Research Article

Existence of Nontrivial Solutions for Perturbed p -Laplacian Equation in $\mathbb{R}^N$ with Critical Nonlinearity

Huixing Zhang and Xiaoqian Liu

Department of Mathematics, China University of Mining and Technology, Xuzhou 221116, China

Correspondence should be addressed to Huixing Zhang; zhx20110906@cumt.edu.cn

Received 5 November 2013; Accepted 17 February 2014; Published 24 March 2014

Academic Editor: Shuangjie Peng

Copyright © 2014 H. Zhang and X. Liu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider a perturbed p -Laplacian equation with critical nonlinearity in $\mathbb{R}^N$. By using variational method, we show that it has at least one positive solution under the proper conditions.

1. Introduction and Main Results

In this paper, we are concerned with the existence of nontrivial solutions for the following nonlinear perturbed p -Laplacian equation with critical nonlinearity:

$$\begin{aligned} & -\varepsilon^p \Delta_p u + V(x) |u|^{p^*-2} u \\ & = K(x) |u|^{p^*-2} u + f(x, u), \quad x \in \mathbb{R}^N, \\ & u(x) > 0, \\ & u(x) \longrightarrow 0 \quad \text{as } |x| \longrightarrow \infty, \end{aligned} \quad (1)$$

where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ is the p -Laplacian operator with $1 < p < N$, $N \geq 3$, $p^* = Np/(N-p)$ denotes the Sobolev critical exponent, $V(x)$ is a nonnegative potential, $K(x)$ is a bounded positive function, and $f(x, u)$ is a superlinear but subcritical function.

For $p = 2$, (1) turns into the following Schrödinger equation of the form

$$-\varepsilon^2 \Delta u + V(x) u = K(x) |u|^{2^*-2} u + f(x, u), \quad x \in \mathbb{R}^N. \quad (2)$$

The equation (2) has been studied extensively under various hypotheses on the potential and nonlinearity by many authors including Ambrosetti and Rabinowitz [1], Bartsch

and Wang [2], Brézis and Lieb [3], Brézis and Nirenberg [4], and Del Pino and Felmer [5] in bounded domains. Meanwhile, we recall some works in unbounded domains which contain Cingolani and Lazzo [6], Clapp and Ding [7], Ding and Lin [8], Floer and Weinstein [9], Grossi [10], Jeanjean and Tanaka [11], Kang and Wei [12], Oh [13], Pistoia [14], Rabinowitz [15], and Tang [16].

For general $p > 1$, most of the work (see [17–19] and the reference therein) dealt with (1) with $\varepsilon = 1$, $K(x) \equiv 0$ and a certain sign potential $V(x)$. Liu and Zheng [20] considered the above mentioned problem with sign-changing potential and subcritical p -superlinear nonlinearity. Cao et al. [21] also studied the similar problem. However, to our best knowledge, it seems that there is almost no work on the existence of semi-classical solutions to the equation in $\mathbb{R}^N$ with critical nonlinearities. This paper will study the critical nonlinearity case in whole space.

Throughout the paper, we make the following assumption:

(H_1) $V \in C(\mathbb{R}^N)$, $V(0) = \inf_{x \in \mathbb{R}^N} V(x) = 0$ and there exists $b > 0$ such that the set $\mathcal{V}^b := \{x \in \mathbb{R}^N : V(x) < b\}$ has finite Lebesgue measure;

(H_2) $K(x) \in C(\mathbb{R}^N, \mathbb{R}^+)$, $0 < \inf K \leq \sup K < \infty$;

(H_3) $f \in C(\mathbb{R}^N \times \mathbb{R}, \mathbb{R})$ and $f(x, t) = o(|t|^{p-2}t)$ uniformly in x as $t \rightarrow 0$;

(H₄) there are $a_0 > 0$ and $p < q < p^*$ such that $|f(x, t)| \leq a_0(1 + |t|^{q-1})$ for all (x, t) ;

(H₅) there exist $b_0 > 0$, $\alpha > p$ and $\mu \in (p, p^*)$ such that $F(x, t) \geq b_0|t|^\alpha$ and $\mu F(x, t) \leq f(x, t)t$ for all (x, t) , where $F(x, t) = \int_0^t f(x, s)ds$.

Our main result reads as follows.

Theorem 1. Assume that (H₁)–(H₅) hold. Then for any $\sigma > 0$, there exists $\varepsilon_\sigma > 0$ such that if $\varepsilon \leq \varepsilon_\sigma$, (1) has at least one positive solution u_ε of least energy which satisfied the following estimate:

$$\frac{\mu - p}{p\mu} \int_{\mathbb{R}^N} (\varepsilon^p |\nabla u_\varepsilon|^p + V(x) |u_\varepsilon|^p) \leq \sigma \varepsilon^N. \quad (3)$$

The main tool used in the proof of Theorem 1 is variational method which was mainly developed in [8]. The main difficulty in the case is to overcome the loss of the compactness of the energy functional related to (1) because of unbounded domain $\mathbb{R}^N$ and critical nonlinearity. Although the energy functional does not satisfy the (PS) condition, we can prove that it possesses (PS)_c condition at some energy level c .

This outline of the paper is organized as follows. In Section 2, we give the variational settings and preliminary results. In Section 3, we show that the corresponding energy functional satisfies (PS)_c condition at the levels less than $\alpha_0 \lambda^{1-N/p}$ with some $\alpha_0 > 0$ independent of λ . Furthermore, it possesses the mountain geometry structure. Section 4 is devoted to the proof of the main result.

2. Preliminaries

Let $\lambda = \varepsilon^{-p}$ in (1). The equation (1) reads as

$$\begin{aligned} -\Delta_p u + \lambda V(x) |u|^{p-2} u \\ = \lambda K(x) |u|^{p^*-2} u + \lambda f(x, u), \quad x \in \mathbb{R}^N, \\ u(x) > 0, \\ u(x) \rightarrow 0, \quad \text{as } |x| \rightarrow \infty. \end{aligned} \quad (4)$$

In order to prove Theorem 1, we are going to prove the following result.

Theorem 2. Assume that (H₁)–(H₅) is satisfied. Then for any $\sigma > 0$, there exists $\lambda_\sigma > 0$ such that if $\lambda > \lambda_\sigma$, (4) has at least one positive solution u_λ satisfying the following estimate:

$$\frac{\mu - p}{p\mu} \int_{\mathbb{R}^N} (|\nabla u_\lambda|^p + \lambda V(x) |u_\lambda|^p) \leq \sigma \lambda^{1-N/p}. \quad (5)$$

Next, we introduce the space

$$\begin{aligned} E_\lambda(\mathbb{R}^N, V) \\ = \left\{ u \in W^{1,p}(\mathbb{R}^N) : \int_{\mathbb{R}^N} \lambda V(x) |u|^p < \infty, \lambda > 0 \right\} \end{aligned} \quad (6)$$

equipped with the norm

$$\|u\|_{E_\lambda} = \left(\int_{\mathbb{R}^N} (|\nabla u|^p + \lambda V(x) |u|^p) \right)^{1/p}. \quad (7)$$

Note that the norm $\|\cdot\|_{E_1}$ is equivalent to the one $\|\cdot\|_{E_\lambda}$ for any $\lambda > 0$. It follows from (H₁) that $E_\lambda(\mathbb{R}^N, V)$ continuously is embedded in $W^{1,p}(\mathbb{R}^N)$. To prove Theorem 2, one considers the C^1 functional $I : W^{1,p}(\mathbb{R}^N) \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} I_\lambda(u) &= \frac{1}{p} \int_{\mathbb{R}^N} (|\nabla u|^p + \lambda V(x) |u|^p) \\ &\quad - \frac{\lambda}{p^*} \int_{\mathbb{R}^N} K(x) |u|^{p^*} - \lambda \int_{\mathbb{R}^N} F(x, u) \\ &= \frac{1}{p} \|u\|_{E_\lambda}^p - \lambda \int_{\mathbb{R}^N} G(x, u), \end{aligned} \quad (8)$$

where $G(x, u) = (1/p^*)K(x)|u|^{p^*} + F(x, u)$.

Under the assumptions of Theorem 2, standard arguments [22] show that $I_\lambda \in C^1(E_\lambda, \mathbb{R})$ and its critical points are weak solutions of (4).

3. Necessary Lemmas

This section will show some lemmas which are important for the proof of the main result.

Lemma 3. Assume that (H₁)–(H₅) is satisfied. For the (PS)_c sequence $\{u_n\} \subset E_\lambda$ for I_λ , we get that $c \geq 0$ and $\{u_n\}$ is bounded in the space E_λ .

Proof. By direct computation and the assumptions (H₂) and (H₅), one has

$$\begin{aligned} I_\lambda(u_n) - \frac{1}{\mu} I'_\lambda(u_n) u_n \\ = \left(\frac{1}{p} - \frac{1}{\mu} \right) \|u_n\|_{E_\lambda}^p + \left(\frac{1}{\mu} - \frac{1}{p^*} \right) \lambda \int_{\mathbb{R}^N} K(x) |u_n|^{p^*} \\ + \lambda \int_{\mathbb{R}^N} \left(\frac{1}{\mu} f(x, u) u - F(x, u) \right). \end{aligned} \quad (9)$$

Together with $I_\lambda(u_n) \rightarrow c$ and $I'_\lambda(u_n) \rightarrow 0$ as $n \rightarrow \infty$, we easily get that the (PS)_c sequence is bounded in E_λ and the energy level $c \geq 0$. $\square$

By Lemma 3, there is $u \in E_\lambda$ such that $u_n \rightharpoonup u$ in E_λ . Furthermore, passing to a subsequence, we have $u_n \rightarrow u$ in $L^b_{\text{loc}}(\mathbb{R}^N)$ for any $b \in [p, p^*)$ and $u_n \rightarrow u$ a.e. in $\mathbb{R}^N$.

Lemma 4. For any $s \in [p, p^*)$, there is a subsequence $\{u_{n_i}\}$ such that, for any $\varepsilon > 0$, there exists $r_\varepsilon > 0$ with

$$\limsup_{i \rightarrow \infty} \int_{B_1 \setminus B_{r_\varepsilon}} |u_{n_i}|^s \leq \varepsilon \quad \text{for any } r \geq r_\varepsilon, \quad (10)$$

where $B_r := \{x \in \mathbb{R}^N : |x| \leq r\}$.

Proof. From $u_n \rightarrow u$ in $L^s_{\text{loc}}(\mathbb{R}^N)$, we have

$$\int_{B_i} |u_n|^s \rightarrow \int_{B_i} |u|^s \quad \text{as } n \rightarrow \infty. \quad (11)$$

Thus, there exists $\tilde{n}_i \in \mathbb{N}$ such that

$$\int_{B_i} (|u_n|^s - |u|^s) < \frac{1}{i}, \quad \forall n = \tilde{n}_i + j, \quad j = 1, 2, \dots \quad (12)$$

In particular, for $n_i = \tilde{n}_i + i$, we have

$$\int_{B_i} (|u_{n_i}|^s - |u|^s) < \frac{1}{i}. \quad (13)$$

Note that there exists $r_\varepsilon > 0$ satisfying

$$\int_{\mathbb{R}^N \setminus B_r} (|u|^s) < \varepsilon \quad \forall r \geq r_\varepsilon. \quad (14)$$

Then

$$\begin{aligned} \int_{B_i \setminus B_r} |u_{n_i}|^s &= \int_{B_i \setminus B_r} (|u_{n_i}|^s - |u|^s) + \int_{B_i \setminus B_r} |u|^s \\ &= \int_{B_i} (|u_{n_i}|^s - |u|^s) + \int_{B_r} (|u|^s - |u_{n_i}|^s) \\ &\quad + \int_{B_i \setminus B_r} |u|^s \\ &\leq \frac{1}{i} + \int_{\mathbb{R}^N \setminus B_r} |u|^s + \int_{B_r} (|u|^s - |u_{n_i}|^s) \\ &\leq \varepsilon, \quad \text{as } i \rightarrow \infty. \end{aligned} \quad (15)$$

This completes the proof of Lemma 4. $\square$

Let $\eta \in C^\infty(\mathbb{R}^+)$ be a smooth function satisfying $0 \leq \eta(t) \leq 1$, $\eta(t) = 1$ if $t \leq 1$ and $\eta(t) = 0$ if $t \geq 2$. Define $\tilde{u}_i(x) = \eta(2|x|/i)u(x)$. It is clear that

$$\|u - \tilde{u}_i\|_{E_\lambda} \rightarrow 0 \quad \text{as } i \rightarrow \infty. \quad (16)$$

Lemma 5. *One has*

$$\limsup_{i \rightarrow \infty} \left| \int_{\mathbb{R}^N} (f(x, u_{n_i}) - f(x, u_{n_i} - \tilde{u}_i) - f(x, \tilde{u}_i)) \varphi \right| = 0 \quad (17)$$

uniformly in $\varphi \in E_\lambda$ with $\|\varphi\|_{E_\lambda} \leq 1$.

Proof. From (16) and the local compactness of Sobolev embedding, for any $r \geq 0$, we have

$$\limsup_{i \rightarrow \infty} \left| \int_{B_r} (f(x, u_{n_i}) - f(x, u_{n_i} - \tilde{u}_i) - f(x, \tilde{u}_i)) \varphi \right| = 0 \quad (18)$$

uniformly in $\|\varphi\|_{E_\lambda} \leq 1$. For any $\varepsilon > 0$, it follows from (14) that

$$\limsup_{i \rightarrow \infty} \int_{B_i \setminus B_r} |\tilde{u}_i|^s \leq \int_{\mathbb{R}^N \setminus B_r} |u|^s \leq \varepsilon \quad (19)$$

for all $r \geq r_\varepsilon$. By Lemma 4 and (H_3) – (H_4) , we obtain

$$\begin{aligned} &\limsup_{i \rightarrow \infty} \left| \int_{\mathbb{R}^N} (f(x, u_{n_i}) - f(x, u_{n_i} - \tilde{u}_i) - f(x, \tilde{u}_i)) \varphi \right| \\ &= \limsup_{i \rightarrow \infty} \left| \int_{B_i \setminus B_r} (f(x, u_{n_i}) - f(x, u_{n_i} - \tilde{u}_i) - f(x, \tilde{u}_i)) \varphi \right| \\ &\leq a_1 \limsup_{i \rightarrow \infty} \int_{B_i \setminus B_r} (|u_{n_i}|^{p-1} + |\tilde{u}_i|^{p-1}) |\varphi| \\ &\quad + a_2 \limsup_{i \rightarrow \infty} \int_{B_i \setminus B_r} (|u_{n_i}|^{q-1} + |\tilde{u}_i|^{q-1}) |\varphi| \\ &\leq a_1 \limsup_{i \rightarrow \infty} \left(\|u_{n_i}\|_{L_p(B_i \setminus B_r)}^{p-1} \right. \\ &\quad \left. + \|\tilde{u}_i\|_{L_p(B_i \setminus B_r)}^{p-1} \right) \|\varphi\|_{L_p(B_i \setminus B_r)} \\ &\quad + a_2 \limsup_{i \rightarrow \infty} \left(\|u_{n_i}\|_{L_q(B_i \setminus B_r)}^{q-1} + \|\tilde{u}_i\|_{L_q(B_i \setminus B_r)}^{q-1} \right) \\ &\quad \times \|\varphi\|_{L_q(B_i \setminus B_r)} \\ &\leq a_3 \varepsilon^{(p-1)/p} + a_4 \varepsilon^{(q-1)/q}. \end{aligned} \quad (20)$$

This shows that the desired conclusion holds. $\square$

Lemma 6. *One has along a subsequence*

$$\begin{aligned} I_\lambda(u_n - \tilde{u}_n) &\rightarrow c - I_\lambda(u), \\ I'_\lambda(u_n - \tilde{u}_n) &\rightarrow 0 \quad \text{in } E_\lambda^{-1} \quad (\text{the dual space of } E_\lambda). \end{aligned} \quad (21)$$

Proof. By Lemma 2.1 of [23] and the arguments of [24], we have

$$\begin{aligned} I_\lambda(u_n - \tilde{u}_n) &= \frac{1}{p} \int_{\mathbb{R}^N} (|\nabla u_n - \nabla \tilde{u}_n|^p + \lambda V(x) |u_n - \tilde{u}_n|^p) \\ &\quad - \frac{\lambda}{p^*} \int_{\mathbb{R}^N} K(x) |u_n - \tilde{u}_n|^{p^*} - \lambda \int_{\mathbb{R}^N} F(x, u_n - \tilde{u}_n) \\ &= I_\lambda(u_n) - I_\lambda(\tilde{u}_n) \\ &\quad + \frac{\lambda}{p^*} \int_{\mathbb{R}^N} K(x) (|u_n|^{p^*} - |u_n - \tilde{u}_n|^{p^*} - |\tilde{u}_n|^{p^*}) \\ &\quad + \lambda \int_{\mathbb{R}^N} (F(x, u_n) - F(x, u_n - \tilde{u}_n) - F(x, \tilde{u}_n)) \\ &\quad + o(1). \end{aligned} \quad (22)$$

By (16) and the similar idea of proving the Brézis-Lieb Lemma [3], we easily get

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x) \left(|u_n|^{p^*} - |u_n - \tilde{u}_n|^{p^*} - |\tilde{u}_n|^{p^*} \right) &= 0, \\ \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} (F(x, u_n) - F(x, u_n - \tilde{u}_n) - F(x, \tilde{u}_n)) &= 0. \end{aligned} \quad (23)$$

Together with the fact $I_\lambda(u_n) \rightarrow c$ and $I_\lambda(\tilde{u}_n) \rightarrow I_\lambda(u)$, one has

$$I_\lambda(u_n - \tilde{u}_n) \rightarrow c - I_\lambda(u). \quad (24)$$

Next, we will check the fact $I'_\lambda(u_n - \tilde{u}_n) \rightarrow 0$ in E_λ^{-1} . For any $\varphi \in E_\lambda$, we have

$$\begin{aligned} I'_\lambda(u_n - \tilde{u}_n) \varphi &= I'_\lambda(u_n) \varphi - I'_\lambda(\tilde{u}_n) \varphi \\ &\quad + \lambda \int_{\mathbb{R}^N} K(x) \left(|u_n|^{p^*-2} u_n - |u_n - \tilde{u}_n|^{p^*-2} (u_n - \tilde{u}_n) \right. \\ &\quad \left. - |\tilde{u}_n|^{p^*-2} \tilde{u}_n \right) \varphi \\ &\quad + \lambda \int_{\mathbb{R}^N} (f(x, u_n) - f(x, u_n - \tilde{u}_n) - f(x, \tilde{u}_n)) \varphi \\ &\quad + o(1). \end{aligned} \quad (25)$$

By the standard argument, it follows that

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x) \left(|u_n|^{p^*-2} u_n - |u_n - \tilde{u}_n|^{p^*-2} (u_n - \tilde{u}_n) \right. \\ \left. - |\tilde{u}_n|^{p^*-2} \tilde{u}_n \right) \varphi = 0 \end{aligned} \quad (26)$$

uniformly in $\|\varphi\|_{E_\lambda} \leq 1$. Together with Lemma 5, we get the desired conclusion. $\square$

Set $u_n^1 = u_n - \tilde{u}_n$; then, $u_n - u = u_n^1 + (\tilde{u}_n - u)$. From (16), it shows that $u_n \rightarrow u$ in E_λ if and only if $u_n^1 \rightarrow 0$ in E_λ .

Furthermore, we have

$$\begin{aligned} I_\lambda(u_n^1) - \frac{1}{p} I'_\lambda(u_n^1) u_n^1 &= \left(\frac{1}{p} - \frac{1}{p^*} \right) \lambda \int_{\mathbb{R}^N} K(x) |u_n^1|^{p^*} \\ &\quad + \lambda \int_{\mathbb{R}^N} \left(\frac{1}{p} u_n^1 f(x, u_n^1) - F(x, u_n^1) \right) \\ &\geq \frac{\lambda}{N} \int_{\mathbb{R}^N} K(x) |u_n^1|^{p^*} \\ &\geq \frac{\lambda}{N} K_{\min} \|u_n^1\|_{p^*}^{p^*}, \end{aligned} \quad (27)$$

where $K_{\min} = \inf_{x \in \mathbb{R}^N} K(x) > 0$.

By the facts that $I_\lambda(u_n^1) \rightarrow c - I_\lambda(u)$ and $I'_\lambda(u_n^1) \rightarrow 0$ in E_λ^{-1} , one has

$$\|u_n^1\|_{p^*}^{p^*} \leq \frac{N(c - I_\lambda(u))}{\lambda K_{\min}} + o(1). \quad (28)$$

Let $V_b(x) := \max\{V(x), b\}$, where b is the positive constant in the assumption (H_1) . Since the set γ_b has finite measure and $u_n^1 \rightarrow 0$ in $L_{\text{loc}}^p(\mathbb{R}^N)$, we get

$$\int_{\mathbb{R}^N} V(x) |u_n^1|^p = \int_{\mathbb{R}^N} V_b(x) |u_n^1|^p + o(1). \quad (29)$$

From (H_2) – (H_5) and Young inequality, there exists $C_b > 0$ such that

$$\int_{\mathbb{R}^N} \left(K(x) |u_n^1|^{p^*} + u_n^1 f(x, u_n^1) \right) \leq b \|u_n^1\|_p^p + C_b \|u_n^1\|_{p^*}^{p^*}. \quad (30)$$

Next, we consider the energy level of the functional I_λ below which the $(PS)_c$ condition held.

Lemma 7. Assume that the assumptions of Theorem 2 are satisfied. There exists $\alpha_0 > 0$ (independent of λ) such that, for any $(PS)_c$ sequence $\{u_n\} \subset E_\lambda$ for I_λ with $u_n \rightarrow u$, either $u_n \rightarrow u$ in E_λ or $c - I_\lambda(u) \geq \alpha_0 \lambda^{1-(N/p)}$.

Proof. Assume that $u_n \not\rightarrow u$; then,

$$\begin{aligned} \liminf_{n \rightarrow \infty} \|u_n^1\|_{E_\lambda} &> 0, \\ c - I_\lambda(u) &> 0. \end{aligned} \quad (31)$$

By the Sobolev inequality, (29), and (30), we get

$$\begin{aligned} S \|u_n^1\|_{p^*}^p &\leq \int_{\mathbb{R}^N} |\nabla u_n^1|^p \\ &= \int_{\mathbb{R}^N} \left(|\nabla u_n^1|^p + \lambda V(x) |u_n^1|^p \right) - \lambda \int_{\mathbb{R}^N} V(x) |u_n^1|^p \\ &= \lambda \int_{\mathbb{R}^N} K(x) |u_n^1|^{p^*} + u_n^1 f(x, u_n^1) - \lambda \\ &\quad \times \int_{\mathbb{R}^N} V_b(x) |u_n^1|^p + o(1) \\ &\leq \lambda b \|u_n^1\|_p^p + \lambda C_b \|u_n^1\|_{p^*}^{p^*} - \lambda b \|u_n^1\|_p^p + o(1) \\ &= \lambda C_b \|u_n^1\|_{p^*}^{p^*} + o(1), \end{aligned} \quad (32)$$

where S is the best Sobolev constant of the immersion

$$S \|u\|_{p^*}^p \leq \int_{\mathbb{R}^N} |\nabla u|^p \quad \forall u \in W^{1,p}(\mathbb{R}^N). \quad (33)$$

This gives

$$\begin{aligned} S &\leq \lambda C_b \|u_n^1\|_{p^*}^{p^*-p} + o(1) \\ &\leq \lambda C_b \left(\frac{N(c - I_\lambda(u))}{\lambda K_{\min}} \right)^{p/N} + o(1) \\ &= \lambda^{1-p/N} C_b \left(\frac{N}{K_{\min}} \right)^{p/N} (c - I_\lambda(u))^{p/N} + o(1). \end{aligned} \quad (34)$$

Set $\alpha_0 = S^{N/p} C_b^{-N/p} N^{-1} K_{\min}$; then,

$$\alpha_0 \lambda^{1-N/p} \leq c - I_\lambda(u) + o(1). \quad (35)$$

This proof is completed. $\square$

From Lemma 7, we will show that I_λ satisfies the following local $(PS)_c$ condition.

Lemma 8. Assume that $(H_1)-(H_5)$ is satisfied. There exists a constant $\alpha_0 > 0$ (independent of λ) such that, if a $(PS)_c$ sequence $\{u_n\} \subset E_\lambda$ for I_λ satisfies $c \leq \alpha_0 \lambda^{1-N/p}$, the sequence $\{u_n\}$ has a strongly convergent subsequence in E_λ .

Proof. By Lemma 7, we easily obtain the required conclusion. $\square$

Now, we consider $\lambda \geq 1$. The following standard arguments show that the energy functional I_λ possesses the mountain-pass structure.

Lemma 9. Under the assumptions of Theorem 2, there exist $\alpha_\lambda, \rho_\lambda > 0$ such that

$$\begin{aligned} I_\lambda(u) &> 0 \quad \text{if } 0 < \|u\|_{E_\lambda} < \rho_\lambda, \\ I_\lambda(u) &\geq \alpha_\lambda \quad \text{if } \|u\|_{E_\lambda} = \rho_\lambda. \end{aligned} \quad (36)$$

Proof. By (30) and (H_5) , for any $\delta > 0$, there is $C_\delta > 0$ such that

$$\int_{\mathbb{R}^N} G(x, u) \leq \delta \|u\|_p^p + C_\delta \|u\|_{p^*}^{p^*}. \quad (37)$$

Thus

$$\begin{aligned} I_\lambda(u) &= \frac{1}{p} \|u\|_{E_\lambda}^p - \lambda \int_{\mathbb{R}^N} G(x, u) \\ &\geq \frac{1}{p} \|u\|_{E_\lambda}^p - \lambda \delta \|u\|_p^p - \lambda C_\delta \|u\|_{p^*}^{p^*}. \end{aligned} \quad (38)$$

Observe that $\|u\|_p^p \leq a_5 \|u\|_{E_\lambda}^p$. Choosing $\delta \leq (2p\lambda a_5)^{-1}$,

$$I_\lambda(u) \geq \frac{1}{2p} \|u\|_{E_\lambda}^p - \lambda C_\delta \|u\|_{p^*}^{p^*}. \quad (39)$$

The fact $p^* > p$ implies the desired conclusion. $\square$

Lemma 10. For any finite dimensional subspace $F \subset E_\lambda$, we have

$$I_\lambda(u) \longrightarrow -\infty, \quad u \in E_\lambda \text{ as } \|u\|_{E_\lambda} \longrightarrow \infty. \quad (40)$$

Proof. By the assumption (H_5) , one has

$$I_\lambda(u) \leq \frac{1}{p} \|u\|_\alpha^p - \lambda b_0 \|u\|_\alpha^\alpha \quad \forall u \in E_\lambda. \quad (41)$$

Since all norms in a finite-dimensional space are equivalent and $\alpha > p$, this implies the desired conclusion. $\square$

Lemma 8 shows that I_λ satisfies $(PS)_{c_\lambda}$ condition for λ large enough and c_λ small sufficiently. In the following, we will find special finite-dimensional subspaces by which we establish sufficiently small minimax levels.

Define the functional

$$\Phi_\lambda(u) = \frac{1}{p} \int_{\mathbb{R}^N} (|\nabla u|^p + \lambda V(x) |u|^p) - \lambda b_0 \int_{\mathbb{R}^N} |u|^\alpha. \quad (42)$$

It is apparent that $\Phi_\lambda \in C^1(E_\lambda)$ and $I_\lambda(u) \leq \Phi_\lambda(u)$ for all $u \in E_\lambda$.

Note that

$$\inf \left\{ \int_{\mathbb{R}^N} |\nabla \phi|^p : \phi \in C_0^\infty(\mathbb{R}^N, \mathbb{R}), \|\phi\|_{L^\alpha(\mathbb{R}^N)} = 1 \right\} = 0. \quad (43)$$

For any $\delta > 0$, there is $\phi_\delta \in C_0^\infty(\mathbb{R}^N, \mathbb{R})$ with $\|\phi_\delta\|_{L^\alpha(\mathbb{R}^N)} = 1$ and $\text{supp } \phi_\delta \subset B_{r_\delta}(0)$ such that $\|\nabla \phi_\delta\|_p^p < \delta$. Let $e_\lambda(x) = \phi_\delta(\sqrt[p]{\lambda}x)$, then $\text{supp } e_\lambda \subset B_{\lambda^{-1/p}r_\delta}(0)$. For any $t \geq 0$, we have

$$\begin{aligned} \Phi_\lambda(te_\lambda) &= \frac{t^p}{p} \|e_\lambda\|_{E_\lambda}^p - b_0 \lambda t^\alpha \int_{\mathbb{R}^N} |\phi_\delta(\sqrt[p]{\lambda}x)|^\alpha \\ &= \lambda^{1-N/p} J_\lambda(t\phi_\delta), \end{aligned} \quad (44)$$

where

$$J_\lambda(u) = \frac{1}{p} \int_{\mathbb{R}^N} (|\nabla u|^p + V(\lambda^{-1/p}x) |u|^p) - b_0 \int_{\mathbb{R}^N} |u|^\alpha. \quad (45)$$

By direct computation, we easily get

$$\begin{aligned} \max_{t \geq 0} J_\lambda(t\phi_\delta) &\leq \frac{\alpha - p}{p\alpha(\alpha b_0)^{p/(\alpha-p)}} \left(\int_{\mathbb{R}^N} (|\nabla \phi_\delta|^p \right. \\ &\quad \left. + V(\lambda^{-1/p}x) |\phi_\delta|^p) \right)^{\alpha/(\alpha-p)}. \end{aligned} \quad (46)$$

In connection with $V(0) = 0$ and $\|\nabla \phi_\delta\|_p^p < \delta$, it shows that there exists $\Lambda_\delta > 0$ such that for all $\lambda \geq \Lambda_\delta$, we have

$$\max_{t \geq 0} J_\lambda(t\phi_\delta) \leq \left(\frac{\alpha - p}{p\alpha(\alpha b_0)^{p/(\alpha-p)}} (2\delta)^{\alpha/(\alpha-p)} \right) \lambda^{1-N/p}. \quad (47)$$

It follows from (47) that

Lemma 11. Assume that (H_1) – (H_5) is satisfied. For any $\sigma > 0$, there is $\Lambda_\sigma > 0$ such that $\lambda \geq \Lambda_\sigma$; there exists $\bar{e}_\lambda \in E_\lambda$ with $\|\bar{e}_\lambda\|_{E_\lambda} > \rho_\lambda$; we have $I_\lambda(\bar{e}_\lambda) \leq 0$ and

$$\max_{t \geq 0} I_\lambda(t\bar{e}_\lambda) \leq \sigma\lambda^{1-N/p}, \quad (48)$$

where ρ_λ is defined in Lemma 9.

Proof. This proof is similar to the one of Lemma 4.3 in [8], so we omit it. $\square$

4. Proof of Theorem 2

In the following, we will give the proof of Theorem 2.

Proof. By Lemma 11, for any $\sigma > 0$ with $0 < \sigma < \alpha_0$, there is $\Lambda_\sigma > 0$ such that for $\lambda \geq \Lambda_\sigma$, we obtain

$$c_\lambda = \inf_{\gamma \in \Gamma_\lambda} \max_{t \in [0,1]} I_\lambda(\gamma(t)) \leq \sigma\lambda^{1-N/p}, \quad (49)$$

where $\Gamma_\lambda = \{\gamma \in C([0,1], E_\lambda) : \gamma(0) = 0, \gamma(1) = \bar{e}_\lambda\}$.

It follows from Lemma 8 that I_λ satisfies $(PS)_{c_\lambda}$ condition. Hence, by the mountain-pass theorem, there exists $u_\lambda \in E_\lambda$ which satisfies $I_\lambda(u_\lambda) = c_\lambda$ and $I'_\lambda(u_\lambda) = 0$. Actually, u_λ is a weak solution of (4). Similar to the argument in [8], we also get that u_λ is a positive least energy solution.

In the end, we show that the solution u_λ satisfies the estimate (5). We easily get

$$\begin{aligned} I_\lambda(u_\lambda) &= I_\lambda(u_\lambda) - \frac{1}{\mu} I'_\lambda(u_\lambda)(u_\lambda) \\ &= \left(\frac{1}{p} - \frac{1}{\mu}\right) \|u_\lambda\|_{E_\lambda}^p + \left(\frac{1}{\mu} - \frac{1}{p^*}\right) \lambda \int_{\mathbb{R}^N} K(x) |u_\lambda|^{p^*} \\ &\quad + \lambda \int_{\mathbb{R}^N} \left(\frac{1}{\mu} u_\lambda f(x, u_\lambda) - F(x, u_\lambda)\right) \\ &\geq \left(\frac{1}{p} - \frac{1}{\mu}\right) \|u_\lambda\|_{E_\lambda}^p. \end{aligned} \quad (50)$$

Note that $I_\lambda(u_\lambda) = c_\lambda$ and $c_\lambda \leq \sigma\lambda^{1-N/p}$ and it implies the required conclusion. The proof is complete. $\square$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors would like to appreciate the referees for their precious comments and suggestions about the original manuscript. This research was supported by the Fundamental Research Funds for the Central Universities (2013XK03) and the National Training Programs of Innovation and Entrepreneurship for Undergraduates (201310290049).

References

- [1] A. Ambrosetti and P. H. Rabinowitz, "Dual variational methods in critical point theory and applications," *Journal of Functional Analysis*, vol. 14, no. 4, pp. 349–381, 1973.
- [2] T. Bartsch and Z. Q. Wang, "Existence and multiplicity results for some superlinear elliptic problems on $\mathbb{R}^N$," *Communications in Partial Differential Equations*, vol. 20, no. 9–10, pp. 1725–1741, 1995.
- [3] H. Brézis and E. Lieb, "A relation between pointwise convergence of functions and convergence of functionals," *Proceedings of the American Mathematical Society*, vol. 88, no. 3, pp. 486–490, 1983.
- [4] H. Brézis and L. Nirenberg, "Some variational problems with lack of compactness," in *Proceedings of Symposia in Pure Mathematics*, vol. 45, pp. 165–201, 1986.
- [5] M. Del Pino and P. L. Felmer, "Multi-peak bound states for nonlinear Schrödinger equations," *Annales de l'Institut Henri Poincaré. Analyse Non Linéaire*, vol. 15, no. 2, pp. 127–149, 1998.
- [6] S. Cingolani and M. Lazzo, "Multiple positive solutions to nonlinear Schrödinger equations with competing potential functions," *Journal of Differential Equations*, vol. 160, no. 1, pp. 118–138, 2000.
- [7] M. Clapp and Y. Ding, "Minimal nodal solutions of a Schrödinger equation with critical nonlinearity and symmetric potential," *Differential and Integral Equations. An International Journal for Theory & Applications*, vol. 16, no. 8, pp. 981–992, 2003.
- [8] Y. Ding and F. Lin, "Solutions of perturbed Schrödinger equations with critical nonlinearity," *Calculus of Variations and Partial Differential Equations*, vol. 30, no. 2, pp. 231–249, 2007.
- [9] A. Floer and A. Weinstein, "Nonspreading wave packets for the cubic Schrödinger equation with a bounded potential," *Journal of Functional Analysis*, vol. 69, no. 3, pp. 397–408, 1986.
- [10] M. Grossi, "Some results on a class of nonlinear Schrödinger equations," *Mathematische Zeitschrift*, vol. 235, no. 4, pp. 687–705, 2000.
- [11] L. Jeanjean and K. Tanaka, "Singularly perturbed elliptic problems with superlinear or asymptotically linear nonlinearities," *Calculus of Variations and Partial Differential Equations*, vol. 21, no. 3, pp. 287–318, 2004.
- [12] X. Kang and J. Wei, "On interacting bumps of semi-classical states of nonlinear Schrödinger equations," *Advances in Differential Equations*, vol. 5, no. 7–9, pp. 899–928, 2000.
- [13] Y.-G. Oh, "On positive multi-lump bound states of nonlinear Schrödinger equations under multiple well potential," *Communications in Mathematical Physics*, vol. 131, no. 2, pp. 223–253, 1990.
- [14] A. Pistoia, "Multi-peak solutions for a class of nonlinear Schrödinger equations," *Nonlinear Differential Equations and Applications*, vol. 9, no. 1, pp. 69–91, 2002.
- [15] P. H. Rabinowitz, "On a class of nonlinear Schrödinger equations," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 43, no. 2, pp. 270–291, 1992.
- [16] Z. Tang, "Least energy solutions for semilinear Schrödinger equations involving critical growth and indefinite potentials," *Communications on Pure and Applied Analysis*, vol. 13, no. 1, pp. 237–248, 2014.
- [17] C. O. Alves and Y. H. Ding, "Multiplicity of positive solutions to a p -Laplacian equation involving critical nonlinearity," *Journal of Mathematical Analysis and Applications*, vol. 279, no. 2, pp. 508–521, 2003.

- [18] X. Fan, “ $p(x)$ -Laplacian equations in $\mathbb{R}^N$ with periodic data and nonperiodic perturbations,” *Journal of Mathematical Analysis and Applications*, vol. 341, no. 1, pp. 103–119, 2008.
- [19] S. Liu, “On ground states of superlinear p -Laplacian equations in $\mathbb{R}^N$,” *Journal of Mathematical Analysis and Applications*, vol. 361, no. 1, pp. 48–58, 2010.
- [20] C. Liu and Y. Zheng, “Existence of nontrivial solutions for p -Laplacian equations in $\mathbb{R}^N$,” *Journal of Mathematical Analysis and Applications*, vol. 380, no. 2, pp. 669–679, 2011.
- [21] D. Cao, S. Peng, and S. Yan, “Infinitely many solutions for p -Laplacian equation involving critical Sobolev growth,” *Journal of Functional Analysis*, vol. 262, no. 6, pp. 2861–2902, 2012.
- [22] J. Mawhin and M. Willem, *Critical Point Theory and Hamiltonian Systems*, Springer, New York, NY, USA, 1989.
- [23] Y. Li, Q. Guo, and P. Niu, “Global compactness results for quasi-linear elliptic problems with combined critical Sobolev-Hardy terms,” *Nonlinear Analysis: Theory, Methods & Applications*, vol. 74, no. 4, pp. 1445–1464, 2011.
- [24] N. Ghoussoub and C. Yuan, “Multiple solutions for quasi-linear PDEs involving the critical Sobolev and Hardy exponents,” *Transactions of the American Mathematical Society*, vol. 352, no. 12, pp. 5703–5743, 2000.