Research Article

Poincaré Inequalities for Composition Operators with L^p -Norm

Ru Fang

Department of Mathematics, Harbin Institute of Technology, Harbin 150001, China

Correspondence should be addressed to Ru Fang; fangr@hit.edu.cn

Received 24 November 2013; Accepted 4 March 2014; Published 1 April 2014

Academic Editor: Shusen Ding

Copyright © 2014 Ru Fang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We establish the Poincaré-type inequalities for the composition of the homotopy operator, exterior derivative operator, and the projection operator with L^p -norm applied to the nonhomogeneous A -harmonic equation in $L^p(\Omega)$ -averaging domains.

1. Introduction

The purpose of the paper is to develop the Poincaré-type inequalities for the composition of the homotopy operator T , exterior derivative operator d , and the projection operator H with L^p -norm. These operators play critical roles in investigating the properties of the solutions to PDEs and in controlling oscillatory behavior of the solutions in domains [1–6]. We first establish the local Poincaré inequalities for the composition $T \circ d \circ H$ in $L^p(\Omega)$ -averaging domains. Then, we prove the global Poincaré inequalities for the composition of $T \circ d \circ H$ in $L^p(\Omega)$ -averaging domains.

In this paper, we assume Ω is a bounded and convex domain in $\mathbb{R}^n$, $n \geq 2$ and $B = B(x_0, r)$ is a ball that is centred at x_0 with r as its radius. For any $\sigma > 0$, we use σB to denote the ball with centred at x_0 with radius σr . We do not distinguish the balls from the cubes in this paper. We use $|E|$ to denote the Lebesgue measure of a set $E \subset \mathbb{R}^n$. We call ω a weight if $\omega \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $\omega > 0$ a.e. For a function u , we denote the average of u over B by $u_B = (1/|B|) \int_B u dx$. Differential forms are extensions of functions in $\mathbb{R}^n$. For example, the function $u(x_1, x_2, \dots, x_n)$ is called a 0-form. Moreover, if $u(x_1, x_2, \dots, x_n)$ is differentiable, it is called a differential 0-form. The 1-form $u(x)$ in $\mathbb{R}^n$ can be written as $u(x) = \sum_{i=1}^n u_i(x_1, x_2, \dots, x_n) dx_i$. If the coefficient functions $u_i(x_1, x_2, \dots, x_n)$, $i = 1, 2, \dots, n$, are differentiable, $u(x)$ is called a differential 1-form. Similarly, a differential k -form $u(x)$ is generated by $\{dx_{i_1} \wedge dx_{i_2} \wedge \dots \wedge dx_{i_k}\}$, $k = 1, 2, \dots, n$, that is, $u(x) = \sum_I u_I(x) dx_I = \sum u_{i_1 i_2 \dots i_k}(x) dx_{i_1} \wedge dx_{i_2} \wedge \dots \wedge dx_{i_k}$, where $I = (i_1, i_2, \dots, i_k)$, $1 \leq i_1 < i_2 < \dots < i_k \leq n$.

Let $\wedge^l = \wedge^l(\mathbb{R}^n)$ be the set of all l -forms in $\mathbb{R}^n$, $D'(\Omega, \wedge^l)$ be the space of all differential l -forms on Ω and $L^p(\Omega, \wedge^l)$ be the l -forms $u(x) = \sum_I u_I(x) dx_I$ on Ω satisfying $\int_\Omega |u_I|^p < \infty$ for all ordered l -tuples I , $l = 1, 2, \dots, n$. We denote the exterior derivative by $d : D'(\Omega, \wedge^l) \rightarrow D'(\Omega, \wedge^{l+1})$ for $l = 0, 1, \dots, n-1$, and define the Hodge star operator $*$: $\wedge^k \rightarrow \wedge^{n-k}$ as follows: if $u = u_{i_1 i_2 \dots i_k}(x_1, x_2, \dots, x_n) dx_{i_1} \wedge dx_{i_2} \wedge \dots \wedge dx_{i_k} = u_I dx_I$, $i_1 < i_2 < \dots < i_k$, is a differential k -form, then $*u = *(u_{i_1 i_2 \dots i_k} dx_{i_1} \wedge dx_{i_2} \wedge \dots \wedge dx_{i_k}) = (-1)^{\sum(I)} u_I dx_J$, where $I = (i_1, i_2, \dots, i_k)$, $J = \{1, 2, \dots, n\} - I$, and $\sum(I) = (k(k+1)/2) + \sum_{i=1}^k i_j$. The Hodge codifferential operator $d^* : D'(\Omega, \wedge^{l+1}) \rightarrow D'(\Omega, \wedge^l)$ is given by $d^* = (-1)^{n-l+1} * d *$ on $D'(\Omega, \wedge^{l+1})$, $l = 0, 1, \dots, n-1$.

We use M to denote a bounded and convex domain on $\mathbb{R}^n$. Let $\wedge^l M$ be the l th exterior power of the cotangent bundle, let $C^\infty(\wedge^l M)$ be the space of smooth l -forms on M , and $\mathcal{W}(\wedge^l M) = \{u \in L^1_{\text{loc}}(\wedge^l M) : u \text{ has generalized gradient}\}$. The harmonic l -fields are defined by $\mathcal{H}(\wedge^l M) = \{u \in \mathcal{W}(\wedge^l M) : du = d^*u = 0, u \in L^p \text{ for some } 1 < p < \infty\}$. The orthogonal complement of $\mathcal{H}$ in L^1 is defined by $\mathcal{H}^\perp = \{u \in L^1 : \langle u, h \rangle = 0 \text{ for all } h \in \mathcal{H}\}$. Then, the Green's operator G is defined as $G : C^\infty(\wedge^l M) \rightarrow \mathcal{H}^\perp \cap C^\infty(\wedge^l M)$ by assigning $G(u)$ as the unique element of $\mathcal{H}^\perp \cap C^\infty(\wedge^l M)$ satisfying Poisson's equation $\Delta G(u) = u - H(u)$, where H is the harmonic projection operator that maps $C^\infty(\wedge^l M)$ onto $\mathcal{H}$ so that $H(u)$ is the harmonic part of u . See [7, 8] for more properties of these operators. The differential forms can

be used to describe various systems of PDEs and to express different geometric structures on manifolds. See [9, 10].

The operator K_y with the case $y = 0$ was first introduced by Cartan in [11]. Then, it was extended to the following version in [12]. To each $y \in \Omega$ there corresponds a linear operator $K_y : C^\infty(\Omega, \wedge^l) \rightarrow C^\infty(\Omega, \wedge^{l-1})$ defined by $(K_y u)(x; \xi_1, \dots, \xi_{l-1}) = \int_0^1 t^{l-1} u(tx + y - ty; x - y, \xi_1, \dots, \xi_{l-1}) dt$ and the decomposition $u = d(K_y u) + K_y(du)$. A homotopy operator $T : C^\infty(\Omega, \wedge^l) \rightarrow C^\infty(\Omega, \wedge^{l-1})$ is defined by averaging K_y over all points $y \in \Omega : Tu = \int_\Omega \phi(y) K_y(du)$, where $\phi \in C_0^\infty(\Omega)$ is normalized so that $\int \phi(y) dy = 1$.

We are particularly interested in a class of differential forms satisfying the well-known nonhomogeneous A -harmonic equation

$$d^* A(x, du) = B(x, du), \quad (1)$$

where $A : \Omega \times \wedge^l(\mathbb{R}^n) \rightarrow \wedge^l(\mathbb{R}^n)$ and $B : \Omega \times \wedge^l(\mathbb{R}^n) \rightarrow \wedge^{l-1}(\mathbb{R}^n)$ satisfy the conditions

$$\begin{aligned} |A(x, \xi)| &\leq a|\xi|^{p-1}, & A(x, \xi) \cdot \xi &\geq |\xi|^p, \\ |B(x, \xi)| &\leq b|\xi|^{p-1} \end{aligned} \quad (2)$$

for almost every $x \in \Omega$ and all $\xi \in \wedge^l(\mathbb{R}^n)$. Here $a > 0$ and $b > 0$ are constants and $1 < p < \infty$ is a fixed exponent associated with (1). A solution to (1) is an element of the Sobolev space $W_{loc}^{1,p}(\Omega, \wedge^{l-1})$ such that $\int_\Omega A(x, du) \cdot d\varphi + B(x, du) \cdot \varphi = 0$ for all $\varphi \in W_{loc}^{1,p}(\Omega, \wedge^{l-1})$ with compact support. If u is a function (0-form) in $\mathbb{R}^n$, (1) reduces to

$$\operatorname{div} A(x, \nabla u) = B(x, \nabla u). \quad (3)$$

If the operator $B = 0$, (1) becomes

$$d^* A(x, du) = 0, \quad (4)$$

which is called the homogeneous A -harmonic equation. Let $A : \Omega \times \wedge^l(\mathbb{R}^n) \rightarrow \wedge^l(\mathbb{R}^n)$ be defined by $A(x, \xi) = \xi|\xi|^{p-2}$ with $p > 1$. Then, A satisfies the required conditions and $d^* A(x, du) = 0$ becomes the p -harmonic equation $d^*(du|du|^{p-2}) = 0$ for differential forms. Some results have been obtained in recent years about different versions of the A -harmonic equation; see [1, 2, 8, 9, 13–15].

2. Main Results and Proofs

Definition 1. Let φ be a continuously increasing convex function on $[0, \infty)$ with $\varphi(0) = 0$, and let Ω be a domain with $\mu(\Omega) < \infty$. If u is a measurable function in Ω , then we define the Orlicz norm of u by

$$\|u\|_{\varphi, \Omega} = \inf \left\{ k > 0 : \int_\Omega \varphi \left(\frac{|u(x)|}{k} \right) dx \leq 1 \right\}. \quad (5)$$

A continuously increasing function $\varphi : [0, \infty)$ with $\varphi(0) = 0$ is called an Orlicz function, and a convex Orlicz function φ is often called a Young function.

From Definition 1, it is easy to see that for any domain $\Omega \subset \mathbb{R}^n$

$$\int_\Omega \varphi \left(\frac{|u(x)|}{\|u\|_{\varphi, \Omega}} \right) dx \leq 1 \quad (6)$$

if $\|u\|_{\varphi, \Omega}$ is finite.

Definition 2. Let φ be an increasing convex function on $[0, \infty)$ with $\varphi(0) = 0$. We call a proper subdomain $\Omega \subset \mathbb{R}^n$ an Orlicz space $L^\varphi(\Omega)$, if $\mu(\Omega) < \infty$ and there exists a constant C such that

$$\|u - u_{B_0}\|_{\varphi, \Omega} \leq C \sup_{B \subset \Omega} \|u - u_B\|_{\varphi, B} \quad (7)$$

for some ball $B_0 \subset \Omega$ and all integrable functions u in Ω , where the supremum is over all balls B with $B \subset \Omega$.

Definition 3 (see [15]). We say that a Young function φ lies in the class $G(p, q, C)$, $1 \leq p < q < \infty$, $C \geq 1$, if (i) $1/C \leq \varphi(t^{1/p})/g(t) \leq C$ and (ii) $1/C \leq \varphi(t^{1/p})/h(t) \leq C$ for all $t > 0$, where g is a convex increasing function and h is a concave increasing function on $[0, \infty)$.

From [15], we know that the class $G(p, q, C)$ contains some very interesting functions, such as $\varphi(t) = t^p$ and $\varphi(t) = t^p \log_+^\alpha(t)$, $p \geq 1$, $\alpha \in \mathbb{R}$, and each of φ, g and h is doubling in the sense that its values at t and $2t$ are uniformly comparable for all $t > 0$, and the consequent fact that

$$C_1 t^q \leq h^{-1}(\varphi(t)) \leq C_2 t^q, \quad C_1 t^p \leq g^{-1}(\varphi(t)) \leq C_2 t^p, \quad (8)$$

where C_1 and C_2 are constants. We will need the following reverse Hölder inequality.

Lemma 4 (see [4]). Let u be a solution of the nonhomogeneous A -harmonic equation (1) in a bounded and convex domain Ω and $0 < s, t < \infty$. Then, there exists a constant C , independent of u , such that

$$\|u\|_{s, B} \leq C |B|^{(t-s)/st} \|u\|_{t, \sigma B} \quad (9)$$

for all balls B with $\sigma B \subset \Omega$ for some $\sigma > 1$.

Lemma 5 (see [1]). Let u be a solution of the nonhomogeneous A -harmonic equation (1) in a bounded and convex domain Ω . Let H be the projection operator, and let $T : C^\infty(\Omega, \wedge^l) \rightarrow C^\infty(\Omega, \wedge^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} \|T(d(H(u))) - (T(d(H(u))))_B\|_{s, B} \\ \leq C |B| \operatorname{diam}(B) \|du\|_{s, B} \end{aligned} \quad (10)$$

for all balls B with $B \subset \Omega$.

Lemma 6 (see [1]). Let u be a solution of the nonhomogeneous A -harmonic equation (1) in a bounded and convex domain Ω . Let H be the projection operator, and let $T : C^\infty(\Omega, \wedge^l) \rightarrow$

$C^\infty(\Omega, \Lambda^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_B\|_{s,B} \\ & \leq C |B| \operatorname{diam}(B) \|u\|_{s,\sigma B} \end{aligned} \quad (11)$$

for all balls B with $\sigma B \subset \Omega$, where $\sigma > 1$ is a constant.

Theorem 7. Let φ be a Young function in the class $G(p, q, C)$, $1 \leq p < q < \infty$, $C \geq 1$, and let Ω be a bounded convex domain. Assume that $\varphi(|u|) \in L^1_{\text{loc}}(\Omega)$ and u is a solution of the nonhomogeneous A -harmonic (1) in Ω , $\varphi(|du|) \in L^1_{\text{loc}}(\Omega)$. Let H be the projection operator, and let $T : C^\infty(\Omega, \Lambda^l) \rightarrow C^\infty(\Omega, \Lambda^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_B\|_{\varphi,B} \\ & \leq C |B| \operatorname{diam}(B) \|du\|_{\varphi,\sigma B} \end{aligned} \quad (12)$$

for some $\sigma > 1$ and all balls B with $\sigma B \subset \Omega$.

Proof. For any constant $k > 0$, from Lemmas 4 and 5, (i) in Definition 3, using the fact that φ is an increasing function, Jensen's inequality, and noticing that φ and g are doubling, we have

$$\begin{aligned} & \varphi\left(\frac{1}{k}\left(\int_B |T(d(H(u))) - (T(d(H(u))))_B|^q dx\right)^{1/q}\right) \\ & \leq \varphi\left(\frac{1}{k} C_1 |B|^{(p-q)/pq} \left(\int_{\sigma B} |T(d(H(u))) - (T(d(H(u))))_B|^p dx\right)^{1/p}\right) \\ & \leq \varphi\left(\frac{1}{k} C_2 |B|^{1+(p-q)/pq} \operatorname{diam}(B) \left(\int_{\sigma B} |du|^p dx\right)^{1/p}\right) \\ & \leq \varphi\left(\left(\frac{1}{k^p} C_2^p |B|^{p+(p-q)/q} (\operatorname{diam}(B))^p \int_{\sigma B} |du|^p dx\right)^{1/p}\right) \\ & \leq C_3 g\left(\frac{1}{k^p} C_2^p |B|^{p+(p-q)/q} (\operatorname{diam}(B))^p \int_{\sigma B} |du|^p dx\right) \\ & = C_3 g\left(\int_{\sigma B} \frac{1}{k^p} C_2^p |B|^{p+(p-q)/q} (\operatorname{diam}(B))^p |du|^p dx\right) \\ & \leq C_3 \int_{\sigma B} g\left(\frac{1}{k^p} C_2^p |B|^{p+(p-q)/q} (\operatorname{diam}(B))^p |du|^p\right) dx. \end{aligned} \quad (13)$$

Since $p \geq 1$, then, $1 + (p - q)/pq > 0$. Hence, we have $|B|^{1+(p-q)/pq} \leq |\Omega|^{1+(p-q)/pq} \leq C_4$. From (i) in Definition 3, we find that $g(t) \leq C_5 \varphi(t^{1/p})$. Thus,

$$\begin{aligned} & \int_{\sigma B} g\left(\frac{1}{k^p} C_2^p |B|^{p+(p-q)/q} (\operatorname{diam}(B))^p |du|^p\right) dx \\ & \leq C_5 \int_{\sigma B} \varphi\left(\frac{1}{k} C_2 |B|^{1+(p-q)/pq} \operatorname{diam}(B) |du|\right) dx \quad (14) \\ & \leq C_5 \int_{\sigma B} \varphi\left(\frac{1}{k} C_2 |B| \operatorname{diam}(B) |du|\right) dx. \end{aligned}$$

Combining (13) and (14) yields

$$\begin{aligned} & \varphi\left(\frac{1}{k}\left(\int_B |T(d(H(u))) - (T(d(H(u))))_B|^q dx\right)^{1/q}\right) \\ & \leq C_6 \int_{\sigma B} \varphi\left(\frac{1}{k} C_2 |B| \operatorname{diam}(B) |du|\right) dx. \end{aligned} \quad (15)$$

Using Jensen's inequality for h^{-1} , (8), and noticing that φ and h are doubling, we obtain

$$\begin{aligned} & \int_B \varphi\left(\frac{|T(d(H(u))) - (T(d(H(u))))_B|}{k}\right) dx \\ & = h\left(h^{-1}\left(\int_B \varphi(|T(d(H(u))) - (T(d(H(u))))_B| \times (k)^{-1}) dx\right)\right) \\ & \leq h\left(\int_B h^{-1}\left(\varphi(|T(d(H(u))) - (T(d(H(u))))_B| \times (k)^{-1})\right) dx\right) \\ & \leq h\left(C_7 \int_B (|T(d(H(u))) - (T(d(H(u))))_B| \times (k)^{-1})^q dx\right)^{1/q} \\ & \leq C_8 \varphi\left(\left(C_7 \int_B (|T(d(H(u))) - (T(d(H(u))))_B| \times (k)^{-1})^q dx\right)^{1/q}\right) \\ & \leq C_8 \varphi\left(\frac{1}{k} \left(C_7 \int_B (|T(d(H(u))) - (T(d(H(u))))_B|^q dx\right)^{1/q}\right) \\ & \leq C_9 \varphi\left(\frac{1}{k} \left(\int_B (|T(d(H(u))) - (T(d(H(u))))_B|^q dx\right)^{1/q}\right). \end{aligned} \quad (16)$$

Substituting (15) into (16) and noticing that φ is doubling, we have

$$\begin{aligned} & \int_B \varphi \left(\frac{|T(d(H(u))) - (T(d(H(u))))_B|}{k} \right) dx \\ & \leq C_{10} \int_{\sigma B} \varphi \left(\frac{1}{k} C_2 |B| \text{diam}(B) |du| \right) dx \quad (17) \\ & \leq C_{11} \int_{\sigma B} \varphi \left(\frac{1}{k} |B| \text{diam}(B) |du| \right) dx. \end{aligned}$$

From Definition 2 and (17), we have the following version of Poincaré inequality with the Orlicz norm:

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_B\|_{\varphi, B} \\ & \leq C |B| \text{diam}(B) \|du\|_{\varphi, \sigma B}. \end{aligned} \quad (18)$$

We have completed the proof of Theorem 7. $\square$

Theorem 8. Let φ be a Young function in the class $G(p, q, C)$, $1 \leq p < q < \infty$, $C \geq 1$, and let Ω be a bounded convex domain. Assume that $\varphi(|u|) \in L^1_{\text{loc}}(\Omega)$ and u is a solution of the non-homogeneous A -harmonic (1) in Ω , $\varphi(|du|) \in L^1_{\text{loc}}(\Omega)$. Let H be the projection operator, and let $T : C^\infty(\Omega, \Lambda^l) \rightarrow C^\infty(\Omega, \Lambda^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_B\|_{\varphi, B} \\ & \leq C |B| \text{diam}(B) \|du\|_{\varphi, B} \end{aligned} \quad (19)$$

for some $\sigma > 1$ and all balls B with $\sigma B \subset \Omega$.

Proof. For any constant $k > 0$, from Lemma 5, (i) in Definition 3, using the fact that φ is an increasing function, Jensen's inequality, and noticing that φ and g are doubling, we have

$$\begin{aligned} & \varphi \left(\frac{1}{k} \left(\int_B |T(d(H(u))) - (T(d(H(u))))_B|^p dx \right)^{1/p} \right) \\ & \leq \varphi \left(\frac{1}{k} C_1 |B| \text{diam}(B) \left(\int_B |du|^p dx \right)^{1/p} \right) \\ & \leq \varphi \left(\left(\frac{1}{k^p} C_1^p |B|^p (\text{diam}(B))^p \int_B |du|^p dx \right)^{1/p} \right) \quad (20) \\ & \leq C_2 g \left(\frac{1}{k^p} C_1^p |B|^p (\text{diam}(B))^p \int_B |du|^p dx \right) \\ & = C_2 g \left(\int_B \frac{1}{k^p} C_1^p |B|^p (\text{diam}(B))^p |du|^p dx \right) \\ & \leq C_2 \int_B g \left(\frac{1}{k^p} C_1^p |B|^p (\text{diam}(B))^p |du|^p \right) dx. \end{aligned}$$

Since $p \geq 1$, then $|B| \leq |\Omega| \leq C_3$. From (i) in Definition 3, we find that $g(t) \leq C_4 \varphi(t^{1/p})$. Thus,

$$\begin{aligned} & \int_B g \left(\frac{1}{k^p} C_1^p |B|^p (\text{diam}(B))^p |du|^p \right) dx \\ & \leq C_4 \int_B \varphi \left(\frac{1}{k} C_1 |B| \text{diam}(B) |du| \right) dx. \end{aligned} \quad (21)$$

Combining (20) and (21) yields

$$\begin{aligned} & \varphi \left(\frac{1}{k} \left(\int_B |T(d(H(u))) - (T(d(H(u))))_B|^p dx \right)^{1/p} \right) \\ & \leq C_5 \int_B \varphi \left(\frac{1}{k} C_1 |B| \text{diam}(B) |du| \right) dx. \end{aligned} \quad (22)$$

Using Jensen's inequality for g^{-1} , (8), and noticing that φ and h are doubling, we obtain

$$\begin{aligned} & \int_B \varphi \left(\frac{|T(d(H(u))) - (T(d(H(u))))_B|}{k} \right) dx \\ & = g \left(g^{-1} \left(\int_B \varphi \left((|T(d(H(u))) - (T(d(H(u))))_B| \times (k)^{-1} \right) dx \right) \right) \\ & \leq g \left(\int_B g^{-1} \left(\varphi \left((|T(d(H(u))) - (T(d(H(u))))_B| \times (k)^{-1} \right) \right) dx \right) \\ & \leq g \left(C_6 \int_B \left((|T(d(H(u))) - (T(d(H(u))))_B| \times (k)^{-1} \right)^p dx \right) \\ & \leq C_7 \varphi \left(\left(C_6 \int_B \left((|T(d(H(u))) - (T(d(H(u))))_B| \times (k)^{-1} \right)^p dx \right)^{1/p} \right) \\ & \leq C_7 \varphi \left(\frac{1}{k} \left(C_6 \int_B (|T(d(H(u))) - (T(d(H(u))))_B|^p dx \right)^{1/p} \right) \\ & \leq C_8 \varphi \left(\frac{1}{k} \left(\int_B (|T(d(H(u))) - (T(d(H(u))))_B|^p dx \right)^{1/p} \right). \end{aligned} \quad (23)$$

Substituting (22) into (23) and noticing that φ is doubling, we have

$$\begin{aligned} & \int_B \varphi \left(\frac{|T(d(H(u))) - (T(d(H(u))))_B|}{k} \right) dx \\ & \leq C_9 \int_B \varphi \left(\frac{1}{k} C_1 |B| \text{diam}(B) |du| \right) dx \\ & \leq C_{10} \int_B \varphi \left(\frac{1}{k} |B| \text{diam}(B) |du| \right) dx. \end{aligned} \quad (24)$$

From Definition 2 and (24), we have the following version of Poincaré inequality with the Orlicz norm:

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_B\|_{\varphi, B} \\ & \leq C |B| \text{diam}(B) \|du\|_{\varphi, B}. \end{aligned} \quad (25)$$

We have completed the proof of Theorem 8. $\square$

Using a similar method to the proof of Theorem 8, we can establish the following version of Poincaré inequality with the Orlicz norm.

Theorem 9. Let φ be a Young function in the class $G(p, q, C)$, $1 \leq p < q < \infty, C \geq 1$, and let Ω be a bounded convex domain. Assume that $\varphi(|u|) \in L^1_{\text{loc}}(\Omega)$ and u is a solution of the non-homogeneous A -harmonic (1) in Ω , $\varphi(|du|) \in L^1_{\text{loc}}(\Omega)$. Let H be the projection operator, and let $T : C^\infty(\Omega, \Lambda^l) \rightarrow C^\infty(\Omega, \Lambda^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_B\|_{\varphi, B} \\ & \leq C |B| \text{diam}(B) \|u\|_{\varphi, \sigma B} \end{aligned} \quad (26)$$

for some $\sigma > 1$ and all balls B with $\sigma B \subset \Omega$.

Theorem 10. Let φ be a Young function in the class $G(p, q, C)$, $1 \leq p < q < \infty, C \geq 1$, and let Ω be a bounded convex domain. Assume that $\varphi(|u|) \in L^1_{\text{loc}}(\Omega)$ and u is a solution of the non-homogeneous A -harmonic (1) in Ω , $\varphi(|du|) \in L^1_{\text{loc}}(\Omega)$. Let H be the projection operator, and let $T : C^\infty(\Omega, \Lambda^l) \rightarrow C^\infty(\Omega, \Lambda^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_{B_0}\|_{\varphi, \Omega} \\ & \leq C |B| \text{diam}(B) \|du\|_{\varphi, \Omega}, \end{aligned} \quad (27)$$

where $B_0 \subset \Omega$ is some fixed ball.

Proof. From definition of the $L^\varphi(\Omega)$ and (12), we have

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_{B_0}\|_{\varphi, \Omega} \\ & \leq C_1 \sup_{B \subset \Omega} \|T(d(H(u))) - (T(d(H(u))))_B\|_{\varphi, B} \\ & \leq C_1 \sup_{B \subset \Omega} (C_2 |B| \text{diam}(B) \|du\|_{\varphi, \sigma B}) \end{aligned}$$

$$\begin{aligned} & \leq C_1 \sup_{B \subset \Omega} (C_2 |B| \text{diam}(B) \|du\|_{\varphi, \Omega}) \\ & \leq C_3 |B| \text{diam}(B) \|du\|_{\varphi, \Omega}. \end{aligned} \quad (28)$$

We have completed the proof of Theorem 10. $\square$

Using a similar method to the proof of Theorem 8, we obtain Theorem 11.

Theorem 11. Let φ be a Young function in the class $G(p, q, C)$, $1 \leq p < q < \infty, C \geq 1$, and let Ω be a bounded convex domain. Assume that $\varphi(|u|) \in L^1_{\text{loc}}(\Omega)$ and u is a solution of the non-homogeneous A -harmonic (1) in Ω , $\varphi(|du|) \in L^1_{\text{loc}}(\Omega)$. Let H be the projection operator, and let $T : C^\infty(\Omega, \Lambda^l) \rightarrow C^\infty(\Omega, \Lambda^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_{B_0}\|_{\varphi, \Omega} \\ & \leq C |B| \text{diam}(B) \|u\|_{\varphi, \Omega}, \end{aligned} \quad (29)$$

where $B_0 \subset \Omega$ is some fixed ball.

It has been proved in [5] that any John domain is special $L^\varphi(\Omega)$ -averaging domain. Hence, we have the following results.

Corollary 12. Let φ be a Young function in the class $G(p, q, C)$, $1 \leq p < q < \infty, C \geq 1$, and let Ω be a bounded John domain. Assume that $\varphi(|u|) \in L^1_{\text{loc}}(\Omega)$ and u is a solution of the non-homogeneous A -harmonic (3) in Ω , $\varphi(|du|) \in L^1_{\text{loc}}(\Omega)$. Let H be the projection operator, and let $T : C^\infty(\Omega, \Lambda^l) \rightarrow C^\infty(\Omega, \Lambda^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_{B_0}\|_{\varphi, \Omega} \\ & \leq C |B| \text{diam}(B) \|du\|_{\varphi, \Omega}, \end{aligned} \quad (30)$$

where $B_0 \subset \Omega$ is some fixed ball.

For some special convex function, we have the following theorems.

Theorem 13. Let $\varphi = t^p$ or $\varphi = t^p \log^\alpha(e + t) \in G(p, q, C)$, $1 \leq p < q < \infty, C \geq 1, \alpha \in \mathbb{R}$ a Young function, and Ω a bounded convex domain. Assume that $\varphi(|u|) \in L^1_{\text{loc}}(\Omega)$ and u is a solution of the nonhomogeneous A -harmonic (1) in Ω , $\varphi(|du|) \in L^1_{\text{loc}}(\Omega)$. Let H be the projection operator, and let $T : C^\infty(\Omega, \Lambda^l) \rightarrow C^\infty(\Omega, \Lambda^{l-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_B\|_{\varphi, B} \\ & \leq C |B| \text{diam}(B) \|du\|_{\varphi, \sigma B} \end{aligned} \quad (31)$$

for some $\sigma > 1$ and all balls B with $\sigma B \subset \Omega$.

Theorem 14. Let $\varphi = t^p$ or $\varphi = t^p \log^\alpha(e+t) \in G(p, q, C)$, $1 \leq p < q < \infty$, $C \geq 1$, $\alpha \in \mathbb{R}$ a Young function, and Ω a bounded convex domain. Assume that $\varphi(|u|) \in L^1_{\text{loc}}(\Omega)$ and u is a solution of the nonhomogeneous A -harmonic (1) in Ω , $\varphi(|du|) \in L^1_{\text{loc}}(\Omega)$. Let H be the projection operator, and let $T : C^\infty(\Omega, \Lambda^1) \rightarrow C^\infty(\Omega, \Lambda^{1-1})$ be the homotopy operator. Then, there exists a constant C , independent of u , such that

$$\begin{aligned} & \|T(d(H(u))) - (T(d(H(u))))_{B_0}\|_{\varphi, \Omega} \\ & \leq C |B| \text{diam}(B) \|u\|_{\varphi, \Omega}, \end{aligned} \quad (32)$$

where $B_0 \subset \Omega$ is some fixed ball.

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of the paper.

Acknowledgment

The author would like to thank Professor Shusen Ding for his precious and thoughtful suggestions.

References

- [1] R. P. Agarwal, S. Ding, and C. Nolder, *Inequalities for Differential Forms*, Springer, New York, NY, USA, 2009.
- [2] S. Ding, “ $L(\varphi, \mu)$ -averaging domains and Poincaré inequalities with Orlicz norms,” *Nonlinear Analysis A: Theory and Methods*, vol. 73, no. 1, pp. 256–265, 2010.
- [3] S. Ding and B. Liu, “A singular integral of the composite operator,” *Applied Mathematics Letters*, vol. 22, no. 8, pp. 1271–1275, 2009.
- [4] S. Ding and C. A. Nolder, “Weighted Poincaré inequalities for solutions to A -harmonic equations,” *Illinois Journal of Mathematics*, vol. 46, no. 1, pp. 199–205, 2002.
- [5] C. A. Nolder, “Global integrability theorems for A -harmonic tensors,” *Journal of Mathematical Analysis and Applications*, vol. 247, no. 1, pp. 236–245, 2000.
- [6] S. Ding, “Weighted Caccioppoli-type estimates and weak reverse Hölder inequalities for A -harmonic tensors,” *Proceedings of the American Mathematical Society*, vol. 127, no. 9, pp. 2657–2664, 1999.
- [7] S. Ding and B. Liu, “Generalized Poincaré inequalities for solutions to the A -harmonic equation in certain domains,” *Journal of Mathematical Analysis and Applications*, vol. 252, no. 2, pp. 538–548, 2000.
- [8] S. Ding, “ $L^p(\mu)$ -averaging domains and the quasi-hyperbolic metric,” *Computers & Mathematics with Applications*, vol. 47, no. 10-11, pp. 1611–1618, 2004.
- [9] G. Bao, “ $A, (\lambda)$ -weighted integral inequalities for A -harmonic tensors,” *Journal of Mathematical Analysis and Applications*, vol. 247, no. 2, pp. 466–477, 2000.
- [10] Y. Wang and C. Wu, “Global Poincaré inequalities for Green’s operator applied to the solutions of the nonhomogeneous A -harmonic equation,” *Computers & Mathematics with Applications*, vol. 47, no. 10-11, pp. 1545–1554, 2004.
- [11] H. Cartan, *Differential Forms*, Houghton Mifflin, Boston, Mass, USA, 1970.
- [12] T. Iwaniec and A. Lutoborski, “Integral estimates for null Lagrangians,” *Archive for Rational Mechanics and Analysis*, vol. 125, no. 1, pp. 25–79, 1993.
- [13] Y. Xing, “Weighted Poincaré-type estimates for conjugate A -harmonic tensors,” *Journal of Inequalities and Applications*, no. 1, pp. 1–6, 2005.
- [14] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and Their Applications*, vol. 88, Academic Press, New York, NY, USA, 1980.
- [15] S. M. Buckley and P. Koskela, “Orlicz-Hardy inequalities,” *Illinois Journal of Mathematics*, vol. 48, no. 3, pp. 787–802, 2004.