Research Article

Limit Cycle Bifurcations by Perturbing a Compound Loop with a Cusp and a Nilpotent Saddle

Huanhuan Tian and Maoan Han

Department of Mathematics, Shanghai Normal University, Shanghai 200234, China

Correspondence should be addressed to Maoan Han; mahan@shnu.edu.cn

Received 13 April 2014; Accepted 14 June 2014; Published 16 July 2014

Academic Editor: Tonghua Zhang

Copyright © 2014 H. Tian and M. Han. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the expansions of the first order Melnikov functions for general near-Hamiltonian systems near a compound loop with a cusp and a nilpotent saddle. We also obtain formulas for the first coefficients appearing in the expansions and then establish a bifurcation theorem on the number of limit cycles. As an application example, we give a lower bound of the maximal number of limit cycles for a polynomial system of Liénard type.

1. Introduction

Consider a planar system of the form

$$\dot{x} = H_y + \epsilon p(x, y, \delta), \quad \dot{y} = -H_x + \epsilon q(x, y, \delta), \quad (1)$$

where ϵ is a small parameter and $H(x, y)$, $p(x, y, \delta)$, and $q(x, y, \delta)$ are C^∞ functions in $(x, y) \in \mathbb{R}^2$ and $\delta \in D \subset \mathbb{R}^m$ with D bounded. For $\epsilon = 0$, (1) becomes

$$\dot{x} = H_y, \quad \dot{y} = -H_x, \quad (2)$$

which is a Hamiltonian system. As we know, the system (1) is said to be a near-Hamiltonian system. For (1), the main task is to study the number of limit cycles which are bifurcated from periodic orbits of the unperturbed system (2). On this aspect, the first order Melnikov function of (1) plays an important role. We can use the expansions of it near Hamiltonian values corresponding to a center or an invariant loop to find its zeros and hence the number of limit cycles. See a survey article [1]. There have been many works on this topic. For the study of general near-Hamiltonian systems, see [2–12]; and especially for the system (2) with the elliptic case, one can see [13–17] and references therein. In [2–4], the number of limit cycles of the system (1) near a homoclinic loop with a cusp of order one or two or a nilpotent saddle of order one (for the definition of an order of a cusp or nilpotent saddle, see [5]) was studied. In the heteroclinic case with two hyperbolic

saddles, a hyperbolic saddle and a cusp of order one, or two cusps of order one or two, the number of limit cycles of the system (1) was studied in [5, 8, 9], respectively. In this paper, we suppose that the unperturbed system (2) has a compound loop consisting of a cusp S_1 of order one, a nilpotent saddle S_2 of order one, a homoclinic loop to S_2 , and two heteroclinic orbits connecting S_1 and S_2 , as shown in Figure 1. We aim to study the number of limit cycles of (1) near the loop for $\epsilon \neq 0$ small.

2. Main Results with Proof

Now consider the C^∞ systems (1) and (2). Suppose that (2) has a compound loop denoted by $L_0 = L_1 \cup L_2 \cup L_3 \cup \{S_1, S_2\}$ and defined by equation $H(x, y) = 0$, where $S_1(x_1, y_1)$ is a cusp and $S_2(x_2, y_2)$ is a nilpotent saddle both having order one, L_1, L_2 are heteroclinic orbits satisfying $\omega(L_1) = \alpha(L_2) = S_2$ and $\omega(L_2) = \alpha(L_1) = S_1$, and L_3 is a homoclinic loop to S_2 . Then, the level curves of $H(x, y)$ define two families of periodic orbits L_{h1} and L_{h2} for h on one side of $h = 0$ and a family of periodic orbits L_{h3} for h on another side of $h = 0$. For the definiteness, let both L_{h1} and L_{h2} exist for $0 < -h \ll 1$ and L_{h3} exist for $0 < h \ll 1$. Thus, we have three Melnikov functions

$$M_i(h, \delta) = \oint_{L_{hi}} q dx - p dy, \quad i = 1, 2, 3. \quad (3)$$

FIGURE 1: Compound loop with a cusp and a nilpotent saddle.

FIGURE 2

Let U_i denote a closed set with diameter $\epsilon_0 > 0$ and with center at S_i , $i = 1, 2$. See Figure 2(a). And further introduce

$$L_{h1}^{(j)} = L_{h1} \cap U_j, \quad j = 1, 2,$$

$$L_{h1}^{(3)} = \text{Cl.} \left(L_{h1} - \bigcup_{j=1}^2 L_{h1}^{(j)} \right), \quad L_{h2}^{(1)} = L_{h2} \cap U_2,$$

$$L_{h2}^{(2)} = L_{h2} - L_{h2}^{(1)}, \quad L_{h3}^{(1)} = L_{h3} \cap U_1, \quad (4)$$

$$L_{h3} \cap U_2 = L_{h3}^{(2)} \cup L_{h3}^{(3)} \quad (\text{as shown in Figure 2(b)}),$$

$$L_{h3}^{(4)} = \text{Cl.} \left(L_{h3} - \bigcup_{j=1}^3 L_{h3}^{(j)} \right).$$

Here the Cl. denotes the closure of a set. Then by (3) and (4), for ϵ_0 sufficiently small we can write

$$M_1(h, \delta) = I_{11}(h, \delta) + I_{12}(h, \delta) + I_{13}(h, \delta) \quad (5)$$

for $0 \leq -h \ll 1$,

$$M_2(h, \delta) = I_{21}(h, \delta) + I_{22}(h, \delta) \quad (6)$$

for $0 \leq -h \ll 1$,

$$M_3(h, \delta) = I_{31}(h, \delta) + I_{32}(h, \delta) + I_{33}(h, \delta) + I_{34}(h, \delta) \quad (7)$$

for $0 \leq h \ll 1$,

where

$$I_{ij}(h, \delta) = \int_{L_{hi}^{(j)}} q dx - p dy, \quad i \in \{1, 2, 3\}, \quad j \in \{1, 2, 3, 4\}. \quad (8)$$

By [5], there exist two transformations of the form

$$(x, y)^T = T_i(u, v)^T + S_i, \quad i = 1, 2, \quad (9)$$

where T_i is a 2×2 matrix satisfying $\det T_i = 1$ such that (1) becomes

$$\dot{u} = H_{iv} + \epsilon p_i(u, v, \delta), \quad \dot{v} = -H_{iu} + \epsilon q_i(u, v, \delta), \quad (10)$$

where

$$H_1(u, v) = \frac{v^2}{2} + \sum_{k+j \geq 3} \tilde{h}_{kj} u^k v^j,$$

$$H_1(u, \varphi_1(u)) = \tilde{h}_3 u^3 + O(u^4), \quad \tilde{h}_3 < 0, \quad (11)$$

$$H_{1v}(u, \varphi_1(u)) = 0, \quad p_1(u, v, \delta) = \sum_{i+j \geq 0} \tilde{a}_{ij} u^i v^j,$$

$$q_1(u, v, \delta) = \sum_{i+j \geq 0} \tilde{b}_{ij} u^i v^j,$$

$$H_2(u, v) = \frac{v^2}{2} + \sum_{k+j \geq 3} \bar{h}_{kj} u^k v^j,$$

$$H_2(u, \varphi_2(u)) = \bar{h}_4 u^4 + O(u^5), \quad \bar{h}_4 < 0,$$

$$H_{2v}(u, \varphi_2(u)) = 0, \quad (12)$$

$$p_2(u, v, \delta) = \sum_{i+j \geq 0} \bar{a}_{ij} u^i v^j,$$

$$q_2(u, v, \delta) = \sum_{i+j \geq 0} \bar{b}_{ij} u^i v^j,$$

for (u, v) near $(0, 0)$. Note that $qdx - pdy = q_1 du - p_1 dv$ for (x, y) near S_1 and $qdx - pdy = q_2 du - p_2 dv$ for (x, y) near S_2 . Then we have

$$I_{i1}(h, \delta) = \int_{\tilde{L}_{hi}^{(1)}} q_1 du - p_1 dv, \quad i = 1, 3, \quad (13)$$

$$I_{21}(h, \delta) = \int_{\tilde{L}_{h2}^{(1)}} q_2 du - p_2 dv, \quad (14)$$

$$I_{12}(h, \delta) = \int_{\tilde{L}_{h1}^{(2)}} q_2 du - p_2 dv, \quad (15)$$

$$I_{3j}(h, \delta) = \int_{\tilde{L}_{h3}^{(j)}} q_2 du - p_2 dv, \quad j = 2, 3,$$

where $\tilde{L}_{hi}^{(1)}$ denote the image of $L_{hi}^{(1)}$ under T_1 and $\tilde{L}_{h2}^{(1)}, \tilde{L}_{h1}^{(2)}$, and $\tilde{L}_{h3}^{(j)}$ denote the image of $L_{h2}^{(1)}, L_{h1}^{(2)}$, and $L_{h3}^{(j)}$ under T_2 , respectively. Then, by using [3, 4] we can obtain the following two lemmas, respectively.

Lemma 1. Consider system (10) with $i = 1$ and suppose (11), (13) hold. Then there are constants $B_{00}, B_{00}^*, B_{10}, B_{10}^*$ satisfying

$$B_{00} = \frac{3}{5} \int_0^1 \frac{dv}{\sqrt{v(1-v^3)}} = \frac{3}{5} \times 2.4286 \cdots > 0,$$

$$B_{00}^* = -\frac{3}{5} \int_{-\infty}^1 \frac{dv}{\sqrt{1-v^3}} = -\frac{3}{5} \times 4.2065 \cdots < 0,$$

$$B_{10} = -\frac{3}{7} \left(\int_0^1 \frac{v^{3/2} dv}{\sqrt{1-v^3}(1+\sqrt{1-v^3})} - 2 \right) > 0,$$

$$\begin{aligned} B_{10}^* &= \frac{3}{7} \left(\int_1^{-1} \frac{v dv}{\sqrt{1-v^3}} - \int_0^1 \frac{v^{3/2} dv}{\sqrt{1+v^3}(1+\sqrt{1+v^3})} - 2 \right) < 0 \end{aligned} \quad (16)$$

such that

$$\begin{aligned} I_{11}(h, \delta) &= B_{00}c_1(S_1, \delta)|h|^{5/6} + B_{10}c_3(S_1, \delta)|h|^{7/6} \\ &\quad - \frac{1}{11}B_{00}c_4(S_1, \delta)|h|^{11/6} + O(h^2) + N_{11}(h, \delta) \end{aligned} \quad (17)$$

for $0 < -h \ll 1$,

$$\begin{aligned} I_{31}(h, \delta) &= B_{00}^*c_1(S_1, \delta)h^{5/6} + B_{10}^*c_3(S_1, \delta)h^{7/6} \\ &\quad + \frac{1}{11}B_{00}^*c_4(S_1, \delta)h^{11/6} + O(h^2) + N_{31}(h, \delta) \end{aligned} \quad (18)$$

for $0 < h \ll 1$, where $N_{i1}(h, \delta) \in C^\omega$ at $h = 0$ with $N_{i1}(0, \delta) = O(\epsilon_0)$, $i = 1, 3$, and

$$\begin{aligned} c_1(S_1, \delta) &= 2\sqrt{2}\tilde{h}_3^{-1/3}(\tilde{a}_{10} + \tilde{b}_{01}), \\ c_3(S_1, \delta) &= 2\sqrt{2}\tilde{h}_3^{-5/3} \\ &\quad \times \left[\tilde{h}_3(2\tilde{a}_{20} + \tilde{b}_{11}) \right. \\ &\quad \left. + \frac{1}{3}(\tilde{h}_{21}^2 - 2\tilde{h}_4 - 3\tilde{h}_3\tilde{h}_{12})(\tilde{a}_{10} + \tilde{b}_{01}) \right], \\ c_4(S_1, \delta) &= 9\mu_1^{-1}\tilde{\alpha}_{01} \\ &\quad - 2\mu_1^{-7} \left[(20\mu_2^3 - 20\mu_1\mu_2\mu_3 + 4\mu_1^2\mu_4)\tilde{\alpha}_{00} \right. \\ &\quad \left. + (4\mu_1^2\mu_3 - 10\mu_1\mu_2^2)\tilde{\alpha}_{10} \right. \\ &\quad \left. + 4\mu_1^2\mu_2\tilde{\alpha}_{20} - \mu_1^3\tilde{\alpha}_{30} \right], \end{aligned} \quad (19)$$

where

$$\begin{aligned} \mu_1 &= \tilde{h}_3^{1/3}, \quad \mu_2 = \frac{1}{3}\tilde{h}_3^{-2/3}\tilde{h}_4, \\ \mu_3 &= \frac{1}{9}\tilde{h}_3^{-5/3}(3\tilde{h}_3\tilde{h}_5 - \tilde{h}_4^2), \\ \mu_4 &= \frac{1}{81}\tilde{h}_3^{-8/3}(27\tilde{h}_3^2\tilde{h}_6 - 18\tilde{h}_3\tilde{h}_4\tilde{h}_5 + 5\tilde{h}_4^3), \\ \tilde{h}_3 &= \tilde{h}_{30}, \quad \tilde{h}_4 = -\frac{1}{2}\tilde{h}_{21}^2 + \tilde{h}_{40}, \\ \tilde{h}_5 &= \tilde{h}_{12}\tilde{h}_{21}^2 - \tilde{h}_{21}\tilde{h}_{31} + \tilde{h}_{50}, \\ \tilde{h}_6 &= -2\tilde{h}_{12}^2\tilde{h}_{21}^2 - \tilde{h}_{03}\tilde{h}_{21}^3 + \tilde{h}_{21}^2\tilde{h}_{22} \\ &\quad + 2\tilde{h}_{12}\tilde{h}_{21}\tilde{h}_{31} - \frac{1}{2}\tilde{h}_{31}^2 - \tilde{h}_{21}\tilde{h}_{41} + \tilde{h}_{60}, \\ \tilde{\alpha}_{00} &= 2\sqrt{2}(\tilde{a}_{10} + \tilde{b}_{01}), \\ \tilde{\alpha}_{10} &= 2\sqrt{2}[-\tilde{h}_{12}(\tilde{a}_{10} + \tilde{b}_{01}) + 2\tilde{a}_{20} + \tilde{b}_{11}], \\ \tilde{\alpha}_{20} &= 2\sqrt{2} \left[(\tilde{a}_{10} + \tilde{b}_{01}) \left(3\tilde{h}_{03}\tilde{h}_{21} - \tilde{h}_{22} + \frac{3}{2}\tilde{h}_{12}^2 \right) - 2\tilde{h}_{12}\tilde{a}_{20} \right. \\ &\quad \left. - \tilde{h}_{12}\tilde{b}_{11} + 3\tilde{a}_{30} + \tilde{b}_{21} - \tilde{a}_{11}\tilde{h}_{21} - 2\tilde{b}_{02}\tilde{h}_{21} \right], \\ \tilde{\alpha}_{30} &= 2\sqrt{2} \left[(\tilde{a}_{10} + \tilde{b}_{01}) \left(3\tilde{h}_{13}\tilde{h}_{21} + 3\tilde{h}_{03}\tilde{h}_{31} + 3\tilde{h}_{12}\tilde{h}_{22} \right) \right. \\ &\quad \left. - 15\tilde{h}_{12}\tilde{h}_{03}\tilde{h}_{21} - \frac{5}{2}\tilde{h}_{12}^3 - \tilde{h}_{32} \right) \\ &\quad + (3\tilde{a}_{11} + 6\tilde{b}_{02})\tilde{h}_{12}\tilde{h}_{21} - 2(\tilde{b}_{12} + \tilde{a}_{21})\tilde{h}_{21} \\ &\quad \left. - (\tilde{a}_{11} + 2\tilde{b}_{02})\tilde{h}_{31} + 4\tilde{a}_{40} + \tilde{b}_{31} \right] \end{aligned}$$

$$\begin{aligned}
& + \left(3\tilde{b}_{11} + 6\tilde{a}_{20} \right) \tilde{h}_{03} \tilde{h}_{21} - \left(2\tilde{a}_{20} + \tilde{b}_{11} \right) \tilde{h}_{22} \\
& + \left(3\tilde{a}_{20} + \frac{3}{2}\tilde{b}_{11} \right) \tilde{h}_{12}^2 - \left(3\tilde{a}_{30} + \tilde{b}_{21} \right) \tilde{h}_{12} \Big], \\
\tilde{\alpha}_{01} = 2\sqrt{2} & \left[\frac{2}{3}\tilde{a}_{12} + 2\tilde{b}_{03} - 2\tilde{h}_{03}\tilde{a}_{11} \right. \\
& \left. - 4\tilde{h}_{03}\tilde{b}_{02} + \left(\tilde{a}_{10} + \tilde{b}_{01} \right) \left(5\tilde{h}_{03}^2 - 2\tilde{h}_{04} \right) \right]. \quad (20)
\end{aligned}$$

Lemma 2. Consider system (10) with $i = 2$ and suppose (12), (15) hold. Then we have

$$\begin{aligned}
I_{12}(h, \delta) = & c_1(S_2, \delta) |h|^{3/4} - c_2(S_2, \delta) h \ln |h| \\
& + c_4(S_2, \delta) |h|^{5/4} + c_5(S_2, \delta) |h|^{7/4} \quad (21) \\
& - c_6(S_2, \delta) h^2 \ln |h| + O(h^2) + N_{12}(h, \delta)
\end{aligned}$$

for $0 < -h \ll 1$,

$$\begin{aligned}
I_{32}(h, \delta) = & \frac{1}{2}c_1^*(S_2, \delta) h^{3/4} + \frac{1}{2}c_3^*(S_2, \delta) h^{5/4} \\
& + \frac{1}{2}c_4^*(S_2, \delta) h^{7/4} + O(h^2) + N_{32}(h, \delta), \quad (22) \\
I_{33}(h, \delta) = & \frac{1}{2}c_1^*(S_2, \delta) h^{3/4} + \frac{1}{2}c_3^*(S_2, \delta) h^{5/4} \\
& + \frac{1}{2}c_4^*(S_2, \delta) h^{7/4} + O(h^2) + N_{33}(h, \delta)
\end{aligned}$$

for $0 < h \ll 1$, where $N_{ij}(h, \delta) \in C^\omega$ at $h = 0$ with $N_{ij}(0, \delta) = O(\epsilon_0)$, $(i, j) \in \{(1, 2), (3, 2), (3, 3)\}$, and

$$\begin{aligned}
c_1(S_2, \delta) = & -2\sqrt{2} |\tilde{A}_0| |\tilde{h}_4|^{-1/4} (\tilde{a}_{10} + \tilde{b}_{01}), \\
c_2(S_2, \delta) = & -\frac{\sqrt{2}}{4} |\tilde{h}_4|^{-1/2} (2\tilde{a}_{20} + \tilde{b}_{11}) + O(\tilde{a}_{10} + \tilde{b}_{01}), \\
c_4(S_2, \delta) = & |\tilde{A}_2| \left[\left(\frac{21}{32} \tilde{h}_5^{-2} - \frac{3}{4} \tilde{h}_4 \tilde{h}_6 \right) |\tilde{h}_4|^{-11/4} \tilde{\alpha}_{00} \right. \\
& \left. + \frac{3}{4} |\tilde{h}_4|^{-7/4} \tilde{h}_5 \tilde{\alpha}_{10} + |\tilde{h}_4|^{-3/4} \tilde{\alpha}_{20} \right], \\
c_5(S_2, \delta) = & \frac{1}{7} |\tilde{A}_0| \left[6d_1^{-1} \tilde{\alpha}_{01} \right. \\
& + d_1^{-9} (105d_1 d_2^2 d_3 - 30d_1^2 d_2 d_4 \\
& - 15d_1^2 d_3^2 + 5d_1^3 d_5 - 70d_2^4) \tilde{\alpha}_{00} \\
& + d_1^{-8} (35d_2^3 - 30d_1 d_2 d_3 + 5d_1^2 d_4) \tilde{\alpha}_{10} \\
& - d_1^{-7} (15d_2^2 - 5d_1 d_3) \tilde{\alpha}_{20} \\
& \left. + 5d_1^{-6} d_2 \tilde{\alpha}_{30} - d_1^{-5} \tilde{\alpha}_{40} \right],
\end{aligned}$$

$$\begin{aligned}
c_6(S_2, \delta) = & -\frac{1}{32} \left[-6d_1^{-3} d_2 \tilde{\alpha}_{01} + 3d_1^{-2} \tilde{\alpha}_{11} \right. \\
& - d_1^{-11} (504d_1 d_2^3 d_3 - 168d_1^2 d_2 d_3^2 \\
& + 42d_1^3 d_3 d_4 - 168d_1^2 d_2^2 d_4 \\
& + 42d_1^3 d_2 d_5 - 252d_2^5 - 6d_1^4 d_6) \tilde{\alpha}_{00} \\
& + d_1^{-10} (168d_1 d_2^2 d_3 - 126d_2^4 \\
& - 42d_1^2 d_2 d_4 - 21d_1^2 d_3^2 + 6d_1^3 d_5) \tilde{\alpha}_{10} \\
& - d_1^{-9} (42d_1 d_2 d_3 - 56d_2^3 - 6d_1^2 d_4) \tilde{\alpha}_{20} \\
& + d_1^{-8} (6d_1 d_3 - 21d_2^2) \tilde{\alpha}_{30} \\
& \left. + 6d_1^{-7} d_2 \tilde{\alpha}_{40} - d_1^{-6} \tilde{\alpha}_{50} \right], \\
c_1^*(S_2, \delta) = & -D_1 c_1(S_2, \delta), \quad c_3^*(S_2, \delta) = -D_2 c_4(S_2, \delta), \\
c_4^*(S_2, \delta) = & D_1 c_5(S_2, \delta), \quad (23)
\end{aligned}$$

where $D_1 = 2|\tilde{A}_0|/|\tilde{A}_0|$, $D_2 = 2|\tilde{A}_1|/|\tilde{A}_2|$, $\tilde{A}_0$, $\tilde{A}_0$, $\tilde{A}_1$, and $\tilde{A}_2$ are constants, given by

$$\begin{aligned}
\tilde{A}_0 = & \frac{2}{3} \int_0^\infty \frac{dv}{\sqrt{1+v^4}} \approx 1.236049785 > 0, \\
\tilde{A}_0 = & -\frac{2}{3} \int_0^1 \frac{dv}{\sqrt{1-v^4}} \\
= & -\frac{\sqrt{2}\pi^{3/2}}{6[\Gamma(3/4)]^2} \approx -0.8740191847 < 0, \\
\tilde{A}_1 = & -\frac{2}{5} \int_0^\infty \frac{dv}{\sqrt{1+v^4} [\sqrt{v^2 + \sqrt{1+v^4}}]} \\
\approx & -0.3388852337 < 0, \\
\tilde{A}_2 = & \frac{2}{5} \left[1 - \int_0^1 \frac{v^2 dv}{\sqrt{1-v^4} (1 + \sqrt{1-v^4})} \right] \\
\approx & 0.2396280470 > 0, \\
d_1 = & |\tilde{h}_4|^{1/4}, \quad d_2 = -\frac{1}{4} |\tilde{h}_4|^{-3/4} \tilde{h}_5, \\
d_3 = & \frac{1}{32} |\tilde{h}_4|^{-7/4} (8\tilde{h}_4 \tilde{h}_6 - 3\tilde{h}_5^2), \\
d_4 = & -\frac{1}{128} |\tilde{h}_4|^{-11/4} (7\tilde{h}_5^3 + 32\tilde{h}_4 \tilde{h}_7 - 24\tilde{h}_4 \tilde{h}_5 \tilde{h}_6),
\end{aligned}$$

$$d_5 = \frac{1}{2048} |\bar{h}_4|^{-15/4} \left(512 \bar{h}_4^3 \bar{h}_8 - 192 \bar{h}_4^2 \bar{h}_6^2 - 384 \bar{h}_4^2 \bar{h}_5 \bar{h}_7 + 336 \bar{h}_4 \bar{h}_5^2 \bar{h}_6 - 77 \bar{h}_5^4 \right),$$

$$d_6 = -\frac{1}{8192} |\bar{h}_4|^{-19/4} \left(2048 \bar{h}_4^4 \bar{h}_9 + 1344 \bar{h}_4^2 \bar{h}_5^2 \bar{h}_6^2 - 1536 \bar{h}_4^3 \bar{h}_5 \bar{h}_8 - 1536 \bar{h}_4^3 \bar{h}_6 \bar{h}_7 + 1344 \bar{h}_4 \bar{h}_5^2 \bar{h}_7 - 1232 \bar{h}_4 \bar{h}_5^3 \bar{h}_6 + 231 \bar{h}_5^5 \right);$$

$$\bar{\alpha}_{00} = 2\sqrt{2} (\bar{a}_{10} + \bar{b}_{01}),$$

$$\bar{\alpha}_{10} = 2\sqrt{2} \left[-\bar{h}_{12} (\bar{a}_{10} + \bar{b}_{01}) + 2\bar{a}_{20} + \bar{b}_{11} \right],$$

$$\bar{\alpha}_{20} = 2\sqrt{2} \left[(\bar{a}_{10} + \bar{b}_{01}) \left(3\bar{h}_{03} \bar{h}_{21} - \bar{h}_{22} + \frac{3}{2} \bar{h}_{12}^2 \right) - 2\bar{h}_{12} \bar{a}_{20} - \bar{h}_{12} \bar{b}_{11} + 3\bar{a}_{30} + \bar{b}_{21} - \bar{a}_{11} \bar{h}_{21} - 2\bar{b}_{02} \bar{h}_{21} \right],$$

$$\begin{aligned} \bar{\alpha}_{30} = 2\sqrt{2} \left[(\bar{a}_{10} + \bar{b}_{01}) \right. \\ \times \left(3\bar{h}_{13} \bar{h}_{21} + 3\bar{h}_{03} \bar{h}_{31} + 3\bar{h}_{12} \bar{h}_{22} \right. \\ \left. - 15\bar{h}_{12} \bar{h}_{03} \bar{h}_{21} - \frac{5}{2} \bar{h}_{12}^3 - \bar{h}_{32} \right) \\ \left. + (3\bar{a}_{11} + 6\bar{b}_{02}) \bar{h}_{12} \bar{h}_{21} - 2(\bar{b}_{12} + \bar{a}_{21}) \bar{h}_{21} \right. \\ \left. - (\bar{a}_{11} + 2\bar{b}_{02}) \bar{h}_{31} + 4\bar{a}_{40} + \bar{b}_{31} \right. \\ \left. + (3\bar{b}_{11} + 6\bar{a}_{20}) \bar{h}_{03} \bar{h}_{21} \right. \\ \left. - (2\bar{a}_{20} + \bar{b}_{11}) \bar{h}_{22} + \left(3\bar{a}_{20} + \frac{3}{2} \bar{b}_{11} \right) \bar{h}_{12}^2 \right. \\ \left. - (3\bar{a}_{30} + \bar{b}_{21}) \bar{h}_{12} \right], \end{aligned}$$

$$\begin{aligned} \bar{\alpha}_{40} = 2\sqrt{2} \left[n_1 (\bar{a}_{10} + \bar{b}_{01}) \right. \\ \left. - \left(\frac{5}{2} \bar{h}_{12}^3 - 3\bar{h}_{12} \bar{h}_{22} + 15\bar{h}_{12} \bar{h}_{03} \bar{h}_{21} + \bar{h}_{32} \right. \right. \\ \left. \left. - 3\bar{h}_{13} \bar{h}_{21} - 3\bar{h}_{03} \bar{h}_{31} \right) (2\bar{a}_{20} + \bar{b}_{11}) \right. \\ \left. + \left(\frac{3}{2} \bar{h}_{12}^2 + 3\bar{h}_{03} \bar{h}_{21} - \bar{h}_{22} \right) \right. \\ \times (3\bar{a}_{30} + \bar{b}_{21} - \bar{a}_{11} \bar{h}_{21} - 2\bar{b}_{02} \bar{h}_{21}) \\ \left. - \bar{h}_{12} (4\bar{a}_{40} + \bar{b}_{31} - 2\bar{a}_{21} \bar{h}_{21} \right. \\ \left. - 2\bar{b}_{12} \bar{h}_{21} + 2\bar{a}_{11} \bar{h}_{12} \bar{h}_{21} + 4\bar{b}_{02} \bar{h}_{12} \bar{h}_{21} \right. \\ \left. - \bar{a}_{11} \bar{h}_{31} - 2\bar{b}_{02} \bar{h}_{31}) + n_3 \right], \end{aligned}$$

$$\begin{aligned} \bar{\alpha}_{50} = 2\sqrt{2} \left[n_2 (\bar{a}_{10} + \bar{b}_{01}) + n_1 (2\bar{a}_{20} + \bar{b}_{11}) \right. \\ \left. - \left(\frac{5}{2} \bar{h}_{12}^3 - 3\bar{h}_{12} \bar{h}_{22} + 15\bar{h}_{12} \bar{h}_{03} \bar{h}_{21} + \bar{h}_{32} \right. \right. \\ \left. \left. - 3\bar{h}_{13} \bar{h}_{21} - 3\bar{h}_{03} \bar{h}_{31} \right) \right. \\ \times (3\bar{a}_{30} + \bar{b}_{21} - \bar{a}_{11} \bar{h}_{21} - 2\bar{b}_{02} \bar{h}_{21}) \\ \left. + \left(\frac{3}{2} \bar{h}_{12}^2 + 3\bar{h}_{03} \bar{h}_{21} - \bar{h}_{22} \right) \right. \\ \times (4\bar{a}_{40} + \bar{b}_{31} - 2\bar{a}_{21} \bar{h}_{21} \\ \left. - 2\bar{b}_{12} \bar{h}_{21} + 2\bar{a}_{11} \bar{h}_{12} \bar{h}_{21} \right. \\ \left. + 4\bar{b}_{02} \bar{h}_{12} \bar{h}_{21} - \bar{a}_{11} \bar{h}_{31} - 2\bar{b}_{02} \bar{h}_{31}) \right. \\ \left. - \bar{h}_{12} n_3 + n_4 \right], \end{aligned}$$

$$\begin{aligned} \bar{\alpha}_{01} = 2\sqrt{2} \left[\frac{2}{3} \bar{a}_{12} + 2\bar{b}_{03} - 2\bar{h}_{03} \bar{a}_{11} - 4\bar{h}_{03} \bar{b}_{02} \right. \\ \left. + (\bar{a}_{10} + \bar{b}_{01}) (5\bar{h}_{03}^2 - 2\bar{h}_{04}) \right], \end{aligned}$$

$$\begin{aligned} \bar{\alpha}_{11} = 2\sqrt{2} \left[(\bar{a}_{10} + \bar{b}_{01}) (10\bar{h}_{04} \bar{h}_{12} - 2\bar{h}_{14} \right. \\ \left. + 10\bar{h}_{03} \bar{h}_{13} - 35\bar{h}_{12} \bar{h}_{03}^2) \right. \\ \left. + (\bar{a}_{11} + 2\bar{b}_{02}) (-2\bar{h}_{13} + 10\bar{h}_{12} \bar{h}_{03}) \right. \\ \left. - 2\bar{h}_{12} (\bar{a}_{12} + 3\bar{b}_{03}) \right. \\ \left. + (2\bar{a}_{20} + \bar{b}_{11}) (5\bar{h}_{03}^2 - 2\bar{h}_{04}) \right. \\ \left. - 4\bar{h}_{03} (\bar{b}_{12} + \bar{a}_{21}) + \frac{4}{3} \bar{a}_{22} + 2\bar{b}_{13} \right]; \end{aligned}$$

$$\bar{h}_4 = -\frac{1}{2} \bar{h}_{21}^2 + \bar{h}_{40}, \quad \bar{h}_5 = \bar{h}_{12} \bar{h}_{21}^2 - \bar{h}_{21} \bar{h}_{31} + \bar{h}_{50},$$

$$\begin{aligned} \bar{h}_6 = -2\bar{h}_{12}^2 \bar{h}_{21}^2 - \bar{h}_{03} \bar{h}_{21}^3 + \bar{h}_{21}^2 \bar{h}_{22} \\ + 2\bar{h}_{12} \bar{h}_{21} \bar{h}_{31} - \frac{1}{2} \bar{h}_{31}^2 - \bar{h}_{21} \bar{h}_{41} + \bar{h}_{60}, \end{aligned}$$

$$\begin{aligned} \bar{h}_7 = -4\bar{h}_{22} \bar{h}_{21}^2 \bar{h}_{12} + \bar{h}_{70} - \bar{h}_{51} \bar{h}_{21} - \bar{h}_{13} \bar{h}_{21}^3 \\ + \bar{h}_{32} \bar{h}_{21}^2 + 4\bar{h}_{12}^3 \bar{h}_{21}^2 - \bar{h}_{31} \bar{h}_{41} + \bar{h}_{12} \bar{h}_{31}^2 \\ + 2\bar{h}_{22} \bar{h}_{21} \bar{h}_{31} + 6\bar{h}_{03} \bar{h}_{21}^3 \bar{h}_{12} - 3\bar{h}_{03} \bar{h}_{21}^2 \bar{h}_{31} \\ - 4\bar{h}_{12}^2 \bar{h}_{21} \bar{h}_{31} + 2\bar{h}_{12} \bar{h}_{21} \bar{h}_{41}, \end{aligned}$$

$$\begin{aligned}
\bar{h}_8 &= \bar{h}_{80} - \bar{h}_{23} \bar{h}_{21}^3 - \bar{h}_{61} \bar{h}_{21} + \bar{h}_{04} \bar{h}_{21}^4 + \bar{h}_{42} \bar{h}_{21}^2 \\
&\quad - 2 \bar{h}_{12}^2 \bar{h}_{31}^2 - 8 \bar{h}_{12}^4 \bar{h}_{21}^2 - 2 \bar{h}_{22}^2 \bar{h}_{21}^2 \\
&\quad - \frac{9}{2} \bar{h}_{03}^2 \bar{h}_{21}^4 - \bar{h}_{31} \bar{h}_{51} + \bar{h}_{22} \bar{h}_{31}^2 - \frac{1}{2} \bar{h}_{41}^2 \\
&\quad + 8 \bar{h}_{12}^3 \bar{h}_{21} \bar{h}_{31} - 4 \bar{h}_{12}^2 \bar{h}_{21} \bar{h}_{41} + 12 \bar{h}_{22} \bar{h}_{21}^2 \bar{h}_{12}^2 \\
&\quad - 24 \bar{h}_{03} \bar{h}_{21}^3 \bar{h}_{12}^2 + 2 \bar{h}_{12} \bar{h}_{31} \bar{h}_{41} + 6 \bar{h}_{12} \bar{h}_{21}^3 \bar{h}_{13} \\
&\quad - 4 \bar{h}_{12} \bar{h}_{21}^2 \bar{h}_{32} + 2 \bar{h}_{12} \bar{h}_{21} \bar{h}_{51} + 6 \bar{h}_{22} \bar{h}_{21}^3 \bar{h}_{03} \\
&\quad + 2 \bar{h}_{22} \bar{h}_{21} \bar{h}_{41} + 2 \bar{h}_{32} \bar{h}_{21} \bar{h}_{31} - 3 \bar{h}_{03} \bar{h}_{21} \bar{h}_{31}^2 \\
&\quad - 3 \bar{h}_{03} \bar{h}_{21}^2 \bar{h}_{41} - 3 \bar{h}_{13} \bar{h}_{21}^2 \bar{h}_{31} \\
&\quad - 8 \bar{h}_{12} \bar{h}_{22} \bar{h}_{21} \bar{h}_{31} + 18 \bar{h}_{12} \bar{h}_{03} \bar{h}_{21}^2 \bar{h}_{31}, \\
\bar{h}_9 &= \bar{h}_{12} \left[\bar{h}_{21}^4 \left(45 \bar{h}_{03}^2 - 8 \bar{h}_{04} \right) + 6 \bar{h}_{21}^3 \left(\bar{h}_{23} - 8 \bar{h}_{03} \bar{h}_{22} \right) \right. \\
&\quad + 2 \bar{h}_{21}^2 \left(9 \bar{h}_{13} \bar{h}_{31} - 2 \bar{h}_{42} + 6 \bar{h}_{22}^2 + 9 \bar{h}_{03} \bar{h}_{41} \right) \\
&\quad + 2 \bar{h}_{21} \left(9 \bar{h}_{03} \bar{h}_{31}^2 - 4 \bar{h}_{41} \bar{h}_{22} - 4 \bar{h}_{31} \bar{h}_{32} \right) \\
&\quad \left. - 4 \bar{h}_{22} \bar{h}_{31}^2 + \bar{h}_{41}^2 + 2 \bar{h}_{31} \bar{h}_{51} + 2 \bar{h}_{21} \bar{h}_{61} \right] \\
&\quad + 4 \bar{h}_{12}^2 \left[\bar{h}_{21} \left(3 \bar{h}_{21} \bar{h}_{32} - 6 \bar{h}_{21}^2 \bar{h}_{13} - \bar{h}_{51} \right. \right. \\
&\quad \left. \left. + 6 \bar{h}_{31} \bar{h}_{22} - 18 \bar{h}_{31} \bar{h}_{03} \bar{h}_{21} \right) - \bar{h}_{31} \bar{h}_{41} \right] \\
&\quad + 4 \bar{h}_{12}^3 \left(20 \bar{h}_{03} \bar{h}_{21}^3 + 2 \bar{h}_{41} \bar{h}_{21} - 8 \bar{h}_{22} \bar{h}_{21}^2 + \bar{h}_{31}^2 \right) \\
&\quad + \bar{h}_{21} \left(2 \bar{h}_{42} \bar{h}_{31} - 16 \bar{h}_{12}^4 \bar{h}_{31} - 4 \bar{h}_{22}^2 \bar{h}_{31} + 2 \bar{h}_{22} \bar{h}_{51} \right. \\
&\quad \left. + 2 \bar{h}_{32} \bar{h}_{41} - 3 \bar{h}_{13} \bar{h}_{31}^2 - \bar{h}_{71} - 6 \bar{h}_{03} \bar{h}_{31} \bar{h}_{41} \right) \\
&\quad + \bar{h}_{21}^2 \left(16 \bar{h}_{12}^5 - 4 \bar{h}_{32} \bar{h}_{22} - 3 \bar{h}_{31} \bar{h}_{23} \right. \\
&\quad \left. - 3 \bar{h}_{03} \bar{h}_{51} + \bar{h}_{52} + 18 \bar{h}_{22} \bar{h}_{31} \bar{h}_{03} \right) \\
&\quad + 2 \bar{h}_{22} \bar{h}_{31} \bar{h}_{41} + 6 \bar{h}_{32} \bar{h}_{21}^3 \bar{h}_{03} + 4 \bar{h}_{04} \bar{h}_{21}^3 \bar{h}_{31} \\
&\quad - \bar{h}_{41} \bar{h}_{51} - \bar{h}_{31} \bar{h}_{61} + \bar{h}_{32} \bar{h}_{31}^2 - \bar{h}_{03} \bar{h}_{31}^3 \\
&\quad - 9 \bar{h}_{03} \bar{h}_{21}^4 \bar{h}_{13} - 18 \bar{h}_{31} \bar{h}_{03}^2 \bar{h}_{21}^3 + \bar{h}_{14} \bar{h}_{21}^4 \\
&\quad - \bar{h}_{33} \bar{h}_{21}^3 + 6 \bar{h}_{22} \bar{h}_{21}^3 \bar{h}_{13} + \bar{h}_{90}; \\
n_1 &= \frac{1}{8} \left(12 \bar{h}_{22}^2 + 35 \bar{h}_{12}^4 + 24 \bar{h}_{12} \bar{h}_{32} \right. \\
&\quad - 60 \bar{h}_{12}^2 \bar{h}_{22} - 72 \bar{h}_{12} \bar{h}_{03} \bar{h}_{31} + 324 \bar{h}_{12}^2 \bar{h}_{03} \bar{h}_{21} \\
&\quad \left. + 108 \bar{h}_{03}^2 \bar{h}_{21}^2 - 72 \bar{h}_{12} \bar{h}_{13} \bar{h}_{21} - 72 \bar{h}_{22} \bar{h}_{03} \bar{h}_{21} \right),
\end{aligned}$$

$$\begin{aligned}
n_2 &= -\frac{1}{8} \left(-24 \bar{h}_{22} \bar{h}_{32} + 60 \bar{h}_{12} \bar{h}_{22}^2 + 60 \bar{h}_{12}^2 \bar{h}_{32} \right. \\
&\quad - 140 \bar{h}_{12}^3 \bar{h}_{22} + 63 \bar{h}_{12}^5 \\
&\quad - 180 \bar{h}_{12}^2 \bar{h}_{03} \bar{h}_{31} + 780 \bar{h}_{12}^3 \bar{h}_{03} \bar{h}_{21} \\
&\quad + 72 \bar{h}_{22} \bar{h}_{03} \bar{h}_{31} - 504 \bar{h}_{12} \bar{h}_{22} \bar{h}_{03} \bar{h}_{21} \\
&\quad - 216 \bar{h}_{03}^2 \bar{h}_{21} \bar{h}_{31} + 972 \bar{h}_{12} \bar{h}_{03}^2 \bar{h}_{21}^2 \\
&\quad + 72 \bar{h}_{03} \bar{h}_{21} \bar{h}_{32} + 72 \bar{h}_{22} \bar{h}_{13} \bar{h}_{21} \\
&\quad \left. - 180 \bar{h}_{12}^2 \bar{h}_{13} \bar{h}_{21} - 216 \bar{h}_{03} \bar{h}_{21}^2 \bar{h}_{13} \right),
\end{aligned}$$

$$\begin{aligned}
n_3 &= 5 \bar{a}_{50} + \bar{b}_{41} - 2 \bar{b}_{22} \bar{h}_{21} - \bar{a}_{11} \bar{h}_{41} \\
&\quad + 2 \bar{a}_{11} \bar{h}_{22} \bar{h}_{21} - 3 \bar{a}_{11} \bar{h}_{03} \bar{h}_{21}^2 + 2 \bar{a}_{11} \bar{h}_{12} \bar{h}_{31} \\
&\quad - 4 \bar{a}_{11} \bar{h}_{12}^2 \bar{h}_{21} - 2 \bar{b}_{02} \bar{h}_{41} + 4 \bar{b}_{02} \bar{h}_{22} \bar{h}_{21} \\
&\quad - 6 \bar{b}_{02} \bar{h}_{03} \bar{h}_{21}^2 + 4 \bar{b}_{02} \bar{h}_{12} \bar{h}_{31} - 8 \bar{b}_{02} \bar{h}_{12}^2 \bar{h}_{21} \\
&\quad - 2 \bar{b}_{12} \bar{h}_{31} + 4 \bar{b}_{12} \bar{h}_{12} \bar{h}_{21} - 2 \bar{a}_{21} \bar{h}_{31} \\
&\quad + 4 \bar{a}_{21} \bar{h}_{12} \bar{h}_{21} - 3 \bar{a}_{31} \bar{h}_{21} + 3 \bar{b}_{03} \bar{h}_{21}^2 + \bar{a}_{12} \bar{h}_{21}^2,
\end{aligned}$$

$$\begin{aligned}
n_4 &= 4 \bar{b}_{12} \bar{h}_{22} \bar{h}_{21} + 4 \bar{b}_{02} \bar{h}_{32} \bar{h}_{21} + 6 \bar{a}_{31} \bar{h}_{12} \bar{h}_{21} \\
&\quad + 2 \bar{a}_{11} \bar{h}_{12} \bar{h}_{41} + 4 \bar{b}_{12} \bar{h}_{12} \bar{h}_{31} - 4 \bar{a}_{11} \bar{h}_{12}^2 \bar{h}_{31} \\
&\quad + 4 \bar{b}_{02} \bar{h}_{12} \bar{h}_{41} - 12 \bar{b}_{03} \bar{h}_{21}^2 \bar{h}_{12} - 4 \bar{a}_{12} \bar{h}_{21}^2 \bar{h}_{12} \\
&\quad + 2 \bar{a}_{11} \bar{h}_{22} \bar{h}_{31} + 6 \bar{b}_{03} \bar{h}_{21} \bar{h}_{31} - 3 \bar{a}_{11} \bar{h}_{13} \bar{h}_{21}^2 \\
&\quad + 6 \bar{a}_{60} + \bar{b}_{51} - 6 \bar{b}_{12} \bar{h}_{03} \bar{h}_{21}^2 + 2 \bar{a}_{11} \bar{h}_{32} \bar{h}_{21} \\
&\quad + 4 \bar{b}_{02} \bar{h}_{22} \bar{h}_{31} + 4 \bar{a}_{21} \bar{h}_{22} \bar{h}_{21} + 2 \bar{a}_{12} \bar{h}_{21} \bar{h}_{31} \\
&\quad + 4 \bar{b}_{22} \bar{h}_{12} \bar{h}_{21} - 8 \bar{b}_{12} \bar{h}_{12}^2 \bar{h}_{21} + 16 \bar{b}_{02} \bar{h}_{12}^3 \bar{h}_{21} \\
&\quad + 8 \bar{a}_{11} \bar{h}_{12}^3 \bar{h}_{21} - 6 \bar{b}_{02} \bar{h}_{13} \bar{h}_{21}^2 - 8 \bar{a}_{21} \bar{h}_{12}^2 \bar{h}_{21} \\
&\quad - 8 \bar{b}_{02} \bar{h}_{12}^2 \bar{h}_{31} - 4 \bar{a}_{41} \bar{h}_{21} - 2 \bar{b}_{32} \bar{h}_{21} \\
&\quad - 6 \bar{a}_{21} \bar{h}_{03} \bar{h}_{21}^2 + 4 \bar{a}_{21} \bar{h}_{12} \bar{h}_{31} + 3 \bar{b}_{13} \bar{h}_{21}^2 \\
&\quad + 2 \bar{a}_{22} \bar{h}_{21}^2 - 6 \bar{a}_{11} \bar{h}_{03} \bar{h}_{21} \bar{h}_{31} + 18 \bar{a}_{11} \bar{h}_{03} \bar{h}_{21}^2 \bar{h}_{12} \\
&\quad - 8 \bar{a}_{11} \bar{h}_{12} \bar{h}_{22} \bar{h}_{21} - 12 \bar{b}_{02} \bar{h}_{03} \bar{h}_{21} \bar{h}_{31} + 36 \bar{b}_{02} \bar{h}_{03} \bar{h}_{21}^2 \bar{h}_{12} \\
&\quad - 16 \bar{b}_{02} \bar{h}_{12} \bar{h}_{22} \bar{h}_{21} - 2 \bar{b}_{22} \bar{h}_{31} - \bar{a}_{11} \bar{h}_{51} - 2 \bar{b}_{12} \bar{h}_{41} \\
&\quad - 2 \bar{a}_{21} \bar{h}_{41} - 2 \bar{b}_{02} \bar{h}_{51} - 3 \bar{a}_{31} \bar{h}_{31}.
\end{aligned}$$

(24)

For convenience, let

$$\begin{aligned} L^* &= L_1 \cup L_2, & L_1^* &= L^* \cap U_1, \\ L_2^* &= L^* \cap U_2, & L_3^* &= \text{Cl.} \left(L^* - \bigcup_{i=1}^2 L_i^* \right), \\ L_0^1 &= L_1^*, & & \\ L_0 \cap U_2 &= L_0^2 \cup L_0^3 \quad (\text{as shown in Figure 2 (b)}), \\ L_0^4 &= \text{Cl.} \left(L_0 - \bigcup_{j=1}^3 L_0^j \right). \end{aligned} \quad (25)$$

Theorem 3. Assume that system (1) has a compound loop L_0 as stated before. Then, the functions $M_i(h, \delta)$ given in (3) at $h = 0$ have the following expansions:

$$\begin{aligned} M_1(h, \delta) &= c_0(\delta) + c_1(\delta) |h|^{3/4} + B_{00} c_2(\delta) |h|^{5/6} \\ &\quad - c_3(\delta) h \ln |h| + c_4(\delta) h + B_{10} c_5(\delta) |h|^{7/6} \\ &\quad + c_6(\delta) |h|^{5/4} + c_7(\delta) |h|^{7/4} \\ &\quad - \frac{1}{11} B_{00} c_8(\delta) |h|^{11/6} - c_9(\delta) h^2 \ln |h| + O(h^2), \end{aligned} \quad (26)$$

$$\begin{aligned} M_2(h, \delta) &= \bar{c}_0(\delta) + c_1(\delta) |h|^{3/4} + c_3(\delta) h \ln |h| \\ &\quad + \bar{c}_3(\delta) h + c_6(\delta) |h|^{5/4} + c_7(\delta) |h|^{7/4} \\ &\quad + c_9(\delta) h^2 \ln |h| + O(h^2) \end{aligned} \quad (27)$$

for $0 < -h \ll 1$, and

$$\begin{aligned} M_3(h, \delta) &= \bar{c}_0(\delta) - D_1 c_1(\delta) h^{3/4} + B_{00}^* c_2(\delta) h^{5/6} \\ &\quad + \bar{c}_2(\delta) h + B_{10}^* c_5(\delta) h^{7/6} - D_2 c_6(\delta) h^{5/4} \\ &\quad + D_1 c_7(\delta) h^{7/4} + \frac{1}{11} B_{00}^* c_8(\delta) h^{11/6} + O(h^2) \end{aligned} \quad (28)$$

for $0 < h \ll 1$, where

$$\begin{aligned} c_1(\delta) &= c_1(S_2, \delta), & c_2(\delta) &= c_1(S_1, \delta), \\ c_3(\delta) &= c_2(S_2, \delta), & c_5(\delta) &= c_3(S_1, \delta), \\ c_6(\delta) &= c_4(S_2, \delta), & c_7(\delta) &= c_5(S_2, \delta), \\ c_8(\delta) &= c_4(S_1, \delta), & c_9(\delta) &= c_6(S_2, \delta), \end{aligned} \quad (29)$$

$$\begin{aligned} c_0(\delta) &= M_1(0, \delta) = \oint_{L^*} q dx - p dy = \sum_{i=1}^2 \int_{L_i} q dx - p dy, \\ \bar{c}_0(\delta) &= M_2(0, \delta) = \oint_{L_3} q dx - p dy, \\ \bar{c}_0(\delta) &= M_3(0, \delta) = c_0(\delta) + \bar{c}_0(\delta), \end{aligned} \quad (30)$$

$$\begin{aligned} c_4(\delta) &= \int_{L_1^*} (p_x + q_y - \sigma) dt \\ &\quad + \int_{L_2^*} [p_x + q_y - \eta_0 - \eta_1(x - x_2)] dt \\ &\quad + \int_{L_3^*} (p_x + q_y) dt + t_2 c_1(\delta) + t_3 c_2(\delta) + t_4 c_3(\delta), \\ \bar{c}_3(\delta) &= \oint_{L_3} [p_x + q_y - \eta_0 - \eta_1(x - x_2)] dt \\ &\quad + t_0 c_1(\delta) + t_1 c_3(\delta), \\ \bar{c}_2(\delta) &= \int_{L_1^*} (p_x + q_y - \sigma) dt + \int_{L_0^2 \cup L_0^3} (p_x + q_y - \eta_0) dt \\ &\quad + \int_{L_0^4} (p_x + q_y) dt + t_5 c_1(\delta) + t_6 c_2(\delta), \end{aligned} \quad (31)$$

where $\sigma = (p_x + q_y)|_{(x_1, y_1)}$, $\eta_0 = (p_x + q_y)|_{(x_2, y_2)}$, $\eta_1 = (p_{xx} + q_{yy})|_{(x_2, y_2)}$. In particular,

$$\begin{aligned} c_4(\delta) &= \sum_{i=1}^2 \int_{L_i} (p_x + q_y) dt, & \bar{c}_3(\delta) &= \oint_{L_3} (p_x + q_y) dt, \\ \bar{c}_2(\delta) &= \sum_{i=1}^3 \int_{L_i} (p_x + q_y) dt \end{aligned} \quad (32)$$

if $c_1(\delta) = c_2(\delta) = c_3(\delta) = 0$. Here, t_i , $i = 0, 1, \dots, 6$, are constants and B_{00} , B_{00}^* , B_{10} , B_{10}^* are given in Lemma 1.

Proof. First, by (6), (10) with $i = 2$, (12), (14), (29), and Theorem 2.2 in [4], we directly obtain (27) with $\bar{c}_0(\delta)$, $\bar{c}_3(\delta)$ given by (30) and (31), respectively. Then we study the expansions of M_1 and M_3 .

By (5), (7), (29), and Lemmas 1 and 2, we have

$$\begin{aligned} M_1(h, \delta) &= c_1(\delta) |h|^{3/4} + B_{00} c_2(\delta) |h|^{5/6} - c_3(\delta) h \ln |h| \\ &\quad + B_{10} c_5(\delta) |h|^{7/6} + c_6(\delta) |h|^{5/4} \\ &\quad + c_7(\delta) |h|^{7/4} - \frac{1}{11} B_{00} c_8(\delta) |h|^{11/6} \\ &\quad - c_9(\delta) h^2 \ln |h| + O(h^2) + N(h, \delta) \end{aligned} \quad (33)$$

for $0 < -h \ll 1$, and

$$\begin{aligned} M_3(h, \delta) = & -D_1 c_1(\delta) h^{3/4} + B_{00}^* c_2(\delta) h^{5/6} + B_{10}^* c_5(\delta) h^{7/6} \\ & - D_2 c_6(\delta) h^{5/4} + D_1 c_7(\delta) h^{7/4} \\ & + \frac{1}{11} B_{00}^* c_8(\delta) h^{11/6} + O(h^2) + N^*(h, \delta) \end{aligned} \quad (34)$$

for $0 < h \ll 1$, where

$$\begin{aligned} N(h, \delta) = & N_{11}(h, \delta) + N_{12}(h, \delta) + I_{13}(h, \delta), \\ N^*(h, \delta) = & N_{31}(h, \delta) + N_{32}(h, \delta) + N_{33}(h, \delta) + I_{34}(h, \delta). \end{aligned} \quad (35)$$

Let

$$\begin{aligned} N(h, \delta) = & c_0(\delta) + c_4(\delta) h + O(h^2), \\ N^*(h, \delta) = & \tilde{c}_0(\delta) + \tilde{c}_2(\delta) h + O(h^2). \end{aligned} \quad (36)$$

It follows further that

$$\begin{aligned} c_0(\delta) = & N_{11}(0, \delta) + N_{12}(0, \delta) + I_{13}(0, \delta) \\ = & \lim_{\epsilon_0 \rightarrow 0} [N_{11}(0, \delta) + N_{12}(0, \delta) + I_{13}(0, \delta)] \\ = & \lim_{\epsilon_0 \rightarrow 0} I_{13}(0, \delta) = \oint_{L_1 \cup L_2} q dx - p dy \\ = & \sum_{i=1}^2 \int_{L_i} q dx - p dy = M_1(0, \delta), \\ \tilde{c}_0(\delta) = & N_{31}(0, \delta) + N_{32}(0, \delta) + N_{33}(0, \delta) + I_{34}(0, \delta) \\ = & \lim_{\epsilon_0 \rightarrow 0} [N_{31}(0, \delta) + N_{32}(0, \delta) \\ & + N_{33}(0, \delta) + I_{34}(0, \delta)] \\ = & \lim_{\epsilon_0 \rightarrow 0} I_{34}(0, \delta) = \oint_{L_0} q dx - p dy \\ = & \sum_{i=1}^3 \int_{L_i} q dx - p dy = M_3(0, \delta), \end{aligned}$$

$$\begin{aligned} c_4(\delta) + O(h) = & N_h(h, \delta) = M_{1h}(h, \delta) \\ & + \frac{3}{4} c_1(\delta) |h|^{-1/4} + \frac{5}{6} B_{00}^* c_2(\delta) |h|^{-1/6} \\ & + c_3(\delta) (\ln |h| + 1) + O(|h|^{1/6}), \end{aligned}$$

$$\begin{aligned} \tilde{c}_2(\delta) + O(h) = & N_h^*(h, \delta) \\ = & M_{3h}(h, \delta) + \frac{3}{4} D_1 c_1(\delta) h^{-1/4} \\ & - \frac{5}{6} B_{00}^* c_2(\delta) h^{-1/6} + O(h^{1/6}). \end{aligned} \quad (37)$$

Then by Lemma 3.1.2 in [5], we have

$$\begin{aligned} c_4(\delta) = & N_h(0, \delta) \\ = & \lim_{h \rightarrow 0^-} \left[\oint_{L_{h1}} (p_x + q_y) dt + \frac{3}{4} c_1(\delta) |h|^{-1/4} \right. \\ & \left. + \frac{5}{6} B_{00}^* c_2(\delta) |h|^{-1/6} + c_3(\delta) (\ln |h| + 1) \right], \\ \tilde{c}_2(\delta) = & N_h^*(0, \delta) \\ = & \lim_{h \rightarrow 0^+} \left[\oint_{L_{h3}} (p_x + q_y) dt \right. \\ & \left. + \frac{3}{4} D_1 c_1(\delta) h^{-1/4} - \frac{5}{6} B_{00}^* c_2(\delta) h^{-1/6} \right]. \end{aligned} \quad (38)$$

It is easy to see that

$$\begin{aligned} \oint_{L_{h1}} (p_x + q_y) dt = & \sum_{i=1}^3 \int_{L_{h1}^{(i)}} (p_x + q_y) dt \\ = & \int_{L_{h1}^{(1)}} (p_x + q_y - \sigma) dt \\ & + \int_{L_{h1}^{(2)}} [p_x + q_y - \eta_0 \\ & - \eta_1(x - x_2)] dt \\ & + \int_{L_{h1}^{(3)}} (p_x + q_y) dt + \sigma \int_{L_{h1}^{(1)}} dt \\ & + \eta_0 \int_{L_{h1}^{(2)}} dt + \eta_1 \int_{L_{h1}^{(2)}} (x - x_2) dt, \end{aligned} \quad (39)$$

$$\begin{aligned} \oint_{L_{h3}} (p_x + q_y) dt = & \sum_{i=1}^4 \int_{L_{h3}^{(i)}} (p_x + q_y) dt = \int_{L_{h3}^{(1)}} (p_x + q_y - \sigma) dt \\ & + \int_{L_{h3}^{(2)} \cup L_{h3}^{(3)}} (p_x + q_y - \eta_0) dt + \int_{L_{h3}^{(4)}} (p_x + q_y) dt \\ & + \sigma \int_{L_{h3}^{(1)}} dt + \eta_0 \int_{L_{h3}^{(2)} \cup L_{h3}^{(3)}} dt. \end{aligned}$$

Noting $\tilde{h}_3 < 0$, $\bar{h}_4 < 0$, by (19), (23), and (29), we have

$$\begin{aligned} c_4(\delta) = & \lim_{h \rightarrow 0^-} \left[\int_{L_{h1}^{(1)}} (p_x + q_y - \sigma) dt \right. \\ & + \int_{L_{h1}^{(2)}} [p_x + q_y - \eta_0 - \eta_1(x - x_2)] dt \\ & \left. + \int_{L_{h1}^{(3)}} (p_x + q_y) dt \right] \end{aligned}$$

$$\begin{aligned}
 & + \lim_{h \rightarrow 0^-} \sigma \left[\int_{L_{h1}^{(1)}} dt - \frac{5\sqrt{2}}{3} B_{00} |\tilde{h}_3|^{-1/3} |h|^{-1/6} \right] \\
 & + \lim_{h \rightarrow 0^-} \eta_0 \left[\int_{L_{h1}^{(2)}} dt + \frac{3\sqrt{2}}{2} \tilde{A}_0 |\tilde{h}_4|^{-1/4} |h|^{-1/4} \right. \\
 & \quad \left. + O(1) (\ln |h| + 1) \right] \\
 & + \lim_{h \rightarrow 0^-} \eta_1 \left[\int_{L_{h1}^{(2)}} (x - x_2) dt \right. \\
 & \quad \left. - \frac{\sqrt{2}}{4} |\tilde{h}_4|^{-1/2} (\ln |h| + 1) \right], \\
 \tilde{c}_2(\delta) & = \lim_{h \rightarrow 0^+} \left[\int_{L_{h3}^{(1)}} (p_x + q_y - \sigma) dt \right. \\
 & \quad + \int_{L_{h3}^{(2)} \cup L_{h3}^{(3)}} (p_x + q_y - \eta_0) dt \\
 & \quad \left. + \int_{L_{h3}^{(4)}} (p_x + q_y) dt \right] \\
 & + \lim_{h \rightarrow 0^+} \sigma \left[\int_{L_{h3}^{(1)}} dt + \frac{5\sqrt{2}}{3} B_{00}^* |\tilde{h}_3|^{-1/3} h^{-1/6} \right] \\
 & + \lim_{h \rightarrow 0^+} \eta_0 \left[\int_{L_{h3}^{(2)} \cup L_{h3}^{(3)}} dt - 3\sqrt{2} \tilde{A}_0 |\tilde{h}_4|^{-1/4} h^{-1/4} \right]. \quad (40)
 \end{aligned}$$

Then by the proof of (3.13) in [3], the following equations hold:

$$\begin{aligned}
 \lim_{h \rightarrow 0^-} \left[\int_{L_{h1}^{(1)}} dt - \frac{5\sqrt{2}}{3} B_{00} |\tilde{h}_3|^{-1/3} |h|^{-1/6} \right] & = T_0, \\
 \lim_{h \rightarrow 0^+} \left[\int_{L_{h3}^{(1)}} dt + \frac{5\sqrt{2}}{3} B_{00}^* |\tilde{h}_3|^{-1/3} h^{-1/6} \right] & = T_0^*, \\
 \lim_{h \rightarrow 0^-} \int_{L_{h1}^{(1)}} (p_x + q_y - \sigma) dt & = \lim_{h \rightarrow 0^+} \int_{L_{h3}^{(1)}} (p_x + q_y - \sigma) dt \\
 & = \int_{L_1^*} (p_x + q_y - \sigma) dt. \quad (41)
 \end{aligned}$$

Here, T_0 and T_0^* are constants. By a similar argument used in Theorems 2.2 and 2.4 in [4], one can obtain

$$\begin{aligned}
 \lim_{h \rightarrow 0^-} \left[\int_{L_{h1}^{(1)}} dt + \frac{3\sqrt{2}}{2} \tilde{A}_0 |\tilde{h}_4|^{-1/4} |h|^{-1/4} + O(1) (\ln |h| + 1) \right] \\
 & = T_1, \\
 \lim_{h \rightarrow 0^-} \left[\int_{L_{h1}^{(2)}} (x - x_2) dt - \frac{\sqrt{2}}{4} |\tilde{h}_4|^{-1/2} (\ln |h| + 1) \right] & = T_1^*, \\
 \lim_{h \rightarrow 0^+} \left[\int_{L_{h3}^{(i)}} dt - \frac{3\sqrt{2}}{2} \tilde{A}_0 |\tilde{h}_4|^{-1/4} h^{-1/4} \right] & = T_i, \quad i = 2, 3,
 \end{aligned}$$

$$\begin{aligned}
 \lim_{h \rightarrow 0^-} \int_{L_{h1}^{(2)}} [p_x + q_y - \eta_0 - \eta_1 (x - x_2)] dt \\
 & = \int_{L_2^*} [p_x + q_y - \eta_0 - \eta_1 (x - x_2)] dt, \\
 \lim_{h \rightarrow 0^+} \int_{L_{h3}^{(i)}} (p_x + q_y - \eta_0) dt & = \int_{L_0^i} (p_x + q_y - \eta_0) dt, \\
 & i = 2, 3. \quad (42)
 \end{aligned}$$

Here $T_1^*, T_i, i = 1, 2, 3$, are constants. Therefore, we can obtain (31) and (32). Thus we have proved Theorem 3. $\square$

In the following we use Theorem 3 to study the problem of limit cycle bifurcation near L_0 . For the sake of convenience, we say that (1) has a distribution $(i, j) + k$ of $i + j + k$ limit cycles if there are i and j limit cycles near the inside of L^* and L_3 , respectively, and k limit cycles near the outside of L_0 . Then we can prove the following theorem.

Theorem 4. Assume that system (1) has a compound loop L_0 as stated before and (26)–(28) hold. Define $c_4^*(\delta) = c_4(\delta)|_{c_1=c_2=c_3=0}$, $\bar{c}_3^*(\delta) = \bar{c}_3(\delta)|_{c_1=c_3=0}$, $\tilde{c}_2^*(\delta) = c_4^*(\delta) + \bar{c}_3^*(\delta)$, $c_3^*(\delta) = c_3(\delta)|_{c_1=0}$. Let there exist $\delta_0 \in \mathbb{R}^m$, such that $(c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*)(\delta_0) = (0, 0, 0, 0, 0, 0, 0)$.

(1) If $c_l(\delta_0) \neq 0$, $c_j(\delta_0) = 0$, $j = 5, \dots, l-1$, and

$$\text{rank} \frac{\partial (c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*, c_5, \dots, c_{l-1})}{\partial (\delta_1, \delta_2, \delta_3, \delta_4, \delta_5, \delta_6, \delta_7, \delta_8, \dots, \delta_m)} = l + 2, \quad (43)$$

then (1) can have $2l + 1$ limit cycles near L_0 for some (ϵ, δ) near $(0, \delta_0)$, where $l = 6, 7$ or 9 .

(2) If $c_5(\delta_0)c_6(\delta_0) < 0$, and

$$\text{rank} \frac{\partial (c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*)}{\partial (\delta_1, \delta_2, \delta_3, \delta_4, \dots, \delta_m)} = 7, \quad (44)$$

then (1) can have 11 limit cycles near L_0 for some (ϵ, δ) near $(0, \delta_0)$.

(3) If $c_8(\delta_0)c_9(\delta_0) > 0$, $c_j(\delta_0) = 0$, $j = 5, 6, 7$, and

$$\text{rank} \frac{\partial (c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*, c_5, c_6, c_7)}{\partial (\delta_1, \delta_2, \delta_3, \delta_4, \delta_5, \delta_6, \delta_7, \delta_8, \dots, \delta_m)} = 10, \quad (45)$$

then (1) can have 18 limit cycles near L_0 for some (ϵ, δ) near $(0, \delta_0)$.

Proof. (1) Because of the similarity in the proof, we only prove the conclusion for $l = 9$ and omit the rest. By our assumptions, there exists $\delta_0 \in \mathbb{R}^m$ such that $\bar{c}_0(\delta_0) = \bar{c}_3^*(\delta_0) = c_3^*(\delta_0) = c_4^*(\delta_0) = 0$, $c_j(\delta_0) = 0$, $j = 0, 1, 2, 5, 6, 7, 8$, $c_9(\delta_0) \neq 0$, and

$$\text{rank} \frac{\partial (c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*, c_5, \dots, c_8)}{\partial (\delta_1, \delta_2, \delta_3, \delta_4, \delta_5, \delta_6, \delta_7, \delta_8, \dots, \delta_m)} = 11. \quad (46)$$

By the implicit function theorem, we can take $\bar{c}_0, \bar{c}_3^*, c_3^*, c_4^*, c_j, j = 0, 1, 2, 5, 6, 7, 8$ as free parameters varying near zero. Obviously, for these parameters varying near zero we have $|c_9| \geq |(1/2)c_9(\delta_0)| > 0$. In the following we proceed the process by 9 steps.

Step 1. Fix $(c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*, c_5, c_6, c_7) = (0, 0, 0, 0, 0, 0, 0, 0, 0, 0)$ and vary c_8 near 0.

First, for $c_8 = 0$, we have by (26) $c_9 M_1 > 0$ for $0 < -h \ll 1$.

Let $0 < |c_8| \ll 1$. Then $c_9 M_1 < 0$ for $0 < -h \ll 1$ if $c_8 c_9 > 0$. Thus, M_1 has a zero. Hence, for $0 < |c_8| \ll 1$,

(1) the condition $c_8 c_9 > 0$ implies a distribution $(1, 0) + 0$ of one limit cycle.

Step 2. Fix $(c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*, c_5, c_6) = (0, 0, 0, 0, 0, 0, 0, 0, 0)$, $c_8 c_9 > 0$ and vary c_7 near 0.

First, for $c_7 = 0$, we have by (26), (27), and (28) $c_8 M_1 < 0$, $c_9 M_2 < 0$ for $0 < -h \ll 1$, and $c_8 M_3 < 0$ for $0 < h \ll 1$.

Let $0 < |c_7| \ll |c_8|$. Then $c_8 M_1 > 0$, $c_9 M_2 > 0$ for $0 < -h \ll 1$, and $c_8 M_3 > 0$ for $0 < h \ll 1$ if $c_7 c_8 > 0, c_7 c_9 > 0$. Thus, M_1, M_2 , and M_3 each gets a zero and the zero of M_1 got in Step 1 still exists. Hence, for $0 < |c_7| \ll |c_8| \ll 1$,

(2) the conditions $c_7 c_8 > 0, c_8 c_9 > 0$ imply a distribution $(2, 1) + 1$ of 4 limit cycles.

Step 3. Fix $(c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*, c_5) = (0, 0, 0, 0, 0, 0, 0, 0)$, $c_7 \neq 0$ and vary c_6 near 0.

First, for $c_6 = 0$, we have by (26), (27), and (28) $c_7 M_1 > 0$, $c_7 M_2 > 0$ for $0 < -h \ll 1$, and $c_7 M_3 > 0$ for $0 < h \ll 1$.

Let $0 < |c_6| \ll |c_7|$. Then $c_7 M_1 < 0$, $c_7 M_2 < 0$ for $0 < -h \ll 1$, and $c_7 M_3 > 0$ for $0 < h \ll 1$ if $c_6 c_7 < 0$. Thus, M_1 and M_2 each gets a new zero and the zeros got in above steps still exist. Hence, for $0 < |c_6| \ll |c_7| \ll |c_8| \ll 1$,

(3) the conditions $c_6 c_7 < 0, c_7 c_8 > 0, c_8 c_9 > 0$ imply a distribution $(3, 2) + 1$ of 6 limit cycles.

Step 4. Fix $(c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*) = (0, 0, 0, 0, 0, 0, 0)$, $c_6 \neq 0$ and vary c_5 near 0.

First, for $c_5 = 0$, we have by (26) and (28) $c_6 M_1 > 0$ for $0 < -h \ll 1$, and $c_6 M_3 < 0$ for $0 < h \ll 1$.

Let $0 < |c_5| \ll |c_6|$. Then $c_6 M_1 < 0$ for $0 < -h \ll 1$, and $c_6 M_3 > 0$ for $0 < h \ll 1$ if $c_5 c_6 < 0$. Thus, M_1 and M_3 each has a new zero and the zeros got in above steps still exist. Hence, for $0 < |c_5| \ll |c_6| \ll |c_7| \ll |c_8| \ll 1$,

(4) the conditions $c_5 c_6 < 0, c_6 c_7 < 0, c_7 c_8 > 0, c_8 c_9 > 0$ imply a distribution $(4, 2) + 2$ of 8 limit cycles.

Step 5. Fix $(c_0, \bar{c}_0, c_1, c_2, c_3^*) = (0, 0, 0, 0, 0)$, $c_5 c_6 < 0$ and vary $(\bar{c}_3^*, c_4^*)$ near $(0, 0)$ with $\bar{c}_2^* (= \bar{c}_3^* + c_4^*) \neq 0$.

First, for $(\bar{c}_3^*, c_4^*) = (0, 0)$, we have by (26), (27), and (28) $c_5 M_1 > 0, c_6 M_2 > 0$ for $0 < -h \ll 1$, and $c_5 M_3 < 0$ for $0 < h \ll 1$.

Let $0 < |\bar{c}_3^*, c_4^*| \ll |c_5|$. Then

$$\begin{aligned} c_5 M_1 &< 0, & c_6 M_2 &< 0 & \text{for } 0 < -h \ll 1, \\ c_5 M_3 &> 0 & & & \text{for } 0 < h \ll 1 \end{aligned} \quad (47)$$

if $c_4^* c_5 > 0, \bar{c}_3^* c_6 > 0, \bar{c}_2^* c_5 > 0$, and

$$\begin{aligned} c_5 M_1 &< 0, & c_6 M_2 &< 0 & \text{for } 0 < -h \ll 1, \\ c_5 M_3 &< 0 & & & \text{for } 0 < h \ll 1 \end{aligned} \quad (48)$$

if $c_4^* c_5 > 0, \bar{c}_3^* c_6 > 0, \bar{c}_2^* c_5 < 0$. Thus, M_1, M_2 , and M_3 each has one more zero in the first case and M_1 and M_2 each has a new zero in the second case. And the zeros got in above steps still exist. Hence, for $0 < |\bar{c}_3^*, c_4^*| \ll |c_5| \ll |c_6| \ll |c_7| \ll |c_8| \ll 1$,

(5i) the conditions $\bar{c}_3^* c_4^* < 0, \bar{c}_3^* \bar{c}_2^* < 0, c_4^* c_5 > 0, c_5 c_6 < 0, c_6 c_7 < 0, c_7 c_8 > 0, c_8 c_9 > 0$ imply a distribution $(5, 3) + 3$ of 11 limit cycles, and

(5ii) the conditions $\bar{c}_3^* c_4^* < 0, \bar{c}_3^* \bar{c}_2^* > 0, c_4^* c_5 > 0, c_5 c_6 < 0, c_6 c_7 < 0, c_7 c_8 > 0, c_8 c_9 > 0$ imply a distribution $(5, 3) + 2$ of 10 limit cycles.

Step 6. Fix $(c_0, \bar{c}_0, c_1, c_2) = (0, 0, 0, 0)$, $c_4^* \bar{c}_3^* < 0$ with $\bar{c}_2^* \neq 0$ and vary c_3^* near 0.

First, for $c_3^* = 0$, we have by (26) and (27) $c_4^* M_1 < 0$ and $\bar{c}_3^* M_2 < 0$ for $0 < -h \ll 1$.

Let $0 < |c_3^*| \ll |c_4^*, \bar{c}_3^*|$. Then $c_4^* M_1 > 0$ and $\bar{c}_3^* M_2 > 0$ for $0 < -h \ll 1$, if $c_3^* c_4^* < 0, \bar{c}_3^* c_3^* > 0$. Thus, M_1 and M_2 each gets a new zero and the zeros got in above steps still exist. Hence, for $0 < |c_3^*| \ll |c_4^*, \bar{c}_3^*| \ll |c_5| \ll |c_6| \ll |c_7| \ll |c_8| \ll 1$,

(6i) the conditions $c_3^* c_4^* < 0, \bar{c}_3^* c_4^* < 0, \bar{c}_3^* \bar{c}_2^* < 0, c_4^* c_5 > 0, c_5 c_6 < 0, c_6 c_7 < 0, c_7 c_8 > 0, c_8 c_9 > 0$ imply a distribution $(6, 4) + 3$ of 13 limit cycles,

(6ii) the conditions $c_3^* c_4^* < 0, \bar{c}_3^* c_4^* < 0, \bar{c}_3^* \bar{c}_2^* > 0, c_4^* c_5 > 0, c_5 c_6 < 0, c_6 c_7 < 0, c_7 c_8 > 0, c_8 c_9 > 0$ imply a distribution $(6, 4) + 2$ of 12 limit cycles.

Step 7. Fix $(c_0, \bar{c}_0, c_1) = (0, 0, 0)$, $c_3^* \bar{c}_2^* \neq 0$ and vary c_2 near 0.

First, for $c_2 = 0$, we have by (26) and (28) $c_3^* M_1 < 0$ for $0 < -h \ll 1$ and $\bar{c}_2^* M_3 > 0$ for $0 < h \ll 1$.

Let $0 < |c_2| \ll |c_3^*| \ll |c_4^*, \bar{c}_3^*|$. Then

$$\begin{aligned} c_3^* M_1 &> 0 & \text{for } 0 < -h \ll 1, \\ \bar{c}_2^* M_3 &< 0 & \text{for } 0 < h \ll 1 \end{aligned} \quad (49)$$

if $c_2 c_3^* > 0, \bar{c}_2^* c_2 > 0$, and

$$\begin{aligned} c_3^* M_1 &> 0 & \text{for } 0 < -h \ll 1, \\ \bar{c}_2^* M_3 &> 0 & \text{for } 0 < h \ll 1 \end{aligned} \quad (50)$$

if $c_2 c_3^* > 0, \bar{c}_2^* c_2 < 0$. Thus, M_1 and M_3 each gets one more zero in the first case and only M_1 has a new zero in the second case. And the zeros got in above steps still exist. Hence, for $0 < |c_2| \ll |c_3^*| \ll |c_4^*, \bar{c}_3^*| \ll |c_5| \ll |c_6| \ll |c_7| \ll |c_8| \ll 1$,

(7i) the conditions $c_2 c_3^* > 0, c_3^* c_4^* < 0, \bar{c}_3^* c_4^* < 0, \bar{c}_3^* \bar{c}_2^* > 0, c_4^* c_5 > 0, c_5 c_6 < 0, c_6 c_7 < 0, c_7 c_8 > 0, c_8 c_9 > 0$ imply a distribution $(7, 4) + 3$ of 14 limit cycles,

(7ii) the conditions $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* < 0$, $c_4^*c_5 > 0$, $c_5c_6 < 0$, $c_6c_7 < 0$, $c_7c_8 > 0$, $c_8c_9 > 0$ imply a distribution (7, 4) + 3 of 14 limit cycles.

Step 8. Fix $(c_0, \bar{c}_0) = (0, 0)$, $c_2c_3^* > 0$ and vary c_1 near 0.

First, for $c_1 = 0$, we have by (26), (27), and (28) $c_2M_1 > 0$, $c_3^*M_2 > 0$ for $0 < -h \ll 1$, and $c_2M_3 < 0$ for $0 < h \ll 1$.

Let $0 < |c_1| \ll |c_2|$. Then $c_2M_1 < 0$, $c_3^*M_2 < 0$ for $0 < -h \ll 1$, and $c_2M_3 > 0$ for $0 < h \ll 1$ if $c_1c_2 < 0$. Thus, M_1 , M_2 , and M_3 each gets a new zero and the zeros got in above steps still exist. Hence, for $0 < |c_1| \ll |c_2| \ll |c_3^*| \ll |c_4^*, \bar{c}_3^*| \ll |c_5| \ll |c_6| \ll |c_7| \ll |c_8| \ll 1$,

(8) the conditions $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$, $c_5c_6 < 0$, $c_6c_7 < 0$, $c_7c_8 > 0$, $c_8c_9 > 0$ imply a distribution (8, 5) + 4 of 17 limit cycles.

Step 9. Fix $c_1 \neq 0$ and vary $(c_0, \bar{c}_0)$ near $(0, 0)$.

First, for $(c_0, \bar{c}_0) = (0, 0)$, we have by (26), (27), and (28) $c_1M_1 > 0$, $c_1M_2 > 0$, for $0 < -h \ll 1$, and $c_1M_3 < 0$ for $0 < h \ll 1$.

Let $0 < |c_0, \bar{c}_0| \ll |c_1|$. Then

$$\begin{aligned} c_1M_1 < 0, \quad c_1M_2 < 0, \quad \text{for } 0 < -h \ll 1, \\ c_1M_3 < 0 \quad \text{for } 0 < h \ll 1 \end{aligned} \quad (51)$$

if $c_0c_1 < 0$, $\bar{c}_0c_1 < 0$,

$$\begin{aligned} c_1M_1 < 0, \quad c_1M_2 > 0, \quad \text{for } 0 < -h \ll 1, \\ c_1M_3 > 0 \quad \text{for } 0 < h \ll 1 \end{aligned} \quad (52)$$

if $c_0c_1 < 0$, $\bar{c}_0c_1 > 0$ and $(c_0 + \bar{c}_0)c_1 > 0$, and

$$\begin{aligned} c_1M_1 > 0, \quad c_1M_2 < 0, \quad \text{for } 0 < -h \ll 1, \\ c_1M_3 > 0 \quad \text{for } 0 < h \ll 1 \end{aligned} \quad (53)$$

if $c_0c_1 > 0$, $\bar{c}_0c_1 < 0$ and $(c_0 + \bar{c}_0)c_1 > 0$. Thus, we have correspondingly (a) M_1 and M_2 each has a new zero, (b) M_1 and M_3 each has a new zero, or (c) M_2 and M_3 each has a new zero. And the zeros got in above steps still exist.

Hence, for $0 < |c_0, \bar{c}_0| \ll |c_1| \ll |c_2| \ll |c_3^*| \ll |c_4^*, \bar{c}_3^*| \ll |c_5| \ll |c_6| \ll |c_7| \ll |c_8| \ll 1$,

(9i) the conditions $c_0\bar{c}_0 > 0$, $c_0c_1 < 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$, $c_5c_6 < 0$, $c_6c_7 < 0$, $c_7c_8 > 0$, $c_8c_9 > 0$ imply a distribution (9, 6) + 4 of 19 limit cycles,

(9ii) the conditions $c_0\bar{c}_0 < 0$, $c_0c_1 < 0$, $(c_0 + \bar{c}_0)c_1 > 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$, $c_5c_6 < 0$, $c_6c_7 < 0$, $c_7c_8 > 0$, $c_8c_9 > 0$ imply a distribution (9, 5) + 5 of 19 limit cycles,

(9iii) the conditions $c_0\bar{c}_0 < 0$, $c_0c_1 > 0$, $(c_0 + \bar{c}_0)c_1 > 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$, $c_5c_6 < 0$, $c_6c_7 < 0$, $c_7c_8 > 0$, $c_8c_9 > 0$ imply a distribution (8, 6) + 5 of 19 limit cycles.

Thus we get the conclusion for $l = 9$.

(2) By our assumptions in case (2) and the implicit function theorem we can take c_0 , $\bar{c}_0$, c_1 , c_2 , c_3^* , c_4^* , $\bar{c}_3^*$ as free parameters varying near zero. Obviously, for these parameters varying near zero we have $c_5c_6 < 0$. By a similar argument in the above proof, we can prove that for $0 < |c_0, \bar{c}_0| \ll |c_1| \ll |c_2| \ll |c_3^*| \ll |c_4^*, \bar{c}_3^*| \ll 1$,

(i) the conditions $c_0\bar{c}_0 > 0$, $c_0c_1 < 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$ imply a distribution (5, 4) + 2 of 11 limit cycles,

(ii) the conditions $c_0\bar{c}_0 < 0$, $c_0c_1 < 0$, $(c_0 + \bar{c}_0)c_1 > 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$ imply a distribution (5, 3) + 3 of 11 limit cycles,

(iii) the conditions $c_0\bar{c}_0 < 0$, $c_0c_1 > 0$, $(c_0 + \bar{c}_0)c_1 > 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$ imply a distribution (4, 4) + 3 of 11 limit cycles.

(3) By our assumptions in case (3) and the implicit function theorem we can take c_0 , $\bar{c}_0$, c_1 , c_2 , c_3^* , c_4^* , $\bar{c}_3^*$, c_5 , c_6 , and c_7 as free parameters varying near zero. Obviously, for these parameters varying near zero we have $c_8c_9 > 0$. By a similar argument used in proving case (1), we can prove that for $0 < |c_0, \bar{c}_0| \ll |c_1| \ll |c_2| \ll |c_3^*| \ll |c_4^*, \bar{c}_3^*| \ll |c_5| \ll |c_6| \ll |c_7| \ll 1$,

(i) the conditions $c_0\bar{c}_0 > 0$, $c_0c_1 < 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$, $c_5c_6 < 0$, $c_6c_7 < 0$, $c_7c_8 > 0$ imply a distribution (8, 6) + 4 of 18 limit cycles,

(ii) the conditions $c_0\bar{c}_0 < 0$, $c_0c_1 < 0$, $(c_0 + \bar{c}_0)c_1 > 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$, $c_5c_6 < 0$, $c_6c_7 < 0$, $c_7c_8 > 0$ imply a distribution (8, 5) + 5 of 18 limit cycles,

(iii) the conditions $c_0\bar{c}_0 < 0$, $c_0c_1 > 0$, $(c_0 + \bar{c}_0)c_1 > 0$, $c_1c_2 < 0$, $c_2c_3^* > 0$, $c_3^*c_4^* < 0$, $\bar{c}_3^*c_4^* < 0$, $\bar{c}_3^*\bar{c}_2^* > 0$ (or $\bar{c}_3^*\bar{c}_2^* < 0$), $c_4^*c_5 > 0$, $c_5c_6 < 0$, $c_6c_7 < 0$, $c_7c_8 > 0$ imply a distribution (7, 6) + 5 of 18 limit cycles.

This completes the proof. $\square$

3. An Application

Consider a Liénard system of the form

$$\dot{x} = y, \quad \dot{y} = -(x+1)^2x^3 \left(x^2 - \frac{1}{4}x - \frac{1}{2} \right) - \epsilon f(x, \delta)y, \quad (54)$$

where

$$f(x, \delta) = \sum_{j=0}^n a_j x^j, \quad \delta = (a_0, a_1, \dots, a_n), \quad (55)$$

$$8 \leq n \leq 12.$$

System (54) $|_{\epsilon=0}$ is Hamiltonian with

$$H(x, y) = \frac{1}{2}y^2 + \frac{1}{8}(x+1)^3x^4(x-1). \quad (56)$$

We have the following theorem.

Theorem 5. Let $C(n)$ denote the maximal number of limit cycles of the system (54) for ϵ small and all δ . Then we have $C(11) \geq 18$, $C(n) \geq 2n - 5$ ($n = 8, 9, 10, 12$).

Proof. It is easy to verify that the unperturbed system has a compound loop $L_0 = L_1 \cup L_2 \cup L_3 \cup \{S_1, S_2\}$ with a cusp $S_1(-1, 0)$ of order one and a nilpotent saddle $S_2(0, 0)$ of order one, L_1, L_2 are heteroclinic orbits satisfying $\omega(L_1) = \alpha(L_2) = S_2$ and $\omega(L_2) = \alpha(L_1) = S_1$, and L_3 is a homoclinic loop to S_2 . Inside $L^* = L_1 + L_2$ (L_3 , resp.), there is a center $C_1((1/8)(1 - \sqrt{33}), 0)$ ($C_2((1/8)(1 + \sqrt{33}), 0)$, resp.).

Because of the similarity in the proof, here we only prove the case for $n = 11$ and omit the rest of the proof.

Let $n = 11$. By Theorem 3, we obtain

$$c_0(\delta) = M_1(0, \delta) = -\oint_{L^*} f(x, \delta) y dx = -\sum_{j=0}^{11} a_j I_1^j, \quad (57)$$

$$\bar{c}_0(\delta) = M_2(0, \delta) = -\oint_{L_3} f(x, \delta) y dx = -\sum_{j=0}^{11} a_j I_2^j,$$

where

$$\begin{aligned} I_1^j &= \oint_{L^*} x^j y dx = 2 \int_{-1}^0 x^j y dx \\ &= \int_{-1}^0 x^{j+2} (x+1) \sqrt{1-x^2} dx, \quad j = 0, 1, \dots, 11, \end{aligned} \quad (58)$$

$$\begin{aligned} I_2^j &= \oint_{L_3} x^j y dx = 2 \int_0^1 x^j y dx \\ &= \int_0^1 x^{j+2} (x+1) \sqrt{1-x^2} dx, \quad j = 0, 1, \dots, 11. \end{aligned}$$

Therefore,

$$\begin{aligned} c_0(\delta) &= -a_0 \left(\frac{\pi}{16} - \frac{2}{15} \right) - a_1 \left(\frac{\pi}{32} - \frac{2}{15} \right) - a_2 \left(\frac{\pi}{32} - \frac{8}{105} \right) \\ &\quad - a_3 \left(\frac{5\pi}{256} - \frac{8}{105} \right) - a_4 \left(\frac{5\pi}{256} - \frac{16}{315} \right) \\ &\quad - a_5 \left(\frac{7\pi}{512} - \frac{16}{315} \right) - a_6 \left(\frac{7\pi}{512} - \frac{128}{3465} \right) \\ &\quad - a_7 \left(\frac{21\pi}{2048} - \frac{128}{3465} \right) - a_8 \left(\frac{21\pi}{2048} - \frac{256}{9009} \right) \\ &\quad - a_9 \left(\frac{33\pi}{4096} - \frac{256}{9009} \right) - a_{10} \left(\frac{33\pi}{4096} - \frac{1024}{45045} \right) \\ &\quad - a_{11} \left(\frac{429\pi}{65536} - \frac{1024}{45045} \right), \end{aligned}$$

$$\begin{aligned} \bar{c}_0(\delta) &= -a_0 \left(\frac{\pi}{16} + \frac{2}{15} \right) - a_1 \left(\frac{\pi}{32} + \frac{2}{15} \right) - a_2 \left(\frac{\pi}{32} + \frac{8}{105} \right) \\ &\quad - a_3 \left(\frac{5\pi}{256} + \frac{8}{105} \right) - a_4 \left(\frac{5\pi}{256} + \frac{16}{315} \right) \\ &\quad - a_5 \left(\frac{7\pi}{512} + \frac{16}{315} \right) - a_6 \left(\frac{7\pi}{512} + \frac{128}{3465} \right) \\ &\quad - a_7 \left(\frac{21\pi}{2048} + \frac{128}{3465} \right) - a_8 \left(\frac{21\pi}{2048} + \frac{256}{9009} \right) \\ &\quad - a_9 \left(\frac{33\pi}{4096} + \frac{256}{9009} \right) - a_{10} \left(\frac{33\pi}{4096} + \frac{1024}{45045} \right) \\ &\quad - a_{11} \left(\frac{429\pi}{65536} + \frac{1024}{45045} \right). \end{aligned} \quad (59)$$

Note that S_2 is a nilpotent saddle of order one and $\bar{h}_4 = -1/8$. By (23), we have

$$\begin{aligned} c_1(\delta) &= c_1(S_2, \delta) = 2^{9/4} |\tilde{A}_0| a_0, \\ c_3(\delta) &= c_2(S_2, \delta) = a_1 + O_1(a_0), \\ c_6(\delta) &= c_4(S_2, \delta) = 2^{3/4} |\tilde{A}_2| (-21a_0 + 12a_1 - 8a_2), \\ c_7(\delta) &= c_5(S_2, \delta) \\ &= 2^{1/4} \frac{|\tilde{A}_0|}{7} \left(\frac{2035}{4} a_0 - 310a_1 + 180a_2 - 80a_3 + 32a_4 \right), \\ c_9(\delta) &= c_6(S_2, \delta) \\ &= \frac{333}{4} a_0 - \frac{207}{4} a_1 + 29a_2 - 15a_3 + 6a_4 - 2a_5. \end{aligned} \quad (60)$$

Making the transformation $x = u - 1$, $y = v$, system (54) becomes

$$\dot{u} = v, \quad \dot{v} = -u^2(u-1)^3 \left(u^2 - \frac{9}{4}u + \frac{3}{4} \right) - \epsilon \tilde{f}(u, \delta) v. \quad (61)$$

Then we have

$$\begin{aligned} H(u, v) &= \frac{v^2}{2} + \frac{1}{8} (u^8 - 6u^7 + 14u^6 - 16u^5 + 9u^4 - 2u^3), \\ \tilde{h}_3 &= -\frac{1}{4}, \\ \tilde{f}(u, \delta) &= \sum_{j=0}^{11} a_j (u-1)^j = \sum_{j=0}^{11} (-1)^j a_j \\ &\quad + \sum_{j=1}^{11} (-1)^{(j-1)} j a_j u + \sum_{j=2}^{11} (-1)^{(j-2)} C_j^2 a_j u^2 \\ &\quad + \sum_{j=3}^{11} (-1)^{(j-3)} C_j^3 a_j u^3 + \sum_{j=4}^{11} (-1)^{(j-4)} C_j^4 a_j u^4 \end{aligned}$$

$$\begin{aligned}
 & + \sum_{j=5}^{11} (-1)^{(j-5)} C_j^5 a_j u^5 + \sum_{j=6}^{11} (-1)^{(j-6)} C_j^6 a_j u^6 \\
 & + \sum_{j=7}^{11} (-1)^{(j-7)} C_j^7 a_j u^7 + \sum_{j=8}^{11} (-1)^{(j-8)} C_j^8 a_j u^8 \\
 & + \sum_{j=9}^{11} (-1)^{(j-9)} C_j^9 a_j u^9 + (a_{10} - 11a_{11}) u^{10} + a_{11} u^{11}.
 \end{aligned} \tag{62}$$

By (19), we have

$$\begin{aligned}
 c_2(\delta) &= c_1(S_1, \delta) = 2^{13/6} \sum_{j=0}^{11} (-1)^j a_j, \\
 c_5(\delta) &= c_3(S_1, \delta) = 2^{17/6} \left[\sum_{j=1}^{11} (-1)^j (j-3) a_j - 3a_0 \right], \\
 c_8(\delta) &= c_4(S_1, \delta) \\
 &= 2^{13/6} \left(-\frac{1316}{3} a_0 + 272a_1 - \frac{460}{3} a_2 + \frac{224}{3} a_3 \right. \\
 &\quad \left. - 28a_4 + \frac{16}{3} a_5 + \frac{4}{3} a_6 - \frac{4}{3} a_8 \right. \\
 &\quad \left. - \frac{16}{3} a_9 + 28a_{10} - \frac{224}{3} a_{11} \right).
 \end{aligned} \tag{63}$$

Note that

$$\begin{aligned}
 c_1(\delta) &= c_1(S_2, \delta) = 2^{9/4} |\widetilde{A}_0| a_0, \\
 c_2(\delta) &= c_1(S_1, \delta) = 2^{13/6} \sum_{j=0}^{11} (-1)^j a_j, \\
 c_3(\delta) &= c_2(S_2, \delta) = a_1 + O_1(a_0).
 \end{aligned} \tag{64}$$

We have $c_1(\delta) = c_2(\delta) = c_3(\delta) = 0$ if and only if $a_0 = a_1 = 0$, and $a_{11} = \sum_{i=2}^{10} (-1)^i a_i$. It implies further that

$$\begin{aligned}
 c_4(\delta) &= \oint_{L^*} (p_x + q_y) dt = -\oint_{L^*} f(x, \delta) dt \\
 &= -\oint_{L^*} \frac{f(x, \delta)}{y} dx = -2 \int_{-1}^0 \frac{f(x, \delta)}{y} dx \\
 &= -2 \int_{-1}^0 \frac{1}{y} \left(\sum_{i=0}^{11} a_i x^i \right) dx \\
 &= -4 \int_{-1}^0 \frac{1}{x^2(x+1)\sqrt{1-x^2}} \\
 &\quad \times \left(\sum_{i=2}^{10} a_i x^i + \sum_{i=2}^{10} (-1)^i a_i x^{11} \right) dx \\
 &= -4 \sum_{i=2}^{10} a_i \int_{-1}^0 f_i(x) dx,
 \end{aligned} \tag{65}$$

where

$$f_i(x) = \frac{x^{i-2} [1 + (-1)^i x^{(11-i)}]}{(x+1)\sqrt{1-x^2}}, \quad i = 2, 3, \dots, 10. \tag{66}$$

Similarly,

$$\begin{aligned}
 \bar{c}_3(\delta) &= \oint_{L_3} (p_x + q_y) dt = -\oint_{L_3} f(x, \delta) dt \\
 &= -\oint_{L_3} \frac{f(x, \delta)}{y} dx = -2 \int_0^1 \frac{f(x, \delta)}{y} dx \\
 &= -4 \sum_{i=2}^{10} a_i \int_0^1 f_i(x) dx.
 \end{aligned} \tag{67}$$

Therefore,

$$\begin{aligned}
 c_4(\delta) &= -4a_2 \left(\frac{315\pi}{256} + \frac{93}{35} \right) + 4a_3 \left(\frac{187\pi}{256} + \frac{93}{35} \right) \\
 &\quad - 4a_4 \left(\frac{187\pi}{256} + \frac{58}{35} \right) + 4a_5 \left(\frac{123\pi}{256} + \frac{58}{35} \right) \\
 &\quad - 4a_6 \left(\frac{123\pi}{256} + \frac{104}{105} \right) + 4a_7 \left(\frac{75\pi}{256} + \frac{104}{105} \right) \\
 &\quad - 4a_8 \left(\frac{75\pi}{256} + \frac{16}{35} \right) + 4a_9 \left(\frac{35\pi}{256} + \frac{16}{35} \right) \\
 &\quad - 4a_{10} \left(\frac{35\pi}{256} \right), \\
 \bar{c}_3(\delta) &= -4a_2 \left(\frac{315\pi}{256} - \frac{93}{35} \right) + 4a_3 \left(\frac{187\pi}{256} - \frac{93}{35} \right) \\
 &\quad - 4a_4 \left(\frac{187\pi}{256} - \frac{58}{35} \right) + 4a_5 \left(\frac{123\pi}{256} - \frac{58}{35} \right) \\
 &\quad - 4a_6 \left(\frac{123\pi}{256} - \frac{104}{105} \right) + 4a_7 \left(\frac{75\pi}{256} - \frac{104}{105} \right) \\
 &\quad - 4a_8 \left(\frac{75\pi}{256} - \frac{16}{35} \right) + 4a_9 \left(\frac{35\pi}{256} - \frac{16}{35} \right) \\
 &\quad - 4a_{10} \left(\frac{35\pi}{256} \right).
 \end{aligned} \tag{68}$$

Let $c_4^*(\delta) = c_4(\delta)|_{c_1=c_2=c_3=0}$, $\bar{c}_3^*(\delta) = \bar{c}_3(\delta)|_{c_1=c_3=0}$, $c_3^*(\delta) = c_3(\delta)|_{c_1(\delta)=0}$. Furthermore, one sees that equations $c_0(\delta) = \bar{c}_0(\delta) = c_1(\delta) = c_2(\delta) = c_3^*(\delta) = c_4^*(\delta) = \bar{c}_3^*(\delta) = c_5(\delta) = c_6(\delta) = c_7(\delta) = 0$ have the solution $a_0 = a_1 = a_2 = 0$, $a_3 = (13/74)a_9 - (1/2)a_7$, $a_4 = (65/148)a_9 - (5/4)a_7$, $a_5 = -(16/37)a_9$, $a_6 = (7/4)a_7 - (255/148)a_9$, $a_8 = (96/37)a_9$, $a_{10} = -(53/37)a_9$, $a_{11} = -(32/37)a_9$, which gives $c_8(\delta) = (55/37)2^{31/6}a_9$, $c_9(\delta) = (32/37)a_9$. And further, $c_8(\delta)c_9(\delta) > 0$ if $a_9 \neq 0$. Thus, fix $a_9 \neq 0$ and take $\delta_0 = (0, 0, 0, (13/74)a_9 - (1/2)a_7, (65/148)a_9 - (5/4)a_7, -(16/37)a_9, (7/4)a_7 - (255/148)a_9, a_7, (96/37)a_9, a_9, -(53/37)a_9, -(32/37)a_9)$.

Then we have

$$\text{rank} \frac{\partial (c_0, \bar{c}_0, c_1, c_2, c_3^*, c_4^*, \bar{c}_3^*, c_5, c_6, c_7)}{\partial (a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, a_{11})} = 10. \tag{69}$$

Hence by Theorem 4(3), we know that there are 18 limit cycles near L_0 for some δ near δ_0 . This ends the proof. $\square$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The project was supported by National Natural Science Foundation of China (11271261) and FP7-PEOPLE-2012-IRSES-316338.

References

- [1] M. Han, "Asymptotic expansions of Melnikov functions and limit cycle bifurcations," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 22, no. 12, Article ID 1250296, 30 pages, 2012.
- [2] A. Atabaigi, H. R. Z. Zangeneh, and R. Kazemi, "Limit cycle bifurcation by perturbing a cuspidal loop of order 2 in a Hamiltonian system," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 75, no. 4, pp. 1945–1958, 2012.
- [3] M. Han, H. Zang, and J. Yang, "Limit cycle bifurcations by perturbing a cuspidal loop in a Hamiltonian system," *Journal of Differential Equations*, vol. 246, no. 1, pp. 129–163, 2009.
- [4] M. Han, J. Yang, and D. Xiao, "Limit cycle bifurcations near a double homoclinic loop with a nilpotent saddle," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 22, no. 8, Article ID 1250189, 33 pages, 2012.
- [5] M. Han, *Bifurcation Theory of Limit Cycles*, Science Press, 2013.
- [6] Z. Hu, B. Gao, and V. G. Romanovski, "Limit cycle bifurcations from centers of symmetric Hamiltonian systems perturbed by cubic polynomials," *International Journal of Bifurcation and Chaos*, vol. 23, no. 3, Article ID 1350043, 20 pages, 2013.
- [7] J. Jiang, "Bifurcation of limit cycles for a quartic near-Hamiltonian system by perturbing a nilpotent center," *Journal of Mathematical Analysis and Applications*, vol. 365, no. 1, pp. 376–384, 2010.
- [8] J. Li, T. Zhang, and M. Han, "Bifurcation of limit cycles from a heteroclinic loop with two cusps," *Chaos, Solitons & Fractals*, vol. 62–63, pp. 44–54, 2014.
- [9] X. Sun, M. Han, and J. Yang, "Bifurcation of limit cycles from a heteroclinic loop with a cusp," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 74, no. 9, pp. 2948–2965, 2011.
- [10] J. Yang, "On the limit cycles of a kind of Liénard system with a nilpotent center under perturbations," *The Journal of Applied Analysis and Computation*, vol. 2, no. 3, pp. 325–339, 2012.
- [11] H. Zang, M. Han, and D. Xiao, "On Melnikov functions of a homoclinic loop through a nilpotent saddle for planar near-Hamiltonian systems," *Journal of Differential Equations*, vol. 245, no. 4, pp. 1086–1111, 2008.
- [12] H. Zang, Z. Wang, and T. Zhang, "Bifurcations and distribution of limit cycles for near-Hamiltonian polynomial systems," *Journal of Mathematical Analysis and Applications*, vol. 348, no. 1, pp. 87–100, 2008.
- [13] A. Atabaigi and H. R. Z. Zangeneh, "Bifurcation of limit cycles in small perturbations of a class of hyper-elliptic Hamiltonian systems of degree 5 with a cusp," *The Journal of Applied Analysis and Computation*, vol. 1, no. 3, pp. 299–313, 2011.
- [14] R. Kazemi, H. R. Z. Zangeneh, and A. Atabaigi, "On the number of limit cycles in small perturbations of a class of hyper-elliptic Hamiltonian systems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 75, pp. 574–587, 2012.
- [15] J. Wang and D. Xiao, "On the number of limit cycles in small perturbations of a class of hyper-elliptic Hamiltonian systems with one nilpotent saddle," *Journal of Differential Equations*, vol. 250, no. 4, pp. 2227–2243, 2011.
- [16] J. Wang, "Estimate of the number of zeros of Abelian integrals for a perturbation of hyperelliptic Hamiltonian system with nilpotent center," *Chaos, Solitons & Fractals*, vol. 45, no. 9–10, pp. 1140–1146, 2012.
- [17] Y. Xiong, "Bifurcation of limit cycles by perturbing a class of hyper-elliptic Hamiltonian systems of degree five," *Journal of Mathematical Analysis and Applications*, vol. 411, no. 2, pp. 559–573, 2014.