Research Article

Landau-Type Theorems for Certain Biharmonic Mappings

Ming-Sheng Liu,¹ Zhen-Xing Liu,¹ and Jun-Feng Xu²

¹ School of Mathematical Sciences, South China Normal University, Guangzhou, Guangdong 510631, China

² Department of Mathematics, Wuyi University, Jiangmen, Guangdong 529020, China

Correspondence should be addressed to Ming-Sheng Liu; liumsh@scnu.edu.cn

Received 2 January 2014; Accepted 2 March 2014; Published 27 March 2014

Academic Editor: Om P. Ahuja

Copyright © 2014 Ming-Sheng Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let $F(z) = |z|^2 g(z) + h(z)$ ($|z| < 1$) be a biharmonic mapping of the unit disk $\mathbb{D}$, where g and h are harmonic in $\mathbb{D}$. In this paper, the Landau-type theorems for biharmonic mappings of the form $L(F)$ are provided. Here L represents the linear complex operator $L = (z\partial/\partial z) - (\bar{z}\partial/\partial \bar{z})$ defined on the class of complex-valued C^1 functions in the plane. The results, presented in this paper, improve the related results of earlier authors.

1. Introduction

Suppose that $f(z) = u(x, y) + iv(x, y)$, $z = x + iy$ is a four times continuously differentiable complex-valued function in a domain $D \subset \mathbb{C}$. If f satisfies the biharmonic equation $\Delta(\Delta f) = 0$, then we call that f is biharmonic, where Δ represents the Laplacian operator:

$$\Delta = 4 \frac{\partial^2}{\partial z \partial \bar{z}} = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}. \quad (1)$$

Biharmonic functions arise in many physical situations, particularly in fluid dynamics and elasticity problems, and have many important applications in engineering (see [1] for details). It is known that a mapping F is biharmonic in a simply connected domain D if and only if F has the following representation:

$$F(z) = |z|^2 g(z) + h(z), \quad (2)$$

where $g(z)$ and $h(z)$ are complex-valued harmonic functions in D [1]. Also, it is known that $g(z)$ and $h(z)$ can be expressed as

$$\begin{aligned} g(z) &= g_1(z) + \overline{g_2(z)}, \quad z \in D, \\ h(z) &= h_1(z) + \overline{h_2(z)}, \quad z \in D, \end{aligned} \quad (3)$$

where g_1, g_2, k_1 , and k_2 are analytic in D [2, 3].

For a continuously differentiable mapping f in D , we define

$$\begin{aligned} \Lambda_f(z) &= \max_{0 \leq \theta \leq 2\pi} |f_z + e^{-2i\theta} f_{\bar{z}}| = |f_z| + |f_{\bar{z}}|, \\ \lambda_f(z) &= \min_{0 \leq \theta \leq 2\pi} |f_z + e^{-2i\theta} f_{\bar{z}}| = ||f_z| - |f_{\bar{z}}||. \end{aligned} \quad (4)$$

We use J_f to denote the Jacobian of f

$$J_f(z) = |f_z(z)|^2 - |f_{\bar{z}}(z)|^2. \quad (5)$$

Then $J_f = \lambda_f \Lambda_f$ if $J_f \geq 0$.

In [4], the authors considered the following differential operator L defined on the class of complex-valued C^1 functions:

$$L = z \frac{\partial}{\partial z} - \bar{z} \frac{\partial}{\partial \bar{z}}. \quad (6)$$

Evidently, L is a complex linear operator and satisfies the usual product rule:

$$\begin{aligned} L(af + bg) &= aL(f) + bL(g), \\ L(fg) &= fL(g) + gL(f), \end{aligned} \quad (7)$$

where a and b are complex constants; f and g are C^1 functions. In addition, the operator L possesses a number of

interesting properties. For instance, it is easy to see that the operator L preserves both harmonicity and biharmonicity. Many other basic properties are stated in [4].

Landau's theorem states that if f is an analytic function on the unit disk $\mathbb{D}$ with $f(0) = f'(0) - 1 = 0$ and $|f(z)| < M$ for $z \in \mathbb{D}$, then f is univalent in the disk $\mathbb{D}_{r_0} = \{z \in \mathbb{C} : |z| < r_0\}$ with $r_0 = 1/(M + \sqrt{M^2 - 1})$, and $f(\mathbb{D}_{r_0})$ contains a disk $\mathbb{D}_{R_0}$ with $R_0 = Mr_0^2$. This result is sharp, with the extremal function $f(z) = Mz((1 - Mz)/(M - z))$. Recently, many authors considered the Landau-type theorems for harmonic mappings [5–9] and biharmonic mappings [1, 4, 10–13]. Chen et al. [10] obtained the Landau-type theorems for biharmonic mappings of the form $L(F)$ as follows.

Theorem A (see [10]). Let $F(z) = |z|^2 g(z) + h(z)$ be a biharmonic mapping of the unit disk $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ such that $F(0) = h(0) = 0$ and $J_h(0) = 1$, where $g(z)$ and $h(z)$ are harmonic in $\mathbb{D}$. Assume that both $|g(z)|$ and $|h(z)|$ are bounded by M . Then there is a constant ρ_1 ($0 < \rho_1 < 1$) such that $L(F)$ is univalent in $\mathbb{D}_{\rho_1}$, where ρ_1 satisfies the following equation:

$$\frac{\pi}{4M} - \frac{6M\rho_1^2}{(1 - \rho_1)^2} - \frac{4M\rho_1^3}{(1 - \rho_1)^3} - \frac{16M}{\pi^2} m_1 \arctan \rho_1 - \frac{4M\rho_1}{(1 - \rho_1)^3} = 0, \quad (8)$$

where $m_1 \approx 6.059$ is the minimum value of the function

$$\frac{2 - x^2 + (4/\pi) \arctan x}{x(1 - x^2)}, \quad (9)$$

for $0 < x < 1$. The minimum is attained at $x \approx 0.588$. Moreover, the range $L(F)(\mathbb{D}_{\rho_1})$ contains a schlicht disk $\mathbb{D}_{R_1}$, where

$$R_1 = \rho_1 \left[\frac{\pi}{4M} - \frac{2M\rho_1^2}{(1 - \rho_1)^2} - \frac{16M}{\pi^2} m_1 \arctan \rho_1 \right]. \quad (10)$$

Theorem B (see [10]). Let $F(z) = |z|^2 g(z)$ be a biharmonic mapping in $\mathbb{D}$ such that $g(0) = 0$, $J_g(0) = 1$, and $|g(z)| < M$, where $g(z)$ is harmonic in $\mathbb{D}$. Then there is a constant ρ_2 ($0 < \rho_2 < 1$) such that $L(F)$ is univalent in $\mathbb{D}_{\rho_2}$, where ρ_2 satisfies the following equation:

$$\frac{\pi}{4M} - \frac{48M}{\pi^2} m_1 \arctan \rho_2 - \frac{2M\rho_2}{(1 - \rho_2)^3} = 0, \quad (11)$$

where m_1 is defined as in Theorem A. Moreover, $L(F)(\mathbb{D}_{\rho_2})$ contains a disk $\mathbb{D}_{R_2}$ with

$$R_2 = \rho_2^3 \left[\frac{\pi}{4M} - \frac{16M}{\pi^2} m_1 \arctan \rho_2 \right]. \quad (12)$$

However, these results are not sharp. The main object of this paper is to improve Theorems A and B. We get three versions of Landau-type theorems for biharmonic mappings of the form $L(F)$, where F belongs to the class of biharmonic

mappings, and Theorems 11 and 14 improve Theorems A and B. In order to establish our main results, we need to recall the following lemmas.

Lemma 1 (see [6, 14]). Suppose that $f(z)$ is a harmonic mapping of the unit disk $\mathbb{D}$ such that $|f(z)| \leq M$ for all $\mathbb{D}$. Then

$$\Lambda_f(z) \leq \frac{4M}{\pi(1 - |z|^2)}, \quad z \in \mathbb{D}. \quad (13)$$

The inequality is sharp.

Lemma 2 (see [9, 12, 15]). Suppose that $f(z) = h(z) + \overline{g(z)}$ is a harmonic mapping of the unit disk $\mathbb{D}$ such that $|f(z)| \leq M$ for all $z \in \mathbb{D}$ with $h(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=1}^{\infty} b_n z^n$. Then $|a_0| \leq M$ and for any $n \geq 1$

$$|a_n| + |b_n| \leq \frac{4M}{\pi}. \quad (14)$$

These estimates are sharp.

Lemma 3 (see [8, 11]). Suppose that f is a harmonic mapping of $\mathbb{D}$ with $f(0) = \lambda_f(0) - 1 = 0$. If $\Lambda_f \leq \Lambda$ for $z \in \mathbb{D}$; then

$$|a_n| + |b_n| \leq \frac{\Lambda^2 - 1}{n\Lambda}, \quad n = 2, 3, \dots \quad (15)$$

These estimates are sharp.

Lemma 4 (see [11]). Suppose that $f(z) = h(z) + \overline{g(z)}$ is a harmonic mapping of the unit disk $\mathbb{D}$ such that $|f(z)| \leq M$ for all $z \in \mathbb{D}$ with $h(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=1}^{\infty} b_n z^n$. If $|J_f(0)| = 1$; then $\lambda_f(0) \geq \lambda_0(M)$, where $M_0 = \pi/2 \sqrt[4]{2\pi^2 - 16} \approx 1.1296$ and

$$\lambda_0(M) = \begin{cases} \frac{\sqrt{2}}{\frac{\sqrt{M^2 - 1}}{\pi} + \sqrt{M^2 + 1}}, & 1 \leq M \leq M_0, \\ \frac{1}{4M}, & M > M_0. \end{cases} \quad (16)$$

Lemma 5 (see [13]). Suppose that $f(z) = h(z) + \overline{g(z)}$ is a harmonic mapping of the unit disk $\mathbb{D}$ with $h(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=1}^{\infty} b_n z^n$. If f satisfies $|f(z)| \leq M$ for all $z \in \mathbb{D}$ and $|J_f(0)| = 1$, then

$$\left(\sum_{n=2}^{\infty} (|a_n| + |b_n|)^2 \right)^{1/2} \leq \sqrt{M^4 - 1} \cdot \lambda_f(0). \quad (17)$$

Lemma 6. Suppose that $M > 0$, $\Lambda \geq 1$. Then the equation

$$\varphi(r) = 1 - \frac{12Mr^2}{\pi(1 - r^2)} - \frac{8Mr^3}{\pi(1 - r^3)} - \frac{\Lambda^2 - 1}{\Lambda} \cdot \frac{2r - r^2}{(1 - r)^2} = 0 \quad (18)$$

has a unique root in $(0, 1)$.

Proof. It is easy to prove that the function φ is continuous and strictly decreasing on $[0, 1)$, $\varphi(0) = 1 > 0$, and $\lim_{r \rightarrow 1^-} \varphi(r) = -\infty$. Hence, the assertion follows from the mean value theorem. This completes the proof. $\square$

Lemma 7. Suppose that $M_1 > 0$, $M_2 \geq 1$, and $\lambda_0(M_2)$ is defined by (16). Then the equation

$$\lambda_0(M_2) - \frac{12M_1r^2}{\pi(1-r^2)} - \frac{8M_1r^3}{\pi(1-r)^3} - \lambda_0(M_2) \sqrt{M_2^4 - 1} \cdot \left[\frac{2r\sqrt{4r^2 + r^4 + 1}}{(1-r^2)^{5/2}} + \frac{r\sqrt{r^4 - 3r^2 + 4}}{(1-r^2)^{3/2}} \right] = 0 \quad (19)$$

has a unique root in $(0, 1)$.

Lemma 8. Let $M \geq 1$. Then the equation

$$1 - \sqrt{M^4 - 1} \cdot \left[\frac{3r\sqrt{r^4 - 3r^2 + 4}}{(1-r^2)^{3/2}} + \frac{2r\sqrt{4r^2 + r^4 + 1}}{(1-r^2)^{5/2}} \right] = 0 \quad (20)$$

has a unique root in $(0, 1)$.

Lemma 9. For any $z_1 \neq z_2$ in $\mathbb{D}_r$ ($0 < r < 1$), we have

$$\int_0^1 |tz_1 + (1-t)z_2|^2 dt \geq \frac{|z_1|^3 + |z_2|^3}{3(|z_1| + |z_2|)} > 0. \quad (21)$$

2. Main Results

We first establish a new version of the Landau-type theorem for biharmonic mappings on the unit disk $\mathbb{D}$ as follows.

Theorem 10. Let $F(z) = |z|^2 g(z) + h(z)$ be a biharmonic mapping of the unit disk $\mathbb{D}$, with $F(0) = h(0) = \lambda_F(0) - 1 = 0$, $|g(z)| \leq M$, and $\Lambda_h(z) \leq \Lambda$ for $z \in \mathbb{D}$, where $M > 0$, $\Lambda \geq 1$. Then $L(F)$ is univalent in the disk $\mathbb{D}_{r_0}$, where r_0 is the unique root in $(0, 1)$ of the equation

$$1 - \frac{12Mr^2}{\pi(1-r^2)} - \frac{8Mr^3}{\pi(1-r)^3} - \frac{\Lambda^2 - 1}{\Lambda} \cdot \frac{2r - r^2}{(1-r)^2} = 0, \quad (22)$$

and $L(F)(\mathbb{D}_{r_0})$ contains a schlicht disk $\mathbb{D}_{\sigma_0}$, where

$$\sigma_0 = r_0 \left[1 - \frac{\Lambda^2 - 1}{\Lambda} \cdot \frac{r_0}{1-r_0} - \frac{4Mr_0^2}{\pi(1-r_0)^2} \right]. \quad (23)$$

Proof. Let $F(z) = |z|^2 g(z) + h(z)$ satisfy the hypothesis of Theorem 10, where

$$g(z) = g_1(z) + \overline{g_2}(z) = \sum_{n=0}^{\infty} a_n z^n + \sum_{n=0}^{\infty} \overline{b_n} \overline{z}^n, \quad (24)$$

$$h(z) = h_1(z) + \overline{h_2}(z) = \sum_{n=1}^{\infty} c_n z^n + \sum_{n=1}^{\infty} \overline{d_n} \overline{z}^n$$

are harmonic in $\mathbb{D}$. As L is linear and $L(|z|^2) = 0$, we may set

$$H := L(F) = |z|^2 L(g) + L(h). \quad (25)$$

Then we have

$$\begin{aligned} H_z &= 2|z|^2 g_z + |z|^2 z g_{zz} - \overline{z}^2 g_{\overline{z}} + h_z + z h_{zz}, \\ H_{\overline{z}} &= -2|z|^2 \overline{g_z} - |z|^2 \overline{z} g_{\overline{z}\overline{z}} + z^2 g_z - h_{\overline{z}} - \overline{z} h_{\overline{z}\overline{z}}. \end{aligned} \quad (26)$$

Note that $\lambda_F(0) = |c_1| - |d_1| = \lambda_h(0) = 1$; by Lemma 3, we have

$$|c_n| + |d_n| \leq \frac{\Lambda^2 - 1}{n\Lambda}, \quad n = 2, 3, \dots \quad (27)$$

Thus, for $z_1 \neq z_2$ in $\mathbb{D}_r$ ($0 < r < r_0$), we have

$$\begin{aligned} |H(z_1) - H(z_2)| &= \left| \int_{[z_1, z_2]} H_z(z) dz + H_{\overline{z}}(z) d\overline{z} \right| \\ &\geq \left| \int_{[z_1, z_2]} h_z(0) dz - h_{\overline{z}}(0) d\overline{z} \right| \\ &\quad - 2 \left| \int_{[z_1, z_2]} |z|^2 (g_z dz - g_{\overline{z}} d\overline{z}) \right| \\ &\quad - \left| \int_{[z_1, z_2]} |z|^2 (z g_{zz} dz - \overline{z} g_{\overline{z}\overline{z}} d\overline{z}) \right| \\ &\quad - \left| \int_{[z_1, z_2]} z h_{zz} dz - \overline{z} h_{\overline{z}\overline{z}} d\overline{z} \right| \\ &\quad - \left| \int_{[z_1, z_2]} z^2 g_z d\overline{z} - \overline{z}^2 g_{\overline{z}} dz \right| \\ &\quad - \left| \int_{[z_1, z_2]} (h_z - h_z(0)) dz - (h_{\overline{z}} - h_{\overline{z}}(0)) d\overline{z} \right|. \end{aligned} \quad (28)$$

Let

$$I_1 = \left| \int_{[z_1, z_2]} h_z(0) dz - h_{\overline{z}}(0) d\overline{z} \right|,$$

$$I_2 = \left| \int_{[z_1, z_2]} |z|^2 (g_z dz - g_{\overline{z}} d\overline{z}) \right|,$$

$$I_3 = \left| \int_{[z_1, z_2]} |z|^2 (z g_{zz} dz - \overline{z} g_{\overline{z}\overline{z}} d\overline{z}) \right|, \quad (29)$$

$$I_4 = \left| \int_{[z_1, z_2]} z h_{zz} dz - \overline{z} h_{\overline{z}\overline{z}} d\overline{z} \right|,$$

$$I_5 = \left| \int_{[z_1, z_2]} z^2 g_z d\overline{z} - \overline{z}^2 g_{\overline{z}} dz \right|,$$

$$I_6 = \left| \int_{[z_1, z_2]} (h_z - h_z(0)) dz - (h_{\overline{z}} - h_{\overline{z}}(0)) d\overline{z} \right|.$$

By Lemmas 1, 2, and 3, elementary calculations yield that

$$\begin{aligned}
I_1 &\geq \int_{[z_1, z_2]} \lambda_h(0) |dz| = \lambda_h(0) |z_1 - z_2| = |z_1 - z_2|, \\
I_2 &\leq \int_{[z_1, z_2]} |z|^2 (|g_z| |dz| + |g_{\bar{z}}| |d\bar{z}|) \\
&\leq r^2 |z_1 - z_2| \Lambda_g(z) \leq |z_1 - z_2| \frac{4Mr^2}{\pi(1-r^2)}, \\
I_3 &\leq |z_1 - z_2| \sum_{n=2}^{\infty} n(n-1) (|a_n| + |b_n|) r^{n+1} \\
&\leq |z_1 - z_2| \frac{8Mr^3}{\pi(1-r)^3}, \\
I_4 &\leq |z_1 - z_2| \sum_{n=2}^{\infty} n(n-1) (|c_n| + |d_n|) r^{n-1} \\
&\leq |z_1 - z_2| \frac{\Lambda^2 - 1}{\Lambda} \cdot \frac{r}{(1-r)^2}, \\
I_5 &\leq \int_{[z_1, z_2]} (|z|^2 |g_z| |d\bar{z}| + |\bar{z}|^2 |g_{\bar{z}}| |dz|) \\
&\leq r^2 |z_1 - z_2| \Lambda_g(z) \leq |z_1 - z_2| \frac{4Mr^2}{\pi(1-r^2)}, \\
I_6 &\leq |z_1 - z_2| \sum_{n=2}^{\infty} n (|c_n| + |d_n|) r^{n-1} \\
&\leq |z_1 - z_2| \frac{\Lambda^2 - 1}{\Lambda} \cdot \frac{r}{1-r}.
\end{aligned} \tag{30}$$

Using these estimates and Lemma 6, we obtain

$$\begin{aligned}
&|H(z_1) - H(z_2)| \\
&\geq I_1 - 2I_2 - I_3 - I_4 - I_5 - I_6 \\
&\geq |z_1 - z_2| \left[1 - \frac{12Mr^2}{\pi(1-r^2)} \right. \\
&\quad \left. - \frac{8Mr^3}{\pi(1-r)^3} - \frac{\Lambda^2 - 1}{\Lambda} \cdot \frac{2r - r^2}{(1-r)^2} \right] > 0,
\end{aligned} \tag{31}$$

which implies $H(z_1) \neq H(z_2)$.

For any z such that $z \in \partial\mathbb{D}_{r_0}$, by Lemmas 2, 4, and 5, we obtain

$$\begin{aligned}
|H(z)| &= |z|^2 (zg_z - \bar{z}g_{\bar{z}}) + (zh_z - \bar{z}h_{\bar{z}})| \\
&\geq |zh_z(0) - \bar{z}h_{\bar{z}}(0)| \\
&\quad - |z(h_z - h_z(0)) - \bar{z}(h_{\bar{z}} - h_{\bar{z}}(0))| \\
&\quad - |z|^2 (zg_z - \bar{z}g_{\bar{z}})| \\
&\geq r_0 \left[1 - \sum_{n=2}^{\infty} (|c_n| + |d_n|) nr_0^{n-1} \right. \\
&\quad \left. - \sum_{n=1}^{\infty} (|a_n| + |b_n|) nr_0^{n+1} \right] \\
&\geq r_0 \left[1 - \frac{\Lambda^2 - 1}{\Lambda} \cdot \frac{r_0}{1-r_0} - \frac{4Mr_0^2}{\pi(1-r_0)^2} \right] = \sigma_0.
\end{aligned} \tag{32}$$

This completes the proof. $\square$

Next we improve Theorem A as follows.

Theorem 11. Let $F(z) = |z|^2 g(z) + h(z)$ be a biharmonic mapping of the unit disk $\mathbb{D}$, with $F(0) = h(0) = J_F(0) - 1 = 0$, $|g(z)| \leq M_1$, and $|h(z)| \leq M_2$ for $z \in \mathbb{D}$, where $M_1 > 0$, $M_2 \geq 1$. Then $L(F)$ is univalent in the disk $\mathbb{D}_{r_3}$, where r_3 is the unique root in $(0, 1)$ of the equation

$$\begin{aligned}
&\lambda_0(M_2) - \frac{12M_1 r^2}{\pi(1-r^2)} - \frac{8M_1 r^3}{\pi(1-r)^3} - \lambda_0(M_2) \sqrt{M_2^4 - 1} \\
&\cdot \left[\frac{2r\sqrt{4r^2 + r^4 + 1}}{(1-r^2)^{5/2}} + \frac{r\sqrt{r^4 - 3r^2 + 4}}{(1-r^2)^{3/2}} \right] = 0,
\end{aligned} \tag{33}$$

and $L(F)(\mathbb{D}_{r_3})$ contains a schlicht disk $\mathbb{D}_{\sigma_3}$, where $\lambda_0(M_2)$ is defined by (16) and

$$\begin{aligned}
\sigma_3 &= r_3 \left[\lambda_0(M_2) - \lambda_0(M_2) \sqrt{M_2^4 - 1} \right. \\
&\quad \left. \cdot \frac{r_3 \sqrt{r_3^4 - 3r_3^2 + 4}}{(1-r_3^2)^{3/2}} - \frac{4M_1 r_3^2}{\pi(1-r_3)^2} \right].
\end{aligned} \tag{34}$$

Proof. Note that $J_F(0) = |c_1|^2 - |d_1|^2 = J_h(0) = 1$; by Lemma 4, we have

$$\lambda_h(0) \geq \lambda_0(M_2). \tag{35}$$

We adopt the same method in Theorem 10, for $z_1 \neq z_2$ in $\mathbb{D}_r (0 < r < r_3)$; by Lemmas 1, 2, and 5, we get

$$\begin{aligned}
 I_1 &\geq \int_{[z_1, z_2]} \lambda_h(0) |dz| = \lambda_h(0) |z_1 - z_2|, \\
 I_2 &\leq \int_{[z_1, z_2]} |z|^2 (|g_z| |dz| + |g_{\bar{z}}| |d\bar{z}|) \\
 &\leq r^2 |z_1 - z_2| \Lambda_g(z) \leq |z_1 - z_2| \frac{4M_1 r^2}{\pi(1-r^2)}, \\
 I_3 &\leq |z_1 - z_2| \sum_{n=2}^{\infty} n(n-1) (|a_n| + |b_n|) r^{n+1} \\
 &\leq |z_1 - z_2| \frac{8M_1 r^3}{\pi(1-r)^3}, \\
 I_4 &\leq |z_1 - z_2| \sum_{n=2}^{\infty} n(n-1) (|c_n| + |d_n|) r^{n-1} \\
 &\leq |z_1 - z_2| \left(\sum_{n=2}^{\infty} (|c_n| + |d_n|)^2 \right)^{1/2} \\
 &\quad \cdot \left(\sum_{n=2}^{\infty} n^2(n-1)^2 r^{2(n-1)} \right)^{1/2} \\
 &\leq |z_1 - z_2| \lambda_h(0) \sqrt{M_2^4 - 1} \cdot \frac{2r \sqrt{4r^2 + r^4 + 1}}{(1-r^2)^{5/2}}, \\
 I_5 &\leq \int_{[z_1, z_2]} (|z|^2 |g_z| |d\bar{z}| + |\bar{z}|^2 |g_{\bar{z}}| |dz|) \\
 &\leq r^2 |z_1 - z_2| \Lambda_g(z) \leq |z_1 - z_2| \frac{4M_1 r^2}{\pi(1-r^2)}, \\
 I_6 &\leq |z_1 - z_2| \sum_{n=2}^{\infty} n (|c_n| + |d_n|) r^{n-1} \\
 &\leq |z_1 - z_2| \left(\sum_{n=2}^{\infty} (|c_n| + |d_n|)^2 \right)^{1/2} \\
 &\quad \cdot \left(\sum_{n=2}^{\infty} n^2 r^{2(n-1)} \right)^{1/2} \\
 &\leq |z_1 - z_2| \lambda_h(0) \sqrt{M_2^4 - 1} \cdot \frac{r \sqrt{r^4 - 3r^2 + 4}}{(1-r^2)^{3/2}}.
 \end{aligned} \tag{36}$$

Using these estimates and Lemma 7, by (35), we obtain

$$\begin{aligned}
 |H(z_1) - H(z_2)| \\
 \geq I_1 - 2I_2 - I_3 - I_4 - I_5 - I_6
 \end{aligned}$$

$$\begin{aligned}
 &\geq |z_1 - z_2| \left(\lambda_h(0) - \frac{12M_1 r^2}{\pi(1-r^2)} \right. \\
 &\quad \left. - \frac{8M_1 r^3}{\pi(1-r)^3} - \lambda_h(0) \sqrt{M_2^4 - 1} \right. \\
 &\quad \left. \cdot \left[\frac{2r \sqrt{4r^2 + r^4 + 1}}{(1-r^2)^{5/2}} + \frac{r \sqrt{r^4 - 3r^2 + 4}}{(1-r^2)^{3/2}} \right] \right) \\
 &\geq |z_1 - z_2| \left(\lambda_0(M_2) - \frac{12M_1 r^2}{\pi(1-r^2)} \right. \\
 &\quad \left. - \frac{8M_1 r^3}{\pi(1-r)^3} - \lambda_0(M_2) \sqrt{M_2^4 - 1} \right. \\
 &\quad \left. \cdot \left[\frac{2r \sqrt{4r^2 + r^4 + 1}}{(1-r^2)^{5/2}} + \frac{r \sqrt{r^4 - 3r^2 + 4}}{(1-r^2)^{3/2}} \right] \right) > 0,
 \end{aligned} \tag{37}$$

which implies $H(z_1) \neq H(z_2)$.

For any z such that $z \in \partial\mathbb{D}_{r_3}$, by (35) and Lemmas 2 and 5, we obtain

$$\begin{aligned}
 |H(z)| &\geq r_3 \left[\lambda_h(0) - \sum_{n=2}^{\infty} (|c_n| + |d_n|) m r_3^{n-1} \right. \\
 &\quad \left. - \sum_{n=1}^{\infty} (|a_n| + |b_n|) n r_3^{n+1} \right] \\
 &\geq r_3 \left[\lambda_h(0) - \lambda_h(0) \sqrt{M_2^4 - 1} \right. \\
 &\quad \left. \cdot \frac{r_3 \sqrt{r_3^4 - 3r_3^2 + 4}}{(1-r_3^2)^{3/2}} - \frac{4M_1 r_3^2}{\pi(1-r_3)^2} \right] \\
 &\geq r_3 \left[\lambda_0(M_2) - \lambda_0(M_2) \sqrt{M_2^4 - 1} \right. \\
 &\quad \left. \cdot \frac{r_3 \sqrt{r_3^4 - 3r_3^2 + 4}}{(1-r_3^2)^{3/2}} - \frac{4M_1 r_3^2}{\pi(1-r_3)^2} \right] = \sigma_3.
 \end{aligned} \tag{38}$$

This completes the proof. $\square$

Setting $M_1 = M_2 = M$ in Theorem 11, we have the following corollary.

Corollary 12. Let $F(z) = |z|^2 g(z) + h(z)$ be a biharmonic mapping of the unit disk $\mathbb{D}$, with $F(0) = h(0) = J_F(0) - 1 = 0$, and both $g(z)$ and $h(z)$ are bounded by M . Then $L(F)$ is

univalent in the disk $\mathbb{D}_{r_1}$, where r_1 is the minimum root of the equation

$$\lambda_0(M) - \frac{12Mr^2}{\pi(1-r^2)} - \frac{8Mr^3}{\pi(1-r)^3} - \lambda_0(M)\sqrt{M^4-1} \cdot \left[\frac{2r\sqrt{4r^2+r^4+1}}{(1-r^2)^{5/2}} + \frac{r\sqrt{r^4-3r^2+4}}{(1-r^2)^{3/2}} \right] = 0, \quad (39)$$

and $L(F)(\mathbb{D}_{r_1})$ contains a schlicht disk $\mathbb{D}_{\sigma_1}$, where

$$\sigma_1 = r_1 \left[\lambda_0(M) - \lambda_0(M)\sqrt{M^4-1} \cdot \frac{r_1\sqrt{r_1^4-3r_1^2+4}}{(1-r_1^2)^{3/2}} - \frac{4Mr_1^2}{\pi(1-r_1)^2} \right]. \quad (40)$$

In order to show Corollary 12 improves Theorem A, we use Mathematica to compute the approximate values for various choices of M as in Table 1.

Remark 13. From Table 1 we can see, for the same M ,

$$r_1 > \rho_1, \quad \sigma_1 > R_1. \quad (41)$$

Finally we improve Theorems B as follows.

Theorem 14. Let $F(z) = |z|^2 g(z)$ be a biharmonic mapping in $\mathbb{D}$ such that $g(0) = 0$, $J_g(0) = 1$ and $|g(z)| < M$, where $M \geq 1$ and $g(z)$ is harmonic in $\mathbb{D}$. Then $L(F)$ is univalent in the disk $\mathbb{D}_{r_2}$, where r_2 is the minimum positive root in $(0, 1)$ of the following equation:

$$1 - \sqrt{M^4-1} \cdot \left[\frac{3r\sqrt{r^4-3r^2+4}}{(1-r^2)^{3/2}} + \frac{2r\sqrt{4r^2+r^4+1}}{(1-r^2)^{5/2}} \right] = 0, \quad (42)$$

and $L(F)(\mathbb{D}_{r_2})$ contains a schlicht disk $\mathbb{D}_{\sigma_2}$ with

$$\sigma_2 = r_2^3 \lambda_0(M) \left[1 - \sqrt{M^4-1} \cdot \frac{r_2\sqrt{r_2^4-3r_2^2+4}}{(1-r_2^2)^{3/2}} \right], \quad (43)$$

where $\lambda_0(M)$ is defined by (16).

Proof. Let

$$g(z) = g_1(z) + \overline{g_2}(z) = \sum_{n=1}^{\infty} a_n z^n + \sum_{n=1}^{\infty} \overline{b_n} \overline{z}^n \quad (44)$$

Let $H(z) := L(F) = |z|^2 L(g)$; then we have

$$\begin{aligned} H_z &= 2|z|^2 g_z - \overline{z}^2 \overline{g_z} + z|z|^2 g_{zz}, \\ H_{\overline{z}} &= -2|z|^2 g_{\overline{z}} + z^2 g_z - \overline{z}|z|^2 g_{z\overline{z}}. \end{aligned} \quad (45)$$

For $z_1 \neq z_2$ in $\mathbb{D}_r$ ($0 < r < r_2$), by Lemmas 4, 5, 8, and 9, we get

$$\begin{aligned} & |H(z_1) - H(z_2)| \\ & \geq |z_1 - z_2| \left(\int_0^1 |tz_1 + (1-t)z_2|^2 dt \right) \\ & \quad \times \left[\lambda_g(0) - 3 \sum_{n=2}^{\infty} n(|a_n| + |b_n|) r^{n-1} \right. \\ & \quad \left. - \sum_{n=2}^{\infty} n(n-1)(|a_n| + |b_n|) r^{n-1} \right] \\ & \geq |z_1 - z_2| \frac{|z_1|^3 + |z_2|^3}{3(|z_1| + |z_2|)} \\ & \quad \times \left[\lambda_g(0) - 3\lambda_g(0)\sqrt{M^4-1} \cdot \frac{r\sqrt{r^4-3r^2+4}}{(1-r^2)^{3/2}} \right. \\ & \quad \left. - \sqrt{M^4-1} \cdot \lambda_g(0) \cdot \frac{2r\sqrt{4r^2+r^4+1}}{(1-r^2)^{5/2}} \right] \\ & \geq |z_1 - z_2| \frac{|z_1|^3 + |z_2|^3}{3(|z_1| + |z_2|)} \lambda_0(M) \\ & \quad \times \left[1 - 3\sqrt{M^4-1} \cdot \frac{r\sqrt{r^4-3r^2+4}}{(1-r^2)^{3/2}} \right. \\ & \quad \left. - \sqrt{M^4-1} \cdot \frac{2r\sqrt{4r^2+r^4+1}}{(1-r^2)^{5/2}} \right] > 0, \end{aligned} \quad (46)$$

which implies $H(z_1) \neq H(z_2)$.

For any z such that $z \in \partial\mathbb{D}_{r_2}$, by Lemmas 4 and 5, we obtain

$$\begin{aligned} |H(z)| &= |L(|z|^2 g)| \\ & \geq |z|^2 |z g_z(0) - \overline{z} g_{\overline{z}}(0)| \\ & \quad - |z|^2 |z(g_z - g_z(0)) - \overline{z}(g_{\overline{z}} - g_{\overline{z}}(0))| \\ & \geq r_2^3 \left[\lambda_g(0) - \sum_{n=2}^{\infty} n(|a_n| + |b_n|) r_2^{n-1} \right] \\ & \geq r_2^3 \lambda_0(M) \left[1 - \sqrt{M^4-1} \cdot \frac{r_2\sqrt{r_2^4-3r_2^2+4}}{(1-r_2^2)^{3/2}} \right] = \sigma_2. \end{aligned} \quad (47)$$

This completes the proof of Theorem 14. $\square$

In order to show Theorem 14 improves Theorem B, we use Mathematica to compute the approximate values for various choices of M as in Table 2.

Remark 15. From Table 2 we can see, for the same M ,

$$r_2 > \rho_2, \quad \sigma_2 > R_2. \quad (48)$$

TABLE 1: The values of r_1, σ_1 are in Corollary 12. The values of ρ_1, R_1 are in Theorem A.

	$M = 1$	$M = 2$	$M = 3$	$M = 4$	$M = 5$
ρ_1	0.0527621	0.0139445	0.00626165	0.00353488	0.0022661
r_1	0.357671	0.0593158	0.0269865	0.015355	0.00988556
R_1	0.013793	0.00164514	0.00048245	0.00020277	0.00010364
σ_1	0.216467	0.0119479	0.00357231	0.00151701	0.00077955

TABLE 2: The values of r_2, σ_2 are in Theorem 14. The values of ρ_2, R_2 are in Theorem B.

	$M = 2$	$M = 3$	$M = 4$	$M = 5$
ρ_2	0.00623234	0.00277176	0.00155948	0.00099817
r_2	0.032209	0.0139701	0.00782686	0.00500376
R_2	6.54254×10^{-8}	3.83564×10^{-9}	5.12297×10^{-10}	1.07466×10^{-10}
σ_2	9.84416×10^{-6}	5.35363×10^{-7}	7.06092×10^{-8}	1.47596×10^{-8}

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The work was financially supported by Foundation for Distinguished Young Talents in Higher Education of Guangdong China (no. 2013LYM0093) and Training plan for the Outstanding Young Teachers in Higher Education of Guangdong (no. Yq 2013159). The authors are grateful to the anonymous referees for making many suggestions that improved the readability of this paper.

References

- [1] Z. AbdulHadi, Y. Abu Muhanna, and S. Khuri, "On univalent solutions of the biharmonic equation," *Journal of Inequalities and Applications*, no. 5, pp. 469–478, 2005.
- [2] J. Clunie and T. Sheil-Small, "Harmonic univalent functions," *Annales Academiæ Scientiarum Fennicæ A*, vol. 9, pp. 3–25, 1984.
- [3] P. Duren, *Harmonic Mappings in the Plane*, Cambridge University Press, New York, NY, USA, 2004.
- [4] Z. AbdulHadi, Y. Abu Muhanna, and S. Khuri, "On some properties of solutions of the biharmonic equation," *Applied Mathematics and Computation*, vol. 177, no. 1, pp. 346–351, 2006.
- [5] H. Chen, P. M. Gauthier, and W. Hengartner, "Bloch constants for planar harmonic mappings," *Proceedings of the American Mathematical Society*, vol. 128, no. 11, pp. 3231–3240, 2000.
- [6] Sh. Chen, S. Ponnusamy, and X. Wang, "Coefficient estimates and Landau-Bloch's constant for planar harmonic mappings," *Bulletin of the Malaysian Mathematical Sciences Society. Second Series*, vol. 34, no. 2, pp. 255–265, 2011.
- [7] X. Z. Huang, "Estimates on Bloch constants for planar harmonic mappings," *Journal of Mathematical Analysis and Applications*, vol. 337, no. 2, pp. 880–887, 2008.
- [8] M. Liu, "Estimates on Bloch constants for planar harmonic mappings," *Science in China A*, vol. 52, no. 1, pp. 87–93, 2009.
- [9] X. Q. Xia and X. Z. Huang, "Estimates on Bloch constants for planar bounded harmonic mappings," *Chinese Annals of Mathematics A*, vol. 31, no. 6, pp. 769–776, 2010 (Chinese).
- [10] Sh. Chen, S. Ponnusamy, and X. Wang, "Landau's theorem for certain biharmonic mappings," *Applied Mathematics and Computation*, vol. 208, no. 2, pp. 427–433, 2009.
- [11] M.-S. Liu, "Landau's theorems for biharmonic mappings," *Complex Variables and Elliptic Equations*, vol. 53, no. 9, pp. 843–855, 2008.
- [12] M. S. Liu, Z. W. Liu, and Y. C. Zhu, "Landau's theorems for certain biharmonic mappings," *Acta Mathematica Sinica. Chinese Series*, vol. 54, no. 1, pp. 69–80, 2011.
- [13] Y.-C. Zhu and M.-S. Liu, "Landau-type theorems for certain planar harmonic mappings or biharmonic mappings," *Complex Variables and Elliptic Equations*, vol. 58, no. 12, pp. 1667–1676, 2013.
- [14] F. Colonna, "The Bloch constant of bounded harmonic mappings," *Indiana University Mathematics Journal*, vol. 38, no. 4, pp. 829–840, 1989.
- [15] Sh. Chen, S. Ponnusamy, and X. Wang, "Bloch constant and Landau's theorem for planar p -harmonic mappings," *Journal of Mathematical Analysis and Applications*, vol. 373, no. 1, pp. 102–110, 2011.