Research Article

A Few Integrable Couplings of Some Integrable Systems and $(2 + 1)$ -Dimensional Integrable Hierarchies

Binlu Feng,¹ Yufeng Zhang,² and Huanhe Dong³

¹ School of Mathematics and Information Sciences, Weifang University, Weifang 261061, China

² College of Sciences, China University of Mining and Technology, Xuzhou 221116, China

³ School of Mathematics, Shandong University of Science and Technology, Qingdao 266590, China

Correspondence should be addressed to Binlu Feng; fbl_1963@126.com

Received 25 April 2014; Accepted 20 July 2014; Published 14 August 2014

Academic Editor: Wen-Xiu Ma

Copyright © 2014 Binlu Feng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Two high-dimensional Lie algebras are presented for which four $(1 + 1)$ -dimensional expanding integrable couplings of the D-AKNS hierarchy are obtained by using the Tu scheme; one of them is a united integrable coupling model of the D-AKNS hierarchy and the AKNS hierarchy. Then $(2 + 1)$ -dimensional DS hierarchy is derived by using the TAH scheme; in particular, the integrable couplings of the DS hierarchy are obtained.

1. Introduction

Seeking new integrable systems has been an important aspect of soliton theory, and the methods for this are also versatile. For example, Magri [1] once proposed the Lax-pair method for generating $(1 + 1)$ -dimensional integrable Hamiltonian equations. Tu [2] took use of some subalgebras of the Lie algebra A_1 to draw a beautiful picture for generating integrable Hamiltonian equation hierarchies with $1 + 1$ dimensions, and in particular, the AKNS hierarchy, the KN hierarchy, the WKI hierarchy, and others were obtained under the frame of zero curvature equations. Ma [3] called the method the Tu scheme, and Ma also obtained a series of interesting results in [4–6]. Hu [7, 8] further generalized the trace identity of Tu scheme to get the supertrace identity for generating superintegrable Hamiltonian systems which has extended the applications of the Tu scheme. Fan [9, 10] applied the Tu scheme not only to have obtained some new integrable hierarchies of equations, but also to have obtained their some geometric properties. Various Lax pairs can be introduced by some reductions of the self-dual Yang-Mills equations. For example, Chakravarty et al. [11] introduced symmetries and gauge transformations as well as isospectral problems by reducing the self-dual Yang-Mills equations to have obtained $(1 + 1)$ - and $(2 + 1)$ -dimensional integrable systems. Ablowitz et al.

[12] introduced some different Lax pairs by reductions of the self-dual Yang-Mills equations and further obtained some celebrated $(1 + 1)$ - and $(2 + 1)$ -dimensional integrable equations, including the KdV equation and the KP equation. It is remarkable that generating $(2 + 1)$ -dimensional integrable hierarchies is more difficult than $(1 + 1)$ -dimensional ones. The papers [11, 12] were proceeded by introducing a Lie algebra $g = \{a_0 + a_1(\partial/\partial y) + a_2(\partial^2/\partial y^2)\}$, where a_i belong to a ring of $n \times n$ complex matrix functions of y , x , and t , while Tu et al. [13] proposed an efficient and direct approach for generating $(2 + 1)$ -dimensional equation hierarchies by introducing a residue operator, which was called by us the TAH scheme. Because few people further discussed the scheme and investigated other new $(2 + 1)$ -dimensional integrable hierarchies, we would like to recall the scheme and apply it to produce new $(2 + 1)$ -dimensional hierarchies of equations along with multipotential functions. First of all, we recall some related notations.

Let $\mathcal{A}$ be an associative algebra over the field $\mathcal{R}$. An operator $\partial : \mathcal{A} \rightarrow \mathcal{A}$ satisfies that

$$\partial(\alpha f + \beta g) = \alpha \partial f + \beta \partial g, \quad \partial(fg) = (\partial f)g + f(\partial g), \quad (1)$$

where $\alpha, \beta \in \mathcal{R}$; $f, g \in \mathcal{A}$.

Introduce an associative algebra $\mathcal{A}[\xi]$ consisting of the pseudodifferential operator $\sum_{-\infty}^N a_i \xi^i$, where the coefficients $a_i \in \mathcal{A}$, and ξ is an operator given by

$$\xi f = f\xi + (\partial_y f), \quad f \in \mathcal{A}. \quad (2)$$

It can be verified that [13]

$$\xi^n f = \sum_{i \geq 0} \binom{n}{i} (\partial^i f) \xi^{n-i}, \quad n \in \mathbf{Z}, \quad (3)$$

$$f \xi^n = \sum_{i \geq 0} (-1)^i \binom{n}{i} \xi^{n-i} (\partial^i f), \quad n \in \mathbf{Z}. \quad (4)$$

Besides, a residue operator is given by

$$R : \mathcal{A}[\xi] \longrightarrow \mathcal{A}, \quad R\left(\sum a_i \xi^i\right) = a_{-1}. \quad (5)$$

Based on the above, we recall the TAH scheme as follows.

- (1) Fix a matrix operator $U = U(\lambda + \xi, u) \in \mathcal{A}[\xi]$, where $u = (u_1, \dots, u_p)^T$.
- (2) Solving the matrix-operator equation

$$V_x = [U, V] \quad (6)$$

leads to a recursion relation among V_n , where $V = \sum V_n \lambda^{-n}$, from which we will obtain a recursion relation among $g^{(n)} \equiv (g_1^{(n)}, \dots, g_p^{(n)})^T$, where $g_i^{(n)}$ comes from the expansion

$$\left\langle V, \frac{\partial U}{\partial u_i} \right\rangle = \sum_n g_i^{(n)} \lambda^{-n}, \quad (7)$$

where $\langle a, b \rangle = \text{tr } R(ab)$.

- (3) Try to find an operator J and form the hierarchy

$$u_{t_n} = Jg^{(n)}. \quad (8)$$

- (4) Employing the trace identity [13]

$$\frac{\delta}{\delta u_i} \left\langle V, \frac{\partial U}{\partial \lambda} \right\rangle = \left(\lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^{\gamma} \right) \left\langle V, \frac{\partial U}{\partial u_i} \right\rangle \quad (9)$$

could obtain the Hamiltonian structure of the $(2+1)$ -dimensional hierarchy (8).

In the paper, we want to employ the TAH scheme to generalize 2×2 matrix operators which are presented in [13, 14] to 4×4 cases by using the constructed higher-dimensional Lie algebras so that $(2+1)$ -dimensional equation hierarchies could be generated, whose Hamiltonian structure is also obtained by the trace identity (9). For that sake, we first consider the D-AKNS hierarchy which is derived by our Lie algebra, and then we expand the D-AKNS hierarchy to obtain four different integrable models with 4, 5, and 6 potential functions, respectively. In particular, one of them can reduce to the well-known AKNS hierarchy.

2. The D-AKNS Hierarchy and Its Expanding Integrable Models

A Lie subalgebra of the Lie algebra A_1 presents that [14]

$$L = \text{span} \{h_1, h_2, h_3, h_4\}, \quad (10)$$

where

$$h_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad h_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad h_3 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad (11)$$

$$h_4 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

A loop algebra is defined by

$$\tilde{L} = \text{span} \{h_1(n), h_2(n), h_3(n), h_4(n)\}, \quad (12)$$

where $h_i(n) = h_i \lambda^n$, $n \in \mathbf{Z}$, which have the commutative relations

$$\begin{aligned} [h_1(m), h_2(n)] &= h_2(m+n), \\ [h_1(m), h_3(n)] &= -h_3(m+n), \quad [h_1(m), h_4(n)] = 0, \\ [h_2(m), h_3(n)] &= h_1(m+n) - h_4(m+n), \\ [h_2(m), h_4(n)] &= h_2(m+n), \\ [h_3(m), h_4(n)] &= -h_3(m+n), \\ m, n &\in \mathbf{Z}. \end{aligned} \quad (13)$$

Paper [14] had derived the D-AKNS hierarchy. Now we employ the loop algebra $\tilde{L}$ to rederive it again by the Tu scheme to make the paper self-contained by following Tu idea, and then two enlarging Lie algebras of the Lie algebra L can be used to generate the expanding integrable models, specially, containing the $(2+1)$ -dimensional expanding equation hierarchies.

Set

$$\begin{aligned} U &= h_4(1) + sh_4(0) + qh_2(0) + rh_3(0), \\ V &= \sum_{m \geq 0} \left(\sum_{i=1}^4 V_{im} h_i(-m) \right). \end{aligned} \quad (14)$$

A solution to (14) for V is given by

$$\begin{aligned} V_{1m,x} &= qV_{3m} - rV_{2m} = -V_{4m,x}, \\ V_{2,m+1} &= -V_{2m,x} - sV_{2m} - qV_{1m} + qV_{4m}, \\ V_{3,m+1} &= V_{3m,x} - sV_{3m} - rV_{1m} + rV_{4m}. \end{aligned} \quad (15)$$

Letting

$$V^{(n)} = \sum_{m=0}^n \left(\sum_{i=1}^4 V_{im} h_i(n-m) \right) + (V_{1n} - V_{4n}) h_4(0), \quad (16)$$

a direct calculation yields that

$$\begin{aligned} -V_x^{(n)} + [U, V^{(n)}] &= (V_{4n,x} - V_{1n,x})h_4(1) + (-V_{2n,x} - sV_{2n}) \\ &\quad \times h_2(0) - (V_{3n,x} - sV_{3n})h_3(0). \end{aligned} \quad (17)$$

According to the Tu scheme, the zero curvature equation

$$U_{t_n} - V_x^{(n)} + [U, V^{(n)}] = 0 \quad (18)$$

admits that

$$\begin{aligned} q_{t_n} &= V_{2n,x} + sV_{2n}, \\ r_{t_n} &= V_{3n,x} - sV_{3n}, \\ s_{t_n} &= -2V_{4n,x}. \end{aligned} \quad (19)$$

Denoting $u = (q, r, s)^T$, then we have

$$u_{t_n} = \begin{pmatrix} 0 & \partial + s & 0 \\ \partial - s & 0 & 0 \\ 0 & 0 & -2\partial \end{pmatrix} \begin{pmatrix} V_{3n} \\ V_{2n} \\ V_{4n} \end{pmatrix} = J_1 \begin{pmatrix} V_{3n} \\ V_{2n} \\ V_{4n} \end{pmatrix}, \quad (20)$$

which is just right the D-AKNS hierarchy. Obviously, J_1 is a Hamiltonian operator. The Hamiltonian structure of (20) can be obtained by the trace identity in [2]:

$$\frac{\delta}{\delta u} (4V_4) = \left(\lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \right) \begin{pmatrix} V_3 \\ V_2 \\ V_4 \end{pmatrix}. \quad (21)$$

Comparing the coefficients of λ^{-n-1} gives

$$\frac{\delta}{\delta u} (4V_{4,n+1}) = (-n + \gamma) \begin{pmatrix} V_{3n} \\ V_{2n} \\ V_{4n} \end{pmatrix}. \quad (22)$$

If set $V_{2,0} = V_{3,0} = V_{4,0} = 0$, $V_{1,0} = \text{constant}$, we can get some explicit values from (15) $V_{2,1} = -\alpha q$, $V_{3,1} = -\alpha r$, and $V_{1,1} = 0, \dots$

In terms of the initial values, we can get $\gamma = 0$. Thus, we obtain the Hamiltonian structure of (20)

$$u_{t_n} = J_1 \frac{\delta H_n}{\delta u}, \quad (23)$$

where $H_n = -4V_{4,n+1}/n$.

In what follows, we discuss four different expanding integrable models of the D-AKNS hierarchy. With the help of the Lie algebra L , we introduce an 8-dimensional Lie algebra

$$G_1 = \text{span} \{f_1, \dots, f_8\}, \quad (24)$$

where

$$\begin{aligned} f_1 &= \begin{pmatrix} h_1 & 0 \\ 0 & h_1 \end{pmatrix}, & f_2 &= \begin{pmatrix} h_4 & 0 \\ 0 & h_4 \end{pmatrix}, & f_3 &= \begin{pmatrix} h_2 & 0 \\ 0 & h_2 \end{pmatrix}, \\ f_4 &= \begin{pmatrix} h_3 & 0 \\ 0 & h_3 \end{pmatrix}, & f_5 &= \begin{pmatrix} 0 & h_1 \\ 0 & h_1 \end{pmatrix}, & f_6 &= \begin{pmatrix} 0 & h_4 \\ 0 & h_4 \end{pmatrix}, \\ f_7 &= \begin{pmatrix} 0 & h_2 \\ 0 & h_2 \end{pmatrix}, & f_8 &= \begin{pmatrix} 0 & h_3 \\ 0 & h_3 \end{pmatrix}, \end{aligned} \quad (25)$$

which possesses the commutative relations

$$\begin{aligned} [f_1, f_2] &= 0, & [f_1, f_3] &= f_3, & [f_1, f_4] &= -f_4, \\ [f_1, f_5] &= [f_1, f_6] = 0, & [f_1, f_7] &= f_7, \\ [f_1, f_8] &= -f_8, & [f_2, f_3] &= -f_3, & [f_2, f_4] &= f_4, \\ [f_2, f_5] &= [f_2, f_6] = 0, & [f_2, f_7] &= -f_7, \\ [f_2, f_8] &= f_8, & [f_3, f_4] &= f_1 - f_2, & [f_3, f_5] &= -f_7, \\ [f_3, f_6] &= f_7, & [f_3, f_7] &= 0, \\ [f_3, f_8] &= f_5 - f_6, & [f_4, f_5] &= f_8, & [f_4, f_6] &= -f_8, \\ [f_4, f_7] &= f_6 - f_5, & [f_4, f_8] &= [f_5, f_6] = 0, \\ [f_5, f_7] &= f_7, & [f_5, f_8] &= -f_8, & [f_6, f_7] &= -f_7, \\ [f_6, f_8] &= f_8, & [f_7, f_8] &= f_5 - f_6. \end{aligned} \quad (26)$$

A loop algebra can be defined as

$$f_i(n) = f_i \lambda^n, \quad i = 1, 2, \dots, 8;$$

$$[f_i(m), f_j(n)] = [f_i, f_j] \lambda^{m+n}, \quad 1 \leq i, j \leq 8; \quad m, n \in \mathbf{Z}. \quad (27)$$

Now we utilize the loop algebra to introduce the following Lax pair:

$$\begin{aligned} U &= f_2(1) + sf_2(0) + qf_3(0) + rf_4(0) + f_6(1) \\ &\quad + u_1 f_7(0) + u_2 f_8(0) + u_3 f_5(0), \end{aligned}$$

$$\begin{aligned} V &= \sum_{m \geq 0} \left(V_{1m} f_1(-m) + V_{2m} f_3(-m) + V_{3m} f_4(-m) \right. \\ &\quad \left. + V_{4m} f_2(-m) + \sum_{i=5}^8 V_{im} f_i(-m) \right). \end{aligned} \quad (28)$$

According to the Tu scheme, we first need to solve the matrix equation for V :

$$V_x = [U, V] \quad (29)$$

which admits the first part equations (15) and the second part ones as follows:

$$\begin{aligned} (V_{5m})_x &= u_1 V_{3m} - u_2 V_{2m} - (r + u_2) V_{7m} + (q + u_1) V_{8m} \\ &= -(V_{6m})_x, \\ (V_{7m})_x &= -2V_{7,m+1} + (-s + u_3) V_{7m} + (q + u_1) V_{6m} \\ &\quad - (q + u_1) V_{5m} - u_1 V_{1m} + u_1 V_{4m} + u_3 V_{2m} - V_{2,m+1}, \\ (V_{8m})_x &= 2V_{8,m+1} + (s - u_3) V_{8m} - (r + u_2) V_{6m} \\ &\quad + (r + u_2) V_{5m} + u_2 V_{1m} - u_2 V_{4m} - u_3 V_{3m} + V_{3,m+1}. \end{aligned} \quad (30)$$

Taking

$$\begin{aligned}
 V^{(n)} = & \sum_{m=0}^n \left(V_{1m} f_1(n-m) + V_{2m} f_3(n-m) + V_{3m} f_4(n-m) \right. \\
 & \left. + V_{4m} f_2(n-m) + \sum_{i=5}^8 V_{im} f_i(-m) \lambda^n \right) \\
 & + (V_{1n} - V_{4n}) f_2(0) + (V_{6n} - V_{5n}) f_5(0), \quad (31)
 \end{aligned}$$

one infers that

$$\begin{aligned}
 -V_x^{(n)} + [U, V^{(n)}] \\
 = & -(V_{1n} - V_{4n})_x f_2(0) + [V_{2,n+1} + q(V_{1n} - V_{4n})] f_3(0) \\
 & - [V_{3,n+1} - r(V_{6n} - V_{5n})] f_4(0) - (V_{6n} - V_{5n})_x f_5(0) \\
 & + [V_{2,n+1} + 2V_{7,n+1} + u_1(V_{1n} - V_{4n}) \\
 & - (q + u_1)(V_{6n} - V_{5n})] f_7(0) \\
 & + [-2V_{8,n+1} - V_{3,n+1} - u_2(V_{1n} - V_{4n}) \\
 & + (r + u_2)(V_{6n} - V_{5n})] f_8(0). \quad (32)
 \end{aligned}$$

Thus, the zero curvature equation

$$U_{t_n} - V_x^{(n)} + [U, V^{(n)}] = 0 \quad (33)$$

gives rise to

$$u_{t_n} = \begin{pmatrix} q \\ r \\ s \\ u_1 \\ u_2 \\ u_3 \end{pmatrix}_{t_n} = \begin{pmatrix} (V_{2n})_x + sV_{2n} \\ (V_{3n})_x - sV_{3n} \\ -2(V_{4n})_x \\ (V_{7n})_x + sV_{7n} - u_3(V_{2n} + V_{7n}) \\ (V_{8n})_x - sV_{8n} + u_3(V_{3n} + V_{8n}) \\ -2(V_{5n})_x \end{pmatrix}. \quad (34)$$

When $u_1 = u_2 = u_3 = 0$, the hierarchy (34) reduces to the D-AKNS hierarchy.

When $s = u_3 = 0$, (34) reduces to the following expanding integrable system of the AKNS hierarchy:

$$\bar{u}_{t_n} = \begin{pmatrix} q \\ r \\ u_1 \\ u_2 \end{pmatrix}_{t_n} = \begin{pmatrix} 0 & \partial_x & 0 & 0 \\ \partial_x & 0 & 0 & 0 \\ 0 & 0 & 0 & \partial_x \\ 0 & 0 & \partial_x & 0 \end{pmatrix} \begin{pmatrix} V_{3n} \\ V_{2n} \\ V_{8n} \\ V_{7n} \end{pmatrix} = J \begin{pmatrix} V_{3n} \\ V_{2n} \\ V_{8n} \\ V_{7n} \end{pmatrix}. \quad (35)$$

In particular, setting $u_1 = u_2 = 0$, (35) again reduces to the known AKNS hierarchy

$$\tilde{u}_{t_n} = \begin{pmatrix} q \\ r \end{pmatrix}_{t_n} = \begin{pmatrix} 0 & \partial_x \\ \partial_x & 0 \end{pmatrix} \begin{pmatrix} V_{3n} \\ V_{2n} \end{pmatrix}. \quad (36)$$

Therefore, the hierarchy (34) is a united expanding integrable model of the D-AKNS hierarchy, the AKNS hierarchy, and

the hierarchy (36). It is easy to see that (34) is different from that presented in [14], which indicates one of the merits for seeking multipotential expanding integrable hierarchies. However, the Hamiltonian structure of (34) cannot be obtained by the variational identity [15], which makes us feel confusions.

In order to get the second expanding integrable model of the D-AKNS hierarchy, we first introduce the following Lie algebra:

$$G_2 = \text{span} \{g_1, \dots, g_8\}, \quad (37)$$

where

$$\begin{aligned}
 g_1 &= f_1, & g_2 &= f_2, & g_3 &= f_3, & g_4 &= f_4, \\
 g_5 &= \begin{pmatrix} 0 & h_1 \\ h_1 & 0 \end{pmatrix}, & g_6 &= \begin{pmatrix} 0 & h_4 \\ h_4 & 0 \end{pmatrix}, & g_7 &= \begin{pmatrix} 0 & h_2 \\ h_2 & 0 \end{pmatrix}, \\
 g_8 &= \begin{pmatrix} 0 & h_3 \\ h_3 & 0 \end{pmatrix}, \quad (38)
 \end{aligned}$$

which has the following commutative relations:

$$\begin{aligned}
 [g_1, g_2] &= 0, & [g_1, g_3] &= g_3, & [g_1, g_4] &= -g_4, \\
 [g_2, g_3] &= -g_3, & [g_2, g_4] &= g_4, & [g_3, g_4] &= g_1 - g_2, \\
 [g_1, g_5] &= [g_1, g_6] = 0, & [g_1, g_7] &= g_7, \\
 [g_1, g_8] &= -g_8, & [g_2, g_5] &= [g_2, g_6] = 0, \\
 [g_2, g_7] &= -g_7, & [g_2, g_8] &= g_8, & [g_3, g_5] &= -g_7, \\
 [g_3, g_6] &= g_7, & [g_3, g_7] &= 0, & [g_3, g_8] &= g_5 - g_6, \\
 [g_4, g_5] &= g_8, & [g_4, g_6] &= -g_8, & [g_4, g_7] &= g_6 - g_5, \\
 [g_4, g_8] &= [g_5, g_6] = 0, & [g_5, g_7] &= g_3, \\
 [g_5, g_8] &= -g_4, & [g_6, g_7] &= -g_3, & [g_6, g_8] &= g_4, \\
 [g_7, g_8] &= g_1 - g_2. \quad (39)
 \end{aligned}$$

Now we let us compare the Lie algebras G_1 and G_2 . Setting

$$\begin{aligned}
 G_{11} &= \text{span} \{f_1, f_2, f_3, f_4\}, \\
 G_{12} &= \text{span} \{f_5, f_6, f_7, f_8\}, \quad (40)
 \end{aligned}$$

we find that

$$G_1 = G_{11} \oplus G_{12}, \quad [G_{11}, G_{12}] \subset G_{12}. \quad (41)$$

Condition (41) is a sufficient condition for generating integrable couplings.

If we set

$$\begin{aligned}
 G_{21} &= \text{span} \{g_1, g_2, g_3, g_4\}, \\
 G_{22} &= \text{span} \{g_5, g_6, g_7, g_8\}, \quad (42)
 \end{aligned}$$

it is easy to see that

$$G_2 = G_{21} + G_{22}, \quad [G_{21}, G_{22}] \text{ not in } G_{22}. \quad (43)$$

Equation (43) is different from (41) because the subalgebra G_{22} is a semisimple Lie algebra, while G_{12} is not. Therefore, if we start the Lie algebra G_2 to introduce Lax pairs, some new integrable hierarchies could be generated by employing the Tu scheme, which were not integrable couplings of some known integrable systems. In what follows, we want to make use of a loop algebra of the Lie algebra G_2 to introduce a Lax pair so that the second expanding hierarchy of the D-AKNS hierarchy could be derived by the Tu scheme. A loop algebra of the Lie algebra G_2 is defined as

$$\tilde{G}_2 = \text{span} \{g_1(n), \dots, g_8(n)\}, \quad (44)$$

where

$$\begin{aligned} g_i(n) &= g_i \lambda^n, \quad i = 1, 2, \dots, 8; \\ [g_i(m), g_j(n)] &= [g_i, g_j] \lambda^{m+n}, \quad 1 \leq i, j \leq 8; \quad m, n \in \mathbb{Z}. \end{aligned} \quad (45)$$

Consider a Lax pair

$$\begin{aligned} U &= g_2(1) + s g_2(0) + q g_3(0) + r g_4(0) + v_1 g_7(0) \\ &\quad + v_2 g_8(0) + v_3 g_5(0), \\ V &= \sum_{m \geq 0} \left(V_{1m} g_1(-m) + V_{2m} g_3(-m) + V_{3m} g_4(-m) \right. \\ &\quad \left. + V_{4m} g_2(-m) + \sum_{i=5}^8 V_{im} g_i(-m) \right). \end{aligned} \quad (46)$$

Solving the stationary zero curvature equation

$$V_x = [U, V] \quad (47)$$

yields that

$$\begin{aligned} (V_{1m})_x &= q V_{3m} - r V_{2m} + v_1 V_{8m} - v_2 V_{7m} = -(V_{4m})_x, \\ V_{2,m+1} &= -(V_{2m})_x - s V_{2m} - q V_{1m} + q V_{4m} - v_1 V_{5m} \\ &\quad + v_1 V_{6m} + v_3 V_{7m}, \\ (V_{3m})_x &= V_{3,m+1} + s V_{3m} + r V_{1m} - r V_{4m} + v_2 V_{5m} \\ &\quad - v_2 V_{6m} - v_3 V_{8m}, \\ (V_{5m})_x &= q V_{8m} - r V_{7m} + v_1 V_{3m} - v_2 V_{2m} = -(V_{6m})_x, \\ V_{7,m+1} &= -(V_{7m})_x - q V_{5m} + q V_{6m} - v_1 V_{1m} + v_1 V_{4m} \\ &\quad + v_3 V_{2m} - s V_{7m}, \\ (V_{8m})_x &= V_{8,m+1} + s V_{8m} + r V_{5m} - r V_{6m} + v_2 V_{1m} \\ &\quad + v_2 V_{1m} - v_2 V_{4m} - v_3 V_{3m}. \end{aligned} \quad (48)$$

Taking

$$\begin{aligned} V^{(n)} &= \sum_{m=0}^n \left(V_{1m} g_1(n-m) + V_{2m} g_3(n-m) + V_{3m} g_4(n-m) \right. \\ &\quad \left. + V_{4m} g_2(n-m) + \sum_{i=5}^8 V_{im} g_i(n-m) \right) \\ &\quad + (V_{1n} - V_{4n}) f_2(0) + (V_{6n} - V_{5n}) f_5(0), \end{aligned} \quad (49)$$

similar to the above discussion, we obtain the integrable hierarchy

$$u_{t_n} = \begin{pmatrix} q \\ r \\ s \\ v_1 \\ v_2 \\ v_3 \end{pmatrix}_{t_n} = \begin{pmatrix} (V_{2n})_x + s V_{2n} - v_3 V_{7n} \\ (V_{3n})_x - s V_{3n} + v_3 V_{8n} \\ -2(V_{4n})_x \\ (V_{7n})_x + s V_{7n} - v_3 V_{2n} \\ (V_{8n})_x - s V_{8n} + v_3 V_{3n} \\ -2(V_{5n})_x \end{pmatrix}. \quad (50)$$

Obviously, (50) is different from (34). However, when $v_3 = u_3 = 0$ in (50) and (34), respectively, they all reduce to the same equation hierarchy (below we will see it). When $v_1 = v_2 = v_3 = 0$, (50) reduces to the D-AKNS hierarchy. Therefore, (50) is the second expanding integrable model of the D-AKNS hierarchy.

In the following, we consider the Hamiltonian structure of (50). Through direct calculation, we get that

$$\begin{aligned} \langle V, U_q \rangle &= 2V_3, & \langle V, U_r \rangle &= 2V_2, & \langle V, U_s \rangle &= 2V_4, \\ \langle V, U_{v_1} \rangle &= 2V_8, & \langle V, U_{v_2} \rangle &= 2V_7, & \langle V, U_{v_3} \rangle &= 2V_5, \\ \langle V, U_\lambda \rangle &= 2V_4, \end{aligned} \quad (51)$$

and here $\langle a, b \rangle = \text{tr}(ab)$, $V_i = \sum_{m \geq 0} V_{im} \lambda^{-m}$, $i = 1, 2, \dots, 8$.

Substituting the above results into the trace identity gives

$$\frac{\delta}{\delta u} (2V_4) = \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \begin{pmatrix} 2V_3 \\ 2V_2 \\ 2V_4 \\ 2V_8 \\ 2V_7 \\ 2V_5 \end{pmatrix}. \quad (52)$$

Comparing the coefficients of λ^{-n-1} leads to

$$\frac{\delta}{\delta u} (2V_{4,n+1}) = (-n + \gamma) \begin{pmatrix} 2V_{3n} \\ 2V_{2n} \\ 2V_{4n} \\ 2V_{8n} \\ 2V_{7n} \\ 2V_{5n} \end{pmatrix}. \quad (53)$$

Similar to the above discussion, we get $\gamma = 0$. Thus, one gets that

$$\begin{pmatrix} 2V_{3n} \\ 2V_{2n} \\ 2V_{4n} \\ 2V_{8n} \\ 2V_{7n} \\ 2V_{5n} \end{pmatrix} = \frac{\delta H_n}{\delta u}, \quad H_n = -\frac{2V_{4n+1}}{n}. \quad (54)$$

Equation (50) can be written as

$$\begin{aligned} u_{t_n} &= \begin{pmatrix} q \\ r \\ s \\ v_1 \\ v_2 \\ v_3 \end{pmatrix}_{t_n} \\ &= \begin{pmatrix} 0 & \frac{\partial+s}{2} & 0 & 0 & -\frac{v_3}{2} & 0 \\ \frac{\partial-s}{2} & 0 & 0 & \frac{v_3}{2} & 0 & 0 \\ 0 & 0 & -\partial & 0 & 0 & 0 \\ 0 & -\frac{v_3}{2} & 0 & 0 & \frac{\partial+s}{2} & 0 \\ \frac{v_3}{2} & 0 & 0 & \frac{\partial-s}{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -\partial \end{pmatrix} \begin{pmatrix} 2V_{3n} \\ 2V_{2n} \\ 2V_{4n} \\ 2V_{8n} \\ 2V_{7n} \\ 2V_{5n} \end{pmatrix} \\ &= J_2 \frac{\delta H_n}{\delta u}. \end{aligned} \quad (55)$$

When $v_3 = 0$, (55) reduces to

$$\begin{aligned} \bar{u}_{t_n} &= \begin{pmatrix} q \\ r \\ s \\ v_1 \\ v_2 \end{pmatrix}_{t_n} = \begin{pmatrix} (V_{2n})_x + sV_{2n} \\ (V - 3n)_x - sV_{3n} \\ -2(V_{4n})_x \\ (V_{7n})_x + sV_{7n} \\ (V_{8n})_x - sV_{8n} \end{pmatrix} \\ &= \begin{pmatrix} 0 & \frac{\partial+s}{2} & 0 & 0 & 0 \\ \frac{\partial-s}{2} & 0 & 0 & 0 & 0 \\ 0 & 0 & -\partial & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{\partial+s}{2} \\ 0 & 0 & 0 & \frac{\partial-s}{2} & 0 \end{pmatrix} \begin{pmatrix} 2V_{3n} \\ 2V_{2n} \\ 2V_{4n} \\ 2V_{8n} \\ 2V_{7n} \end{pmatrix} \\ &= J_3 \begin{pmatrix} 2V_{3n} \\ 2V_{2n} \\ 2V_{4n} \\ 2V_{8n} \\ 2V_{7n} \end{pmatrix}, \end{aligned} \quad (56)$$

which is also an expanding integrable model of the D-AKNS hierarchy.

3. The (2 + 1)-Dimensional D-AKNS Hierarchy and Its Expanding Models

In the section, we will apply the TAH scheme to deduce the (2 + 1)-dimensional D-AKNS hierarchy and its various expanding models, from which we obtain the united (2 + 1)-dimensional model of the (2 + 1)-dimensional D-AKNS hierarchy and the Davey-Stewartson (DS) hierarchy. The key idea presents that 4×4 matrix operators are introduced by our Lie algebras which extend the 2×2 cases in [13]. The work is worth going on because there are few people applying the method to generate new (2 + 1)-dimensional integrable hierarchies. First of all, we deduce the (2 + 1)-dimensional D-AKNS hierarchy. Set

$$U = \begin{pmatrix} 0 & q \\ r & \lambda + \xi + s \end{pmatrix}, \quad V = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \quad (57)$$

where

$$\begin{aligned} A &= \sum_{m \geq 0} A_m \lambda^{-m}, & B &= \sum_{m \geq 0} B_m \lambda^{-m}, & C &= \sum_{m \geq 0} C_m \lambda^{-m}, \\ D &= \sum_{m \geq 0} D_m \lambda^{-m}. \end{aligned} \quad (58)$$

According to the TAH scheme, we solve the equation of the matrix operators (6) which gives rise to

$$\begin{aligned} A_x &= qC - Br, \\ B_x &= qD - Aq - \lambda B - B\xi - Bs, \\ C_x &= rA - Dr + \lambda C + \xi C + sC, \\ D_x &= rB - Cq + \xi D + sD - D\xi - Ds. \end{aligned} \quad (59)$$

Inserting (58) into (59) reads

$$\begin{aligned} A_{nx} &= qC_n - B_n r, \\ B_{n+1} &= -B_{nx} + qD_n - A_n q - B_n \xi - D_n s, \\ C_{n+1} &= C_{nx} - rA_n + D_n r - \xi C_n - sC_n, \\ D_{nx} - D_{ny} &= rB_n - C_n q + sD_n - D_n s. \end{aligned} \quad (60)$$

Taking a few initial values as follows

$$B_0 = C_0 = D_0 = 0, \quad A_0 = -\xi^{-1}, \quad (61)$$

one infers from (60) that

$$\begin{aligned} B_1 &= q\xi^{-1} - q_y \xi^{-2} + q_{yy} \xi^{-3} + O(\xi^{-4}), \\ C_1 &= r\xi^{-1}, \end{aligned} \quad (62)$$

$$A_1 = \bar{A}_1 \xi^{-2} + \tilde{A}_1 \xi^{-3} + O(\xi^{-4}),$$

where $\partial_x \bar{A}_1 = (qr)_y$, $\partial_x \bar{A}_1 = -q_{yy}$, and

$$\begin{aligned} D_{1x} - D_{1y} + [D_1, s] &= 0 \implies D_1 = 0, \\ B_2 &= -q + (-q_x + q_y - qs) \xi^{-1} + O(\xi^{-2}), \\ C_2 &= -r + (r_x - r_y - sr) \xi^{-1} + O(\xi^{-2}), \\ D_{2x} - D_{2y} + [D_2, s] &= [-(rq)_x + (rq)_y + [-rq, s]] \xi^{-1} \\ &\quad + O(\xi^{-2}) \\ \implies D_2 &= -(rq) \xi^{-1} + O(\xi^{-2}). \end{aligned} \quad (63)$$

Thus, in terms of step (3) of the TAH scheme and (20), we obtain the $(2+1)$ -dimensional D-AKNS hierarchy

$$u_{t_n} = \begin{pmatrix} q \\ r \\ s \end{pmatrix}_{t_n} = \begin{pmatrix} (\partial + s) R(V_{2n}) \\ (\partial - s) R(V_{3n}) \\ -2R(V_{4n,x}) \end{pmatrix}. \quad (64)$$

When $s = 0$, (64) reduces to the DS hierarchy.

When $n = 2$, $t_2 = t$, (64) reduces to

$$\begin{aligned} q_t &= -q_{xx} + q_{xy} + (qs)_x - s(q_x + q_y + qs), \\ r_t &= r_{xx} - r_{xy} - (sr)_x - s(r_x - r_y - sr), \\ s_t &= 2(rq)_x. \end{aligned} \quad (65)$$

Now we consider the Hamiltonian structure of (64). A direct computation gives

$$\begin{aligned} \frac{\partial U}{\partial q} &= U_q = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad U_r = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \\ U_s &= \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad U_\lambda = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \\ \langle V, U_q \rangle &= R(V_3), \quad \langle V, U_r \rangle = R(V_2), \\ \langle V, U_s \rangle &= R(V_4), \quad \langle V, U_\lambda \rangle = 4R(V_4). \end{aligned} \quad (66)$$

Substituting the above results into the trace identity (9) yields that

$$\frac{\delta}{\delta u} (4R(V_4)) = \left(\lambda^{-\gamma} \gamma \frac{\partial}{\partial \lambda} \lambda^\gamma \right) \begin{pmatrix} R(V_3) \\ R(V_2) \\ R(V_4) \end{pmatrix}. \quad (67)$$

Comparing the coefficients of λ^{-n-1} reads

$$\frac{\delta}{\delta u} (4R(V_{4,n+1})) = (-n + \gamma) \begin{pmatrix} R(V_{3n}) \\ R(V_{2n}) \\ R(V_{4n}) \end{pmatrix}. \quad (68)$$

Inserting the initial values in (60) gives $\gamma = 0$. Thus, we have

$$\begin{pmatrix} R(V_{3n}) \\ R(V_{2n}) \\ R(V_{4n}) \end{pmatrix} = \frac{\delta}{\delta u} \left(-\frac{4R(V_{4,n+1})}{n} \right) \equiv \frac{\delta H_n}{\delta u}. \quad (69)$$

Therefore, the $(2+1)$ -dimensional D-AKNS hierarchy (64) can be written as the Hamiltonian form

$$u_{t_n} = \begin{pmatrix} q \\ r \\ s \end{pmatrix}_{t_n} = J \frac{\delta H_n}{\delta u}, \quad (70)$$

where

$$J = \begin{pmatrix} 0 & \partial + s & 0 \\ \partial - s & 0 & 0 \\ 0 & 0 & -2\partial \end{pmatrix}. \quad (71)$$

In what follows, we want to discuss the $(2+1)$ -dimensional expanding models of the $(2+1)$ -dimensional D-AKNS hierarchy (64). Employing the loop algebra of the Lie algebra G_2 introduces the following Lax pair:

$$\begin{aligned} U &= \begin{pmatrix} 0 & q & 0 & v_1 \\ r & \lambda + \xi + s & v_2 & 0 \\ 0 & v_1 & 0 & q \\ v_2 & 0 & r & \lambda + \xi + s \end{pmatrix}, \\ V &= \begin{pmatrix} A & B \\ B & A \end{pmatrix}, \end{aligned} \quad (72)$$

where

$$\begin{aligned} A &= \begin{pmatrix} V_1 & V_2 \\ V_3 & V_4 \end{pmatrix}, \quad B = \begin{pmatrix} V_5 & V_7 \\ V_8 & V_6 \end{pmatrix}, \\ V_i &= \sum_{m \geq 0} V_{im} \lambda^{-m}, \quad i = 1, 2, \dots, 8. \end{aligned} \quad (73)$$

Solving the matrix-operator equation (6) yields that

$$\begin{aligned} V_{1x} &= qV_3 + v_3V_5 + v_1V_8 - V_2r - V_5v_3 - V_7v_2, \\ V_{2x} &= qV_4 + v_3V_7 + v_1V_6 - V_1q - \lambda V_2 - V_2\xi - V_2s - V_5v_1, \\ V_{5x} &= qV_8 + v_3V_1 + v_1V_3 - V_1v_3 - V_2v_2 - V_7r, \\ V_{7x} &= qV_6 + v_3V_2 + v_1V_4 - V_1v_1 - V_5q - V_7\lambda - V_7\xi - V_7s, \\ V_{3x} &= rV_1 + \lambda V_3 + \xi V_3 + sV_3 + v_2V_5 - V_4r - V_8v_3 - V_6v_2, \\ V_{4x} &= rV_2 + \xi V_4 + sV_4 + v_1V_7 - V_3q - V_4\xi - V_4s - V_8v_1, \\ V_{8x} &= rV_5 + \lambda V_8 + \xi V_8 + sV_8 + v_2V_1 - V_3v_3 - V_4v_2 - V_6r, \\ V_{6x} &= rV_7 + \xi V_6 + sV_6 + v_2V_2 - V_3v_1 - V_8q - V_6\xi - V_6s. \end{aligned} \quad (74)$$

Substituting (73) into (74) gets that

$$\begin{aligned} V_{1n,x} &= qV_{3n} + v_1V_{8n} - V_{2n}r - V_{7n}v_2, \\ V_{2,n+1} &= -V_{2n,x} + qV_{4n} + v_1V_{6n} - V_{1n}q - V_{2n}\xi - V_{2n}s - V_{5n}v_1, \end{aligned}$$

$$\begin{aligned}
V_{5n,x} &= qV_{8n} + v_1V_{3n} - V_{2n}v_2 - V_{7n}r, \\
V_{7,n+1} &= -V_{7n,x} + qV_{6n} + v_1V_{4n} - V_{1n}v_1 - V_{5n}q - V_{7n}\xi - V_{7n}s, \\
V_{3,n+1} &= \partial_- V_{3n} - rV_{1n} - V_{3n}\xi - sV_{3n} - v_2V_{5n} + V_{4n}r + V_{6n}v_2, \\
V_{4n,x} &= rV_{2n} + V_{4n,y} + sV_{4n} + v_2V_{7n} - V_{3n}q - V_{4n}s - V_{8n}v_1, \\
V_{8,n+1} &= \partial_- V_{8n} - rV_{5n} - V_{8n}\xi - sV_{8n} - v_2V_{1n} + V_{4n}v_2 + V_{6n}r, \\
V_{6n,x} &= rV_{7n} + V_{6n,y} + sV_{6n} + v_2V_{2n} - V_{3n}v_1 - V_{8n}q - V_{6n}s,
\end{aligned} \tag{75}$$

where $\partial_- = \partial_x - \partial_y$.

Taking

$$\begin{aligned}
V_{2,0} = V_{3,0} = V_{4,0} = V_{6,0} = V_{7,0} = V_{8,0} &= 0, \\
V_{1,0} = V_{5,0} &= -\xi^{-1},
\end{aligned} \tag{76}$$

then one infers from (60) that

$$\begin{aligned}
V_{2,1} &= (q + v_1)\xi^{-1} - (v_{1y} + q_y)\xi^{-2} + (q_{yy} + v_{1yy})\xi^{-3} \\
&\quad + O(\xi^{-4}), \\
V_{3,1} &= V_{8,1} = (r + v_2)\xi^{-1}, \\
V_{7,1} &= (q + v_1)\xi^{-1} - (v_{1y} + q_y)\xi^{-2} + (v_{1yy} + q_{yy})\xi^{-3} \\
&\quad + O(\xi^{-4}), \\
V_{1,1} &= \bar{V}_{1,1}\xi^{-2} + O(\xi^{-3}),
\end{aligned} \tag{77}$$

where $\partial_x \bar{V}_{1,1} = (qr + v_1r + v_1v_2 + qv_2)_y$,

$$V_{5,1} = \bar{V}_{5,1}\xi^{-2} + O(\xi^{-3}), \tag{78}$$

where $\partial_x \bar{V}_{5,1} = (v_1v_2 + v_1r + qr + qv_2)_y$,

$$\begin{aligned}
(V_{4,1})_x + [V_{4,1}, s] &= 0 \implies V_{4,1} = 0, \\
(V_{6,1})_x + [V_{6,1}, s] &= 0 \implies V_{6,1} = 0, \\
V_{2,2} &= -q - v_1 - (q_x - q_y + v_{1x} - v_{1y} - qs - v_1s)\xi^{-1} \\
&\quad + O(\xi^{-2}), \\
V_{7,2} &= (-v_{1x} - q_x + v_{1y} + q_y - v_1s - qs)\xi^{-1} + O(\xi^{-2}), \\
V_{3,2} = V_{8,2} &= -r - v_2 + (r_x + v_{2x} - r_y - v_{2y} - sr - sv_2)\xi^{-1} \\
&\quad + O(\xi^{-2}),
\end{aligned}$$

$$\begin{aligned}
&(V_{4,2})_x - (V_{4,2})_y + [V_{4,2}, s] \\
&= -(rq + rv_1 + v_2v_1 + v_2q)_x \xi^{-1} \\
&\quad + (rq + rv_1 + v_2v_1 + v_2q)_y \xi^{-1} \\
&\quad + [(rq + rv_1 + v_1v_1 + v_2q)s \\
&\quad \quad - s(rq + v_2q + rv_1 + v_2v_1)] \xi^{-1} + O(\xi^{-2}) \\
&\implies V_{4,2} = -(rq + rv_1 + v_2v_1 + v_2q) \xi^{-1}.
\end{aligned} \tag{79}$$

According to step (3) of the TAH scheme and (56), we obtain the $(2+1)$ -dimensional equation hierarchy

$$u_{t_n} = \begin{pmatrix} q \\ r \\ s \\ v_1 \\ v_2 \end{pmatrix}_{t_n} = J_3 \begin{pmatrix} 2R(V_{3n}) \\ 2R(V_{2n}) \\ 2R(V_{4n}) \\ 2R(V_{8n}) \\ 2R(V_{7n}) \end{pmatrix}, \tag{80}$$

where J_3 appears in (56).

When $n = 2$, $t_2 = t$, (80) reduces to the following equations:

$$\begin{aligned}
q_t &= -q_{xx} + q_{xy} - v_{1xx} + v_{1xy} + (qs + v_1s)_x - sq_x - sq_y \\
&\quad - sv_{1x} + sv_{1y} + sqs + sv_1s, \\
r_t &= r_{xx} - r_{xy} + v_{2xx} - v_{2xy} - (sr + sv_2)_x - sr_x - sv_{2x} + sr_y \\
&\quad + sv_{2y} + s^2r + s^2v_2, \\
s_t &= 2(rq + rv_1 + v_2v_1 + v_2q)_x, \\
(v_1)_t &= -q_{xx} + q_{xy} - v_{1xx} + v_{1xy} - (v_1s + qs)_x - sv_{1x} - sq_x \\
&\quad + sv_{1y} + sq_y - sv_1s - sqs, \\
(v_2)_t &= r_{xx} - r_{xy} + v_{2xx} - v_{2xy} - (sr + sv_2)_x - sr_x - sv_{2x} \\
&\quad + sr_y + sv_{2y} + s^2r + s^2v_2.
\end{aligned} \tag{81}$$

We can regard (81) as a kind of expanding model of (65). That is, when letting $s = v_1 = v_2 = 0$, the previous three equations of (81) are just right (65). Therefore, when $s = v_1 = v_2 = 0$, (80) can be regarded as a united model of the $(2+1)$ -dimensional D-AKNS hierarchy and the DS hierarchy.

Remark. Seeking the Hamiltonian structure of (80) is our interesting problem. However, if we apply the trace identity (9) to deduce its Hamiltonian structure, we find that it is the same with that in (70). We guess this is not a right result. In forthcoming days, we will go on to consider the problem.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

This work was supported by the Natural Science Foundation of Shandong Province (Grant no. ZR2013AL016) and the National Natural Science Foundation of China (Grant no. 11371361) as well as the Fundamental Research Funds for the Central Universities (2013XK03).

References

- [1] F. Magri, "A simple model of the integrable Hamiltonian equation," *Journal of Mathematical Physics*, vol. 19, no. 5, pp. 1156–1162, 1978.
- [2] G. Z. Tu, "The trace identity, a powerful tool for constructing the Hamiltonian structure of integrable systems," *Journal of Mathematical Physics*, vol. 30, no. 2, pp. 330–338, 1989.
- [3] W. X. Ma, "A hierarchy of Liouville integrable generalized Hamiltonian equations and its reduction," *Chinese Journal of Contemporary Mathematics*, vol. 13, no. 1, pp. 79–89, 1992.
- [4] W. X. Ma, "A hierarchy of Liouville integrable finite-dimensional Hamiltonian systems," *Applied Mathematics and Mechanics*, vol. 13, no. 4, pp. 369–377, 1992.
- [5] W. Ma, "An approach for constructing non-isospectral hierarchies of evolution equations," *Journal of Physics A: Mathematical and General*, vol. 25, no. 12, pp. L719–L726, 1992.
- [6] W. Ma, "A Hamiltonian structure associated with a matrix spectral problem of arbitrary-order," *Physics Letters. A*, vol. 367, no. 6, pp. 473–477, 2007.
- [7] X. B. Hu, "A powerful approach to generate new integrable systems," *Journal of Physics A: Mathematical and General*, vol. 27, no. 7, pp. 2497–2514, 1994.
- [8] X. Hu, "An approach to generate superextensions of integrable systems," *Journal of Physics A: Mathematical and General*, vol. 30, no. 2, pp. 619–632, 1997.
- [9] E. Fan, "A family of completely integrable multi-Hamiltonian systems explicitly related to some celebrated equations," *Journal of Mathematical Physics*, vol. 42, no. 9, pp. 4327–4344, 2001.
- [10] E. G. Fan, "Integrable evolution systems based on Gerdjikov-Ivanov equations, bi-Hamiltonian structure, finite-dimensional integrable systems and N -fold Darboux transformation," *Journal of Mathematical Physics*, vol. 41, no. 11, pp. 7769–7782, 2000.
- [11] S. Chakravarty, S. L. Kent, and E. T. Newman, "Some reductions of the self-dual Yang-Mills equations to integrable systems in $2 + 1$ dimensions," *Journal of Mathematical Physics*, vol. 36, no. 2, pp. 763–772, 1995.
- [12] M. J. Ablowitz, S. Chakravarty, and L. A. Takhtajan, "A self-dual Yang-Mills hierarchy and its reductions to integrable systems in $1 + 1$ and $2 + 1$ dimensions," *Communications in Mathematical Physics*, vol. 158, no. 2, pp. 289–314, 1993.
- [13] G. Z. Tu, R. I. Andrushkiw, and X. C. Huang, "A trace identity and its application to integrable systems of $1 + 2$ dimensions," *Journal of Mathematical Physics*, vol. 32, no. 7, pp. 1900–1907, 1991.
- [14] G. Z. Tu and D. Z. Meng, "The trace identity, a powerful tool for constructing the Hamiltonian structure of integrable systems (II)," *Acta Mathematicae Applicatae Sinica*, vol. 5, no. 1, pp. 89–96, 1989.
- [15] W. Ma and M. Chen, "Hamiltonian and quasi-Hamiltonian structures associated with semi-direct sums of Lie algebras," *Journal of Physics A: Mathematical and General*, vol. 39, no. 34, pp. 10787–10801, 2006.