Research Article

On Convergence in L -Valued Fuzzy Topological Spaces

A. A. Ramadan,¹ M. El-Dardery,² and Hu Zhao³

¹Department of Mathematics, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt

²Department of Mathematics, Faculty of Science, Fayoum University, Fayoum, Egypt

³School of Science, Xian Polytechnic University, Xian 710048, China

Correspondence should be addressed to M. El-Dardery; mdardery6@gmail.com

Received 30 November 2014; Revised 28 December 2014; Accepted 29 December 2014

Academic Editor: Abdelghani Bellouquid

Copyright © 2015 A. A. Ramadan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce the concept of L -fuzzy neighborhood systems using complete MV -algebras and present important links with the theory of L -fuzzy topological spaces. We investigate the relationships among the degrees of L -fuzzy r -adherent points (r -convergent, r -cluster, and r -limit, resp.) in an L -fuzzy topological spaces. Also, we investigate the concept of LF -continuous functions and their properties.

1. Introduction

Šostak [1–3] introduced a new definition of L -fuzzy topology as the concept of the degree of the openness of fuzzy set. It is an extension of $I = [0, 1]$ -fuzzy topology defined by Chang [4]. It has been developed in many directions [5–11]. The study of neighborhood systems and convergence of nets in Chang fuzzy topology was initiated by Pao-Ming and Ying-Ming [11] and Liu and Luo [12]. In [13] Ying introduced the degree to which a fuzzy point x_t belongs to a fuzzy subset λ by $m(x_t, \lambda) = \min(1, 1 - t + \lambda(x))$ and gave the idea of graded neighborhood on fuzzy topological spaces. This plays an important role in the theory of convergence in Chang fuzzy topology see also [14–18]. Following Ying [13], Demirci [5] introduced the idea of graded neighborhood systems in smooth topological spaces [19] (a smooth topology is similar to fuzzy topology as defined by Šostak [1], Hazra and Samanta [6]) in a different approach but restricted himself to the I -valued fuzzy sets.

In this paper, we study the concept of L -fuzzy neighborhood systems and present important links with the theory of L -fuzzy topological spaces and investigate some of their properties. We investigate the relationships among the degrees of L -fuzzy r -adherent points (r -convergent, r -cluster, and r -limit, resp.) nets in an L -fuzzy topological spaces. Also, we give some related examples to illustrate some

of the introduced notions. In the end, we characterize LF -continuous functions in terms of some of the various notions introduced in this paper.

2. Preliminaries

Throughout the text we consider $(L, \leq, \wedge, \vee, 0, 1)$ as a completely distributive lattice with 0 and 1, respectively, being the universal upper and lower bound and $L_0 = L - \{0\}$. A lattice L is called order dense if for each $a, b \in L$ such that $a < b$, there exist $c \in L$ such that $a < c < b$. If L is a completely distributive lattice and $x \triangleleft \bigvee_{i \in \Gamma} y_i$, then there must be $i_0 \in \Gamma$ such that $x \triangleleft y_{i_0}$, where $x \triangleleft a$ means $K \subset L$, $a \leq \bigvee K \Rightarrow \exists y \in K$ such that $x \leq y$. If $a \triangleleft b$ and $c \triangleleft d$, we always assume $a \wedge c \triangleleft b \wedge d$ [20] and some properties of $\triangleleft$ can be found in [12].

A completely distributive lattice $L = (L, \leq, \wedge, \vee, \odot, \rightarrow, 0, 1)$ (or L , in short) is called a residuated lattice [9, 21–23] if it satisfies the following conditions: for each $x, y, z \in L$,

- (R1) $(L, \odot, 1)$ is a commutative monoid,
- (R2) if $x \leq y$, then $x \odot z \leq y \odot z$ ($\odot$ is isotone operation),
- (R3) (Galois correspondence) $x \leq y \rightarrow z \Leftrightarrow x \odot y \leq z$.

In a residuated lattice L , $x' = x \rightarrow 0$ is called complement of $x \in L$.

A residuated lattice L is called a BL -algebra [9, 21, 23] if it satisfies the following conditions: for each $x, y, z \in L$,

- (B1) $x \wedge y = x \odot (x \rightarrow y)$,
- (B2) $x \vee y = [(x \rightarrow y) \rightarrow y] \wedge [(y \rightarrow x) \rightarrow x]$,
- (B3) $(x \rightarrow y) \vee (y \rightarrow x) = 1$.

A BL -algebra is called an MV -algebra if $x = x''$, for each $x \in L$.

Lemma 1 (see [9, 21, 23]). *Let L be a complete MV -algebra. For each $x, y, z \in L$, $\{y_i, x_i \mid i \in \Gamma\} \subset L$, one has the following properties:*

- (1) $x \odot y \leq x \wedge y \leq x \vee y$,
- (2) $x \odot y \leq x, y$,
- (3) If $y \leq z$, $(x \odot y) \leq (x \odot z)$, $x \rightarrow y \leq x \rightarrow z$ and $z \rightarrow x \leq y \rightarrow x$,
- (4) $x \odot y = (x \rightarrow y')'$,
- (5) $x \leq y$ iff $x' \geq y'$,
- (6) $x \rightarrow y = y' \rightarrow x'$,
- (7) $\bigwedge_{i \in \Gamma} (x \odot y_i) = x \odot (\bigwedge_{i \in \Gamma} y_i)$,
- (8) $\bigvee_{i \in \Gamma} (x \odot y_i) = x \odot (\bigvee_{i \in \Gamma} y_i)$,
- (9) $x \rightarrow 1 = 1, 0 \rightarrow x = 1, x \rightarrow x = 1$,
- (10) $x \leq y \Leftrightarrow x \rightarrow y = 1$ and $1 \rightarrow x = x$,
- (11) $x \rightarrow \bigwedge_{i \in \Gamma} y_i = \bigwedge_{i \in \Gamma} (x \rightarrow y_i)$,
- (12) $(\bigvee_{i \in \Gamma} y_i) \rightarrow x = \bigwedge_{i \in \Gamma} (y_i \rightarrow x)$,
- (13) $x \rightarrow \bigvee_{i \in \Gamma} y_i = \bigvee_{i \in \Gamma} (x \rightarrow y_i)$,
- (14) $\bigwedge_{i \in \Gamma} y_i \rightarrow x = \bigvee_{i \in \Gamma} (y_i \rightarrow x)$,
- (15) $\bigwedge_{i \in \Gamma} y_i' = (\bigvee_{i \in \Gamma} y_i)'$ and $\bigvee_{i \in \Gamma} y_i' = (\bigwedge_{i \in \Gamma} y_i)'$.

In this paper, we always assume that L is a complete MV -algebra. Let X be a nonempty set, and the family L^X denotes the set of all L -fuzzy subsets of a given set X . For $\alpha \in L, \lambda \in L^X$, we denote $(\alpha \rightarrow \lambda), (\alpha \odot \lambda)$, and $\alpha_X \in L^X$ as $(\alpha \rightarrow \lambda)(x) = \alpha \rightarrow \lambda(x), (\alpha \odot \lambda)(x) = \alpha \odot \lambda(x)$, and $\alpha_X(x) = \alpha$.

A fuzzy point x_t for $t \in L_0$ is an element of L^X such that

$$x_t(y) = \begin{cases} t, & \text{if } y = x, \\ 0, & \text{if } y \neq x. \end{cases} \quad (1)$$

The set of all fuzzy points in X is denoted by $\text{Pt}(X)$. For $\lambda \in L^X$ and $x_t \in \text{Pt}(X)$, $x_t \in \lambda$ if and only if $t \leq \lambda(x)$.

Given a mapping $\phi : X \rightarrow Y$, we write $\phi^{\leftarrow}$ for the mapping $L^Y \rightarrow L^X$ defined by $\phi^{\leftarrow}(\mu) = \mu \circ \phi$; we write $\phi^{\rightarrow}$ for the mapping $L^X \rightarrow L^Y$ defined by $\phi^{\rightarrow}(\mu)(y) = \bigvee \{\mu(x) \mid \phi(x) = y\}$ for all $\mu \in L^X, y \in Y$.

For a given set X , define a binary mapping $S(\cdot) : L^X \times L^X \rightarrow L$ as

$$S(\lambda, \mu) = \bigwedge_{x \in X} (\lambda(x) \rightarrow \mu(x)), \quad \forall (\lambda, \mu) \in L^X \times L^X. \quad (2)$$

For each $\lambda, \mu \in L^X$, $S(\lambda, \mu)$ can be interpreted as the degree to which λ is fuzzy included in μ . It is called the L -fuzzy inclusion order [24].

Lemma 2 (see [24]). *For each $\lambda, \mu, \rho, \mu_i \in L^X, i \in \Gamma$ and $e, x_t \in \text{Pt}(X)$, the following properties hold:*

- (1) $\lambda \leq \mu \Leftrightarrow S(\lambda, \mu) = 1$,
- (2) $\lambda \leq \mu \Rightarrow S(\rho, \lambda) \leq S(\rho, \mu)$ and $S(\lambda, \rho) \geq S(\mu, \rho)$, for any $\rho \in L^X$,
- (3) $S(x, \lambda) = \lambda(x)$, for any $\lambda \in L^X$,
- (4) $S(x_t, \lambda) = 0$ if and only if $t = 1$ and $\lambda(x) = 0$,
- (5) $S(e, \lambda) \wedge S(e, \mu) = S(e, \lambda \wedge \mu)$,
- (6) $S(x_t, \bigwedge_{i \in \Gamma} \mu_i) = \bigwedge_{i \in \Gamma} S(x_t, \mu_i)$, for any $\{\mu_i\}_{i \in \Gamma} \subset L^X$,
- (7) $S(x_t, \bigvee_{i \in \Gamma} \mu_i) = \bigvee_{i \in \Gamma} S(x_t, \mu_i)$, for any $\{\mu_i\}_{i \in \Gamma} \subset L^X$.

Lemma 3 (see [16]). *Let $f : X \rightarrow Y$ be a mapping. Then the following statement hold:*

- (1) $S(\lambda, \mu) \leq S(f^{\rightarrow}(\lambda), f^{\rightarrow}(\mu))$, for each $\lambda, \mu \in L^X$
- (2) $S(\rho, \nu) \leq S(f^{\leftarrow}(\rho), f^{\leftarrow}(\nu))$, for each $\rho, \nu \in L^Y$.

In particular, if the mapping $f : X \rightarrow Y$ is bijective, and then the equalities hold.

Definition 4 (see [1, 9]). A map $\mathcal{T} : L^X \rightarrow L$ is called an L -fuzzy topology on X if it satisfies the following conditions:

- (LO1) $\mathcal{T}(1_X) = \mathcal{T}(0_X) = 1$,
- (LO2) $\mathcal{T}(\mu_1 \wedge \mu_2) \geq \mathcal{T}(\mu_1) \wedge \mathcal{T}(\mu_2)$, for all $\mu_1, \mu_2 \in L^X$,
- (LO3) $\mathcal{T}(\bigvee_{i \in \Lambda} \mu_i) \geq \bigwedge_{i \in \Lambda} \mathcal{T}(\mu_i)$, for any $\{\mu_i\}_{i \in \Lambda} \subset L^X$.

The pair $(X, \mathcal{T})$ is called an L -fuzzy topological space.

Let $\mathcal{T}_1$ and $\mathcal{T}_2$ be L -fuzzy topologies on X . We say that $\mathcal{T}_1$ is finer than $\mathcal{T}_2$ ($\mathcal{T}_2$ is coarser than $\mathcal{T}_1$), denoted by $\mathcal{T}_2 \leq \mathcal{T}_1$, if $\mathcal{T}_2(\lambda) \leq \mathcal{T}_1(\lambda)$ for all $\lambda \in L^X$. Let $(X, \mathcal{T}_1)$ and $(Y, \mathcal{T}_2)$ be L -fuzzy topological space spaces. A map $f : (X, \mathcal{T}_1) \rightarrow (Y, \mathcal{T}_2)$ is L -fuzzy continuous (LF -continuous, for short) if $\mathcal{T}_2(\lambda) \leq \mathcal{T}_1(f^{\leftarrow}(\lambda)), \forall \lambda \in L^Y$.

Theorem 5 (see [7, 9]). *Let $(X, \mathcal{T})$ be an L -fuzzy topological space. For each $r \in L_0$ and $\lambda \in L^X$, one defines operators $I_{\mathcal{T}}, C_{\mathcal{T}} : L^X \times L_0 \rightarrow L^X$ as follows:*

$$\begin{aligned} I_{\mathcal{T}}(\lambda, r) &= \bigvee \{\rho \in L^X \mid \rho \leq \lambda, \mathcal{T}(\rho) \geq r\}, \\ C_{\mathcal{T}}(\lambda, r) &= \bigwedge \{\nu \in L^X \mid \lambda \leq \nu, \mathcal{T}(\nu') \geq r\}. \end{aligned} \quad (3)$$

For each $\lambda, \mu \in L^X$ and $r, s \in L_0$, one has the following properties:

- (I1) $\mathcal{I}_{\mathcal{T}}(1_X, r) = 1_X$,
- (I2) $\mathcal{I}_{\mathcal{T}}(\lambda, r) \leq \lambda$,
- (I3) if $\lambda \leq \mu$ and $r \leq s$, then $\mathcal{I}_{\mathcal{T}}(\lambda, s) \leq \mathcal{I}_{\mathcal{T}}(\mu, r)$,
- (I4) $\mathcal{I}_{\mathcal{T}}(\lambda \wedge \mu, r \wedge s) \geq \mathcal{I}_{\mathcal{T}}(\lambda, r) \wedge \mathcal{I}_{\mathcal{T}}(\mu, s)$,
- (I5) $\mathcal{I}_{\mathcal{T}}(\mathcal{I}_{\mathcal{T}}(\lambda, r), r) = \mathcal{I}_{\mathcal{T}}(\lambda, r)$,
- (I6) $\mathcal{I}_{\mathcal{T}}(\lambda', r) = (C_{\mathcal{T}}(\lambda, r))'$.

Definition 6 (see [12]). Let D be a directed set. A function $T : D \rightarrow \text{Pt}(X)$ is called a fuzzy net in X . Let $\lambda \in L^X$, and one says that T is a fuzzy net in λ if $T(n) \in \lambda$ for every $n \in D$.

Definition 7 (see [12, 25]). Let T be a fuzzy net and $\lambda \in L^X$.

- (1) T is often in λ if for each $n \in D$, there exists $n_0 \in D$ such that $n_0 \geq n$ and $T(n_0) \in \lambda$.
- (2) T is finally in λ if there exists $n_0 \in D$ such that for each $n \in D$ with $n \geq n_0$, one has $T(n) \in \lambda$.

Definition 8 (see [12, 25]). Let $T : D \rightarrow \text{Pt}(X)$ and $U : E \rightarrow \text{Pt}(X)$ be two fuzzy nets. A fuzzy net U is called a subnet of T if there exists a function $N : E \rightarrow D$, called by a cofinal selection on T , such that

- (1) $U = T \circ N$;
- (2) for every $n_0 \in D$, there exists $m_0 \in E$ such that $N(m) \geq n_0$, for $m \geq m_0$.

3. L-Fuzzy Neighborhood Systems

Definition 9. Let $\lambda \in L^X$ and $x_t \in \text{Pt}(X)$. Then the degree to which x_t belongs to λ is

$$S(x_t, \lambda) = \bigwedge_{x \in X} (t \rightarrow \lambda(x)). \quad (4)$$

Definition 10. Let $(X, \mathcal{T})$ be an L -fuzzy topological space, $\lambda \in L^X$, $e \in \text{Pt}(X)$, and $r \in L_0$. The degree to which λ is a r -neighborhood of e is defined by

$$(\mathcal{N}^{\mathcal{T}})_e(\lambda, r) = \bigvee \{S(e, \mu) \mid \mu \leq \lambda, r \triangleleft \mathcal{T}(\mu)\}. \quad (5)$$

A mapping $(\mathcal{N}^{\mathcal{T}})_e : L^X \times L_0 \rightarrow L$ is called the L -fuzzy neighborhood system of e .

Theorem 11. Let $(X, \mathcal{T})$ be an L -fuzzy topological space and let $(\mathcal{N}^{\mathcal{T}})_e$ be the fuzzy neighborhood system of e . For all $\lambda, \mu \in L^X$ and $r, s \in L_0$, the following properties hold:

- (1) $(\mathcal{N}^{\mathcal{T}})_e(0_X, r) = S(e, 0_X)$ and $(\mathcal{N}^{\mathcal{T}})_e(1_X, r) = 1$,
- (2) $(\mathcal{N}^{\mathcal{T}})_e(\lambda, r) \leq S(e, \lambda)$,
- (3) $(\mathcal{N}^{\mathcal{T}})_e(\lambda, r) \geq (\mathcal{N}^{\mathcal{T}})_e(\lambda, s)$, if $r \leq s$,
- (4) $(\mathcal{N}^{\mathcal{T}})_e(\lambda, r) \leq (\mathcal{N}^{\mathcal{T}})_e(\mu, r)$, if $\lambda \leq \mu$,
- (5) $(\mathcal{N}^{\mathcal{T}})_e(\lambda_1, r) \wedge (\mathcal{N}^{\mathcal{T}})_e(\lambda_2, s) \leq (\mathcal{N}^{\mathcal{T}})_e(\lambda_1 \wedge \lambda_2, r \wedge s)$,
- (6) $(\mathcal{N}^{\mathcal{T}})_e(\lambda, r) \leq \bigvee \{(\mathcal{N}^{\mathcal{T}})_e(\mu, r) \mid \mu \leq \lambda, S(d, \mu) \leq (\mathcal{N}^{\mathcal{T}})_d(\mu, r) \forall d \in \text{Pt}(X)\}$,
- (7) $(\mathcal{N}^{\mathcal{T}})_{x_t}(\lambda, r) = \bigwedge_{x \in X} (t \rightarrow (\mathcal{N}^{\mathcal{T}})_{x_t}(\lambda, r))$.

Proof. (1), (3), and (4) are easily proved.

(2) is proved from the following:

$$\begin{aligned} (\mathcal{N}^{\mathcal{T}})_e(\lambda, r) &= \bigvee \{S(e, \mu_i) \mid \mu_i \leq \lambda, r \triangleleft \mathcal{T}(\mu_i)\} \\ &\leq \bigvee \left\{ S\left(e, \bigvee \mu_i\right) \mid \mu_i \leq \lambda, r \triangleleft \mathcal{T}(\mu_i) \right\} \\ &\quad \text{(by Lemma 2 (2))} \\ &\leq \left\{ S\left(e, \bigvee \mu_i\right) \mid \bigvee \mu_i \leq \lambda, r \leq \mathcal{T}\left(\bigvee \mu_i\right) \right\} \\ &\leq S(e, \lambda). \end{aligned} \quad (6)$$

In (5) if $a \triangleleft (\mathcal{N}^{\mathcal{T}})_e(\lambda_1, r) \wedge (\mathcal{N}^{\mathcal{T}})_e(\lambda_2, s)$, then $a \triangleleft (\mathcal{N}^{\mathcal{T}})_e(\lambda_1, r)$ and $a \triangleleft (\mathcal{N}^{\mathcal{T}})_e(\lambda_2, s)$, and there exists $\rho_1 \in L^X$ with $\rho_1 \leq \lambda_1$ and $r \triangleleft \mathcal{T}(\rho_1)$ such that $a \triangleleft S(e, \rho_1)$. Again, there exists $\rho_2 \in L^X$ with $\rho_2 \leq \lambda_2$ and $s \triangleleft \mathcal{T}(\rho_2)$ such that $a \triangleleft S(e, \rho_2)$. So, $\rho_1 \wedge \rho_2 \leq \lambda_1 \wedge \lambda_2$, $r \wedge s \triangleleft \mathcal{T}(\rho_1) \wedge \mathcal{T}(\rho_2)$, and $a \leq S(e, \rho_1) \wedge S(e, \rho_2) = S(e, \rho_1 \wedge \rho_2) \leq (\mathcal{N}^{\mathcal{T}})_e(\lambda_1 \wedge \lambda_2, r \wedge s)$. Hence,

$$(\mathcal{N}^{\mathcal{T}})_e(\lambda_1 \wedge \lambda_2, r \wedge s) \geq (\mathcal{N}^{\mathcal{T}})_e(\lambda_1, r) \wedge (\mathcal{N}^{\mathcal{T}})_e(\lambda_2, s). \quad (7)$$

In (6) if $r \triangleleft \mathcal{T}(\mu)$, then $S(d, \mu) = (\mathcal{N}^{\mathcal{T}})_d(\mu, r)$, for each $d \in \text{Pt}(X)$. It implies

$$\begin{aligned} (\mathcal{N}^{\mathcal{T}})(\lambda, r) &= \bigvee \{S(e, \mu) \mid \mu \leq \lambda, r \triangleleft \mathcal{T}(\mu)\} \\ &= \bigvee \{(\mathcal{N}^{\mathcal{T}})_e(\mu, r) \mid \mu \leq \lambda, \\ &\quad S(d, \mu) = (\mathcal{N}^{\mathcal{T}})_d(\mu, r), \\ &\quad \forall d \in \text{Pt}(X)\} \\ &\leq \bigvee \{(\mathcal{N}^{\mathcal{T}})_e(\mu, r) \mid \mu \leq \lambda, \\ &\quad S(d, \mu) \leq (\mathcal{N}^{\mathcal{T}})_d(\mu, r), \\ &\quad \forall d \in \text{Pt}(X)\}. \end{aligned} \quad (8)$$

(7) is proved from

$$\begin{aligned} (\mathcal{N}^{\mathcal{T}})_{x_t}(\lambda, r) &= \bigvee \{S(x_t, \mu) \mid \mu \leq \lambda, \mathcal{T}(\mu) \geq r\} \\ &= \bigvee \left\{ \bigwedge_{x \in X} (t \rightarrow \mu(x)) \mid \mu \leq \lambda, \mathcal{T}(\mu) \geq r \right\} \end{aligned}$$

$$\begin{aligned}
&= \bigwedge_{x \in X} \left\{ t \longrightarrow \bigvee \{ \mu(x) \mid \mu \leq \lambda, \mathcal{T}(\mu) \geq r \} \right\} \\
&\quad \text{(by Lemma 2 (7))} \\
&= \bigwedge_{x \in X} \left(t \longrightarrow (\mathcal{N}^{\mathcal{T}})_{x_1}(\lambda, r) \right). \\
\end{aligned} \tag{9}$$

Theorem 12. Let X be a nonempty set. Let for each $e \in \text{Pt}(X)$, and $\mathcal{N}_e : L^X \times L_0 \rightarrow L$ satisfying the above conditions (1)–(5). Define $\mathcal{T}_{\mathcal{N}} : L^X \rightarrow L$ by

$$\mathcal{T}_{\mathcal{N}}(\lambda) = \bigvee \{ r \in L_0 \mid S(e, \lambda) = \mathcal{N}_e(\lambda, r), \forall e \in \text{Pt}(X) \}. \tag{10}$$

Then one has the following:

- (a) $\mathcal{T}_{\mathcal{N}}$ is an L -fuzzy topology on X ;
- (b) if $(\mathcal{N}^{\mathcal{T}})_e$ is the L -fuzzy neighborhood system of e induced by $(X, \mathcal{T})$, then $\mathcal{T}_{\mathcal{N}^{\mathcal{T}}} = \mathcal{T}$;
- (c) if $\mathcal{N}_e$'s satisfy the conditions (6) and (7), then

$$\mathcal{T}_{\mathcal{N}}(\lambda) = \bigvee \{ r \in L_0 \mid S(x, \lambda) = \mathcal{N}_x(\lambda, r), \forall x \in X \}; \tag{11}$$

- (d) $\mathcal{N}_{\mathcal{T}_{\mathcal{N}}} = \mathcal{N}$.

Proof. (a) (LO1) It is easily proved from Theorem 11(1).

(LO2) It is proved from the following:

$$\begin{aligned}
&\mathcal{T}_{\mathcal{N}}(\lambda_1) \wedge \mathcal{T}_{\mathcal{N}}(\lambda_2) \\
&= \left(\bigvee \{ r \in L_0 \mid S(e, \lambda_1) = \mathcal{N}_e(\lambda_1, r) \} \right) \\
&\quad \wedge \left(\bigvee \{ s \in L_0 \mid S(e, \lambda_2) = \mathcal{N}_e(\lambda_2, s) \} \right) \\
&= \bigvee \{ r \wedge s \in L_0 \mid S(e, \lambda_1) \wedge S(e, \lambda_2) \\
&\quad = \mathcal{N}_e(\lambda_1, r) \wedge \mathcal{N}_e(\lambda_2, s) \} \\
&\leq \bigvee \{ r \wedge s \in L_0 \mid S(e, \lambda_1) \wedge S(e, \lambda_2) \\
&\quad \leq \mathcal{N}_e(\lambda_1 \wedge \lambda_2, r \wedge s) \} \\
&\leq \bigvee \{ r \wedge s \in L_0 \mid S(e, \lambda_1 \wedge \lambda_2) \leq \mathcal{N}_e(\lambda_1 \wedge \lambda_2, r \wedge s) \} \\
&\quad \text{(by Lemma 2 (5))} \\
&\leq \mathcal{T}_{\mathcal{N}}(\lambda_1 \wedge \lambda_2). \\
\end{aligned} \tag{12}$$

(LO3) If $a \triangleleft \bigwedge_{i \in \Gamma} \mathcal{T}_{\mathcal{N}}(\lambda_i)$, then $a \triangleleft \mathcal{T}_{\mathcal{N}}(\lambda_i)$ for each $i \in \Gamma$, and note that

$$\begin{aligned}
\mathcal{T}_{\mathcal{N}}(\lambda_i) &= \bigvee \{ r_i \in L_0 \mid S(e, \lambda_i) = \mathcal{N}_e(\lambda_i, r_i), \\
&\quad \forall e \in \text{Pt}(X) \}, \\
\end{aligned} \tag{13}$$

so there exists $r_i \in L_0$, with $S(e, \lambda_i) = \mathcal{N}_e(\lambda_i, r_i)$ such that $a \triangleleft r_i$. Put $r = \bigwedge_{i \in \Gamma} r_i$, and then $a \leq r$. By Theorem 11, we have

$$S(e, \lambda_i) \leq \mathcal{N}_e(\lambda_i, r_i) \leq \mathcal{N}_e(\lambda_i, r) \leq S(e, \lambda_i). \tag{14}$$

It implies $S(e, \lambda_i) = \mathcal{N}_e(\lambda_i, r)$. Furthermore, by Lemma 2(7), we have

$$\begin{aligned}
&S\left(e, \bigvee_{i \in \Gamma} \lambda_i\right) \\
&= \bigvee_{i \in \Gamma} S(e, \lambda_i) = \bigvee_{i \in \Gamma} \mathcal{N}_e(\lambda_i, r_i) \\
&\leq \bigvee_{i \in \Gamma} \mathcal{N}_e(\lambda_i, r) \leq \mathcal{N}_e\left(\bigvee_{i \in \Gamma} \lambda_i, r\right) \leq S\left(e, \bigvee_{i \in \Gamma} \lambda_i\right). \\
\end{aligned} \tag{15}$$

So $\mathcal{N}_e(\bigvee_{i \in \Gamma} \lambda_i, r) = S(e, \bigvee_{i \in \Gamma} \lambda_i)$. Hence, $\mathcal{T}_{\mathcal{N}}(\bigvee_{i \in \Gamma} \lambda_i) \geq r \geq a$. Therefore, $\mathcal{T}_{\mathcal{N}}(\bigvee_{i \in \Gamma} \lambda_i) \geq \bigwedge_{i \in \Gamma} \mathcal{T}_{\mathcal{N}}(\lambda_i)$.

(b) If $a \triangleleft \mathcal{T}_{\mathcal{N}}(\lambda)$, then there exists $r_0 \in L_0$ with $S(e, \lambda) = \mathcal{N}_e(\lambda, r_0)$ such that $r_0 \triangleleft \mathcal{T}(\lambda)$. Since

$$S(e, \lambda) = \mathcal{N}_e(\lambda, r_0) = \bigvee \{ S(e, \mu_i) \mid \mu_i \leq \lambda, r_0 \triangleleft \mathcal{T}(\mu_i) \}, \tag{16}$$

then, for each $x_1 \in \text{Pt}(X)$,

$$\begin{aligned}
\lambda(x) &= S(x_1, \lambda) \\
&= \bigvee \{ S(x_1, \mu_i) \mid \mu_i \leq \lambda, r_0 \triangleleft \mathcal{T}(\mu_i) \} \\
\end{aligned} \tag{17}$$

$$= S\left(x_1, \bigvee_{i \in \Gamma} \mu_i\right) = \bigvee_{i \in \Gamma} \mu_i(x).$$

Thus, $\lambda = \bigvee \mu_i$. So $\mathcal{T}(\lambda) \geq r_0 \geq a$. Hence, $\mathcal{T}_{\mathcal{N}}(\lambda) \leq \mathcal{T}(\lambda)$. We can easily obtain $\mathcal{T}_{\mathcal{N}}(\lambda) \geq \mathcal{T}(\lambda)$.

(c) We only show that $S(x_t, \lambda) = \mathcal{N}_{x_t}(\lambda, r)$, $\forall x_t \in \text{Pt}(X)$ if and only if $S(x, \lambda) = \lambda(x) = \mathcal{N}_x(\lambda, r)$, $\forall x \in X$.

($\Rightarrow$) It is trivial.

($\Leftarrow$) From condition (7),

$$\begin{aligned}
\mathcal{N}_{x_t}(\lambda, r) &= \bigwedge_{x \in X} (t \longrightarrow \mathcal{N}_{x_t}(\lambda, r)) \\
&= \bigwedge_{x \in X} (t \longrightarrow S(x_1, \lambda)) \\
&= \bigwedge_{x \in X} (t \longrightarrow \lambda(x)) \\
&= S(x_t, \lambda). \\
\end{aligned} \tag{18}$$

(d) From the proof of Theorem 11(6), we easily obtain $\mathcal{N}_{\mathcal{T}_{\mathcal{N}}} \geq \mathcal{N}$.

If $a \triangleleft (\mathcal{N}_{\mathcal{T}_{\mathcal{N}}})_e(\lambda, r) = \bigvee \{ S(e, \mu) \mid \mu \leq \lambda, r \triangleleft \mathcal{T}_{\mathcal{N}}(\mu) \}$, there exists μ_0 with $\mu_0 \leq \lambda$, $r \triangleleft \mathcal{T}_{\mathcal{N}}(\mu_0)$ such that $a \triangleleft S(e, \mu_0)$. Note that

$$\mathcal{T}_{\mathcal{N}}(\mu_0) = \bigvee \{ t \in L_0 \mid S(e, \mu_0) = \mathcal{N}_e(\mu_0, t), \forall e \in \text{Pt}(X) \}, \tag{19}$$

and there exists $t_0 \in L_0$ with $S(e, \mu_0) = \mathcal{N}_e(\mu_0, t_0)$ such that $r \triangleleft t_0$ (thus $r \leq t_0$). So $a \triangleleft \mathcal{N}_e(\mu_0, t_0) \leq \mathcal{N}_e(\mu_0, r) \leq \mathcal{N}_e(\lambda, r)$. Therefore, $\mathcal{N}_{\mathcal{T}, \mathcal{N}} \leq \mathcal{N}$. $\square$

By Theorem 12, we have the following corollary.

Corollary 13. *The set of all L-fuzzy topologies on X and the set of all L-fuzzy neighborhood systems on X are in one to one correspondence.*

Example 14. Let $L = [0, 1]$, $X = \{a, b\}$ be a set, $x \rightarrow y = \min(1 - x + y, 1)$, and let $\mu \in L^X$ be defined as follows:

$$\mu(a) = 0.3, \quad \mu(b) = 0.4. \quad (20)$$

We define an L-fuzzy topology on X as

$$\mathcal{T}(\lambda) = \begin{cases} 1, & \text{if } \lambda = 0_X \text{ or } 1_X, \\ \frac{1}{2}, & \text{if } \lambda = \mu, \\ 0, & \text{otherwise.} \end{cases} \quad (21)$$

From Definition 10, $\mathcal{N}_{a_1}, \mathcal{N}_{b_1} : L^X \times L_0 \rightarrow L$ as follows:

$$\mathcal{N}_{a_1}(\lambda, r) = \begin{cases} 1, & \text{if } \lambda = 1_X, \quad r \in L_0, \\ 0.3, & \text{if } 1_X \neq \lambda \geq \mu, \quad 0 < r \leq \frac{1}{2}, \\ 0, & \text{otherwise,} \end{cases} \quad (22)$$

$$\mathcal{N}_{b_1}(\lambda, r) = \begin{cases} 1, & \text{if } \lambda = 1_X, \quad r \in L_0, \\ 0.4, & \text{if } 1_X \neq \lambda \geq \mu, \quad 0 < r \leq \frac{1}{2}, \\ 0, & \text{otherwise.} \end{cases}$$

From Theorem 12(c), we have

$$\mathcal{T}_{\mathcal{N}}(\lambda) = \begin{cases} 1, & \text{if } \lambda = 0_X \text{ or } 1_X, \\ \frac{1}{2}, & \text{if } \lambda = \mu, \\ 0, & \text{otherwise.} \end{cases} \quad (23)$$

4. R-Convergence

Definition 15. Let $(X, \mathcal{T})$ be an L-fuzzy topological space, $\lambda \in L^X$, $e \in \text{Pt}(X)$, and $r \in L_0$. The degree to which a fuzzy net T in X is r -convergent to e and T is r -cluster to e are defined, respectively, as follows:

$$\text{Con}_e(T, r) = \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid T \text{ is often in } \lambda' \},$$

$$\text{Cl}_e(T, r) = \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid T \text{ is finally in } \lambda' \}. \quad (24)$$

Definition 16. Let $(X, \mathcal{T})$ be an L-fuzzy topological space, $\lambda \in L^X$, $e \in \text{Pt}(X)$, and $r \in L_0$. The degree to which e is r -adherent point of e is defined by

$$\text{Ad}_e(\lambda, r) = \mathcal{N}'_e(\lambda', r). \quad (25)$$

Proposition 17. Let $(X, \mathcal{T})$ be an L-fuzzy topological space. For each $\lambda \in L^X$, $e, x_t \in \text{Pt}(X)$ and $r \in L_0$, one has

- (1) $S(e, I_{\mathcal{T}}(\lambda, r)) = \mathcal{N}_e(\lambda, r)$,
- (2) $S(e, C'_{\mathcal{T}}(\lambda, r)) = \text{Ad}'_e(\lambda, r)$,
- (3) $\text{Ad}_{x_t}(\lambda, r) = \bigvee_{x \in X} (t \odot \text{Ad}_x(\lambda, r))$.

Proof. (1) From Lemma 2(7), we have

$$\begin{aligned} S(e, I_{\mathcal{T}}(\lambda, r)) &= S\left(e, \bigvee \{ \mu_i \mid \mu_i \leq \lambda, \mathcal{T}(\mu_i) \geq r \}\right) \\ &= \bigvee \{ S(e, \mu_i) \mid \mu_i \leq \lambda, \mathcal{T}(\mu_i) \geq r \} \\ &= \mathcal{N}_e(\lambda, r). \end{aligned} \quad (26)$$

(2) From Theorem 5, we have

$$\begin{aligned} S(e, C'_{\mathcal{T}}(\lambda, r)) &= S(e, I_{\mathcal{T}}(\lambda', r)) \\ &= \mathcal{N}_e(\lambda', r) \quad (\text{by (1)}) \\ &= \text{Ad}'_e(\lambda, r). \end{aligned} \quad (27)$$

(3) From Theorem 11(7), we have

$$\begin{aligned} \text{Ad}_{x_t}(\lambda, r) &= \mathcal{N}'_{x_t}(\lambda', r) \\ &= \left(\bigwedge_{x \in X} (t \rightarrow \mathcal{N}_{x_t}(\lambda', r)) \right)' \\ &= \bigvee_{x \in X} (t \rightarrow \mathcal{N}_{x_t}(\lambda', r))' \\ &= \bigvee_{x \in X} (t \odot \mathcal{N}'_{x_t}(\lambda', r)) \\ &\quad (\text{by Lemma 2 (4)}) \\ &= \bigvee_{x \in X} (t \odot \text{Ad}_x(\lambda, r)). \end{aligned} \quad (28)$$

$\square$

Theorem 18. Let $(X, \mathcal{T})$ be an L-fuzzy topological space. Let $T : D \rightarrow \text{Pt}(X)$ be fuzzy net and let $U : E \rightarrow \text{Pt}(X)$ be a subnet of S . For $r, s \in L_0$, the following properties hold:

- (1) if $r_1 \leq r_2$, $\text{Con}_e(T, r_1) \leq \text{Con}_e(T, r_2)$, and $\text{Cl}_e(T, r_1) \leq \text{Cl}_e(T, r_2)$,
- (2) $\text{Con}_e(T, r) \leq \text{Cl}_e(T, r)$,
- (3) $\text{Cl}_e(U, r) \leq \text{Cl}_e(T, r)$,
- (4) $\text{Con}_e(T, r) \leq \text{Con}_e(U, r)$,
- (5) $\text{Con}_{x_t}(T, r) = \bigvee_{x \in X} (t \odot \text{Con}_x(T, r))$, and $\text{Cl}_{x_t}(T, r) = \bigvee_{x \in X} (t \odot \text{Cl}_x(T, r))$.

Proof. (1) is easily proved.

In (2) if T is finally in λ' , T is often in λ' . Hence

$$\begin{aligned} \text{Con}_e(T, r) &= \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid T \text{ is often in } \lambda' \} \\ &\leq \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid T \text{ is finally in } \lambda' \} \\ &= \text{Cl}_e(T, r). \end{aligned} \quad (29)$$

In (3) if T is finally in λ' , U is finally in λ' . Hence

$$\begin{aligned} \text{Cl}_e(U, r) &= \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid U \text{ is finally in } \lambda' \} \\ &\leq \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid T \text{ is finally in } \lambda' \} \\ &= \text{Cl}_e(T, r). \end{aligned} \quad (30)$$

In (4) let U be often in λ' . We will show that T is often in λ' . Let $n \in D$. Since $U : E \rightarrow \text{Pt}(X)$ is a subnet of T , there exists a cofinal selection $N : E \rightarrow D$. For each $n \in D$, there exists $m \in E$ such that $N(k) \geq n$ for $k \geq m$. Since U is often in λ' , for $m \in E$, there exists $m_0 \in E$ such that $m_0 \geq m$ for $U(m_0) \in \lambda'$. Put $n_0 = N(m_0)$. Then $n_0 \geq n$ and $T(n_0) = T(N(m_0)) = T(n_0) \in \lambda'$. Thus, U is often in λ' . Hence

$$\begin{aligned} \text{Con}_e(T, r) &= \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid T \text{ is often in } \lambda' \} \\ &\leq \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid U \text{ is often in } \lambda' \} \\ &= \text{Con}_e(U, r). \end{aligned} \quad (31)$$

In (5) one has

$$\begin{aligned} \text{Con}_{x_i}(T, r) &= \bigwedge \{ \mathcal{N}'_{x_i}(\lambda, r) \mid T \text{ is often in } \lambda' \} \\ &= \bigwedge \left\{ \left(\bigwedge_{x \in X} (t \rightarrow \mathcal{N}_{x_i}(\lambda, r)) \right)' \mid \right. \\ &\quad \left. T \text{ is finally in } \lambda' \right\} \\ &\quad (\text{by Theorem 11 (7)}) \\ &= \bigvee_{x \in X} \bigwedge \left\{ (t \rightarrow \mathcal{N}_{x_i}(\lambda, r))' \mid \right. \\ &\quad \left. T \text{ is finally in } \lambda' \right\} \\ &= \bigvee_{x \in X} \bigwedge \{ t \odot \mathcal{N}'_{x_i}(\lambda, r) \mid T \text{ is finally in } \lambda' \} \\ &\quad (\text{by Lemma 1 (4)}) \\ &= \bigvee_{x \in X} (t \odot \bigwedge \{ \mathcal{N}'_{x_i}(\lambda, r) \mid T \text{ is finally in } \lambda' \}) \\ &= \bigvee_{x \in X} (t \odot \text{Con}_x(T, r)). \end{aligned} \quad (32)$$

The other case is the same. $\square$

Proposition 19. Let $(X, \mathcal{T})$ be an L -fuzzy topological space, let T be a fuzzy net, $e \in \text{Pt}(X)$, and $r \in L_0$. Then one has

$$\begin{aligned} \text{Ad}_e(\lambda, r) &= \bigvee \{ \text{Con}_e(T, r) \mid T \text{ is a fuzzy net in } \lambda \} \\ &= \bigvee \{ \text{Cl}_e(T, r) \mid T \text{ is a fuzzy net in } \lambda \}. \end{aligned} \quad (33)$$

Proof. Since T is finally in λ , T is often in λ . We easily show that

$$\begin{aligned} \text{Ad}_e(\lambda, r) &= \mathcal{N}'_e(\lambda', r) \\ &\geq \bigvee \{ \text{Cl}_e(T, r) \mid T \text{ is a fuzzy net in } \lambda \} \\ &\geq \bigvee \{ \text{Con}_e(T, r) \mid T \text{ is a fuzzy net in } \lambda \}. \end{aligned} \quad (34)$$

We only show that

$$\text{Ad}_e(\lambda, r) \leq \bigvee \{ \text{Con}_e(T, r) \mid T \text{ is a fuzzy net in } \lambda \}. \quad (35)$$

Let $\text{Ad}_e(\lambda, r) = t$. If $t > 0$, then $\mathcal{N}'_e(\lambda', r) = t$. Put $D = \{ \mu \in L^X \mid \mathcal{N}_e(\mu, r) > t' \}$. Define a relation on D by

$$\mu_1 \leq \mu_2 \quad \text{iff} \quad \mu_1 \geq \mu_2, \quad \forall \mu_1, \mu_2 \in D. \quad (36)$$

For each $\mu_1, \mu_2 \in D$, since by Theorem 11(5),

$$\mathcal{N}_e(\mu_1 \wedge \mu_2, r) \geq \mathcal{N}_e(\mu_1, r) \wedge \mathcal{N}_e(\mu_2, r) > t'. \quad (37)$$

Hence, $\mu_1 \wedge \mu_2 \in D$ and $\mu_1, \mu_2 \leq \mu_1 \wedge \mu_2$. Thus, $(D, \leq)$ is a directed set. For each $\mu \in D$, that is, $\mathcal{N}_e(\mu, r) > t'$, we have $\mu \notin \lambda'$; that is, there exists $x \in X$ such that $\lambda(x) > \mu'(x)$. Thus, we can define a fuzzy net $T_0 : D \rightarrow \text{Pt}(X)$ by $T_0(\mu) = x_{\lambda(x)}$ where $T_0(\mu) \in \lambda$ and $\lambda(x) = T_0(\mu)(x) > \mu'(x)$.

We will show that if $\mu \in D$, then T_0 is not often in μ' . Suppose that T_0 is often in μ' . For $\mu \in D$, there exists $\rho \in D$ such that $\mu \leq \rho$ such that

$$T_0(\rho) = y_{\lambda(y)} \in \mu', \quad (38)$$

and $\lambda(y) = T_0(\rho)(y) > \rho'(y)$. Since $\mu \leq \rho$ implies $\mu \geq \rho$, it implies

$$\lambda(y) \leq \mu'(y) \leq \rho'(y), \quad (39)$$

It is contradiction for the definition of T_0 . Thus, if T_0 is often in μ' , then $\mu \notin D$; that is, $\mathcal{N}_e(\mu, r) \leq t'$. Therefore,

$$\begin{aligned} &\bigvee \{ \text{Con}_e(T, r) \mid T \text{ is a fuzzy net in } \lambda \} \\ &\geq \text{Con}_e(T, r) \\ &= \bigwedge \{ \mathcal{N}'_e(\mu, r) \mid T_0 \text{ is often in } \mu' \} \\ &\geq t = \text{Ad}_e(\lambda, r). \end{aligned} \quad (40)$$

$\square$

Theorem 20. Let $(X, \mathcal{T})$ be L -fuzzy topological space and let $T, U : D \rightarrow \text{Pt}(X)$ be fuzzy nets such that $T(n) \vee U(n), T(n) \wedge U(n) \in \text{Pt}(X)$ for each $n \in D$. Define fuzzy nets $T \vee U, T \wedge U : D \rightarrow \text{Pt}(X)$ by, for each $n \in D$,

$$\begin{aligned} (T \vee U)(n) &= T(n) \vee U(n), \\ (T \wedge U)(n) &= T(n) \wedge U(n). \end{aligned} \quad (41)$$

For each $r \in L_0$, the following properties hold:

(1) if $T(n) \leq U(n)$ for all $n \in D$, then

$$Cl_e(T, r) \leq Cl_e(U, r), \quad Con_e(T, r) \leq Con_e(U, r), \quad (42)$$

$$(2) Cl_e(T \wedge U, r) \leq Cl_e(T, r) \wedge Cl_e(U, r),$$

$$(3) Con_e(T \vee U, r) \geq Con_e(T, r) \vee Con_e(U, r),$$

$$(4) Con_e(T \wedge U, r) \leq Con_e(T, r) \wedge Con_e(U, r),$$

$$(5) \text{ if } L \text{ is order dense, then } Cl_e(T \vee U, r) = Cl_e(T, r) \vee Cl_e(U, r).$$

Proof. In (1) let U be finally (often) in λ . Then let T be finally (often) in λ , respectively. Thus it is trivial. (2), (3), and (4) are easily proved.

In (5) since $T \leq T \vee U$ and $U \leq T \vee U$, by (1), we have

$$Cl_e(T \vee U, r) \geq Cl_e(T, r) \vee Cl_e(U, r). \quad (43)$$

Suppose that $Cl_e(T \vee U, r) \not\geq Cl_e(T, r) \vee Cl_e(U, r)$. Since L is order dense, then there exist $t \in L_0$ and a fuzzy point $e \in Pt(X)$ such that

$$Cl_e(T \vee U, r) > t > Cl_e(T, r) \vee Cl_e(U, r). \quad (44)$$

Since $Cl_e(T, r) < t$ and $Cl_e(U, r) < t$, by the definition Cl_e , there exist $\lambda, \mu \in L^X$ such that T and U are finally in λ' and μ' , respectively, with

$$Cl_e(T, r) \vee Cl_e(U, r) \leq \mathcal{N}'_e(\lambda, r) \vee \mathcal{N}'_e(\mu, r) < t. \quad (45)$$

Since T is finally in λ' , there exists $n_1 \in D$ such that $T(n) \in \lambda'$ for every $n \in D$ with $n \geq n_1$. Since U is finally in μ' , there exists $n_2 \in D$ such that $T(n) \in \mu'$ for every $n \in D$ with $n \geq n_2$. Let $n_3 \in D$ such that $n_3 \geq n_1$ and $n_3 \geq n_2$. For $n \geq n_3$, we have

$$(T \vee U)(n) \leq \lambda' \vee \mu' = (\lambda \wedge \mu)'. \quad (46)$$

Thus, $(T \vee U)$ is finally in $(\lambda \wedge \mu)'$. It implies

$$\begin{aligned} Cl_e(T \vee U, r) &\leq \mathcal{N}'_e(\lambda \wedge \mu, r) \\ &\leq \mathcal{N}'_e(\lambda, r) \vee \mathcal{N}'_e(\mu, r) < t. \end{aligned} \quad (47)$$

It is a contradiction. Hence, we have

$$Cl_e(T \vee U, r) \leq Cl_e(T, r) \vee Cl_e(U, r). \quad (48)$$

□

Example 21. Let $(L = [0, 1], \rightarrow)$ be defined as Example 14. Let $X = \{a, b\}$ be a set and $\mu \in I^X$ as follows:

$$\mu(x) = 0.3, \quad \mu(y) = 0.4. \quad (49)$$

We define L -fuzzy topology $\mathcal{T} : I^X \rightarrow I$ as follows:

$$\mathcal{T}(\lambda) = \begin{cases} 1, & \text{if } \lambda = 0_X \text{ or } 1_X, \\ \frac{1}{2}, & \text{if } \lambda = \mu, \\ 0, & \text{otherwise.} \end{cases} \quad (50)$$

(1) In general, $Cl_e(T \wedge U, r) \neq Cl_e(T, r) \wedge Cl_e(U, r)$.

Let N be a natural numbers. Define fuzzy nets $T, U : N \rightarrow Pt(X)$ by

$$\begin{aligned} T(n) &= x_{a_n}, \quad a_n = 0.8 + (-1)^n 0.2, \\ U(n) &= x_{b_n}, \quad b_n = 0.8 + (-1)^{n+1} 0.2. \end{aligned} \quad (51)$$

From Theorem 20, $(T \wedge U)(n) = x_{0.6}$ is a fuzzy net. Let $e = x_{0.3}$. From Definition 15, we have for $0 < r \leq 1/2$,

$$Cl_e(x_{0.6}, r) = 1 - \mathcal{N}_e(\mu, r) = 1 - m(x_{0.3}, \mu) = 0. \quad (52)$$

Since T or U is finally in 1_X ,

$$Cl_e(T, r) = 1 - \mathcal{N}_e(0_X, r) = 1 - m(x_{0.3}, 0_X) = 0.3. \quad (53)$$

Similarly, $Cl_e(U, r) = 0.3$. For $0 < r \leq 1/2$,

$$0 = Cl_e(T \wedge U, r) \neq Cl_e(T, r) \wedge Cl_e(U, r) = 0.3. \quad (54)$$

(2) In general, $Con_e(T \vee U, r) \neq Con_e(T, r) \vee Con_e(U, r)$. Define fuzzy nets $T, U : N \rightarrow Pt(X)$ by

$$\begin{aligned} T(n) &= x_{a_n}, \quad a_n = 0.6 + (-1)^n 0.2, \\ U(n) &= x_{b_n}, \quad b_n = 0.6 + (-1)^{n+1} 0.2. \end{aligned} \quad (55)$$

From Theorem 20, $(T \vee U)(n) = x_{0.8}$ is a fuzzy net. Let $e = x_{0.3}$. For all $r \in I_0$,

$$Ad_e(x_{0.8}, r) = 1 - \mathcal{N}_e(0_X, r) = 1 - m(x_{0.3}, 0_X) = 0.3. \quad (56)$$

Since T or U is often in μ' , for $0 < r \leq 1/2$,

$$Cl_e(T, r) = 1 - \mathcal{N}_e(\mu, r) = 1 - m(x_{0.3}, \mu) = 0. \quad (57)$$

Similarly, $Cl_e(U, r) = 0$. For $0 < r \leq 1/2$

$$0.3 = Con_e(T \vee U, r) > (Con_e(T, r) \vee Con_e(U, r)) = 0. \quad (58)$$

5. Fuzzy r -Limit Nets and LF -Continuous Mappings

Definition 22. Let $(X, \mathcal{T})$ be an L -fuzzy topological space. Let $T : D \rightarrow Pt(X)$ be fuzzy net in X , $e \in Pt(X)$, and $r \in L_0$. Then the degree to which T is r -limit to e is defined, denoted by $\lim_e(T, r) = t$, if $Cl_e(T, r) = Con_e(T, r) = t$.

Theorem 23. Let $(X, \mathcal{T})$ be L -fuzzy topological space and let $T, U : D \rightarrow Pt(X)$ be fuzzy nets such that $T(n) \vee U(n) \in Pt(X)$ for each $n \in D$. If L is order dense, $Cl_e(T, r) = Con_e(T, r)$, and $Cl_e(U, r) = Con_e(U, r)$, then

$$\lim_e(T \vee U, r) = \lim_e(T, r) \vee \lim_e(U, r). \quad (59)$$

Proof. From Theorem 20, $T \vee U$ is a fuzzy net. We easily proved it from the following:

$$\begin{aligned}
 \text{Cl}_e(T \vee U, r) &= \text{Cl}_e(T, r) \vee \text{Cl}_e(U, r) \quad (\text{by Theorem 20 (2)}) \\
 &(\text{since } \text{Cl}_e(T, r) = \text{Con}_e(T, r), \text{Cl}_e(U, r) = \text{Con}_e(U, r)) \\
 &= \text{Con}_e(T, r) \vee \text{Con}_e(U, r) \\
 &\leq \text{Con}_e(T \vee U, r) \quad (\text{by Theorem 20 (4)}) \\
 &\leq \text{Cl}_e(T \vee U, r) \quad (\text{by Theorem 20 (2)}).
 \end{aligned} \tag{60}$$

□

Theorem 24. Let $(X, \mathcal{F})$ be L -fuzzy topological space. Let T be a fuzzy net and $\mathcal{H} = \{U \mid U \text{ is a subnet of } T\}$. Then, if L is an order dense, the following statements hold:

- (1) $\text{Con}_e(T, r) = \bigwedge_{U \in \mathcal{H}} \text{Cl}_e(U, r)$;
- (2) $\text{Cl}_e(T, r) = \bigvee_{U \in \mathcal{H}} \text{Con}_e(U, r)$.

Proof. (1) For each $U \in \mathcal{H}$, by Theorem 18, we have

$$\text{Con}_e(T, r) \leq \text{Con}_e(U, r) \leq \text{Cl}_e(U, r) \leq \text{Cl}_e(T, r). \tag{61}$$

Hence

$$\text{Con}_e(T, r) \leq \bigwedge_{U \in \mathcal{H}} \text{Cl}_e(U, r). \tag{62}$$

Suppose

$$\text{Con}_e(T, r) \not\leq \bigwedge_{U \in \mathcal{H}} \text{Cl}_e(U, r). \tag{63}$$

Then there exist $x_p \in \text{Pt}(X)$ and $t \in L_0$ such that

$$\text{Con}_{x_p}(T, r) < t < \bigwedge_{U \in \mathcal{H}} \text{Cl}_{x_p}(U, r). \tag{64}$$

Since $\text{Con}_{x_p}(T, r) < t$, there exists $\mu \in L^X$ with T is often in μ' such that

$$\text{Con}_{x_p}(T, r) \leq \mathcal{N}'_{x_p}(\mu, r) < \bigwedge_{U \in \mathcal{H}} \text{Cl}_{x_p}(U, r). \tag{65}$$

Since T is often in μ' , for each $n \in D$ there exists $N(n) \in D$ with $N(n) \geq n$ and $T(N(n)) \in \mu'$. Hence there exists a cofinal selection $N : E \rightarrow D$ such that $U = T \circ N$. Thus U is a subnet of T and U is finally in μ' . It is a contradiction.

(2) From (1), we have

$$\bigvee_{U \in \mathcal{H}} \text{Con}_e(U, r) \leq \text{Cl}_e(T, r). \tag{66}$$

Conversely, let $\text{Cl}_e(T, r) = t > 0$. Then $\mathcal{N}_e(\lambda, r) \leq t'$, for T is finally in λ' . Let $F = \{\mu \mid \mathcal{N}_e(\mu, r) > t'\}$. Define a relation on $E = D \times F$ by

$$(m, \mu_1) \leq (n, \mu_2) \quad \text{iff } m \leq n, \mu_1 \geq \mu_2. \tag{67}$$

Then $(E, \leq)$ is a directed set. If $\mu \in F$, then T is not finally in μ' . For each $(n, \mu) \in E$, there exists $N(n, \mu) \in D$ with $N(n, \mu) \geq n$ such that $T(N(n, \mu)) \notin \lambda'$. So, we can define $N : E \rightarrow D$. For each $n_0 \in D$ and $\mu_0 \in F$, there exists $N(n_0, \mu_0) \in D$ with $N(n_0, \mu_0) \geq n_0$ such that $T(N(n_0, \mu_0)) \notin \mu'_0$. Hence for every $(n, \mu) \geq (n_0, \mu_0)$, since $n \geq n_0$, we have $N(n, \mu) \geq n \geq n_0$. Therefore N is a cofinal selection on T . So $U = T \circ N$ is a fuzzy subnet of T and U is finally to every member of F . If U is often in λ' , then U is not finally of λ ; that is, $\lambda \notin F$. Thus

$$\bigvee_{U \in \mathcal{H}} \text{Con}_e(T, r) = \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid U \text{ is often in } \lambda' \} \geq t. \tag{68}$$

Since t is arbitrary, we complete the proof. □

Theorem 25. Let L be an order dense, let $(X, \mathcal{F})$ be L -fuzzy topological space, and let T be a fuzzy net. If every subnet U of T has a subnet K of U such that $\lim_e(K, r) = t$, then $\lim_e(T, r) = t$.

Proof. Let $\mathcal{H} = \{U \mid U \text{ is a subnet of } T\}$. For each $U \in \mathcal{H}$, since U has a subnet K with $\lim_{\mathcal{F}}(K, r) = t$, by Theorem 18(4), we have

$$\text{Con}_e(U, r) \leq \text{Con}_e(K, r) = \text{Cl}_e(K, r) = t. \tag{69}$$

Hence, by Theorem 24(2),

$$\text{Cl}_e(T, r) = \bigvee_{U \in \mathcal{H}} \text{Con}_e(U, r) \leq t. \tag{70}$$

Conversely, by Theorem 18(2),

$$t = \text{Con}_e(K, r) = \text{Cl}_e(K, r) \leq \text{Cl}_e(U, r). \tag{71}$$

Hence, by Theorem 24(1),

$$t \leq \bigwedge_{U \in \mathcal{H}} \text{Cl}_e(U, r) = \text{Con}_e(T, r). \tag{72}$$

Hence, $\text{Cl}_e(T, r) \leq \text{Con}_e(T, r)$. Since $\text{Con}_e(T, r) \leq \text{Cl}_e(T, r)$ from Theorem 18(2), $\text{Cl}_e(T, r) = \text{Con}_e(T, r)$; that is, $\lim_e(T, r) = t$. □

Example 26. Let $(L = [0, 1], \rightarrow)$ be defined as in Example 21. Let N be a natural number set. Define a fuzzy net $T : N \rightarrow \text{Pt}(X)$ by

$$T(n) = x_{a_n}, \quad a_n = 0.6 + (-1)^n 0.2. \tag{73}$$

Let $e = x_{0.3}$. Since T is often in μ' , for $0 < r \leq 1/2$,

$$\text{Con}_e(T, r) = 1 - \mathcal{N}_e(\mu, r) = 1 - m(x_{0.3}, \mu) = 0. \tag{74}$$

Since T is finally in 1_X , for each $r \in I_0$,

$$\text{Cl}_e(T, r) = 1 - \mathcal{N}_e(0_X, r) = 1 - m(x_{0.3}, 0_X) = 0.3. \tag{75}$$

Thus, since $\text{Con}_e(T, r) \neq \text{Cl}_e(T, r)$ for $0 < r \leq 1/2$, $\lim_e(T, r)$ does not exist.

Since $\text{Con}_e(T, r) = \text{Cl}_e(T, r) = 0.3$ for $1/2 < r \leq 1$, $\lim_e(T, r) = 0.3$.

Theorem 27. Let $(X, \mathcal{T}_1)$ and $(Y, \mathcal{T}_2)$ be L -fuzzy topological spaces. For every fuzzy net T in X , $x_t \in \text{Pt}(X)$, $r \in L_0$, and $\lambda \in L^X$, the following statements are equivalent:

- (1) $f : (X, \mathcal{T}_1) \rightarrow (Y, \mathcal{T}_2)$ is LF -continuous;
- (2) $\mathcal{N}_{f \rightarrow (e)}(\mu, r) \leq \bigvee \{ \mathcal{N}_e(\lambda, r) \mid f \rightarrow (\lambda) \leq \mu \}$;
- (3) $Cl_e(T, r) \leq Cl_{f \rightarrow (e)}(f \circ T, r)$;
- (4) $Con_e(T, r) \leq Con_{f \rightarrow (e)}(f \circ T, r)$;
- (5) $f \rightarrow (C_{\mathcal{T}_1}(\lambda, r)) \leq C_{\mathcal{T}_2}(f \rightarrow (\lambda), r)$;
- (6) $C_{\mathcal{T}_1}(f \leftarrow (\mu), r) \leq f \leftarrow (C_{\mathcal{T}_2}(\mu), r)$;
- (7) $f \leftarrow (I_{\mathcal{T}_2}(\mu, r)) \leq I_{\mathcal{T}_1}(f \leftarrow (\mu), r)$.

Proof. (1) $\Rightarrow$ (2) For any $\rho \in L^Y$ such that $\mathcal{T}_2(\rho) \geq r$ and $\rho \leq \mu$. Since f is LF -continuous, then $\mathcal{T}_1(f \leftarrow (\rho)) \geq \mathcal{T}_2(\rho) \geq r$, and we have by Lemma 3(2)

$$\begin{aligned} S(f \rightarrow (e), \rho) &\leq S(e, f \leftarrow (\rho)) \quad (e = x_t, f \rightarrow (e) = f(x)_t) \\ &= \mathcal{N}_e(f \leftarrow (\rho), r) \quad (\mathcal{T}_1(f \rightarrow (f \leftarrow (\rho))) \geq r) \\ &\leq \bigvee \{ \mathcal{N}_e(\lambda, r) \mid f \rightarrow (\lambda) \leq \mu \} \\ &\quad (f \rightarrow (f \leftarrow (\rho)) \leq \rho \leq \mu). \end{aligned} \quad (76)$$

Thus, $\mathcal{N}_{f \rightarrow (e)}(\mu, r) \leq \bigvee \{ \mathcal{N}_e(\lambda, r) \mid f \rightarrow (\lambda) \leq \mu \}$.

(2) $\Rightarrow$ (3) If $f \rightarrow (\lambda) \leq \mu$ and $f \circ T$ is finally in μ' , there exists $n_0 \in D$ such that, for all $n \geq n_0$, $f(T(n)) \in \mu'$. Let $T(n) = x_t$. Then

$$t \leq \mu'(f(x)) \leq (f(\lambda))'(f(x)) \leq \lambda'(x). \quad (77)$$

It implies $T(n) \in \lambda'$. Therefore, T is finally in λ' . One has

$$\begin{aligned} Cl_e(T, r) &= \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid T \text{ is finally in } \lambda' \} \\ &\leq \bigwedge \{ \mathcal{N}'_e(\lambda, r) \mid \exists \mu, f \rightarrow (\lambda) \leq \mu, \\ &\quad f \circ T \text{ is finally in } \mu' \} \\ &= \bigwedge \{ \bigvee \{ \mathcal{N}_e(\lambda, r) \mid f \rightarrow (\lambda) \leq \mu \}' , \\ &\quad f \circ T \text{ is finally in } \mu' \} \\ &\leq \bigwedge \{ \mathcal{N}'_{f \rightarrow (e)}(\mu, r), f \circ T \text{ is finally in } \mu' \} \\ &= Cl_e(f \circ T, r) \quad (\text{by (2)}). \end{aligned} \quad (78)$$

(3) $\Rightarrow$ (4) Every subnet $U : E \rightarrow \text{Pt}(Y)$ of $f(T)$, and there exists a cofinal selection $N : E \rightarrow D$ such that $U =$

$f(T) \circ N = f \circ (T \circ N)$. Put $K = T \circ N$. Then K is a subnet of T . We can prove it from the following:

$$\begin{aligned} Con_e(T, r) &\leq Con_e(K, r) \quad (\text{by Theorem 18 (5)}) \\ &\leq Cl_e(K, r) \quad (\text{by Theorem 18 (2)}) \\ &\leq Cl_{f \rightarrow (e)}(f \circ K, r) \quad (\text{by (3)}) \\ &= Cl_{f \rightarrow (e)}(f \circ (T \circ N), r) \\ &= Cl_{f \rightarrow (e)}(U, r). \end{aligned} \quad (79)$$

From Theorem 18(2), we have $Con_e(T, r) \leq Con_{f \rightarrow (e)}(f \circ T, r)$.

(4) $\Rightarrow$ (5) From Theorem 5 and Proposition 17(2),

$$S(x_1, C'_{\mathcal{T}_1}(\lambda, r)) = C'_{\mathcal{T}_1}(\lambda, r)(x) = \text{Ad}'_x(\lambda, r). \quad (80)$$

It implies

$$C_{\mathcal{T}_1}(\lambda, r)(x) = \text{Ad}_x(\lambda, r). \quad (81)$$

Thus, we have

$$\begin{aligned} f \rightarrow (C_{\mathcal{T}_1}(\lambda, r))(y) &= \bigvee \{ C_{\mathcal{T}_1}(\lambda, r)(x) \mid f(x) = y \} \\ &= \bigvee \{ \text{Ad}_x(\lambda, r) \mid f(x) = y \} \quad (\text{by (81)}) \\ &= \bigvee_{f(x)=y} \bigvee \{ \text{Con}_x(T, r) \mid T \text{ is fuzzy net in } \lambda \} \\ &\quad (\text{by Proposition 19}) \\ &\leq \bigvee_{f(x)=y} \bigvee \{ \text{Con}_y(f \circ T, r) \mid T \text{ is fuzzy net in } \lambda \} \\ &\quad (\text{by (4)}) \end{aligned} \quad (82)$$

$$\begin{aligned} &= \bigvee \{ \text{Con}_y(f \circ T, r) \mid T \text{ is fuzzy net in } \lambda \} \\ &\leq \bigvee \{ \text{Con}_y(T, r) \mid T \text{ is fuzzy net in } f \rightarrow (\lambda) \} \\ &= \text{Ad}_y(f \rightarrow (\lambda), r) \quad (\text{by Proposition 19}) \\ &= C_{\mathcal{T}_2}(f \rightarrow (\lambda), r)(y) \quad (\text{by (81)}). \end{aligned}$$

(5) $\Rightarrow$ (6) and (6) $\Rightarrow$ (7) are easily proved.

(7) $\Rightarrow$ (1) We will show that $\mathcal{T}_1(f \leftarrow (\mu)) \geq \mathcal{T}_2(\mu)$, for all $\mu \in L^Y$.

Let $\mathcal{T}_2(\mu) = 0$. It is trivial.

Let $\mathcal{T}_2(\mu) = r > 0$. Since $\mathcal{T}_N = \mathcal{T}_2$ from Theorem 12(b), we have for all $y \in Y$,

$$S(y, \mu) = \mathcal{N}_y(\mu, r). \quad (83)$$

It implies, for all $x \in X$,

$$S(f(x), \mu) = S(x, f \leftarrow (\mu)) = \mathcal{N}_{f(x)}(\mu, r). \quad (84)$$

Since $f^{\leftarrow}(I_{\mathcal{T}_2}(\mu, r)) = f^{\leftarrow}(\mu)$,

$$\begin{aligned} S(x, f^{-1}(\mu)) &= S(x, f^{\leftarrow}(I_{\mathcal{T}_2}(\mu, r))) \\ &(\text{since } f^{\leftarrow}(I_{\mathcal{T}_2}(\mu, r)) \leq I_{\mathcal{T}_1}(f^{\leftarrow}(\mu), r)) \\ &\leq S(x, I_{\mathcal{T}_1}(f^{\leftarrow}(\mu), r)) \\ &= \mathcal{N}_x(f^{\leftarrow}(\mu), r) \quad (\text{by Proposition 17(1)}). \end{aligned} \quad (85)$$

Thus, by Theorem 11(2), we have

$$S(x, f^{\leftarrow}(\mu)) = \mathcal{N}_x(f^{\leftarrow}(\mu), r). \quad (86)$$

Hence, $\mathcal{T}_1(f^{\leftarrow}(\mu)) \geq r$. $\square$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

The authors would like to thank the anonymous referee for his comments, which helped us to improve the final version of this paper.

References

- [1] A. P. Šostak, "On a fuzzy topological structure," *Rendiconti del Circolo Matematico di Palermo II*, vol. 11, pp. 89–103, 1985.
- [2] A. P. Šostak, "On the neighborhood structure of a fuzzy topological spaces," *Zbornik Radova*, no. 4, pp. 7–14, 1990.
- [3] A. P. Šostak, "Basic structures of fuzzy topology," *Journal of Mathematical Sciences*, vol. 78, no. 6, pp. 662–701, 1996.
- [4] C. L. Chang, "Fuzzy topological spaces," *Journal of Mathematical Analysis and Applications*, vol. 24, pp. 182–190, 1968.
- [5] M. Demirci, "Neighborhood structures of smooth topological spaces," *Fuzzy Sets and Systems*, vol. 92, no. 1, pp. 123–128, 1997.
- [6] R. N. Hazra, S. K. Samanta, and K. C. Chattopadhyay, "Fuzzy topology redefined," *Fuzzy Sets and Systems*, vol. 45, no. 1, pp. 79–82, 1992.
- [7] K. C. Chattopadhyay, R. N. Hazra, and S. K. Samanta, "Gradation of openness: fuzzy topology," *Fuzzy Sets and Systems*, vol. 49, no. 2, pp. 237–242, 1992.
- [8] U. Höhle, *Many Valued Topology and Its Application*, Kluwer Academic, Boston, Mass, USA, 2001.
- [9] U. Höhle and S. E. Rodabaugh, *Mathematics of Fuzzy Sets, Logic, Topology and Measure Theory*, vol. 3 of *The Handbooks of Fuzzy Sets Series*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1999.
- [10] U. Höhle and A. P. Šostak, "Axiomatic foundations of fixed-basis fuzzy topology," in *Mathematics of Fuzzy Sets*, vol. 3 of *The Handbooks of Fuzzy Sets Series*, chapter 3, pp. 123–272, Kluwer Academic, Boston, Mass, USA, 1999.
- [11] P. Pao-Ming and L. Ying-Ming, "Fuzzy topology. I. Neighborhood structure of a fuzzy point and Moore-Smith convergence," *Journal of Mathematical Analysis and Applications*, vol. 76, no. 2, pp. 571–599, 1980.
- [12] Y.-M. Liu and M.-K. Luo, *Fuzzy Topology*, World Scientific, Singapore, 1997.
- [13] M. S. Ying, "On the method of neighborhood systems in fuzzy topology," *Fuzzy Sets and Systems*, vol. 68, no. 2, pp. 227–238, 1994.
- [14] S. L. Chen and J. S. Cheng, "On convergence of nets of L -fuzzy sets," *Journal of Fuzzy Mathematics*, vol. 2, no. 3, pp. 517–524, 1994.
- [15] S. L. Chen and J. S. Cheng, " θ -convergence of nets of L -fuzzy sets and its applications," *Fuzzy Sets and Systems*, vol. 86, pp. 235–240, 1997.
- [16] J. Fang, "Relationships between L -ordered convergence structures and strong L -topologies," *Fuzzy Sets and Systems*, vol. 161, no. 22, pp. 2923–2944, 2010.
- [17] D. N. Georgiou and B. K. Papadopoulos, "Convergences in fuzzy topological spaces," *Fuzzy Sets and Systems*, vol. 101, no. 3, pp. 495–504, 1999.
- [18] W. Yao, "On many-valued stratified L -fuzzy convergence spaces," *Fuzzy Sets and Systems*, vol. 159, no. 19, pp. 2503–2519, 2008.
- [19] A. A. Ramadan, "Smooth topological spaces," *Fuzzy Sets and Systems*, vol. 48, no. 3, pp. 371–375, 1992.
- [20] G. Gierz and K. H. Hofmann, *Continuous Lattices and 305 Domains*, Cambridge University Press, Cambridge, UK, 2003.
- [21] P. Hájek, *Metamathematics of Fuzzy Logic*, vol. 4 of *Trends in Logic*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1998.
- [22] E. Turunen, "Algebraic structures in fuzzy logic," *Fuzzy Sets and Systems*, vol. 52, no. 2, pp. 181–188, 1992.
- [23] E. Turunen, *Mathematics Behind Fuzzy Logic*, Springer, New York, NY, USA, 1999.
- [24] F. Jinming, "Stratified L -ordered convergence structures," *Fuzzy Sets and Systems*, vol. 161, no. 16, pp. 2130–2149, 2010.
- [25] H. Lai and D. Zhang, "Fuzzy preorder and fuzzy topology," *Fuzzy Sets and Systems*, vol. 157, no. 14, pp. 1865–1885, 2006.