Research Article

A Simplified Proof of Uncertainty Principle for Quaternion Linear Canonical Transform

Mawardi Bahri¹ and Ryuichi Ashino²

¹Department of Mathematics, Hasanuddin University, Makassar 90245, Indonesia

²Division of Mathematical Sciences, Osaka Kyoiku University, Osaka 582-8582, Japan

Correspondence should be addressed to Ryuichi Ashino; ashino@cc.osaka-kyoiku.ac.jp

Received 20 August 2015; Accepted 9 December 2015

Academic Editor: Jozef Banas

Copyright © 2016 M. Bahri and R. Ashino. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We provide a short and simple proof of an uncertainty principle associated with the quaternion linear canonical transform (QLCT) by considering the fundamental relationship between the QLCT and the quaternion Fourier transform (QFT). We show how this relation allows us to derive the inverse transform and Parseval and Plancherel formulas associated with the QLCT. Some other properties of the QLCT are also studied.

1. Introduction

It is well-known that the traditional linear canonical transform (LCT) plays an important role in many fields of optics and signal processing. It can be regarded as a generalization of many mathematical transforms such as the Fourier transform, Laplace transform, the fractional Fourier transform, and the Fresnel transform. Many fundamental properties of this extended transform are already known, including shift, modulation, convolution, and correlation and uncertainty principle, for example, in [1–6].

Recently, there are so many studies in the literature that are concerned with the generalization of the LCT within the context of quaternion algebra, which is the so-called quaternion linear canonical transform (QLCT) (see, e.g., [7–10]). They also established some important properties of the QLCT such as inversion formula and the uncertainty principle. An application of the QLCT to study of generalized swept-frequency filters was presented in [11]. In this paper, we will focus on the two-dimensional case and provide a new proof of uncertainty principle associated with the QLCT, the ones proposed in [8], the proof of which is much simpler using the component-wise and directional uncertainty principles for the QFT [12, 13]. Therefore, before proving this main result, we first derive the fundamental relationship between the QLCT and QFT. Using the relation, we obtain

useful properties of the QLCT such as inverse transform and Parseval formula associated with the QLCT.

The quaternion algebra over $\mathbb{R}$, denoted by $\mathbb{H}$, is an associative noncommutative four-dimensional algebra:

$$\mathbb{H} = \{q = q_0 + \mathbf{i}q_1 + \mathbf{j}q_2 + \mathbf{k}q_3; q_0, q_1, q_2, q_3 \in \mathbb{R}\}, \quad (1)$$

which obeys the following multiplication rules:

$$\begin{aligned} \mathbf{i}\mathbf{j} &= -\mathbf{j}\mathbf{i} = \mathbf{k}, \\ \mathbf{j}\mathbf{k} &= -\mathbf{k}\mathbf{j} = \mathbf{i}, \\ \mathbf{k}\mathbf{i} &= -\mathbf{i}\mathbf{k} = \mathbf{j}, \\ \mathbf{i}^2 &= \mathbf{j}^2 = \mathbf{k}^2 = \mathbf{i}\mathbf{j}\mathbf{k} = -1. \end{aligned} \quad (2)$$

For a quaternion $q = q_0 + \mathbf{i}q_1 + \mathbf{j}q_2 + \mathbf{k}q_3 \in \mathbb{H}$, q_0 is called the *scalar* part of q denoted by $\text{Sc}(q)$ and $\mathbf{i}q_1 + \mathbf{j}q_2 + \mathbf{k}q_3$ is called the *vector* (or *pure*) part of q . The vector part of q is conventionally denoted by $\mathbf{q}$. Let $p, q \in \mathbb{H}$ and $\mathbf{p}, \mathbf{q}$ be their vector parts, respectively. Equation (2) yields the quaternionic multiplication qp as

$$qp = q_0p_0 - \mathbf{q} \cdot \mathbf{p} + q_0\mathbf{p} + p_0\mathbf{q} + \mathbf{q} \times \mathbf{p}, \quad (3)$$

where $\mathbf{q} \cdot \mathbf{p} = (q_1p_1 + q_2p_2 + q_3p_3)$ and $\mathbf{q} \times \mathbf{p} = \mathbf{i}(q_2p_3 - q_3p_2) + \mathbf{j}(q_3p_1 - q_1p_3) + \mathbf{k}(q_1p_2 - q_2p_1)$.

The quaternion conjugate of q , given by

$$\bar{q} = q_0 - \mathbf{i}q_1 - \mathbf{j}q_2 - \mathbf{k}q_3, \quad q_0, q_1, q_2, q_3 \in \mathbb{R}, \quad (4)$$

is an anti-involution; that is,

$$\bar{q}\bar{p} = \overline{p\bar{q}}. \quad (5)$$

From (4) we obtain the norm or modulus of $q \in \mathbb{H}$ defined as

$$|q| = \sqrt{q\bar{q}} = \sqrt{q_0^2 + q_1^2 + q_2^2 + q_3^2}. \quad (6)$$

It is not difficult to see that

$$|qp| = |q||p|, \quad \forall p, q \in \mathbb{H}. \quad (7)$$

Furthermore, it is easily seen that

$$|pqr| = |rqp|, \quad \forall p, q, r \in \mathbb{H}. \quad (8)$$

Using conjugate (4) and the modulus of q , we can define the inverse of $q \in \mathbb{H} \setminus \{0\}$ as

$$q^{-1} = \frac{\bar{q}}{|q|^2}, \quad (9)$$

which shows that $\mathbb{H}$ is a normed division algebra.

It is convenient to introduce an inner product for quaternion-valued (in the rest of the paper, we will always consider quaternion function) functions $f, g : \mathbb{R}^2 \rightarrow \mathbb{H}$ as

$$(f, g) = \int_{\mathbb{R}^2} f(\mathbf{x}) \overline{g(\mathbf{x})} d\mathbf{x}, \quad d\mathbf{x} = dx_1 dx_2, \quad (10)$$

with symmetric real scalar part

$$\langle f, g \rangle = \frac{1}{2} [(f, g) + (g, f)] = \text{Sc} \int_{\mathbb{R}^2} f(\mathbf{x}) \overline{g(\mathbf{x})} d\mathbf{x}. \quad (11)$$

In particular, for $f = g$, we obtain the $L^2(\mathbb{R}^2; \mathbb{H})$ -norm:

$$\|f\| = \sqrt{\langle f, f \rangle} = \left(\int_{\mathbb{R}^2} |f(\mathbf{x})|^2 d^2 \mathbf{x} \right)^{1/2}. \quad (12)$$

2. Quaternion Linear Canonical Transform

In this section we begin by defining the two-sided QFT (for simplicity of notation we write the QFT instead of the two-sided QFT in the next section). We discuss some properties, which will be used to prove the uncertainty principle.

Definition 1. The QFT of $f \in L^1(\mathbb{R}^2; \mathbb{H})$ is the transform $\mathcal{F}_q\{f\} : \mathbb{R}^2 \rightarrow \mathbb{H}$ given by the integral

$$\mathcal{F}_q\{f\}(\omega) = \frac{1}{\sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{-i\omega_1 x_1} f(\mathbf{x}) e^{-j\omega_2 x_2} d\mathbf{x}, \quad (13)$$

where $\mathbf{x} = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2$, $\omega = \omega_1 \mathbf{e}_1 + \omega_2 \mathbf{e}_2$, and the quaternion exponential product $e^{-i\omega_1 x_1} e^{-j\omega_2 x_2}$ is the quaternion Fourier kernel. Here $\mathcal{F}_q$ is called the quaternion Fourier transform operator.

Definition 2. If $f \in L^1(\mathbb{R}^2; \mathbb{H})$ and $\mathcal{F}_q\{f\} \in L^1(\mathbb{R}^2; \mathbb{H})$, then the inverse transform of the QFT is given by

$$\begin{aligned} f(\mathbf{x}) &= \mathcal{F}_q^{-1} [\mathcal{F}_q\{f\}](\mathbf{x}) \\ &= \frac{1}{\sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{i\omega_1 x_1} \mathcal{F}_q\{f\}(\omega) e^{j\omega_2 x_2} d\omega, \end{aligned} \quad (14)$$

where $\mathcal{F}_q^{-1}$ is called the inverse QFT operator.

An important property of the QFT is stated in the following lemma, which is needed to prove Parseval formula of the QLCT. For more details of the QFT, see [12–16].

Lemma 3 (QFT Parseval). *The quaternion product of $f, g \in L^1(\mathbb{R}^2; \mathbb{H}) \cap L^2(\mathbb{R}^2; \mathbb{H})$ and its QFT are related by*

$$\langle f, g \rangle_{L^2(\mathbb{R}^2; \mathbb{H})} = \langle \mathcal{F}_q\{f\}, \mathcal{F}_q\{g\} \rangle_{L^2(\mathbb{R}^2; \mathbb{H})}. \quad (15)$$

In particular, with $f = g$, we get the quaternion version of the Plancherel formula; that is,

$$\|f\|_{L^2(\mathbb{R}^2; \mathbb{H})}^2 = \|\mathcal{F}_q\{f\}\|_{L^2(\mathbb{R}^2; \mathbb{H})}^2. \quad (16)$$

Based on the definition of the QFT mentioned above, we consider the two-sided QLCT which is defined as follows.

Definition 4 (QLCT). Let $A_1 = (a_1, b_1, c_1, d_1)$ and $A_2 = (a_2, b_2, c_2, d_2)$ be two matrix parameters satisfying $\det(A_s) = a_s d_s - b_s c_s = 1$, $s = 1, 2$. The QLCT of a quaternion signal $f \in L^1(\mathbb{R}^2; \mathbb{H})$ is defined by

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}}\{f\}(\omega) &= \begin{cases} \int_{\mathbb{R}^2} K_{A_1}(x_1, \omega_1) f(\mathbf{x}) K_{A_2}(x_2, \omega_2) d\mathbf{x}, & b_1 b_2 \neq 0 \\ \sqrt{d_1} e^{i(c_1 d_1/2)\omega_1^2} f(d_1 \omega_1, x_2) K_{A_2}(x_2, \omega_2), & b_1 = 0, b_2 \neq 0 \\ \sqrt{d_2} e^{j(c_2 d_2/2)\omega_2^2} f(x_1, d_2 \omega_2) K_{A_1}(x_1, \omega_1), & b_1 \neq 0, b_2 = 0 \\ \sqrt{d_1 d_2} e^{i(c_1 d_1/2)\omega_1^2} f(d_1 \omega_1, d_2 \omega_2) e^{j(c_2 d_2/2)\omega_2^2}, & b_1 = b_2 = 0, \end{cases} \end{aligned} \quad (17)$$

where the kernel functions of the QLCT are given by, respectively,

$$\begin{aligned} K_{A_1}(x_1, \omega_1) &= \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1 \omega_1 + (d_1/b_1)\omega_1^2 - (\pi/2))}, \\ & \quad b_1 \neq 0, \end{aligned} \quad (18)$$

$$\begin{aligned} K_{A_2}(x_2, \omega_2) &= \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2 \omega_2 + (d_2/b_2)\omega_2^2 - (\pi/2))}, \\ & \quad b_2 \neq 0. \end{aligned} \quad (19)$$

From the definition of the QLCT, we can see easily that when $b_1 b_2 = 0$ and $b_1 = b_2 = 0$, the QLCT of a signal

is essentially a quaternion chirp multiplication. Therefore, in this work we always assume that $b_1 b_2 \neq 0$. As a special case, when $A_1 = A_2 = (a_i, b_i, c_i, d_i) = (0, 1, -1, 0)$ for $i = 1, 2$, LCT definition (17) reduces to the QFT definition. That is,

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega) &= \int_{\mathbb{R}^2} \frac{e^{-i(\pi/4)}}{\sqrt{2\pi}} f(\mathbf{x}) e^{-i\omega_1 x_1} \frac{e^{-j(\pi/4)}}{\sqrt{2\pi}} e^{-j\omega_2 x_2} d\mathbf{x} \\ &= e^{-i(\pi/4)} \mathcal{F}_q \{f\}(\omega) e^{-j(\pi/4)}, \end{aligned} \quad (20)$$

where $\mathcal{F}_q \{f\}$ is the QFT of f given by (13).

We need the following important result (compare to [17, 18]), which will be useful in proving Theorem 15.

Theorem 5. *The QLCT of a quaternion function $f \in L^1(\mathbb{R}^2; \mathbb{H})$ with matrix parameters $A_1 = (a_1, b_1, c_1, d_1)$ and $A_2 = (a_2, b_2, c_2, d_2)$ can be reduced to the QFT*

$$\mathcal{F}_q \{g_f\}(\omega) = \frac{1}{\sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{-i\mathbf{x}_1 \omega_1} g_f(\mathbf{x}) e^{-j\mathbf{x}_2 \omega_2} d\mathbf{x}, \quad (21)$$

where

$$\begin{aligned} \mathcal{F}_q \{g_f\}(\omega) &= \widetilde{\mathcal{F}}(\mathbf{b}\omega), \\ g_f(\mathbf{x}) &= \frac{e^{-i(\pi/4)}}{\sqrt{b_1}} \tilde{f}(\mathbf{x}) \frac{e^{-j(\pi/4)}}{\sqrt{b_2}}, \\ \tilde{f}(\mathbf{x}) &= e^{i(a_1/2b_1)x_1^2} f(\mathbf{x}) e^{j(a_2/2b_2)x_2^2}, \end{aligned} \quad (22)$$

with

$$\widetilde{\mathcal{F}}(\omega) = \frac{1}{\sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{-i\mathbf{x}_1(\omega_1/b_1)} g_f(\mathbf{x}) e^{-j\mathbf{x}_2(\omega_2/b_2)} d\mathbf{x}, \quad (23)$$

$$\widetilde{\mathcal{F}}(\omega) = e^{-i(d_1/2b_1)\omega_1^2} L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega) e^{-j(d_2/2b_2)\omega_2^2}.$$

Proof. Simple computations using Definition 4 show that

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega) &= \frac{1}{\sqrt{2\pi b_1}} \\ &\cdot \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega_1 + (d_1/b_1)\omega_1^2 - (\pi/2))} f(\mathbf{x}) \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega_2 + (d_2/b_2)\omega_2^2 - (\pi/2))} d\mathbf{x} \\ &= \frac{e^{-i(\pi/4)}}{\sqrt{2\pi b_1}} e^{i(d_1/2b_1)\omega_1^2} \int_{\mathbb{R}^2} e^{-i\mathbf{x}_1(\omega_1/b_1)} \left(e^{i(a_1/2b_1)x_1^2} f(\mathbf{x}) \right. \\ &\cdot e^{j(a_2/2b_2)x_2^2} \left. \right) \frac{e^{-j(\pi/4)}}{\sqrt{2\pi b_2}} e^{-j\mathbf{x}_2(\omega_2/b_2)} e^{j(d_2/2b_2)\omega_2^2} d\mathbf{x} \\ &= \frac{e^{-i(\pi/4)}}{\sqrt{2\pi b_1}} e^{i(d_1/2b_1)\omega_1^2} \int_{\mathbb{R}^2} e^{-i\mathbf{x}_1(\omega_1/b_1)} \tilde{f}(\mathbf{x}) \\ &\cdot e^{-j\mathbf{x}_2(\omega_2/b_2)} d\mathbf{x} \frac{e^{-i(\pi/4)}}{\sqrt{2\pi b_2}} e^{-j(d_2/2b_2)\omega_2^2}. \end{aligned} \quad (24)$$

Then, multiplying both sides of (24) by $e^{-i(d_1/2b_1)\omega_1^2} e^{-j(d_2/2b_2)\omega_2^2}$ results in

$$\begin{aligned} e^{-i(d_1/2b_1)\omega_1^2} L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega) e^{-j(d_2/2b_2)\omega_2^2} &= \frac{e^{-i(\pi/4)}}{\sqrt{2\pi b_1}} \\ &\cdot \int_{\mathbb{R}^2} e^{-i\mathbf{x}_1(\omega_1/b_1)} \tilde{f}(\mathbf{x}) e^{-j\mathbf{x}_2(\omega_2/b_2)} d\mathbf{x} \frac{e^{-j(\pi/4)}}{\sqrt{2\pi b_2}} = \frac{1}{\sqrt{2\pi}} \\ &\cdot \int_{\mathbb{R}^2} e^{-i\mathbf{x}_1(\omega_1/b_1)} \frac{e^{-i(\pi/4)}}{\sqrt{b_1}} \tilde{f}(\mathbf{x}) \frac{e^{-i(\pi/4)}}{\sqrt{b_2}} e^{-j\mathbf{x}_2(\omega_2/b_2)} d\mathbf{x} \\ &\cdot \frac{1}{\sqrt{2\pi}} = \frac{1}{\sqrt{(2\pi)^2}} \\ &\cdot \int_{\mathbb{R}^2} e^{-i\mathbf{x}_1(\omega_1/b_1)} g_f(\mathbf{x}) e^{-j\mathbf{x}_2(\omega_2/b_2)} d\mathbf{x} = \mathcal{F}_q \{g_f\} \\ &\cdot \left(\frac{\omega}{\mathbf{b}} \right). \end{aligned} \quad (25)$$

This is the desired result. $\square$

Theorem 6. *If $f \in L^1(\mathbb{R}^2; \mathbb{H})$ and $L_{A_1, A_2}^{\mathbb{H}} \{f\} \in L^1(\mathbb{R}^2; \mathbb{H})$, then the inverse transform of the QLCT can be derived from that of the QFT.*

Proof. Indeed, we have

$$\begin{aligned} g_f(\mathbf{x}) &= \frac{1}{\sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{i\mathbf{x}_1 \omega_1} \mathcal{F}_q \{g_f\}(\omega) e^{j\mathbf{x}_2 \omega_2} d\omega \\ &= \frac{1}{\sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{i\mathbf{x}_1 \omega_1} \widetilde{\mathcal{F}}(\mathbf{b}\omega) e^{j\mathbf{x}_2 \omega_2} d\omega \\ &= \frac{1}{b_1 b_2 \sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{i\mathbf{x}_1(\omega_1/b_1)} \widetilde{\mathcal{F}}(\omega) e^{j\mathbf{x}_2(\omega_2/b_2)} d\omega \\ &= \frac{1}{b_1 b_2 \sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{i\mathbf{x}_1(\omega_1/b_1)} e^{-i(d_1/2b_1)\omega_1^2} L_{A_1, A_2}^{\mathbb{H}} \{f\} \\ &\cdot (\omega) e^{-j(d_2/2b_2)\omega_2^2} e^{j\mathbf{x}_2(\omega_2/b_2)} d\omega. \end{aligned} \quad (26)$$

It means that

$$\begin{aligned} \frac{e^{-i(\pi/4)}}{\sqrt{b_1}} e^{i(a_1/2b_1)x_1^2} f(\mathbf{x}) e^{j(a_2/2b_2)x_2^2} \frac{e^{-j(\pi/4)}}{\sqrt{b_2}} \\ &= \frac{1}{b_1 b_2 \sqrt{(2\pi)^2}} \int_{\mathbb{R}^2} e^{i\mathbf{x}_1(\omega_1/b_1)} e^{-i(d_1/2b_1)\omega_1^2} L_{A_1, A_2}^{\mathbb{H}} \{f\} \\ &\cdot (\omega) e^{-j(d_2/2b_2)\omega_2^2} e^{j\mathbf{x}_2(\omega_2/b_2)} d\omega. \end{aligned} \quad (27)$$

Or, equivalently,

$$\begin{aligned}
 f(\mathbf{x}) &= \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{-i(a_1/2b_1)x_1^2} e^{ix_1(\omega_1/b_1)} e^{-i(d_1/2b_1)\omega_1^2} e^{i(\pi/4)} L_{A_1, A_2}^{\mathbb{H}} \{f\} \\
 &\cdot (\omega) \frac{1}{\sqrt{2\pi b_2}} e^{-j(a_2/2b_2)x_2^2} e^{jx_2(\omega_2/b_2)} e^{-j(d_2/2b_2)\omega_2^2} e^{-i(\pi/4)} d\omega \\
 &= \int_{\mathbb{R}^2} \overline{K_{A_1}(x_1, \omega_1)} L_{A_1, A_2}^{\mathbb{H}} \{f\} (\omega) \overline{K_{A_2}(x_2, \omega_2)} d\omega,
 \end{aligned} \quad (28)$$

which is inverse transform of the QLCT. This proves the theorem. $\square$

In following we give an alternative proof of Parseval formula for the QLCT (cf. [8]).

Theorem 7 (QLCT Parseval). *Two quaternion functions $f, h \in L^1(\mathbb{R}^2; \mathbb{H}) \cap L^2(\mathbb{R}^2; \mathbb{H})$ are related to their QLCT via the Parseval formula, given as*

$$\langle f, h \rangle_{L^2(\mathbb{R}^2; \mathbb{H})} = \langle L_{A_1, A_2}^{\mathbb{H}} \{f\}, L_{A_1, A_2}^{\mathbb{H}} \{h\} \rangle_{L^2(\mathbb{R}^2; \mathbb{H})}. \quad (29)$$

For $f = h$, one has

$$\|f\|_{L^2(\mathbb{R}^2; \mathbb{H})}^2 = \|L_{A_1, A_2}^{\mathbb{H}} \{f\}\|_{L^2(\mathbb{R}^2; \mathbb{H})}^2. \quad (30)$$

Proof. From Parseval formula (15), it follows that

$$\begin{aligned}
 \langle g_f, g_h \rangle &= \langle \mathcal{F}_q \{g_f\}, \mathcal{F}_q \{g_h\} \rangle \\
 &= \text{Sc} \int_{\mathbb{R}^2} \mathcal{F}_q \{g_f\} (\omega) \overline{\mathcal{F}_q \{g_h\} (\omega)} d\omega \\
 &= \frac{1}{|b_1 b_2|} \text{Sc} \int_{\mathbb{R}^2} \mathcal{F}_q \{g_f\} \left(\frac{\omega}{b} \right) \overline{\mathcal{F}_q \{g_h\} \left(\frac{\omega}{b} \right)} d\omega.
 \end{aligned} \quad (31)$$

Applying the cyclic multiplication symmetry, we get

$$\begin{aligned}
 \langle g_f, g_h \rangle &= \frac{1}{|b_1 b_2|} \text{Sc} \int_{\mathbb{R}^2} e^{-i(d_1/2b_1)\omega_1^2} L_{A_1, A_2}^{\mathbb{H}} \{f\} (\omega) \\
 &\cdot e^{-j(d_2/2b_2)\omega_2^2} e^{-i(d_1/2b_1)\omega_1^2} \overline{L_{A_1, A_2}^{\mathbb{H}} \{h\} (\omega)} e^{-j(d_2/2b_2)\omega_2^2} d\omega \\
 &= \frac{1}{|b_1 b_2|} \text{Sc} \int_{\mathbb{R}^2} e^{-i(d_1/2b_1)\omega_1^2} L_{A_1, A_2}^{\mathbb{H}} \{f\} (\omega) \\
 &\cdot e^{-j(d_2/2b_2)\omega_2^2} e^{-j(d_2/2b_2)\omega_1^2} \overline{L_{A_1, A_2}^{\mathbb{H}} \{h\} (\omega)} e^{-i(d_1/2b_1)\omega_1^2} d\omega \\
 &= \frac{1}{|b_1 b_2|} \text{Sc} \int_{\mathbb{R}^2} L_{A_1, A_2}^{\mathbb{H}} \{f\} (\omega) \overline{L_{A_1, A_2}^{\mathbb{H}} \{h\} (\omega)} d\omega.
 \end{aligned} \quad (32)$$

On the other hand,

$$\begin{aligned}
 \langle g_f, g_h \rangle &= \text{Sc} \int_{\mathbb{R}^2} g_f(\mathbf{x}) \overline{g_h(\mathbf{x})} d\mathbf{x} \\
 &= \text{Sc} \int_{\mathbb{R}^2} \frac{e^{-i(\pi/4)}}{\sqrt{b_1}} \tilde{f}(\mathbf{x}) \\
 &\cdot \frac{e^{-j(\pi/4)}}{\sqrt{b_2}} \frac{e^{-i(\pi/4)}}{\sqrt{b_1}} \tilde{h}(\mathbf{x}) \frac{e^{-j(\pi/4)}}{\sqrt{b_2}} d\mathbf{x} \\
 &= \frac{1}{|b_1 b_2|} \text{Sc} \int_{\mathbb{R}^2} \tilde{f}(\mathbf{x}) \overline{\tilde{h}(\mathbf{x})} d\mathbf{x} \\
 &= \frac{1}{|b_1 b_2|} \text{Sc} \int_{\mathbb{R}^2} e^{i(a_1/2b_1)x_1^2} f(\mathbf{x}) \\
 &\cdot e^{j(a_2/2b_2)x_2^2} e^{i(a_1/2b_1)x_1^2} \overline{h(\mathbf{x})} e^{j(a_2/2b_2)x_2^2} d\mathbf{x} \\
 &= \frac{1}{|b_1 b_2|} \text{Sc} \int_{\mathbb{R}^2} f(\mathbf{x}) \overline{h(\mathbf{x})} d\mathbf{x}.
 \end{aligned} \quad (33)$$

The proof is complete. $\square$

It is interesting to describe the relationship between the QLCT and QFT as shown in the following example.

Example 8. Let us now compute the QLCT of the two-dimensional Gaussian function $f(\mathbf{x}) = e^{-(k_1 x_1^2 + k_2 x_2^2)}$ with $k_1, k_2 > 0$.

From the definition of QLCT (17), we easily obtain

$$\begin{aligned}
 L_{A_1, A_2}^{\mathbb{H}} \{f\} (\omega) &= \frac{1}{\sqrt{2\pi b_1}} \\
 &\cdot \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega + (d_1/b_1)\omega^2 - (\pi/2))} f(\mathbf{x}) \\
 &\cdot e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega + (d_2/b_2)\omega^2 - (\pi/2))} \frac{1}{\sqrt{2\pi b_2}} d\mathbf{x} \\
 &= \frac{1}{\sqrt{2\pi b_1}} \\
 &\cdot \int_{\mathbb{R}} e^{-k_1 x_1^2} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega + (d_1/b_1)\omega^2 - (\pi/2))} dx_1 \\
 &\cdot \frac{1}{\sqrt{2\pi b_2}} \\
 &\cdot \int_{\mathbb{R}} e^{-k_2 x_2^2} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega + (d_2/b_2)\omega^2 - (\pi/2))} dx_2 \\
 &= \frac{1}{\sqrt{2\pi b_1}} \\
 &\cdot e^{i(1/2)(d_1/b_1)\omega^2} \int_{\mathbb{R}} e^{-(1/2b_1)(2k_1 b_1 - ia_1)x_1^2 - i(\omega/b_1)x_1} e^{-i(\pi/2)} dx_1 \\
 &\cdot \frac{1}{\sqrt{2\pi b_2}} \\
 &\cdot e^{j(1/2)(d_2/b_2)\omega^2} \int_{\mathbb{R}} e^{-(1/2b_2)(2k_2 b_2 - ja_2)x_2^2 - j(\omega/b_2)x_2} e^{-j(\pi/2)} dx_2.
 \end{aligned} \quad (34)$$

Using the QFT of the Gaussian function,

$$\begin{aligned}\mathcal{F}_q\{f\}(\omega) &= \int_{\mathbb{R}^2} e^{-i\omega_1 x_1} e^{-(k_1 x_1^2 + k_2 x_2^2)} e^{-j\omega_2 x_2} d\mathbf{x} \\ &= \frac{\pi}{\sqrt{k_1 k_2}} e^{-(\omega_1^2/4k_1 + \omega_2^2/4k_2)}.\end{aligned}\quad (35)$$

We immediately obtain

$$\begin{aligned}L_{A_1, A_2}^{\mathbb{H}}\{f\}(\omega) &= \frac{1}{\sqrt{2\pi b_1 i}} \\ &\cdot e^{i(1/2)(d_1/b_1)\omega_1^2} \sqrt{\frac{2\pi b_1}{(2k_1 b_1 - i a_1)}} e^{-\omega_1^2/2b_1(2k_1 b_1 - i a_1)} \\ &\cdot \frac{1}{\sqrt{2\pi b_1}} \\ &\cdot e^{j(1/2)(d_2/b_2)\omega_2^2} \sqrt{\frac{2\pi b_2}{(2k_2 b_2 - j a_2)}} e^{-\omega_2^2/2b_2(2k_2 b_2 - j a_2)} \\ &= \frac{1}{\sqrt{a_1 + 2k_1 b_1}} e^{(\omega_1^2/2)((c_1 + 2k_1 i d_1)/(2k_1 b_1 - i a_1))} \\ &\cdot \frac{1}{\sqrt{a_2 + 2k_2 b_2}} e^{(\omega_2^2/2)((c_2 + 2k_2 j d_2)/(2k_2 b_2 - j a_2))}.\end{aligned}\quad (36)$$

3. Properties of the QLCT

In this section we present useful properties of the QLCT in detail. We see that the results are generalizations of the properties of the LCT [5, 19]. In [9], the authors derived the asymptotic behavior of the QLCT. In the following, we shall provide a different proof of the results using the QLCT kernel properties.

3.1. Asymptotic Behavior of the QLCT. Like the classical Fourier transform, the Riemann-Lebesgue lemma is also valid for the QLCT, expressed as follows.

Theorem 9 (Riemann-Lebesgue lemma). *Suppose that $f \in L^1(\mathbb{R}^2; \mathbb{H})$. Then*

$$\begin{aligned}\lim_{|\omega_1| \rightarrow \infty} |L_{A_1, A_2}^{\mathbb{H}}\{f\}(\omega)| &= 0, \\ \lim_{|\omega_2| \rightarrow \infty} |L_{A_1, A_2}^{\mathbb{H}}\{f\}(\omega)| &= 0.\end{aligned}\quad (37)$$

Proof. It is not difficult to see that

$$\begin{aligned}e^{-i(\omega_1 x_1/b_1)} &= -e^{-i(\omega_1/b_1)(x_1 + b_1 \pi/\omega_1)}, \\ e^{-j(\omega_2 x_2/b_2)} &= -e^{-j(\omega_2/b_2)(x_2 + b_2 \pi/\omega_2)}.\end{aligned}\quad (38)$$

Now applying (38) gives

$$\begin{aligned}L_{A_1, A_2}^{\mathbb{H}}\{f\}(\omega) &= \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f(\mathbf{x}) \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{x} \\ &= - \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)(x_1 + b_1 \pi/\omega_1)\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f(\mathbf{x}) \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{x}.\end{aligned}\quad (39)$$

Therefore, by making the change of variable $x_1 + b_1 \pi/\omega_1 = t_1$ in the above identity, we immediately obtain

$$\begin{aligned}L_{A_1, A_2}^{\mathbb{H}}\{f\}(\omega) &= - \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)(t_1 - b_1 \pi/\omega_1)^2 - (2/b_1)t_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f\left(t_1 - \frac{b_1 \pi}{\omega_1}, t_2\right) \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)t_2^2 - (2/b_2)t_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} dt = \frac{1}{2} \left[\int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)t_1^2 - (2/b_1)t_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f(\mathbf{t}) \right. \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)t_2^2 - (2/b_2)t_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{x} \\ &\left. - \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)(t_1 - b_1 \pi/\omega_1)^2 - (2/b_1)t_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f\left(t_1 - \frac{b_1 \pi}{\omega_1}, t_2\right) \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)t_2^2 - (2/b_2)t_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} dt \right]\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \left[\int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)t_1^2 - (2/b_1)t_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} \left(f(\mathbf{t}) - e^{i(a_1/2b_1)(-2t_1b_1\pi/\omega_1 + (b_1\pi/\omega_1)^2)} f\left(t_1 - \frac{b_1\pi}{\omega_1}, t_2\right) \right) \right. \\
&\quad \left. \cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)t_2^2 - (2/b_2)t_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{t} \right].
\end{aligned} \tag{40}$$

This means that

$$\begin{aligned}
\lim_{|\omega_1| \rightarrow \infty} |L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)| &\leq \frac{1}{|4\pi b_1 b_2|} \lim_{|\omega_1| \rightarrow \infty} \int_{\mathbb{R}^2} \left| f(\mathbf{t}) \right. \\
&\quad \left. - e^{i(a_1/2b_1)(-2t_1b_1\pi/\omega_1 + (b_1\pi/\omega_1)^2)} f\left(t_1 - \frac{b_1\pi}{\omega_1}, t_2\right) \right| d\mathbf{t} \\
&= 0.
\end{aligned} \tag{41}$$

Similarly we can prove

$$\lim_{|\omega_2| \rightarrow \infty} |L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)| = 0. \tag{42}$$

Theorem 10 (continuity). *If $f \in L^1(\mathbb{R}^2; \mathbb{H})$, then $L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)$ is continuous on $\mathbb{R}^2$.*

Proof. Simple computations show that

$$\begin{aligned}
|L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega + \mathbf{h}) - L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)| &= \left| \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)(x_1+h_1)^2 - (2/b_1)(x_1+h_1)\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f(\mathbf{x}) \right. \\
&\quad \cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)(x_2+h_2)^2 - (2/b_2)(x_2+h_2)\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{x} - \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f(\mathbf{x}) \frac{1}{\sqrt{2\pi b_2}} \\
&\quad \cdot e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{x} \left| \leq \int_{\mathbb{R}^2} \left| \frac{1}{\sqrt{2\pi b_1}} \right. \right. \\
&\quad \cdot e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} \left(e^{i(1/2)((a_1/b_1)(2x_1h_1+h_1^2) - (2/b_1)\omega_1h_1)} f(\mathbf{x}) e^{j(1/2)((a_2/b_2)(2x_2h_2+h_2^2) - (2/b_2)\omega_2h_2)} - f(\mathbf{x}) \right) \\
&\quad \cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{x} = \frac{1}{|2\pi b_1 b_2|} \\
&\quad \cdot \int_{\mathbb{R}^2} \left| \left(e^{i(1/2)((a_1/b_1)(2x_1h_1+h_1^2) - (2/b_1)\omega_1h_1)} f(\mathbf{x}) e^{j(1/2)((a_2/b_2)(2x_2h_2+h_2^2) - (2/b_2)\omega_2h_2)} - f(\mathbf{x}) \right) \right| d\mathbf{x} \leq \frac{1}{|\pi b_1 b_2|} \int_{\mathbb{R}^2} |f(\mathbf{x})| d\mathbf{x}.
\end{aligned} \tag{43}$$

By the Lebesgue dominated convergence theorem, we may conclude that

$$|L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega + \mathbf{h}) - L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)| \longrightarrow 0 \tag{44}$$

when $\mathbf{h} \rightarrow 0$. This proves that $L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)$ is continuous on $\mathbb{R}^2$. Again since (43) is independent of ω , $L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)$ is, in fact, uniformly continuous on $\mathbb{R}^2$. $\square$

3.2. Useful Properties of the QLCT. Due to the noncommutativity of the kernel of the QLCT, we only have a left linearity property with specific constants

$$\alpha, \beta \in \{q \mid q = q_0 + \mathbf{i}q_1, q_0, q_1 \in \mathbb{R}\}, \tag{45}$$

which is

$$\begin{aligned}
L_{A_1, A_2}^{\mathbb{H}} \{\alpha f + \beta g\}(\omega) &= \alpha L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega) \\
&\quad + \beta L_{A_1, A_2}^{\mathbb{H}} \{g\}(\omega),
\end{aligned} \tag{46}$$

and a right linearity property with specific constants

$$\alpha', \beta' \in \{q \mid q = q_0 + \mathbf{j}q_2, q_0, q_2 \in \mathbb{R}\}. \tag{47}$$

Theorem 11 (shift property). *Given a quaternion function $f \in L^2(\mathbb{R}^2; \mathbb{H})$, let $\tau_{\mathbf{k}} f(\mathbf{x})$ denote the shifted (translated) function defined by $\tau_{\mathbf{k}} f(\mathbf{x}) = f(\mathbf{x} - \mathbf{k})$, where $\mathbf{k} \in \mathbb{R}^2$. Then one gets*

$$\begin{aligned}
L_{A_1, A_2}^{\mathbb{H}} \{\tau_{\mathbf{k}} f\}(\omega) &= e^{-\mathbf{i}a_1c_1k_1^2/2 + \mathbf{i}c_1k_1\omega_1} L_{A_1, A_2}^{\mathbb{H}} \{f\} \\
&\quad \cdot (\omega_1 - a_1k_1, \omega_2 - a_2k_2) \\
&\quad \cdot e^{-\mathbf{j}a_2c_2k_2^2/2 + \mathbf{j}c_2k_2\omega_2}.
\end{aligned} \tag{48}$$

Proof. Taking into account the definition of QLCT (17), we get

$$L_{A_1, A_2}^{\mathbb{H}} \{ \tau_{\mathbf{k}} f \} (\omega) = \frac{1}{\sqrt{2\pi b_1}} \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f(\mathbf{x} - \mathbf{k}) d\mathbf{x}. \quad (49)$$

By making the change of a variable $\mathbf{x} - \mathbf{k} = \mathbf{m}$, we easily obtain

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \tau_{\mathbf{k}} f \} (\omega) &= \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)(m_1+k_1)^2 - (2/b_1)(m_1+k_1)\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} f(\mathbf{m}) \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)(m_2+k_2)^2 - (2/b_2)(m_2+k_2)\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{m} = \frac{1}{\sqrt{2\pi b_1}} \\ &\cdot \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)m_1^2 - (2/b_1)m_1(\omega_1 - k_1 a_1) + (d_1/b_1)\omega_1^2 - \pi/2)} e^{i((1/2)(a_1/b_1)k_1^2)} e^{i(-(1/2)(2k_1\omega_1/b_1))} f(\mathbf{m}) \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)m_2^2 - (2/b_2)m_2(\omega_2 - k_2 a_2) + (d_2/b_2)\omega_2^2 - \pi/2)} e^{j((1/2)(a_2/b_2)k_2^2)} e^{j(-(1/2)(2k_2\omega_2/b_2))} d\mathbf{m}. \end{aligned} \quad (50)$$

Therefore, we further get

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \tau_{\mathbf{k}} f \} (\omega) &= \frac{1}{\sqrt{2\pi b_1}} \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)m_1^2 - (2/b_1)m_1(\omega_1 - k_1 a_1) + (d_1/b_1)(\omega_1 - k_1 a_1 + k_1 a_1)^2 - \pi/2)} e^{i((1/2)(a_1/b_1)k_1^2)} e^{i(-(1/2)(2k_1\omega_1/b_1))} f(\mathbf{m}) \\ &\cdot \sqrt{2\pi b_2} e^{j(1/2)((a_2/b_2)m_2^2 - (2/b_2)m_2(\omega_2 - k_2 a_2) + (d_2/b_2)(\omega_2 - k_2 a_2 + k_2 a_2)^2 - \pi/2)} e^{j((1/2)(a_2/b_2)k_2^2)} e^{j(-(1/2)(2k_2\omega_2/b_2))} d\mathbf{m} \\ &= e^{i(1/2)(d_1/b_1)(2(\omega_1 - k_1 a_1)k_1 a_1 + (k_1 a_1)^2)} e^{i((1/2)(a_1/b_1)k_1^2)} e^{i(-(1/2)(2k_1\omega_1/b_1))} \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)m_1^2 - (2/b_1)m_1(\omega_1 - k_1 a_1) + (d_1/b_1)(\omega_1 - k_1 a_1)^2)} f(\mathbf{m}) \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)m_2^2 - (2/b_2)m_2(\omega_2 - k_2 a_2) + (d_2/b_2)(\omega_2 - k_2 a_2)^2)} d\mathbf{m} e^{j((1/2)(d_2/b_2)(2(\omega_2 - k_2 a_2)k_2 a_2 + (k_2 a_2)^2))} e^{j((1/2)(a_2/b_2)k_2^2)} e^{j(-(1/2)(2k_2\omega_2/b_2))}. \end{aligned} \quad (51)$$

Applying the definition of the QLCT (17), the above expression can be rewritten in the form

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \tau_{\mathbf{k}} f \} (\omega) &= e^{i(1/2)(d_1/b_1)(2(\omega_1 - k_1 a_1)k_1 a_1 + (k_1 a_1)^2)} e^{i((1/2)(a_1/b_1)k_1^2)} e^{i(-(1/2)(2k_1\omega_1/b_1))} L_{A_1, A_2}^{\mathbb{H}} \{ f \} (\omega_1 - a_1 k_1, \omega_2 - a_2 k_2) \\ &\cdot e^{j(1/2)(d_2/b_2)(2(\omega_2 - k_2 a_2)k_2 a_2 + (k_2 a_2)^2)} e^{j((1/2)(a_2/b_2)k_2^2)} e^{j(-(1/2)(2k_2\omega_2/b_2))}. \end{aligned} \quad (52)$$

We notice that

$$\begin{aligned} &e^{i(1/2)(d_1/b_1)(2(\omega_1 - k_1 a_1)k_1 a_1 + (k_1 a_1)^2)} e^{i((1/2)(a_1/b_1)k_1^2)} e^{i(-(1/2)(2k_1\omega_1/b_1))} \\ &= e^{ik_1\omega_1(d_1 a_1/b_1 - 1/b_1)} e^{-i(1/2)k_1^2 a_1(d_1 a_1/b_1 - 1/b_1)}, \\ &e^{j(1/2)(d_2/b_2)(2(\omega_2 - k_2 a_2)k_2 a_2 + (k_2 a_2)^2)} e^{j((1/2)(a_2/b_2)k_2^2)} e^{j(-(1/2)(2k_2\omega_2/b_2))} \\ &= e^{jk_2\omega_2(d_2 a_2/b_2 - 1/b_2)} e^{-j(1/2)k_2^2 a_2(d_2 a_2/b_2 - 1/b_2)}. \end{aligned} \quad (53)$$

Because $a_i d_i - b_i c_i = 1$, then $d_i a_i/b_i - 1/b_i = c_i$ for $i = 1, 2$. It means that we get

$$\begin{aligned} &e^{i(1/2)(d_1/b_1)(2(\omega_1 - k_1 a_1)k_1 a_1 + (k_1 a_1)^2)} e^{i((1/2)(a_1/b_1)k_1^2)} e^{i(-(1/2)(2k_1\omega_1/b_1))} \\ &= e^{ik_1\omega_1 c_1} e^{-i(a_1 k_1^2/2)c_1}. \end{aligned} \quad (54)$$

By the above equalities, we finally arrive at

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \tau_{\mathbf{k}} f \} (\boldsymbol{\omega}) &= e^{i k_1 \omega_1 c_1} e^{-i(a_1 k_1^2/2) c_1} L_{A_1, A_2}^{\mathbb{H}} \{ f \} \\ &\cdot (\omega_1 - a_1 k_1, \omega_2 - a_2 k_2) \\ &\cdot e^{j k_2 \omega_2 c_2} e^{-j(a_2 k_2^2/2) c_2}. \end{aligned} \quad (55)$$

This completes the proof of theorem. $\square$

Next, we are concerned with the behavior of the QLCT under modulation.

Theorem 12 (modulation property). *Let $\mathbb{M}_{\boldsymbol{\omega}_0} f$ be modulation operator defined by $\mathbb{M}_{\boldsymbol{\omega}_0} f(\mathbf{x}) = e^{i \mathbf{x}_1 u_0} f(\mathbf{x}) e^{j \mathbf{x}_2 v_0}$ with $\boldsymbol{\omega}_0 = u_0 \mathbf{e}_1 + v_0 \mathbf{e}_2$. Then*

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \mathbb{M}_{\boldsymbol{\omega}_0} f \} (\boldsymbol{\omega}) &= L_{A_1, A_2}^{\mathbb{H}} \{ e^{i \mathbf{x}_1 u_0} f(\mathbf{x}) e^{j \mathbf{x}_2 v_0} \} (\boldsymbol{\omega}) \\ &= e^{-i b_1 d_1 u_0^2/2 + i d_1 u_0 \omega_1} L_{A_1, A_2}^{\mathbb{H}} \{ f \} \\ &\cdot (\omega_1 - u_0 b_1, \omega_2 - v_0 b_2) e^{-j b_2 d_2 v_0^2/2 + j d_2 v_0 \omega_2}. \end{aligned} \quad (56)$$

Proof. From Definition 4, it follows that

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \mathbb{M}_{\boldsymbol{\omega}_0} f \} (\boldsymbol{\omega}) &= \int_{\mathbb{R}^2} \frac{1}{\sqrt{2\pi b_1}} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega_1 + (d_1/b_1)\omega_1^2 - \pi/2)} e^{i \mathbf{x}_1 u_0} f(\mathbf{x}) \\ &\cdot e^{j \mathbf{x}_2 v_0} \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega_2 + (d_2/b_2)\omega_2^2 - \pi/2)} d\mathbf{x} \\ &= \frac{1}{\sqrt{2\pi b_1}} \\ &\cdot \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1\omega_1 + (d_1/b_1)\omega_1^2 + 2u_0x_1 - \pi/2)} f(\mathbf{x}) \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2\omega_2 + (d_2/b_2)\omega_2^2 + 2v_0x_2 - \pi/2)} d\mathbf{x} \\ &= \frac{1}{\sqrt{2\pi b_1}} \\ &\cdot \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1(\omega_1 - u_0 b_1) + (d_1/b_1)(\omega_1 - u_0 b_1)^2 - \pi/2)} f(\mathbf{x}) \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2(\omega_2 - v_0 b_2) + (d_2/b_2)(\omega_2 - v_0 b_2)^2 - \pi/2)} d\mathbf{x}. \end{aligned} \quad (57)$$

Subsequent calculations reveal that

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \mathbb{M}_{\boldsymbol{\omega}_0} f \} (\boldsymbol{\omega}) &= \frac{1}{\sqrt{2\pi b_1}} \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1(\omega_1 - u_0 b_1) + (d_1/b_1)(\omega_1 - u_0 b_1)^2 - \pi/2)} f(\mathbf{x}) \\ &\cdot \frac{1}{\sqrt{2\pi b_2}} e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2(\omega_2 - v_0 b_2) + (d_2/b_2)((\omega_2 - v_0 b_2) + v_0 b_2)^2 - \pi/2)} d\mathbf{x} = \frac{1}{\sqrt{2\pi b_1}} \\ &\cdot \int_{\mathbb{R}^2} e^{i(1/2)((a_1/b_1)x_1^2 - (2/b_1)x_1(\omega_1 - u_0 b_1) + (d_1/b_1)((\omega_1 - u_0 b_1)^2 + 2(\omega_1 - u_0 b_1)u_0 b_1 + u_0^2 b_1^2) - \pi/2)} f(\mathbf{x}) \frac{1}{\sqrt{2\pi b_2}} \\ &\cdot e^{j(1/2)((a_2/b_2)x_2^2 - (2/b_2)x_2(\omega_2 - v_0 b_2) + (d_2/b_2)((\omega_2 - v_0 b_2)^2 + 2(\omega_2 - v_0 b_2)v_0 b_2 + v_0^2 b_2^2) - \pi/2)} d\mathbf{x}. \end{aligned} \quad (58)$$

Hence,

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \mathbb{M}_{\boldsymbol{\omega}_0} f \} (\boldsymbol{\omega}) &= e^{i(\omega_1 - u_0 b_1)u_0 d_1 + i d_1 u_0^2 b_1/2} L_{A_1, A_2}^{\mathbb{H}} (\omega_1 \\ &- u_0 b_1, \omega_2 - v_0 b_2) e^{j(\omega_2 - v_0 b_2)v_0 d_2 + j d_2 v_0^2 b_2/2} \\ &= e^{i \omega_1 u_0 d_1} e^{-i((2b_1 d_1 u_0^2 - b_1 d_1 u_0^2)/2)} L_{A_1, A_2}^{\mathbb{H}} (\omega_1 - u_0 b_1, \omega_2 \\ &- v_0 b_2) e^{j \omega_2 v_0 d_2} e^{-j((2b_2 d_2 v_0^2 - b_2 d_2 v_0^2)/2)} \\ &= e^{i \omega_1 u_0 d_1} e^{-i(b_1 d_1 u_0^2/2)} L_{A_1, A_2}^{\mathbb{H}} (\omega_1 - u_0 b_1, \omega_2 - v_0 b_2) \\ &\cdot e^{j \omega_2 v_0 d_2} e^{-j(b_2 d_2 v_0^2/2)}. \end{aligned} \quad (59)$$

This is desired result. $\square$

Theorem 13 (time-frequency shift). *If quaternion function $f \in L^2(\mathbb{R}^2; \mathbb{H})$, then one gets*

$$\begin{aligned} L_{A_1, A_2}^{\mathbb{H}} \{ \mathbb{M}_{\boldsymbol{\omega}_0} \tau_{\mathbf{k}} f \} (\boldsymbol{\omega}) &= L_{A_1, A_2}^{\mathbb{H}} \{ e^{i \mathbf{x}_1 u_0} f(\mathbf{x} - \mathbf{k}) e^{j \mathbf{x}_2 v_0} \} \\ &\cdot (\boldsymbol{\omega}) \\ &= e^{-i(a_1 c_1 k_1^2 + b_1 d_1 u_0^2)/2 + i(c_1 k_1 + d_1 u_0)\omega_1 - i b_1 c_1 k_1 u_0} L_{A_1, A_2}^{\mathbb{H}} \{ f \} \\ &\cdot (\omega_1 - a_1 k_1 - u_0 b_1, \omega_2 - a_2 k_2 - v_0 b_2) \\ &\cdot e^{-j(a_2 c_2 k_2^2 + b_2 d_2 v_0^2)/2 + j(c_2 k_2 + d_2 v_0)\omega_2 - j b_2 c_2 k_2 v_0}. \end{aligned} \quad (60)$$

Proof. The proof directly follows from two previous theorems. $\square$

The above properties of the QLCT are summarized in Table 1.

TABLE 1: Properties of the QLCT of $f, g \in L^2(\mathbb{R}^2; \mathbb{H})$, where $\alpha, \beta, \alpha', \beta' \in \mathbb{H}$ are constants and $\omega_0 = u_0 \mathbf{e}_1 + v_0 \mathbf{e}_2 \in \mathbb{R}^2$.

Property	Quaternion func.	QLCT
Left linearity	$\alpha f + \beta g$	$\alpha L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega) + \beta L_{A_1, A_2}^{\mathbb{H}} \{g\}(\omega)$
Right linearity	$f \alpha' + g \beta'$	$L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega) \alpha' + L_{A_1, A_2}^{\mathbb{H}} \{g\}(\omega) \beta'$
Shift	$f(\mathbf{x} - \mathbf{k})$	$e^{-i a_1 c_1 k_1^2/2 + i c_1 k_1 \omega_1} L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega_1 - a_1 k_1, \omega_2 - a_2 k_2) e^{-j a_2 c_2 k_2^2/2 + j c_2 k_2 \omega_2}$
Modulation	$e^{i x_1 u_0} f(\mathbf{x}) e^{j x_2 v_0}$	$e^{-i b_1 d_1 u_0^2/2 + i d_1 u_0 \omega_1} L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega_1 - u_0 b_1, \omega_2 - v_0 b_2) e^{-j b_2 d_2 v_0^2/2 + j d_2 v_0 \omega_2}$
Time-frequency	$e^{i x_1 u_0} f(\mathbf{x} - \mathbf{k}) e^{j x_2 v_0}$	$e^{-i(a_1 c_1 k_1^2 + b_1 d_1 u_0^2)/2 + i(c_1 k_1 + d_1 u_0)\omega_1 - i b_1 c_1 k_1 u_0} L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega_1 - a_1 k_1 - u_0 b_1, \omega_2 - a_2 k_2 - v_0 b_2) e^{-j(a_2 c_2 k_2^2 + b_2 d_2 v_0^2)/2 + j(c_2 k_2 + d_2 v_0)\omega_2 - j b_2 c_2 k_2 v_0}$
Gaussian function	$e^{-(k_1 x_1^2 + k_2 x_2^2)}$	$\frac{1}{\sqrt{a_1 + 2k_1} b_1 i} e^{(\omega_1^2/2)/((c_1 + 2k_1 i d_1)/(2k_1 b_1 - i a_1))} \frac{1}{\sqrt{a_2 + 2k_2} b_2 j} e^{(\omega_2^2/2)/((c_2 + 2k_2 j d_2)/(2k_2 b_2 - j a_2))}$

4. Heisenberg Uncertainty Principle for QLCT

The classical uncertainty principle of harmonic analysis states that a nontrivial function and its Fourier transform cannot be sharply localized simultaneously. In quantum mechanics, the uncertainty principle asserts that one cannot at the same time be certain of the position and of the velocity of an electron (or any particle) [20]. Let us now give an alternative proof of the Heisenberg type uncertainty principle for the QLCT, which is recently studied in [8] (the uncertainty principle of the QCT was proved using the exponential form of a 2D quaternion function and proposed proof of this paper uses the relationship between the QFT and QLCT). However, before proceeding with the statement of this main result, we need to introduce the component-wise uncertainty principle for the QFT as follows (see [12] for more details).

Theorem 14 (the QFT component-wise uncertainty principle). *Suppose that $f \in L^1(\mathbb{R}^2; \mathbb{H}) \cap L^2(\mathbb{R}^2; \mathbb{H})$. If $\partial f / \partial x_k$ and $\omega_k(\partial f / \partial x_k) \in L^2(\mathbb{R}^2; \mathbb{H})$, then one has*

$$\int_{\mathbb{R}^2} x_k^2 |f(\mathbf{x})|^2 d\mathbf{x} \int_{\mathbb{R}^2} \omega_k^2 |\mathcal{F}_q \{f\}(\omega)|^2 d\omega \geq \frac{1}{4} \left(\int_{\mathbb{R}^2} |f(\mathbf{x})|^2 d\mathbf{x} \right)^2, \quad k = 1, 2. \quad (61)$$

The generalization of the above uncertainty principle to the QLCT domain is given by the following theorem (for more detailed information, see [8]).

Theorem 15 (the QLFT component-wise uncertainty principle). *Assume that $f \in L^1(\mathbb{R}^2; \mathbb{H}) \cap L^2(\mathbb{R}^2; \mathbb{H})$, $\partial f / \partial x_k \in L^2(\mathbb{R}^2; \mathbb{H})$ and that $L_{A_1, A_2}^{\mathbb{H}} \{f\}$, $\omega_k L_{A_1, A_2}^{\mathbb{H}} \{f\} \in L^2(\mathbb{R}^2; \mathbb{H})$, $k = 1, 2$. Then, the following inequality holds:*

$$\int_{\mathbb{R}^2} x_k^2 |f(\mathbf{x})|^2 d\mathbf{x} \int_{\mathbb{R}^2} \omega_k^2 |L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)|^2 d\omega \geq \frac{b_k^2}{4} \left(\int_{\mathbb{R}^2} |f(\mathbf{x})|^2 d\mathbf{x} \right)^2, \quad k = 1, 2. \quad (62)$$

Proof. Substituting the quaternion function f by g_f defined by (21) on both sides of (62), we easily obtain

$$\int_{\mathbb{R}^2} x_k^2 |g_f(\mathbf{x})|^2 d\mathbf{x} \int_{\mathbb{R}^2} \omega_k^2 |\mathcal{F}_q \{g_f\}(\omega)|^2 d\omega \geq \frac{1}{4} \left(\int_{\mathbb{R}^2} |g_f(\mathbf{x})|^2 d\mathbf{x} \right)^2. \quad (63)$$

Now setting $\omega = \omega/\mathbf{b}$, we further have

$$\int_{\mathbb{R}^2} x_k^2 \left| \frac{e^{-i(\pi/4)}}{\sqrt{b_1}} \tilde{f}(\mathbf{x}) \frac{e^{-j(\pi/4)}}{\sqrt{b_2}} \right|^2 d\mathbf{x} \cdot \int_{\mathbb{R}^2} \frac{\omega_k^2}{b_k^2} |\mathcal{F}_q \{g_f\}(\frac{\omega}{\mathbf{b}})|^2 d\frac{\omega}{\mathbf{b}} \geq \frac{1}{4} \left(\int_{\mathbb{R}^2} \left| \frac{e^{-i(\pi/4)}}{\sqrt{b_1}} \tilde{f}(\mathbf{x}) \frac{e^{-j(\pi/4)}}{\sqrt{b_2}} \right|^2 d\mathbf{x} \right)^2, \quad (64)$$

and thus

$$\int_{\mathbb{R}^2} \frac{x_k^2}{|b_1 b_2|^2} |\tilde{f}(\mathbf{x})|^2 d\mathbf{x} \int_{\mathbb{R}^2} \frac{\omega_k^2}{b_k^2} |\mathcal{F}_q \{g_f\}(\frac{\omega}{\mathbf{b}})|^2 d\omega \geq \frac{1}{4 |b_1 b_2|^2} \left(\int_{\mathbb{R}^2} |\tilde{f}(\mathbf{x})|^2 d\mathbf{x} \right)^2. \quad (65)$$

Hence,

$$\int_{\mathbb{R}^2} \frac{x_k^2}{|b_1 b_2|^2} |e^{i(a_1/2b_1)x_1^2} f(\mathbf{x}) e^{j(a_2/2b_2)x_2^2}|^2 d\mathbf{x} \cdot \int_{\mathbb{R}^2} \frac{\omega_k^2}{b_k^2} |\mathcal{F}_q \{g_f\}(\frac{\omega}{\mathbf{b}})|^2 d\omega \geq \frac{1}{4 |b_1 b_2|^2} \left(\int_{\mathbb{R}^2} |e^{i(a_1/2b_1)x_1^2} f(\mathbf{x}) e^{j(a_2/2b_2)x_2^2}|^2 d\mathbf{x} \right)^2. \quad (66)$$

By inserting (23) into (66), we immediately obtain

$$\begin{aligned} & \int_{\mathbb{R}^2} \frac{x_k^2}{|b_1 b_2|^2} \left| e^{i(a_1/2b_1)x_1^2} f(\mathbf{x}) e^{j(a_2/2b_2)x_2^2} \right|^2 d\mathbf{x} \\ & \cdot \int_{\mathbb{R}^2} \frac{\omega_k^2}{b_k^2} \left| e^{-i(d_1/2b_1)\omega_1^2} L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega) \right. \\ & \cdot e^{-j(d_2/2b_2)\omega_2^2} \left. \right|^2 d\omega \\ & \geq \frac{1}{4|b_1 b_2|^2} \left(\int_{\mathbb{R}^2} \left| e^{i(a_1/2b_1)x_1^2} f(\mathbf{x}) e^{j(a_2/2b_2)x_2^2} \right|^2 d\mathbf{x} \right)^2. \end{aligned} \quad (67)$$

Simplifying it gives

$$\begin{aligned} & \int_{\mathbb{R}^2} x_k^2 |f(\mathbf{x})|^2 d\mathbf{x} \int_{\mathbb{R}^2} \omega_k^2 |L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)|^2 d\omega \\ & \geq \frac{b_k^2}{4} \left(\int_{\mathbb{R}^2} |f(\mathbf{x})|^2 d\mathbf{x} \right)^2. \end{aligned} \quad (68)$$

This finishes the proof of theorem. $\square$

It is not difficult to check that directional uncertainty principle for the QFT takes the following form (cf. [21, 22]).

Theorem 16. Suppose that $f \in L^1(\mathbb{R}^2; \mathbb{H}) \cap L^2(\mathbb{R}^2; \mathbb{H})$. If $\partial f / \partial x_k$ and $\omega_k(\partial f / \partial x_k) \in L^2(\mathbb{R}^2; \mathbb{H})$, then one has

$$\begin{aligned} & \int_{\mathbb{R}^2} |x|^2 |f(\mathbf{x})|^2 d\mathbf{x} \int_{\mathbb{R}^2} |\omega|^2 |\mathcal{F}_q \{f\}(\omega)|^2 d\omega \\ & \geq \left(\int_{\mathbb{R}^2} |f(\mathbf{x})|^2 d\mathbf{x} \right)^2. \end{aligned} \quad (69)$$

Proceeding as in the proof of Theorem 15, we obtain the QLCT directional uncertainty principle as follows.

Theorem 17. Suppose that $f \in L^1(\mathbb{R}^2; \mathbb{H}) \cap L^2(\mathbb{R}^2; \mathbb{H})$ and $L_{A_1, A_2}^{\mathbb{H}} \{f\}$ and $|\omega|^2 L_{A_1, A_2}^{\mathbb{H}} \{f\} \in L^2(\mathbb{R}^2; \mathbb{H})$. Then the following inequality is satisfied:

$$\begin{aligned} & \int_{\mathbb{R}^2} |x|^2 |f(\mathbf{x})|^2 d\mathbf{x} \int_{\mathbb{R}^2} |\omega|^2 |L_{A_1, A_2}^{\mathbb{H}} \{f\}(\omega)|^2 d\omega \\ & \geq |b|^2 \left(\int_{\mathbb{R}^2} |f(\mathbf{x})|^2 d\mathbf{x} \right)^2. \end{aligned} \quad (70)$$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The first author is partially supported by Hibah Penelitian Kompetisi Internal Tahun 2014 (no. 1454/UN4.20/PL.9/2014) from the Hasanuddin University, Indonesia. The second author is partially supported by JSPS. KAKENHI (C)25400202 of Japan.

References

- [1] N. Goel and K. Singh, "A modified convolution and product theorem for the linear canonical transform derived by representation transformation in quantum mechanics," *International Journal of Applied Mathematics and Computer Science*, vol. 23, no. 3, pp. 685–695, 2013.
- [2] P. Dang, G.-T. Deng, and T. Qian, "A tighter uncertainty principle for linear canonical transform in terms of phase derivative," *IEEE Transactions on Signal Processing*, vol. 61, no. 21, pp. 5153–5164, 2013.
- [3] H. M. Ozaktas, Z. Zalevsky, and M. A. Kutay, *The Fractional Fourier Transform with Application in Optics and Signal Processing*, Wiley, New York, NY, USA, 2001.
- [4] J. Zhao, R. Tao, Y.-L. Li, and Y. Wang, "Uncertainty principles for linear canonical transform," *IEEE Transactions on Signal Processing*, vol. 57, no. 7, pp. 2856–2858, 2009.
- [5] X. Guanlai, W. Xiaotong, and Y. Xiaogang, "On uncertainty principle for the linear canonical transform of complex signals," *IEEE Transactions on Signal Processing*, vol. 58, no. 9, pp. 4916–4918, 2010.
- [6] J. Shi, X. Liu, and N. Zhang, "On uncertainty principles for linear canonical transform of complex signals via operator methods," *Signal, Image and Video Processing*, vol. 8, no. 1, pp. 85–93, 2014.
- [7] B. Hu, Y. Zhou, L. D. Lie, and J. Y. Zhang, "Polar linear canonical transform in quaternion domain," *Journal of Information Hiding and Multimedia Signal Processing*, vol. 6, no. 6, pp. 1185–1193, 2015.
- [8] K. I. Kou, J.-Y. Ou, and J. Morais, "On uncertainty principle for quaternionic linear canonical transform," *Abstract and Applied Analysis*, vol. 2013, Article ID 725952, 14 pages, 2013.
- [9] K. I. Kou and J. Morais, "Asymptotic behaviour of the quaternion linear canonical transform and the Bochner-Minlos theorem," *Applied Mathematics and Computation*, vol. 247, no. 15, pp. 675–688, 2014.
- [10] Y. Yang and K. I. Kou, "Uncertainty principles for hypercomplex signals in the linear canonical transform domains," *Signal Processing*, vol. 95, pp. 67–75, 2014.
- [11] M. Bahri, "Quaternion linear canonical transform application," *Global Journal of Pure and Applied Mathematics*, vol. 11, no. 1, pp. 19–24, 2015.
- [12] M. Bahri, E. S. Hitzler, A. Hayashi, and R. Ashino, "An uncertainty principle for quaternion Fourier transform," *Computers & Mathematics with Applications*, vol. 56, no. 9, pp. 2398–2410, 2008.
- [13] E. M. Hitzler, "Directional uncertainty principle for quaternion Fourier transform," *Advances in Applied Clifford Algebras*, vol. 20, no. 2, pp. 271–284, 2010.
- [14] T. Bülöw, *Hypercomplex spectral signal representations for the processing and analysis of images [Ph.D. thesis]*, University of Kiel, Kiel, Germany, 1999.
- [15] M. Bahri, R. Ashino, and R. Vaillancourt, "Convolution theorems for quaternion Fourier transform: properties and applications," *Abstract and Applied Analysis*, vol. 2013, Article ID 162769, 10 pages, 2013.
- [16] E. M. Hitzler, "Quaternion Fourier transform on quaternion fields and generalizations," *Advances in Applied Clifford Algebras*, vol. 17, no. 3, pp. 497–517, 2007.
- [17] Q. Xiang and K.-Y. Qin, "On the relationship between the linear canonical transform and the Fourier transform," in *Proceedings*

of the 4th International Congress on Image and Signal Processing (CISP '11), pp. 2214–2217, IEEE, Shanghai, China, October 2011.

- [18] A. I. Zayed, “On the relationship between the Fourier and fractional Fourier transforms,” *IEEE Signal Processing Letters*, vol. 3, no. 12, pp. 310–311, 1996.
- [19] M. Bahri, Zulfajar, and R. Ashino, “Convolution and correlation theorem for linear canonical transform and properties,” *Information B*, vol. 17, no. 6, pp. 2509–2521, 2014.
- [20] W. Weyl, *The Theory of Groups and Quantum Mechanics*, Dover, New York, NY, USA, 2nd edition, 1950.
- [21] L.-P. Chen, K. I. Kou, and M.-S. Liu, “Pitt’s inequality and the uncertainty principle associated with the quaternion Fourier transform,” *Journal of Mathematical Analysis and Applications*, vol. 423, no. 1, pp. 681–700, 2015.
- [22] Y. Yang, P. Dang, and T. Qian, “Tighter uncertainty principles based on quaternion Fourier transform,” *Advances in Applied Clifford Algebras*, 2015.