Research Article

On Solvability Theorems of Second-Order Ordinary Differential Equations with Delay

Nai-Sher Yeh

Department of Mathematics, Fu Jen Catholic University, Xinzhuang District, New Taipei City 24205, Taiwan

Correspondence should be addressed to Nai-Sher Yeh; 038300@mail.fju.edu.tw

Received 24 October 2017; Accepted 28 January 2018; Published 1 April 2018

Academic Editor: Chun-Lei Tang

Copyright © 2018 Nai-Sher Yeh. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

For each $x_0 \in [0, 2\pi)$ and $k \in \mathbb{N}$, we obtain some existence theorems of periodic solutions to the two-point boundary value problem $u''(x) + k^2 u(x - x_0) + g(x, u(x - x_0)) = h(x)$ in $(0, 2\pi)$ with $u(0) - u(2\pi) = u'(0) - u'(2\pi) = 0$ when $g : (0, 2\pi) \times \mathbb{R} \rightarrow \mathbb{R}$ is a Caratheodory function which grows linearly in u as $|u| \rightarrow \infty$, and $h \in L^1(0, 2\pi)$ may satisfy a generalized Landesman-Lazer condition $(1 + \text{sign}(\beta)) \int_0^{2\pi} h(x)v(x)dx < \int_{v(x)>0} g_\beta^+(x)|v(x)|^{1-\beta}dx + \int_{v(x)<0} g_\beta^-(x)|v(x)|^{1-\beta}dx$ for all $v \in N(L) \setminus \{0\}$. Here $N(L)$ denotes the subspace of $L^1(0, 2\pi)$ spanned by $\sin kx$ and $\cos kx$, $-1 < \beta \leq 0$, $g_\beta^+(x) = \liminf_{u \rightarrow \infty} (g(x, u)u/|u|^{1-\beta})$, and $g_\beta^-(x) = \liminf_{u \rightarrow -\infty} (g(x, u)u/|u|^{1-\beta})$.

1. Introduction

Let $x_0 \in [0, 2\pi)$ and $k \in \mathbb{N}$ be fixed. We consider the following two-point boundary value problems:

$$u''(x) + k^2 u(x - x_0) + g(x, u(x - x_0)) = h(x) \quad \text{in } (0, 2\pi), \quad (1)$$

$$u(0) - u(2\pi) = u'(0) - u'(2\pi) = 0,$$

$$u''(x) + k^2 u(x) - g(x, u(x - x_0)) = -h(x) \quad \text{in } (0, 2\pi), \quad (2)$$

$$u(0) - u(2\pi) = u'(0) - u'(2\pi) = 0,$$

where $h \in L^1(0, 2\pi)$ is given and $g : (0, 2\pi) \times \mathbb{R} \rightarrow \mathbb{R}$ is a Caratheodory function; that is, $g(x, u)$ is continuous in $u \in \mathbb{R}$, for a.e. $x \in (0, 2\pi)$, is measurable in $x \in (0, 2\pi)$ for all $u \in \mathbb{R}$, and satisfies, for each $r > 0$, the fact that there exists an $a_r \in L^1(0, 2\pi)$ such that

$$|g(x, u)| \leq a_r(x) \quad (3)$$

for a.e. $x \in (0, 2\pi)$ and all $|u| \leq r$. Concerning the growth condition of the nonlinear term g to (1) and (2), we assume that

(H) there exist constants $-1 < \beta \leq 0$, $r_0 > 0$, and $a, b, c, d \in L^1(0, 2\pi)$, $a, b \geq 0$ and $a(x) \leq 2k + 1$ for a.e. $x \in (0, 2\pi)$ with strict inequality on a positive measurable subset of $(0, 2\pi)$, such that for a.e. $x \in (0, 2\pi)$ and all $u \geq r_0$

$$c(x)|u|^{-\beta} \leq g(x, u) \leq a(x)|u| + b(x); \quad (4)$$

and for a.e. $x \in (0, 2\pi)$ and all $u \leq -r_0$

$$-a(x)|u| - b(x) \leq g(x, u) \leq d(x)|u|^{-\beta}; \quad (5)$$

(G) there exist constants $-1 < \beta \leq 0$, $r_0 > 0$, and $a, b, c, d \in L^1(0, 2\pi)$, $a, b \geq 0$ and $a(x) \leq 2k - 1$ for a.e. $x \in (0, 2\pi)$ with strict inequality on a positive measurable subset of $(0, 2\pi)$, such that for a.e. $x \in (0, 2\pi)$ and all $u \geq r_0$

$$c(x)|u|^{-\beta} \leq g(x, u) \leq a(x)|u| + b(x); \quad (6)$$

and for a.e. $x \in (0, 2\pi)$ and all $u \leq -r_0$

$$-a(x)|u| - b(x) \leq g(x, u) \leq d(x)|u|^{-\beta}; \quad (7)$$

respectively, and a generalized Landesman-Lazer condition

$$0 < \int_{v(x)>0} g_{\beta}^{+}(x) |v(x)|^{1-\beta} dx + \int_{v(x)<0} g_{\beta}^{-}(x) |v(x)|^{1-\beta} dx, \quad (8)$$

for all $v \in N(L) \setminus \{0\}$, may be satisfied. Here $N(L)$ denotes the subspace of $L^1(0, 2\pi)$ spanned by $\sin kx$ and $\cos kx$, $\beta \in \mathbf{R}$, $g_{\beta}^{+}(x) = \liminf_{u \rightarrow \infty} (g(x, u)u/|u|^{1-\beta})$, and $g_{\beta}^{-}(x) = \liminf_{u \rightarrow -\infty} (g(x, u)u/|u|^{1-\beta})$. Under assumptions and either with or without the Landesman-Lazer condition

$$\int_0^{2\pi} h(x) v(x) dx < \int_{v(x)>0} g_0^{+} |v(x)| dx + \int_{v(x)<0} g_0^{-} |v(x)| dx \quad (9)$$

for all $v \in N(L) \setminus \{0\}$, the solvability of the problem (1) has been extensively studied if the nonlinearity $g(x, u)$ has at most linear growth in u as $|u| \rightarrow \infty$ (see [1–13] for the case $x_0 = 0$ and [14–16] for the general case) or grows superlinearly in u in one of directions $u \rightarrow \infty$ and $u \rightarrow -\infty$ and may be bounded in the other (see [8, 17] for the case $x_0 = 0$ and [14] for the general case when $k = 0$). Based on the well-known Leray-Schauder continuation method (see [18, 19]), we obtain solvability theorems to (1) (resp., (2)) when $g(x, u)$ satisfies (H) (resp., (G)) and either (8) with $-1 < \beta < 0$ or (9) with $\beta = 0$ is satisfied, which extends the results of [15] for the nonresonance case, and has been established in [9] for the case $x_0 = 0$ and $g(x, u)$ grows sublinearly in u as $|u| \rightarrow \infty$ with $-1 < \beta \leq 1$. Unfortunately, it is still unknown when $k \in \mathbf{N}$, $g(x, u)$ grows linearly in u as $|u| \rightarrow \infty$ and the assumption of (8) is replaced by

$$\int_0^{2\pi} h(x) v(x) dx = 0 < \int_{v(x)>0} g_{\beta}^{+}(x) |v(x)|^{1-\beta} dx + \int_{v(x)<0} g_{\beta}^{-}(x) |v(x)|^{1-\beta} dx \quad (10)$$

for all $v \in N(L) \setminus \{0\}$ with $\beta > 0$. In the following we will make use of real Banach spaces $L^p(0, 2\pi)$, $C[0, 2\pi]$ and Sobolev spaces $W^{2,1}(0, 2\pi)$ and $H^1(0, 2\pi)$. The norms of $L^p(0, 2\pi)$, $C[0, 2\pi]$ and $H^1(0, 2\pi)$ are denoted by $\|u\|_{L^p}$, $\|u\|_C$ and $\|u\|_{H^1}$, respectively. By a solution of (1), we mean a periodic function $u : \mathbf{R} \rightarrow \mathbf{R}$ of period 2π which belongs to $W^{2,1}(0, 2\pi)$ and satisfies the differential equation in (1) a.e. $x \in (0, 2\pi)$.

2. Existence Theorems

For each $v \in W^{2,1}(0, 2\pi)$ with $v(0) - v(2\pi) = v'(0) - v'(2\pi) = 0$ and $k \in \mathbf{N}$, we write $\bar{v} = \sum_{0 \leq j \leq k} P_j v$, $\tilde{v} = \sum_{j > k} P_j v$, and $v^{\perp} = \sum_{0 \leq j \neq k} P_j v$. Here $P_j v$ denotes the projection of v on the

eigenspace of d^2/dx^2 spanned by $\sin jx$ and $\cos jx$ for $j \in \mathbf{N} \cup \{0\}$. Just as an application of [11, Lemma 2] or [1, Lemma 2.2], we can modify slightly the proof of [15, Lemma 1] to obtain the next lemma.

Lemma 1. *Let $k \in \mathbf{N} \cup \{0\}$ and Γ be a nonnegative $L^1(0, 2\pi)$ -function such that for a.e. $x \in (0, 2\pi)$, $\Gamma(x) \leq 2k+1$ with strict inequality on a positive measurable subset of $(0, 2\pi)$. Then there exists a constant $K_1 > 0$ such that*

$$\begin{aligned} & \int_0^{2\pi} (\bar{u}(x - x_0) - \bar{u}(x)) \cdot (u''(x) + k^2 u(x - x_0) + p(x) u(x - x_0)) dx \\ & \geq K_1 \|u^{\perp}\|_{H^1}^2 \end{aligned} \quad (11)$$

whenever $p \in L^1(0, 2\pi)$ with $0 \leq p(x) \leq \Gamma(x)$ for a.e. $x \in (0, 2\pi)$ and $u \in W^{2,1}(0, 2\pi)$ is a periodic function of period 2π with $u(0) - u(2\pi) = u'(0) - u'(2\pi) = 0$.

Proof. Just as in [20, Lemma 1], we can modify slightly the proof of [11, Lemma 2] or [1, Lemma 2.2] to obtain the fact that there exists a constant $K_1 > 0$ such that

$$\begin{aligned} & \int_0^{2\pi} (\bar{u}'(x))^2 - (k^2 + p(x)) (\bar{u}(x))^2 dx \\ & + \int_0^{2\pi} (k^2 + p(x)) (\bar{u}(x))^2 - (\bar{u}'(x))^2 dx \\ & \geq 2K_1 \|u^{\perp}\|_{H^1}^2 \end{aligned} \quad (12)$$

whenever $p \in L^1(0, 2\pi)$ with $0 \leq p(x) \leq \Gamma(x)$ for a.e. $x \in (0, 2\pi)$ and $u \in W^{2,1}(0, 2\pi)$ with $u(0) - u(2\pi) = u'(0) - u'(2\pi) = 0$. Let us extend $u(x)$ and $p(x)$ 2π periodically in x to all of $\mathbf{R}$ and then use the same notations for the periodic extensions as for the original functions. In this case, we have $\int_0^{2\pi} (\bar{u}'(x))^2 dx = \int_0^{2\pi} (\bar{u}'(x - x_0))^2 dx$ and

$$\begin{aligned} & \int_0^{2\pi} [u''(x) + (k^2 + p(x)) u(x - x_0)] \cdot (\bar{u}(x - x_0) - \bar{u}(x)) dx = \int_0^{2\pi} (\bar{u}'(x))^2 dx \\ & - \int_0^{2\pi} \bar{u}'(x) \bar{u}'(x - x_0) dx + \frac{1}{2} \int_0^{2\pi} (k^2 + p(x)) \cdot [(\bar{u}(x - x_0))^2 - (\bar{u}(x))^2 - (\bar{u}(x - x_0))^2] dx \\ & + \frac{1}{2} \int_0^{2\pi} (k^2 + p(x)) \cdot [\bar{u}(x - x_0) + \bar{u}(x - x_0) - \bar{u}(x)]^2 dx \end{aligned}$$

$$\begin{aligned}
&\geq \int_0^{2\pi} (\bar{u}'(x))^2 dx - \frac{1}{2} \int_0^{2\pi} (\bar{u}'(x))^2 \\
&+ (\bar{u}'(x-x_0))^2 dx + \frac{1}{2} \int_0^{2\pi} (k^2 + p(x)) \\
&\cdot [(\bar{u}(x-x_0))^2 - (\bar{u}(x))^2 - (\bar{u}(x-x_0))^2] dx \\
&+ \frac{1}{2} \int_0^{2\pi} (k^2 + p(x)) \\
&\cdot [\bar{u}(x-x_0) + \bar{u}(x-x_0) - \bar{u}(x)]^2 dx \geq \frac{1}{2} \\
&\cdot \int_0^{2\pi} (\bar{u}'(x))^2 \\
&- (k^2 + p(x)) (\bar{u}(x))^2 dx + \frac{1}{2} \\
&\cdot \int_0^{2\pi} (\bar{u}'(x-x_0))^2 - (k^2 + p(x)) \\
&\cdot (\bar{u}(x-x_0))^2 dx + \frac{1}{2} \int_0^{2\pi} (k^2 + p(x)) \\
&\cdot (\bar{u}(x-x_0))^2 - (\bar{u}'(x-x_0))^2 dx - \frac{1}{2} \\
&\cdot \int_0^{2\pi} (\bar{u}'(x))^2 dx + \frac{1}{2} \int_0^{2\pi} (k^2 + p(x)) \\
&\cdot [\bar{u}(x-x_0) + \bar{u}(x-x_0) - \bar{u}(x)]^2 dx.
\end{aligned} \tag{13}$$

Since $\int_0^{2\pi} (\bar{u}(x))^2 dx = \int_0^{2\pi} (\bar{u}(x-x_0))^2 dx$, $p(x) \geq 0$ for a.e. $x \in (0, 2\pi)$, and $\int_0^{2\pi} \bar{v}(x)\bar{w}(x) dx = 0$ for all $v, w \in W^{2,1}(0, 2\pi)$ with $v(0) - v(2\pi) = v'(0) - v'(2\pi) = 0$ and $w(0) - w(2\pi) = w'(0) - w'(2\pi) = 0$, we have $\int_0^{2\pi} (\bar{u}'(x))^2 - (k^2 + p(x))(\bar{u}(x))^2 dx \geq 0$ and

$$\begin{aligned}
&-\frac{1}{2} \int_0^{2\pi} (\bar{u}'(x))^2 dx + \frac{1}{2} \int_0^{2\pi} (k^2 + p(x)) \\
&\cdot [\bar{u}(x-x_0) + \bar{u}(x-x_0) - \bar{u}(x)]^2 dx \geq -\frac{1}{2} \\
&\cdot \int_0^{2\pi} (\bar{u}'(x))^2 dx + \frac{1}{2} \\
&\cdot \int_0^{2\pi} k^2 [\bar{u}(x-x_0) + \bar{u}(x-x_0) - \bar{u}(x)]^2 dx \quad (14) \\
&= \frac{1}{2} \left[- \int_0^{2\pi} (\bar{u}'(x))^2 dx \right. \\
&+ k^2 \int_0^{2\pi} (\bar{u}(x-x_0))^2 dx \left. + \frac{1}{2} \right. \\
&\cdot k^2 \int_0^{2\pi} [\bar{u}(x-x_0) - \bar{u}(x)]^2 dx \geq 0.
\end{aligned}$$

Combining (12) with (13), we have

$$\begin{aligned}
&\int_0^{2\pi} (\bar{u}(x-x_0) - \bar{u}(x)) \\
&\cdot (u''(x) + k^2 u(x-x_0) + p(x)u(x-x_0)) dx \\
&\geq \frac{1}{2} \int_0^{2\pi} (\bar{u}'(x-x_0))^2 - (k^2 + p(x)) \\
&\cdot (\bar{u}(x-x_0))^2 dx + \frac{1}{2} \int_0^{2\pi} (k^2 + p(x)) \\
&\cdot (\bar{u}(x-x_0))^2 - (\bar{u}'(x-x_0))^2 dx \\
&\geq K_1 \|u^\perp\|_{H^1(x_0, x_0+2\pi)}^2 = K_1 \|u^\perp\|_{H^1}^2.
\end{aligned} \tag{15}$$

□

Lemma 2. Let $k \in \mathbf{N}$ and Γ be a nonnegative $L^1(0, 2\pi)$ -function such that for a.e. $x \in (0, 2\pi)$, $\Gamma(x) \leq 2k-1$ with strict inequality on a positive measurable subset of $(0, 2\pi)$. Then there exists a constant $K_2 > 0$ such that

$$\begin{aligned}
&\int_0^{2\pi} (\bar{\bar{u}}(x-x_0) - \bar{\bar{u}}(x)) \\
&\cdot (u''(x) + k^2 u(x) - p(x)u(x-x_0)) dx \quad (16) \\
&\geq K_2 \|u^\perp\|_{H^1}^2
\end{aligned}$$

whenever $p \in L^1(0, 2\pi)$ with $0 \leq p(x) \leq \Gamma(x)$ for a.e. $x \in (0, 2\pi)$ and $u \in W^{2,1}(0, 2\pi)$ is a periodic function of period 2π with $u(0) - u(2\pi) = u'(0) - u'(2\pi) = 0$. Here $\bar{\bar{v}} = \sum_{0 \leq j < k} P_j v$ and $\bar{\bar{v}} = \sum_{j \geq k} P_j v$ for each $v \in W^{2,1}(0, 2\pi)$ with $v(0) - v(2\pi) = v'(0) - v'(2\pi) = 0$.

Theorem 3. Let $k \in \mathbf{N} \cup \{0\}$ and $g : (0, 2\pi) \times \mathbf{R} \rightarrow \mathbf{R}$ be a Caratheodory function satisfying (H). Then for each $h \in L^1(0, 2\pi)$ problem (1) has a solution u , provided that either (8) with $-1 < \beta < 0$ or (9) with $\beta = 0$ holds.

Proof. Let $\alpha \in \mathbf{R}$ be fixed and $0 < \alpha < 2k+1$. We consider the boundary value problems

$$\begin{aligned}
&u''(x) + k^2 u(x-x_0) + (1-t)\alpha u(x-x_0) \\
&+ tg(x, u(x-x_0)) = th(x) \quad \text{in } (0, 2\pi), \quad (17) \\
&u(0) - u(2\pi) = u'(0) - u'(2\pi) = 0
\end{aligned}$$

for $0 \leq t \leq 1$, which becomes the original problem when $t = 1$. Since $0 < \alpha < 2k+1$, we observe from Lemma 1 that (17) has only a trivial solution when $t = 0$. To apply the Leray-Schauder continuation method, it suffices to show that solutions to (17) for $0 < t < 1$ have an a priori bound in $H^1(0, 2\pi)$. To this end, let $\theta : \mathbf{R} \rightarrow \mathbf{R}$ be a continuous function

such that $0 \leq \theta \leq 1$, $\theta(u) = 0$ for $|u| \leq r_0$, and $\theta(u) = 1$ for $|u| \geq 2r_0$. We define $e(x) = \max\{a_{r_0}(x), b(x), |c(x)|, |d(x)|\}$,

$$g_1(x, u) = \begin{cases} \min\{g(x, u) + e(x)|u|^{-\beta}, a(x)u\} \theta(u) & \text{if } u \geq 0 \\ \max\{g(x, u) - e(x)|u|^{-\beta}, a(x)u\} \theta(u) & \text{if } u \leq 0, \end{cases} \quad (18)$$

and $g_2(x, u) = g(x, u) - g_1(x, u)$. Then $g_1, g_2 : (0, 2\pi) \times \mathbf{R} \rightarrow \mathbf{R}$ are Caratheodory functions, such that for a.e. $x \in (0, 2\pi)$ and $u \in \mathbf{R}$, $u \neq 0$

$$0 \leq \frac{g_1(x, u)}{u} \leq a(x), \quad (19)$$

$$|g_2(x, u)| \leq e(x)|u|^{-\beta} + e(x). \quad (20)$$

If u is a possible solution to (17) for some $0 < t < 1$, then using (19), (20), and Lemma 1, we have

$$\begin{aligned} 0 &= \int_0^{2\pi} (\bar{u}(x) - \bar{u}(x - x_0)) [u''(x) + k^2 u(x - x_0) \\ &\quad + (1 - t)\alpha u(x - x_0) + t g_1(x, u(x - x_0)) \\ &\quad + t g_2(x, u(x - x_0)) - t h(x)] dx \\ &\geq K_1 \|u^\perp\|_{H^1}^2 - (\|e\|_{L^1} \|u\|_C^{-\beta} + \|e\|_{L^1} + \|h\|_{L^1}) (\|\bar{u}\|_C \\ &\quad + \|\bar{u}\|_C) \geq K_1 \|u^\perp\|_{H^1}^2 - C_1 (\|u\|_C^{-\beta} + 1) (\|\bar{u}\|_{H^1} \\ &\quad + \|\bar{u}\|_{H^1}), \end{aligned} \quad (21)$$

which implies that

$$\begin{aligned} \|u^\perp\|_{H^1}^2 &\leq \frac{C_1}{K_1} (\|u\|_C^{-\beta} + 1) (\|\bar{u}\|_{H^1} + \|\bar{u}\|_{H^1}) \\ &\leq C_2 (\|u\|_{H^1} + \|u\|_{H^1}^{1-\beta}) \end{aligned} \quad (22)$$

for some constants $C_1, C_2 > 0$ independent of u . It remains to show that solutions to (17) for $0 < t < 1$ have an a priori bound in $H^1(0, 2\pi)$. We argue by contradiction and suppose that there exists a sequence $\{u_n\}$ of periodic functions with period 2π and a corresponding sequence $\{t_n\}$ in $(0, 1)$ such that u_n is a solution to (17) with $t = t_n$ and $\|u_n\|_{H^1} \geq n$ for all n . Let $v_n = u_n / \|u_n\|_{H^1}$; then $\|v_n\|_{H^1} = 1$ for all $n \in \mathbf{N}$, and by (22) we have $\|v_n^\perp\|_{H^1} \rightarrow 0$ as $n \rightarrow \infty$. Since $\|v_n\|_{H^1} = 1$ and $\|P_k v_n\|_{H^1} \leq \|v_n\|_{H^1} + \|v_n^\perp\|_{H^1}$ for all $n \in \mathbf{N}$, we have a bounded sequence $\{P_k v_n\}$ in $H^1(0, 2\pi)$. For simplicity, we may assume that v_n converges to v in $H^1(0, 2\pi)$ for some $v \in N(L)$ with $\|v\|_{H^1} = 1$. In particular, $v_n \rightarrow v$ in $C[0, 2\pi]$. Clearly, $v(\cdot - x_0) \in N(L)$ and $\|v(\cdot - x_0)\|_{H^1} = \|v\|_{H^1}$. It follows that $u_n(x) \rightarrow \infty$ for each $x \in \mathbf{R}$ with $v(x) > 0$, and $u_n(x) \rightarrow -\infty$ for each $x \in \mathbf{R}$ with $v(x) < 0$. Since $\int_0^{2\pi} u_n''(x) P_k u_n(x -$

$x_0) dx + \int_0^{2\pi} k^2 u_n(x) P_k u_n(x - x_0) dx = 0$ and $\|P_k u_n(\cdot)\|_{L^2}^2 = \|P_k u_n(\cdot - x_0)\|_{L^2}^2$, we have

$$\begin{aligned} &\int_0^{2\pi} u_n''(x) P_k u_n(x - x_0) dx + \int_0^{2\pi} k^2 u_n(x - x_0) \\ &\quad \cdot P_k u_n(x - x_0) dx = \int_0^{2\pi} u_n''(x) \\ &\quad \cdot P_k u_n(x - x_0) dx + \int_0^{2\pi} k^2 u_n(x) \\ &\quad \cdot P_k u_n(x - x_0) dx \\ &\quad + \int_0^{2\pi} k^2 [u_n(x - x_0) - u_n(x)] \\ &\quad \cdot P_k u_n(x - x_0) dx \\ &= \int_0^{2\pi} k^2 [u_n(x - x_0) - u_n(x)] \\ &\quad \cdot P_k u_n(x - x_0) dx \\ &= \int_0^{2\pi} k^2 [P_k u_n(x - x_0) - P_k u_n(x)] \\ &\quad \cdot P_k u_n(x - x_0) dx \geq 0. \end{aligned} \quad (23)$$

Multiplying each side of (17) by $P_k v_n(x - x_0)$, and then integrating them over $[0, 2\pi]$ when $u = u_n$ and $t = t_n$, we get

$$\begin{aligned} &t_n \int_0^{2\pi} g(x, u_n(x - x_0)) P_k v_n(x - x_0) dx \\ &\leq (1 - t_n) \alpha \int_0^{2\pi} u_n(x - x_0) P_k v_n(x - x_0) dx \\ &\quad + t_n \int_0^{2\pi} g(x, u_n(x - x_0)) P_k v_n(x - x_0) dx \\ &\leq t_n \int_0^{2\pi} h(x) P_k v_n(x - x_0) dx. \end{aligned} \quad (24)$$

By (19) and the assumption of $-1 < \beta \leq 0$, we have

$$\begin{aligned} &g_1(x, u_n(x - x_0)) P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta \\ &= \frac{g_1(x, u_n(x - x_0))}{u_n(x - x_0)} u_n(x - x_0) P_k v_n(x - x_0) \\ &\quad \cdot \|u_n\|_{H^1}^\beta \geq \frac{g_1(x, u_n(x - x_0))}{u_n(x - x_0)} \\ &\quad \cdot \frac{-1}{2} [u_n(x - x_0) - P_k u_n(x - x_0)]^2 \|u_n\|_{H^1}^{\beta-1} \geq \frac{-1}{2} \\ &\quad \cdot a(x) [u_n^\perp(x - x_0)]^2 \|u_n\|_{H^1}^{\beta-1} \end{aligned} \quad (25)$$

for a.e. $x \in (0, 2\pi)$. Combining (22) with (25), we get that $g_1(x, u_n(x - x_0)) P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta$ is bounded from below

by an $L^1(0, 2\pi)$ -function independent of n . By (20) and the assumption of $-1 < \beta \leq 0$, we have

$$\begin{aligned} & |g_2(x, u_n(x - x_0)) P_k v_n(x - x_0)| \|u_n\|_{H^1}^\beta \\ & \leq [e(x) |u_n(x - x_0)|^{-\beta} + e(x)] |P_k v_n(x - x_0)| \\ & \cdot \|u_n\|_{H^1}^\beta \leq [e(x) |v_n(x - x_0)|^{-\beta} + e(x)] \\ & \cdot |P_k v_n(x - x_0)| \end{aligned} \quad (26)$$

for a.e. $x \in (0, 2\pi)$. In particular, $g_2(x, u_n(x - x_0)) P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta$ is bounded from below by an $L^1(0, 2\pi)$ -function independent of n , which implies that $g(x, u_n(x - x_0)) P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta$ is also so, $\int_{v(x-x_0)=0} \liminf_{n \rightarrow \infty} g(x, u_n(x - x_0)) P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta dx = 0$, and

$$\begin{aligned} & g(x, u_n(x - x_0)) P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta \\ & = \frac{g(x, u_n(x - x_0)) u_n(x - x_0)}{|u_n(x - x_0)|^{1-\beta}} \\ & \cdot \frac{P_k v_n(x - x_0) \operatorname{sgn}(u_n(x - x_0))}{|v_n(x - x_0)|^\beta} \end{aligned} \quad (27)$$

for all $n \in \mathbb{N}$ with $u_n(x - x_0) \neq 0$. Here $\operatorname{sign}(w) = 1$ if $w > 0$, $\operatorname{sign}(w) = 0$ if $w = 0$, and $\operatorname{sign}(w) = -1$ if $w < 0$. Applying Fatou's lemma to the integral $\int_0^{2\pi} g(x, u_n(x - x_0)) P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta dx$, we have

$$\begin{aligned} & \int_{v(x-x_0)>0} g_\beta^+(x) |v(x - x_0)|^{1-\beta} dx + \int_{v(x-x_0)<0} g_\beta^-(x) \\ & \cdot |v(x - x_0)|^{1-\beta} dx \\ & = \int_{v(x-x_0)>0} \liminf_{n \rightarrow \infty} \frac{g(x, u_n(x - x_0)) u_n(x - x_0)}{|u_n(x - x_0)|^{1-\beta}} \\ & \cdot \lim_{n \rightarrow \infty} \frac{P_k v_n(x - x_0) \operatorname{sgn}(u_n(x - x_0))}{|v_n(x - x_0)|^\beta} dx \\ & + \int_{v(x-x_0)<0} \liminf_{n \rightarrow \infty} \frac{g(x, u_n(x - x_0)) u_n(x - x_0)}{|u_n(x - x_0)|^{1-\beta}} \\ & \cdot \lim_{n \rightarrow \infty} \frac{P_k v_n(x - x_0) \operatorname{sgn}(u_n(x - x_0))}{|v_n(x - x_0)|^\beta} dx \\ & = \int_{v(x-x_0)>0} \liminf_{n \rightarrow \infty} g(x, u_n(x - x_0)) P_k v_n(x - x_0) \\ & \cdot \|u_n\|_{H^1}^\beta dx \\ & + \int_{v(x-x_0)<0} \liminf_{n \rightarrow \infty} g(x, u_n(x - x_0)) P_k v_n(x - x_0) \end{aligned}$$

$$\begin{aligned} & \cdot \|u_n\|_{H^1}^\beta dx \\ & = \int_{v(x-x_0)>0} \liminf_{n \rightarrow \infty} g(x, u_n(x - x_0)) P_k v_n(x - x_0) \\ & \cdot \|u_n\|_{H^1}^\beta dx \\ & + \int_{v(x-x_0)<0} \liminf_{n \rightarrow \infty} g(x, u_n(x - x_0)) P_k v_n(x - x_0) \\ & \cdot \|u_n\|_{H^1}^\beta dx + \int_{v(x-x_0)=0} \lim_{n \rightarrow \infty} g(x, u_n(x - x_0)) \\ & \cdot P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta dx \\ & = \int_0^{2\pi} \liminf_{n \rightarrow \infty} g(x, u_n(x - x_0)) P_k v_n(x - x_0) \\ & \cdot \|u_n\|_{H^1}^\beta dx \leq \liminf_{n \rightarrow \infty} \int_0^{2\pi} g(x, u_n(x - x_0)) \\ & \cdot P_k v_n(x - x_0) \|u_n\|_{H^1}^\beta dx \leq (1 + \operatorname{sign}(\beta)) \\ & \cdot \int_0^{2\pi} h(x) v(x - x_0) dx, \end{aligned} \quad (28)$$

which is a contradiction when either (8) with $-1 < \beta < 0$ or (9) with $\beta = 0$ is satisfied. Hence, the proof of this theorem is complete. $\square$

By slightly modifying the proof of Theorem 3, we can apply Lemma 2 to obtain an existence theorem to (2) when condition (H) is replaced by (G) and either (8) with $-1 < \beta < 0$ or (9) with $\beta = 0$ is satisfied, which has been established in [20] for the case $x_0 = 0$ when (9) with $\beta = 0$ is satisfied and in [9] for the case $x_0 = 0$ when (8) with $\beta = -1$ is satisfied.

Theorem 4. Let $k \in \mathbb{N}$ and $g : (0, 2\pi) \times \mathbb{R} \rightarrow \mathbb{R}$ be a Caratheodory function satisfying (G). Then for each $h \in L^1(0, 2\pi)$ problem (2) has a solution u , provided that either (8) with $-1 < \beta < 0$ or (9) with $\beta = 0$ holds.

Conflicts of Interest

There are no conflicts of interest involved.

References

- [1] P. Drábek, "Landesman-lazer condition for nonlinear problem with jumping nonlinearity," *Journal of Differential Equations*, vol. 85, no. 1, pp. 186–199, 1990.
- [2] P. Drábek and S. Invernizzi, "On the periodic BVP for the forced duffing equation with jumping nonlinearity," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 10, no. 7, pp. 643–650, 1986.
- [3] J.-P. Gossez and P. Omari, "Nonresonance with respect to the Fucik spectrum for periodic solutions of second order ordinary differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 14, no. 12, pp. 1079–1104, 1990.

- [4] C. P. Gupta, J. J. Nieto, and L. Sanchez, "Periodic solutions of some lienard and duffing equations," *Journal of Mathematical Analysis and Applications*, vol. 140, no. 1, pp. 67–82, 1989.
- [5] C. P. Gupta and J. Mawhin, "Asymptotic conditions at the two first eigenvalues for the periodic solutions of lineard differential equations and an inequality of E. Schmidt," *Zeitschrift Für Analysis Und Ihre Anwendungen*, vol. 3, no. 1, pp. 33–42, 1984.
- [6] C. P. Gupta, "Solvability of a forced autonomous duffing's equation with boundary conditions in the presence of damping," *Applications of Mathematics*, vol. 38, no. 3, pp. 195–203, 1993.
- [7] R. Iannacci and M. N. Nkashama, "Nonlinear two-point boundary value problems at resonance without landesman-lazer condition," *Proceedings of the American Mathematical Society*, vol. 106, no. 4, pp. 943–952, 1989.
- [8] C.-C. Kuo, "Periodic solutions of a two-point boundary value problem at resonance," *Osaka Journal of Mathematics*, vol. 37, no. 2, pp. 345–353, 2000.
- [9] C.-C. Kuo, "On the solvability of semilinear differential equations at resonance," *Proceedings of the Edinburgh Mathematical Society*, vol. 43, no. 1, pp. 103–112, 2000.
- [10] M. Martelli and J. D. Schuur, "Periodic solutions of linear type second order ordinary differential equations," *Tohoku Mathematical Journal*, vol. 32, pp. 337–351, 1983.
- [11] J. Mawhin and J. Ward, "Periodic solutions of some forced lineard differential equations at resonance," *Archiv der Mathematik*, vol. 41, no. 4, pp. 337–351, 1983.
- [12] J. Mawhin and J. R. Ward, "Nonuniform nonresonance conditions at the two first eigenvalues for periodic solutions of forced lineard and duffing equations," *Rocky Mountain Journal of Mathematics*, vol. 12, no. 4, pp. 643–654, 1982.
- [13] J. R. Ward Jr., "Periodic solutions for systems of second order differential equations," *Journal of Mathematical Analysis and Applications*, vol. 81, no. 1, pp. 92–98, 1981.
- [14] R. Iannacci and M. N. Nkashama, "Nonresonance conditions for periodic solutions of lineard and duffing equations with delay," *Annales de la Société scientifique de Bruxelles Série I*, vol. 99, no. 1, pp. 29–43, 1985.
- [15] R. Iannacci and M. N. Nkashama, "On periodic solutions of forced second order differential equations with a deviating argument," in *Lecture Notes in Mathematics*, vol. 1151, pp. 224–232, Springer, Berlin, Germany, 1985.
- [16] E. De Pascale and R. Iannacci, "Periodic solutions of generalized lineard equations with delay," in *Lecture Notes in Mathematics*, vol. 1017, pp. 148–156, Springer, Berlin, Germany, 1983.
- [17] R. Kannan and R. Ortega, "Periodic solutions of pendulum-type equations," *Journal of Differential Equations*, vol. 59, no. 1, pp. 123–144, 1985.
- [18] K. Deimling, *Nonlinear Functional Analysis*, Springer, New York, NY, USA, 1985.
- [19] J. Mawhin, *Topological Degree Methods in Nonlinear Boundary Value Problems*, vol. 40 of *CBMS Regional Conference Series in Mathematics*, American Mathematical Society, Providence, RI, USA, 1979.
- [20] R. Iannacci and M. N. Nkashama, "Nonlinear boundary value problems at resonance," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 11, no. 4, pp. 455–473, 1987.