Review Article

Impact of Autologous and Allogeneic Stem Cell Transplantation in Peripheral T-Cell Lymphomas

Peter Reimer

Klinik für Hämatologie, Internistische Onkologie und Stammzelltransplantation, Kliniken Essen Süd, Evangelisches Krankenhaus Essen-Werden gGmbH, Pattbergstraße 1-3, 45239 Essen, Germany

Correspondence should be addressed to Peter Reimer, p.reimer@kliniken-essen-sued.de

Received 25 August 2010; Revised 12 October 2010; Accepted 18 October 2010

Academic Editor: Brian Bolwell

Copyright © 2010 Peter Reimer. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peripheral T/NK-cell lymphomas (PTCLs) are rare malignancies characterized by poor prognosis. So far, no standard therapy has been established, due to the lack of randomised studies. High-dose therapy and autologous stem cell transplantation (HDT-autoSCT) have shown good feasibility with low toxicity in retrospective studies. In relapsing and refractory PTCL several comparison analyses suggest similar efficacy for PTCL when compared with aggressive B-cell lymphoma. In the upfront setting, prospective data show promising results with a long-lasting overall survival in a relevant subset of patients. Achieving a complete remission at transplantation seems to be the most important prognostic factor. Allogeneic stem cell transplantation (alloSCT) has been investigated only as salvage treatment. Especially when using reduced intensity conditioning regimen, eligible patients seem to benefit from this approach. To define the role for upfront stem cell transplantation a randomised trial by the German High-Grade Non-Hodgkin Lymphoma Study Group comparing HDT-autoSCT and alloSCT will be initiated this year.

1. Introduction

Peripheral T-cell lymphomas (PTCLs) represent approximately 10%–15% of all non-Hodgkin's lymphomas (NHLs) in Western countries [1–3]. Although the clinical appearance and the manifestation sites vary widely between the different subgroups, most PTCL share some characteristics. Most patients are of older age (median age >60 years) and usually present with advanced stage disease [4, 5]. PTCLs in general show an aggressive course and most studies detect the T-cell phenotype as an independent negative prognostic factor [6–9]. Both, the international prognostic index (IPI) and the prognostic index for T-cell lymphomas (PITs) that also include the bone marrow involvement, have shown prognostic value in PTCL and determine the outcome of patients with nodal PTCL [10–13]. In addition, in retrospective studies further parameters like the expression of Ki-67, the level of β_2 -microglobulin, and the detection of the Epstein-Barr virus (EBV) have been found to have some prognostic relevance in PTCL [14–16].

The prognosis of PTCL is poor with the exception of the ALK (anaplastic lymphoma kinase) expressing anaplastic large cell lymphoma (ALCL) with a more favourable outcome after conventional chemotherapy and the primary cutaneous T-cell lymphomas (CTCL) that usually show an indolent clinical course [17, 18]. In contrast, for the remaining PTCL the outcome following anthracycline-based chemotherapy is worse compared to aggressive B-cell lymphomas even regarding the pre-rituximab era with a median overall survival (OS) of 9 to 42 months [19–21].

So far, no accepted standard treatment could be defined for PTCL. This mainly results from a lack of PTCL-restricted randomised trials and the heterogeneity of most published series. Although the CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) and CHOP-like regimen are widely used first-line, these protocols have never been established prospectively in PTCL and are rather adopted from treatment strategies for aggressive B-cell lymphomas. Even the role of anthracyclines in the first-line treatment of PTCL is controversial since a large retrospective

international survey did not reveal a significant impact on OS [5].

To improve the treatment results in PTCL more aggressive strategies such as high-dose therapy with autologous stem cell transplantation (HDT-autoSCT) and allogeneic stem cell transplantation (alloSCT) seem attractive strategies in PTCL. In this paper the data on stem cell transplantation for PTCL will be discussed.

2. Autologous Stem Cell Transplantation

2.1. Second-Line Therapy. HDT-autoSCT has become the standard of care in relapsing and primary refractory high-grade B-cell lymphomas. In PTCL, prospective randomized studies on salvage HDT-autoSCT are lacking. To date, at least 16 retrospective studies, each including more than 15 patients have addressed this issue and are listed in Table 1 [22–37].

The cited studies were heterogeneous in terms of histological subgroups, patient characteristics, prognostic factors, myeloablative regimen, and duration of follow-up. In addition, some series included patients receiving upfront autoSCT and did not provide separate analyses for the patients treated in second line. Taken together, this strategy is feasible and safe with a low morbidity and mortality rate. The OS in these series ranged from 35% at 2 years to 70% at 5 years, respectively, and the disease-free survival (DFS) or event-free survival (EFS) from 28% at 2 years to 56% at 5 years, respectively. Although the earlier reports tend to show somewhat better results than the series published recently, when subgroup or matched control analyses were performed, the OS results for PTCL were equivalent to the long-term outcome in patients with aggressive B-cell lymphomas [22, 26, 32]. So far, it is unclear whether histology impacts the outcome of PTCL after salvage autoSCT. In some series ALCL showed a favorable outcome compared to other pathological subtypes [25, 26, 30, 31]. However, the ALK status was not determined in all series and furthermore, Zamkoff et al. could not demonstrate a long-term DFS for recurrent (ALK-negative) ALCL following HDT-autoSCT [29]. The encouraging results for patients with ALCL by Fanin et al., (5-year OS and PFS of 70% and 56%, resp.) were probably biased by age and inclusion of patients in first complete remission (CR), who showed a significantly better outcome in a subgroup analysis [23].

The disease status at the time of transplantation often correlates with the outcome after salvage HDT-autoSCT. In fact, several authors found a better long-term survival in patients transplanted in CR than in patients with other disease status at transplantation [23, 27, 32, 33, 35–37]. Other authors could not confirm this finding in their survey [24, 25, 31]. However, since all data in this setting are generated retrospectively, the value of this observation needs further observation.

In summary, second-line HDT-autoSCT in PTCL is feasible and seems an effective approach for a considerable subgroup of patients.

2.2. First-Line Therapy. Some retrospective studies on upfront HDT-autoSCT have been published and are summarized in Table 2(a) [38–43]. Like in the salvage setting, a comparison of the cited series is hampered by their variety. Some series reported mainly on patients with a low or intermediate low IPI, whereas others predominantly included patients with an unfavorable prognostic index. In addition, most studies contained patients receiving HDT-autoSCT in second line, but not all of them show a subgroup analysis for the upfront setting. The OS in these retrospective studies ranged from 53% at 3 years to 62–68% at 5 years. Interestingly, the DFS/EFS did not appear to be much lower than the OS in most cohorts that might indicate a substantial curative potential for this approach in previously untreated PTCL. The EBMT (European Group for Blood and Marrow Transplantation) published the largest study in this setting. Kyriakou et al. reported data on 146 patients with angioimmunoblastic T-cell lymphoma (AITL) showing an actuarial OS of 67% at 2 years and 59% at 4 years, respectively, after a median observation of 31 months. About two thirds of the patients were transplanted in first CR or PR. Interestingly, patients who received a TBI- (total body irradiation-) based conditioning regimen had a significantly lower relapse rate in this study [41].

In these retrospective studies chemotherapy-sensitive disease was the major factor predicting OS and PFS [39–41, 43]. Patients transplanted in CR or PR showed a superior long-term outcome compared to patients with chemotherapy-refractory disease. Other parameters (e.g., age, PIT, IPI) could not consistently be detected as being of prognostic value.

Although mainly showing promising results, the cited retrospective studies are limited by focussing on patients, only, who actually proceed to transplantation leading to superior results due to patient selection.

Prospective randomized PTCL-restricted studies assessing the value of upfront high-dose therapy in PTCL are lacking. Two French trials by the GELA (Groupe d'Etude des Lymphomes de l'Adulte) published data on autoSCT as frontline strategy in poor-risk, aggressive NHL, including PTCL [57–59]. In the LNH87-2 study, patients were treated with either consolidative sequential chemotherapy or HDT-autoSCT [57, 58]. The LNH93-3 trial compared a high-dose arm with shortened first-line myeloablative chemotherapy with a sequential consolidation chemotherapy arm. In the intent-to-treat analysis, none of these studies demonstrated a significant benefit for the high-dose arm [59]. In addition, a pooled data matched control analysis failed to show a significant advantage for upfront HDT-autoSCT [44, 60]. However, the limited number of patients in the high-dose group and the restriction to high-risk patients, only, do not allow to definitely clarify the impact of first-line HDT-autoSCT in PTCL from these data. In another subgroup analysis, Nickelsen et al. reported results of 33 patients with PTCL from a single-arm study by the German High-Grade Non-Hodgkin Lymphoma Study Group [45]. Patients with high-risk aggressive lymphomas were treated with dose-escalated CHOP plus etoposide necessitating repeated autoSCT. Compared to B-cell NHL, PTCL showed a

TABLE 1: Studies on high-dose therapy and autologous stem cell transplantation in PTCL as second-line therapy.

Author	Year	n	Age	Histologies (WHO)	Retrospective data					Followup (months)	Comment
					High-dose regimen	IPI	Status at Tx	DFS/EFS/PFS/RFS	OS		
Vose et al. [22]	1990	17	33	No data*	No data	No data	CR 42% PR 26%	28% (2 y)	35% (2 y)	28	
Fanin et al. [23]	1999	64	25	ALCL 100%	Diverse	aallPI 0/1 48% 2/3 22% Unknown 30%	CR 47%	56% (5 y)	70% (5 y)	43	34/64 transplanted 2.line
Rodríguez et al. [24]	2001	29	43	No data*	Diverse	0/1 31% 2 41% 3 21% 4/5 7%	CR 38% PR 48%	32% (3 y)	39% (3 y)	43	
Blystad et al. [25]	2001	40	42	PTCLu 50% ALCL 35% Other 15%	Diverse	aallPI 0/1 42% 2/3 45% Unknown 13%	CR 70% PR 30%	56% (3 y)	58% (3 y)	25	23/40 transplanted 2.line
Song et al. [26]	2002	36	46	PTCLu 56% ALCL 25% NK/T 11% Other 8%	Mel/Eto	No data	CR 42% PR 50%	37% (3 y)	48% (3 y)	42	
Rodríguez et al. [27]	2003	115	41	PTCLu 63% ALCL 22% NK/T 15%	Diverse	aallP 0/1 40% 2/3 60%	CR 56% PR 38%	60% (5 y) 49% (5 y)**	56% (5 y) 45% (5 y)**	37	78/115 transplanted 2.line
Schetelig et al. [28]	2003	29	51	AITL 100%	Diverse	aallPI 0/1 21% 2/3 79%	No data	37% (5 y) 39% (5 y)**	60% (5 y) 44% (5 y)**	60	15/29 transplanted 2.line
Zamkoff et al. [29]	2004	16	52	ALK ⁻ ALCL 100%	Diverse	aallPI 0/1 67% 2/3 33%	CR 60% PR 40%	12 w (median)	72 w (median)	No data	
Jantunen et al. [30]	2004	37	46	PTCLu 38% ALCL 38% EATL 14% Other 11%	BEAC/BEAM	0/1 46% 2 22% 3 19% 4/5 14% Unknown 3%	CR/PR 87%	44% (5 y) 28% (5 y)**	54% (5 y) 45% (5 y)**	24	19/37 transplanted 2.line TRM 16%
Jagasia et al. [31]	2004	28	39	ALCL 57% PTCLu 21% AITL 11% NK/T 11%	Cy/Eto/TBI or CBV	aallPI 0/1 29% 2/3 50% Unknown 21%	CR 39% PR 46%	50% (3 y)	69% (3 y)	44	7/28 underwent alloSCT

TABLE 1: Continued.

Author	Year	n	Age	Histologies (WHO)	Retrospective data				Followup (months)	Comment
					High-dose regimen	Status at Tx	DFS/EFS/PFS/RFS	OS		
Kewalramani et al. [32]	2006	24	48	PTCLu 58%	Diverse	CR 63% PR 37%	24% (5 y)	33% (5 y)	72	
				ALCL 17%						
				AITL 17%						
				Other 8%						
Kim et al. [33]	2007	40	44	PTCLu 50%	Diverse	CR 28% PR 52%	No data	11.5 m (median)	16	29/40 transplanted 2.line
				NK/T 25%						
				ALCL 13%						
				Other 13%						
Smith et al. [34]	2007	32	44	ALCL 66%	BEC	No data	18% (5 y)	34% (5 y)	30	26/32 transplanted 2.line
				PTCLu 34%						
Chen et al. [35]	2008	53	45	ALCL 34%	Diverse	CR/PR 89%	25% (5 y) 9% (5 y)**	48% (5 y) 37% (5 y)**	60	38/53 transplanted 2.line
				PTCLu 30%						
				AITL 17%						
				Other 19%						
Lee et al. [36]	2008	47	42	NK/T 100%	CVB/BEAM/ MCEC (72%)	CR 58%	No data	No data	117	33/47 transplanted 2.line
Yang et al. [37]	2009	64	44	PTCLu 100%	BEAM/CVB (70%)	CR 33% PR 58%	44% (3 y) 33% (3 y)**	53% (3 y) 46% (3 y)**	30	36/64 transplanted 2.line

Studies including both patients receiving HDT-autoSCT 1.line and 2.line are listed in the table representing the predominant group

* classified according to the Working Formulation

** for the subgroup of patients transplanted 2.line.

TABLE 2: Studies on high-dose therapy and autologous stem cell transplantation in PTCL as first-line therapy.

Author	Year	n	Age	Histologies (WHO)	IPI	High-dose regimen	Status at Tx	DFS/EFS/PFS/RFS	OS	Followup (months)	Comment	
Rodríguez et al. [38]	2007	19	46	AITL 100%	aalPI 0/1 37% 2/3 63%	BEAM/BEAC (79%)	CR1 42% PR1 26%	55% (3 y)	60% (3 y)	25	15/19 transplanted 1.line	
Rodríguez et al. [39]	2007	74	46	PTCLu 50% ALCL 31% AITL 11% Other 7%	aalPI 0/1 35% 2/3 65%	BEAM/BEAC (91%)	No data	63% (5 y)	68% (5 y)	67		
Feyler et al. [40]	2007	64	45	PTCLu 47% ALCL 31% AITL 8% Other 14%	0/1 34% 2 11% 3 11% 4/5 2% Unknown 42%	Diverse	CR1 48% PR1 23%	50% (3 y)	53% (3 y)	48	Incl. CTCL, T-cell leukemia/lymphoma 46/64 transplanted 1.line	
Kyriakou et al. [41]	2008	146	53	AITL 100%	No data	BEAM (74%)	CR1 33% PR 36%	49% (4 y)	59% (4 y)	31	101/146 transplanted 1.line	
Prochazka et al. [42]	2009	18	59	PTCLu 56% ALCL 39% AITL 6%	0/1 28% 2 28% ≥3 44%	BEAM	No data	52% (2 y)	71% (2 y)	26		
Numata et al. [43]	2010	39	53	PTCLu 31% AITL 28% ALCL 23% NK/T 18%	aalPI 0/1 33% 2/3 59% Unknown 8%	MCEC (n = 32) TBI-based (n = 7)	CR1 69%	61% (5 y)	62% (5 y)	78	23/39 transplanted 1.line	
Studies including both patients receiving HDT-autoSCT 1.line and 2.line are listed in the table representing the predominant group.												
(b) Prospective data												
Author	Year	n	Age	Histologies (WHO)	IPI	High-dose regimen	Status at Tx	Tx rate	DFS/EFS/PFS	OS	Followup (months)	Comment
Mounier et al. [44]	2004	28	36	PTCLu 56% Precursor 44% ALK ⁺ ALCL 39%	0/1 25% ≥2 75%	BEAM/CBV	CR 100%	No data	44% (5 y)	54% (5 y)	78'	Incl. precursor T-cell lymphoma
Nickelsen et al. [45]	2009	33	48	PTCLu 33% AITL 12% Other 16%	aalPI 0/1 9% 2/3 91%	MegaCHOEP	CR 49% PR 6% after therapy	67%	26% (3 y)	45% (3 y)	53	Sequential HDT-autoSCT Subgroup analysis

significantly worse OS and EFS at 3 years in an intent-to-treat analysis.

So far, five larger prospective PTCL-restricted trials have published data on 372 patients with frontline HDT-autoSCT [46–50]. Compared to the cited retrospective studies, these prospective series are much more homogeneous. The median age ranged between 43 and 57 years, PTCL unspecified; AITL and ALCL accounted for 77 to 100% of all histological subtypes; the age-adjusted IPI was high or intermediate high in 46 to 72%; the most commonly used myeloablative regimen was the BEAM protocol ($n = 228$), and the disease status at transplantation was CR or PR in 59 to 76%. Only one study included ALK-positive ALCL [46]. In these trials the OS ranged from 48 to 73% at 3 years to 34% at 12 years. The DFS/EFS or the progression-free survival (PFS) was between 36 to 53% at 3 years and 30% at 12 years. One consistently found problem of upfront HDT-autoSCT is early progressive disease leading to about one third of patients in intent-to-treat analyses, who finally fail to achieve transplantation. Mercadal et al. reported a still significant lower transplantation rate of 41%. Of note, in this trial poor stem cell mobilization was the second most frequent cause of failing HDT-autoSCT [48].

With regard to these prospective data, again the remission status at the time of transplantation was a significantly prognostic factor in most studies that provided this analysis [46–48]. In addition, the IPI and PIT also show prognostic value in some [48, 49] but not in all series [47]. Other parameters, for example, histological subtype, age, sex, stage have not been concordantly been found to impact the outcome. The prospective series on upfront autoSCT are listed in Table 2(b).

3. Allogeneic Stem Cell Transplantation

In contrast to the cytotoxic effect of HDT-autoSCT, allogeneic SCT (alloSCT) could add a graft-versus-lymphoma (GVL) effect to the myeloablative or reduced intensity conditioning (RIC) regimen, potentially improving the therapeutic outcome. However, the experience with alloSCT for PTCL is limited. To date, no relevant data for the upfront setting are available. Besides some case reports, five retrospective series with at least 10 patients have been reported in patients with relapsing and refractory PTCL (Table 3(a)) [40, 51–54].

The largest series was published by the Société Française de Greffe de Moëlle et de Thérapie Cellulaire. In 77 pretreated patients who mainly had a myeloablative conditioning regimen the 5-year OS and PFS were 57% and 53%, respectively, after a median followup of 43 months. The treatment-related mortality (TRM) was 33% at 5 years. In a multivariate analysis, chemotherapy-resistant disease at transplantation and grade 3/4 acute graft-versus-host disease (GVHD) were the strongest adverse prognostic factors for OS. The TRM was similar in both conditioning groups [53]. Most studies could reveal a GVL effect [52–54]. However, the TRM/NRM (nonrelapse mortality) had a relevant impact on outcome and was increasing over time up to 69% at 3 years in the series

by Hamadani et al [52]. The OS ranged from 40% at 2 years to 57% at 5 years.

Two prospective studies have been published so far (Table 3(b)) [55, 56]. In the Italian phase II trial by Corradini et al. 17 patients underwent RIC and alloSCT as salvage therapy [55]. Eight out of 17 patients had failed front-line HDT-autoSCT. After a median followup of 28 months, 14 of 17 patients were alive. The estimated 3-year OS and PFS rates were 81% and 64%, respectively. The TRM was impressively low with only 6%. Notably, donor lymphocyte infusions given at the time of progression resulted in a disease response in two out of four patients, indicating evidence of a GVL effect. In the German study by Wulf et al. 10 patients were treated with chemotherapy combined with the humanized antiCD52 monoclonal antibody, alemtuzumab, followed by RIC and alloSCT. Two patients had prior HDT-autoSCT. The OS was 70% with six patients in CR after a median followup of 7 months [56].

4. Summary

Due to their generally poor prognosis after conventional chemotherapy more effective treatment strategies in PTCL are urgently needed. Although randomised trials are lacking, HDT-autoSCT can be regarded as feasible and safe in PTCL. In the salvage setting, several subgroup analysis and comparisons show similar results compared to diffuse large-cell B-cell lymphoma. This finding could recently be confirmed by Sohn et al. [61]. Therefore, taken together the existing data, HDT-autoSCT seems a reasonable approach in relapsing and refractory PTCL particularly in those with chemotherapy-sensitive disease.

The value of upfront HDT-autoSCT remains to be definitely established. A recently published retrospective comparison did not find a significant benefit of this strategy compared to conventional treatment [62]. However, in this study the high-dose group was heterogeneous ranging from high-dose CHOP to alloSCT. In contrast, other retrospective studies revealed a significant better outcome when HDT-autoSCT was compared to chemotherapy, alone [5, 36]. Taken together, the prospective data mainly show promising results, especially for patients achieving a good remission status prior to transplantation, that has been reported as independent prognostic factor in most series. Therefore, these patients should mainly be regarded as candidates for upfront autoSCT. Since one major obstacle of this approach is early progressive disease, novel treatment concepts incorporating new agents and/or dose-dense regimen should be further investigated to improve remission status prior to transplantation. In a subanalysis of several trials by the DSHNHL (German High-Grade Non-Hodgkin Lymphoma Study Group), the addition of etoposide to the CHOP-protocol improved the outcome in younger patients with PTCL [63]. Furthermore, alemtuzumab has shown efficacy especially in untreated PTCL [64–67]. However, this agent can cause significant infectious and hematologic toxicities that have led to early closure of some trials [65–67]. Recently, EBV-associated B-cell lymphomas have been reported complicating alemtuzumab therapy especially when

TABLE 3: Studies on High-Dose Therapy and Allogeneic Stem Cell Transplantation in PTCL.

(a) Retrospective data														
Author	Year	n	Age	Histologies	IPI	Previous autoSCT	Regimen (MA versus RIC)	Status at Tx	DFS/EFS/PFS	OS	TRM/NRM	Followup (months)	GVHD	
Feyler et al. [40]	2007	18	28	PTCLu 50%										
				T-cell leuk. 28%	0/1 33%	11%	100% MA	No data	33% (3 y)	39% (3 y)	38%*	57	Acute GVHD °3/4: 28%	
				ALCL 17%	≥2 66%								Ext. chronic GVHD: 6%	
Murashige et al. [51]	2005	28	38	NK/T 79%										
				Blastic NK 11%	No data	32%	82% versus 18%	CR 57%	34% (2 y)	40% (2 y)	29%	34	Acute GVHD °3/4: 29%	
				NK-leukemia 11%									Ext. chronic GVHD: 11%	
Hamadani et al. [52]	2008	14	43	PTCLu 36%										
				AITL 28%	aaIPI	14%	57% versus 43%	CR 21%	31% (3 y)	35% (3 y)	28%*	34	Acute GVHD °3/4: 21%	
				ALCL 14%	0/1 57%			PR 35%					Chronic GVHD: 50%	
				NK/T 14%	2/3 42%									
				Other 7%										
Le Gouill et al. [53]	2008	77	36	PTCLu 35%										
				ALCL 35%	0/1 61%	25%	74% versus 26%	CR 40%	53% (5 y)	57% (5 y)	21%*	43	Acute GVHD °3/4: 21%	
				AITL 14%	≥2 32%			PR 30%						
Kyriakou et al. [54]	2009	45	48	Other 16%	Unknown 6%									
				AITL 100%	No data	33%	56% versus 45%	CR 27%	53% (3 y)	64% (3 y)	18%*	29	Acute GVHD °3/4: 11%	
								PR 22%					Ext. chronic GVHD: 24%	
(b) Prospective data														
Author	Year	n	Age	Histologies	IPI	Previous autoSCT	Regimen (MA versus RIC)	Status at Tx	DFS/EFS/PFS	OS	TRM/NRM	Followup (months)	GVHD	
Corradini et al. [55]	2004	17	41	PTCLu 53%										
				ALCL 24%	aaIPI	47%	100% RIC	CR 12%	64% (3 y)	81% (3 y)	6%	28	Acute GVHD °3/4 : 12%	
				AITL 24%	0/1 24%			PR 71%					Ext. chronic GVHD: 6%	
Wulff et al. [56]	2005	10	45	PTCLu 40%										
				ALCL 30%	No data	20%	100% RIC	CR 10%	60% (7 m)	70% (7 m)	30%	7	Acute GVHD °3/4 : 10%	
				AITL 20%				PR 50%					Ext. chronic GVHD: 50%	
				T-PLL 10%										

* at day 100.

given in higher dosages [68, 69]. To better define the role of chemoimmunotherapy in the concept of HDT-autoSCT, the Nordic Lymphoma Group is conducting a multicenter randomized trial using dose-dense chemotherapy induction with or without alemtuzumab.

Allogeneic stem cell transplantation could offer a curative option in younger patients. However, the experience with this approach is sparse and limited to relapsed and refractory PTCL. In the prospective series nonmyeloablative conditioning protocols were used with very encouraging results especially in the Italian study. Furthermore, a GVL effect could be demonstrated. Taken together, the current data support the concept of alloSCT in eligible patients with relapsing chemosensitive PTCL, especially after failing prior HDT-autoSCT.

To further investigate the role of stem cell transplantation in previously untreated PTCL, this year the DSHNHL in cooperation with other groups will initiate a prospective randomised multicenter trial comparing upfront autoSCT versus alloSCT following dose-dense induction chemotherapy.

Abbreviations

AaIPI:	Age-adjusted IPI
ALCL:	Anaplastic large cell lymphoma
ALK:	Anaplastic lymphoma kinase
AlloSCT:	Allogeneic stem cell transplantation
AITL:	Angioimmunoblastic T-cell lymphoma
BEAC:	BCNU, etoposide, cytarabin, cyclophosphamide
BEAM:	BCNU, etoposide, cytarabin, melphalan
BCNU:	Carmustine
BEC:	Busulfan, etoposide, cyclophosphamide
CR:	Complete remission
CVB:	BCNU, etoposide, cyclophosphamide
DSF:	Disease-free survival
EATL:	Enteropathy-associated T-cell lymphoma
EFS:	Event-free survival
Eto:	Etoposide
IPI:	International prognostic index
M:	Months
Mel:	Melphalan
NK/T:	Natural killer-cell/T-cell leukemia/lymphoma
OS:	Overall survival
PFS:	Progression-free survival
PR:	Partial remission
PTCLu:	Peripheral T-cell lymphoma, unspecified
Thio:	Thiotepa
TRM:	Treatment-related mortality
Tx:	Transplantation
W:	Weeks
Y:	Year
CHOEP:	Cyclophosphamide, vincristine, doxorubicin, etoposide, prednisone
CTCL:	Cutaneous T-cell lymphoma
CVB/CBV:	BCNU, etoposide, cyclophosphamide
Cy:	Cyclophosphamide

HDT-autoSCT:	High-dose therapy with autologous stem cell transplantation
HSTL:	Hepatosplenic T-cell lymphoma
MCEC:	Ranimustine, cyclophosphamide, etoposide, carboplatin
Mito:	Mitoxantrone
TBI:	Total body irradiation
AutoSCT:	Autologous stem cell transplantation
Ext:	Extensive
GVHD:	Graft-versus-host disease
Leuk:	Leukemia
MA:	Myeloablative conditioning
NK:	Natural killer cell
NRM:	Nonrelapse mortality
RIC:	Reduced intensity conditioning
T-PLL:	T-cell prolymphocytic leukaemia.

Conflict of Interests

The author indicated no potential conflicts of interest.

References

- [1] S. Ascani, P. L. Zinzani, F. Gherlinzoni et al., "Peripheral T-cell lymphomas. Clinico-pathologic study of 168 cases diagnosed according to the R.E.A.L. classification," *Annals of Oncology*, vol. 8, no. 6, pp. 583–592, 1997.
- [2] J. O. Armitage, "A clinical evaluation of the International Lymphoma Study Group classification of non-Hodgkin's lymphoma," *Blood*, vol. 89, no. 11, pp. 3909–3918, 1997.
- [3] T. Rüdiger, D. D. Weisenburger, J. R. Anderson et al., "Peripheral T-cell lymphoma (excluding anaplastic large-cell lymphoma): results from the non-Hodgkin's lymphoma classification project," *Annals of Oncology*, vol. 13, no. 1, pp. 140–149, 2002.
- [4] B. Coiffier, N. Brousse, M. Peuchmaur et al., "Peripheral T-cell lymphomas have a worse prognosis than B-cell lymphomas: a prospective study of 361 immunophenotyped patients treated with the LNH-84 regimen," *Annals of Oncology*, vol. 1, no. 1, pp. 45–50, 1990.
- [5] J. Vose, J. Armitage, and D. Weisenburger, "International peripheral T-cell and natural killer/T-cell lymphoma study: pathology findings and clinical outcomes international T-cell lymphoma project," *Journal of Clinical Oncology*, vol. 26, no. 25, pp. 4124–4130, 2008.
- [6] J. O. Armitage, J. M. Vose, J. Linder et al., "Clinical significance of immunophenotype in diffuse aggressive non-Hodgkin's lymphoma," *Journal of Clinical Oncology*, vol. 7, no. 12, pp. 1783–1790, 1989.
- [7] C. Gisselbrecht, P. Gaulard, E. Lepage et al., "Prognostic significance of T-cell phenotype in aggressive non-Hodgkin's lymphomas. Groupe d'Etudes des Lymphomes de l'Adulte (GELA)," *Blood*, vol. 92, no. 1, pp. 76–82, 1998.
- [8] A. Melnyk, A. Rodríguez, W. C. Pugh, and F. Cabannillas, "Evaluation of the revised European-American lymphoma classification confirms the clinical relevance of immunophenotype in 560 cases of aggressive non-Hodgkin's lymphoma," *Blood*, vol. 89, no. 12, pp. 4514–4520, 1997.
- [9] S. M. Lippman, T. P. Miller, C. M. Spier, D. J. Slymen, and T. M. Grogan, "The prognostic significance of the immunotype

- in diffuse large-cell lymphoma: a comparative study of the T-cell and B-cell phenotype," *Blood*, vol. 72, no. 2, pp. 436–441, 1988.
- [10] S. M. Ansell, T. M. Habermann, P. J. Kurtin et al., "Predictive capacity of the International Prognostic Factor Index in patients with peripheral T-cell lymphoma," *Journal of Clinical Oncology*, vol. 15, no. 6, pp. 2296–2301, 1997.
 - [11] F. Zaja, D. Russo, F. Silvestri et al., "Retrospective analysis of 23 cases with peripheral T-cell lymphoma, unspecified: clinical characteristics and outcome," *Haematologica*, vol. 82, no. 2, pp. 171–177, 1997.
 - [12] R. Sonnen, W.-P. Schmidt, H. Konrad Müller-Hermelink, and N. Schmitz, "The International Prognostic Index determines the outcome of patients with nodal mature T-cell lymphomas," *British Journal of Haematology*, vol. 129, no. 3, pp. 366–372, 2005.
 - [13] A. Gallamini, C. Stelitano, R. Calvi et al., "Peripheral T-cell lymphoma unspecified (PTCL-U): a new prognostic model from a retrospective multicentric clinical study," *Blood*, vol. 103, no. 7, pp. 2474–2479, 2004.
 - [14] P. Went, C. Agostinelli, A. Gallamini et al., "Marker expression in peripheral T-cell lymphoma: a proposed clinical-pathologic prognostic score," *Journal of Clinical Oncology*, vol. 24, no. 16, pp. 2472–2479, 2006.
 - [15] J. Rodríguez, E. Conde, A. Gutiérrez et al., "The adjusted International Prognostic Index and β -2-microglobulin predict the outcome after autologous stem cell transplantation in relapsing/refractory peripheral T-cell lymphoma," *Haematologica*, vol. 92, no. 8, pp. 1067–1074, 2007.
 - [16] J. Dupuis, J.-F. Emile, N. Mounier et al., "Prognostic significance of Epstein-Barr virus in nodal peripheral T-cell lymphoma, unspecified: a Groupe d'Etude des Lymphomes de l'Adulte (GELA) study," *Blood*, vol. 108, no. 13, pp. 4163–4169, 2006.
 - [17] R. D. Gascoyne, P. Aoun, D. Wu et al., "Prognostic significance of anaplastic lymphoma kinase (ALK) protein expression in adults with anaplastic large cell lymphoma," *Blood*, vol. 93, no. 11, pp. 3913–3921, 1999.
 - [18] B. Falini, S. Pileri, P. L. Zinzani et al., "ALK⁺ lymphoma: clinico-pathological findings and outcome," *Blood*, vol. 93, no. 8, pp. 2697–2706, 1999.
 - [19] H. Q. Huang, Y. L. Peng, X. B. Lin et al., "Clinical outcomes of 106 patients with peripheral T-cell lymphoma treated by standard CHOP regimen," *Ai Zheng*, vol. 23, no. 11, pp. 1443–1447, 2004.
 - [20] K. Kim, W. S. Kim, C. W. Jung et al., "Clinical features of peripheral T-cell lymphomas in 78 patients diagnosed according to the Revised European-American lymphoma (REAL) classification," *European Journal of Cancer*, vol. 38, no. 1, pp. 75–81, 2002.
 - [21] A. López-Guillermo, J. Cid, A. Salar et al., "Peripheral T-cell lymphomas: initial features, natural history, and prognostic factors in a series of 174 patients diagnosed according to the R.E.A.L. Classification," *Annals of Oncology*, vol. 9, no. 8, pp. 849–855, 1998.
 - [22] J. M. Vose, C. Peterson, P. J. Bierman et al., "Comparison of high-dose therapy and autologous bone marrow transplantation for T-cell and B-cell non-Hodgkin's lymphomas," *Blood*, vol. 76, no. 2, pp. 424–431, 1990.
 - [23] R. Fanin, M. C. Ruiz de Elvira, A. Sperotto, M. Baccarani, and A. Goldstone, "Autologous stem cell transplantation for T and null cell CD30-positive anaplastic large cell lymphoma: analysis of 64 adult and paediatric cases reported to the European Group for Blood and Marrow Transplantation (EBMT)," *Bone Marrow Transplantation*, vol. 23, no. 5, pp. 437–442, 1999.
 - [24] J. Rodríguez, M. Munsell, S. Yazji et al., "Impact of high-dose chemotherapy on peripheral T-cell lymphomas," *Journal of Clinical Oncology*, vol. 19, no. 17, pp. 3766–3770, 2001.
 - [25] A. K. Blystad, G. Enblad, S. Kvaløy et al., "High-dose therapy with autologous stem cell transplantation in patients with peripheral T cell lymphomas," *Bone Marrow Transplantation*, vol. 27, no. 7, pp. 711–716, 2001.
 - [26] K. W. Song, P. Mollee, A. Keating, and M. Crump, "Autologous stem cell transplant for relapsed and refractory peripheral T-cell lymphoma: variable outcome according to pathological subtype," *British Journal of Haematology*, vol. 120, no. 6, pp. 978–985, 2003.
 - [27] J. Rodríguez, M. D. Caballero, A. Gutiérrez et al., "High-dose chemotherapy and autologous stem cell transplantation in peripheral T-cell lymphoma: the GEL-TAMO experience," *Annals of Oncology*, vol. 14, no. 12, pp. 1768–1775, 2003.
 - [28] J. Schetelig, S. Fetscher, A. Reichle et al., "Long-term disease-free survival in patients with angioimmunoblastic T-cell lymphoma after high-dose chemotherapy and autologous stem cell transplantation," *Haematologica*, vol. 88, no. 11, pp. 1272–1278, 2003.
 - [29] K. W. Zamkoff, M. D. Matulis, A. C. Mehta, M. W. Beaty, R. E. Hutchison, and T. C. Gentile, "High-dose therapy and autologous stem cell transplant does not result in long-term disease-free survival in patients with recurrent chemotherapy-sensitive ALK-negative anaplastic large-cell lymphoma," *Bone Marrow Transplantation*, vol. 33, no. 6, pp. 635–638, 2004.
 - [30] E. Jantunen, T. Wiklund, E. Juvonen et al., "Autologous stem cell transplantation in adult patients with peripheral T-cell lymphoma: a nation-wide survey," *Bone Marrow Transplantation*, vol. 33, no. 4, pp. 405–410, 2004.
 - [31] M. Jagasia, D. Morgan, S. Goodman et al., "Histology impacts the outcome of peripheral T-cell lymphomas after high dose chemotherapy and stem cell transplant," *Leukemia and Lymphoma*, vol. 45, no. 11, pp. 2261–2267, 2004.
 - [32] T. Kewalramani, A. D. Zelenetz, J. Teruya-Feldstein et al., "Autologous transplantation for relapsed or primary refractory peripheral T-cell lymphoma," *British Journal of Haematology*, vol. 134, no. 2, pp. 202–207, 2006.
 - [33] M. K. Kim, S. Kim, S. S. Lee et al., "High-dose chemotherapy and autologous stem cell transplantation for peripheral T-cell lymphoma: complete response at transplant predicts survival," *Annals of Hematology*, vol. 86, no. 6, pp. 435–442, 2007.
 - [34] S. D. Smith, B. J. Bolwell, L. A. Rybicki et al., "Autologous hematopoietic stem cell transplantation in peripheral T-cell lymphoma using a uniform high-dose regimen," *Bone Marrow Transplantation*, vol. 40, no. 3, pp. 239–243, 2007.
 - [35] A. I. Chen, A. McMillan, R. S. Negrin, S. J. Horning, and G. G. Laport, "Long-term results of autologous hematopoietic cell transplantation for peripheral T cell lymphoma: the Stanford experience," *Biology of Blood and Marrow Transplantation*, vol. 14, no. 7, pp. 741–747, 2008.
 - [36] J. Lee, W.-Y. Au, M. J. Park et al., "Autologous hematopoietic stem cell transplantation in extranodal natural killer/T cell lymphoma: a multinational, multicenter, matched controlled study," *Biology of Blood and Marrow Transplantation*, vol. 14, no. 12, pp. 1356–1364, 2008.
 - [37] D.-H. Yang, W. S. Kim, S. J. Kim et al., "Prognostic factors and clinical outcomes of high-dose chemotherapy followed by autologous stem cell transplantation in patients with

- peripheral T cell lymphoma, unspecified: complete remission at transplantation and the prognostic index of peripheral T cell lymphoma are the major factors predictive of outcome," *Biology of Blood and Marrow Transplantation*, vol. 15, no. 1, pp. 118–125, 2009.
- [38] J. Rodríguez, E. Conde, A. Gutiérrez et al., "Prolonged survival of patients with angioimmunoblastic T-cell lymphoma after high-dose chemotherapy and autologous stem cell transplantation. The GELTAMO experience," *European Journal of Haematology*, vol. 78, no. 4, pp. 290–296, 2007.
 - [39] J. Rodríguez, E. Conde, A. Gutiérrez et al., "The results of consolidation with autologous stem-cell transplantation in patients with peripheral T-cell lymphoma (PTCL) in first complete remission: the Spanish Lymphoma and Autologous Transplantation Group experience," *Annals of Oncology*, vol. 18, no. 4, pp. 652–657, 2007.
 - [40] S. Feyler, H. M. Prince, R. Pearce et al., "The role of high-dose therapy and stem cell rescue in the management of T-cell malignant lymphomas: a BSBMT and ABMTRR study," *Bone Marrow Transplantation*, vol. 40, no. 5, pp. 443–450, 2007.
 - [41] C. Kyriakou, C. Canals, A. Goldstone et al., "High-dose therapy and autologous stem-cell transplantation in angioimmunoblastic lymphoma: complete remission at transplantation is the major determinant of outcome- Lymphoma Working Party of the European Group for blood and marrow transplantation," *Journal of Clinical Oncology*, vol. 26, no. 2, pp. 218–224, 2008.
 - [42] V. Prochazka, E. Faber, L. Raida et al., "Prolonged survival of patients with peripheral T-cell lymphoma after first-line intensive sequential chemotherapy with autologous stem cell transplantation," *Biomedical Papers*, vol. 153, no. 1, pp. 63–66, 2009.
 - [43] A. Numata, T. Miyamoto, Y. Ohno et al., "Long-term outcomes of autologous PBSCT for peripheral T-cell lymphoma: retrospective analysis of the experience of the Fukuoka BMT group," *Bone Marrow Transplantation*, vol. 45, no. 2, pp. 311–316, 2010.
 - [44] N. Mounier, C. Gisselbrecht, J. Brière et al., "Prognostic factors in patients with aggressive non-Hodgkin's lymphoma treated by front-line autotransplantation after complete remission: a cohort study by the Groupe d'Etude des Lymphomes de l'Adulte," *Journal of Clinical Oncology*, vol. 22, no. 14, pp. 2826–2834, 2004.
 - [45] M. Nickelsen, M. Ziepert, S. Zeinalova et al., "High-dose CHOP plus etoposide (MegaCHOEP) in T-cell lymphoma: a comparative analysis of patients treated within trials of the German High-Grade Non-Hodgkin Lymphoma Study Group (DSHNHL)," *Annals of Oncology*, vol. 20, no. 12, pp. 1977–1984, 2009.
 - [46] P. Corradini, C. Tarella, F. Zallio et al., "Long-term follow-up of patients with peripheral T-cell lymphomas treated up-front with high-dose chemotherapy followed by autologous stem cell transplantation," *Leukemia*, vol. 20, no. 9, pp. 1533–1538, 2006.
 - [47] J. Rodríguez, E. Conde, A. Gutiérrez et al., "Frontline autologous stem cell transplantation in high-risk peripheral T-cell lymphoma: a prospective study from the Gel-Tamo Study Group," *European Journal of Haematology*, vol. 79, no. 1, pp. 32–38, 2007.
 - [48] S. Mercadal, J. Briones, B. Xicoy et al., "Intensive chemotherapy (high-dose CHOP/ESHAP regimen) followed by autologous stem-cell transplantation in previously untreated patients with peripheral T-cell lymphoma," *Annals of Oncology*, vol. 19, no. 5, pp. 958–963, 2008.
 - [49] P. Reimer, T. Rüdiger, E. Geissinger et al., "Autologous stem-cell transplantation as first-line therapy in peripheral t-cell lymphomas: results of a prospective multicenter study," *Journal of Clinical Oncology*, vol. 27, no. 1, pp. 106–113, 2009.
 - [50] F. D'Amore, T. Relander, G. Lauritszen et al., "Dose-dense induction followed by autologous stem cell transplant (ASCT) as 1st line treatment in peripheral T-cell lymphomas (PTCL)—a phase II study of the Nordic Lymphoma Group (NLG)," *Haematologica*, vol. 94, supplement 2, p. 437, 2009.
 - [51] N. Murashige, M. Kami, Y. Kishi et al., "Allogeneic haematopoietic stem cell transplantation as a promising treatment for natural killer-cell neoplasms," *British Journal of Haematology*, vol. 130, no. 4, pp. 561–567, 2005.
 - [52] M. Hamadani, F. T. Awan, P. Elder et al., "Allogeneic hematopoietic stem cell transplantation for peripheral T cell lymphomas; evidence of graft-versus-T cell lymphoma effect," *Biology of Blood and Marrow Transplantation*, vol. 14, no. 4, pp. 480–483, 2008.
 - [53] S. Le Gouill, N. Milpied, A. Buzyn et al., "Graft-versus-lymphoma effect for aggressive T-cell lymphomas in adults: a study by the Société Française de Greffe de Moëlle et de Thérapie Cellulaire," *Journal of Clinical Oncology*, vol. 26, no. 14, pp. 2264–2271, 2008.
 - [54] C. Kyriakou, C. Canals, J. Finke et al., "Allogeneic stem cell transplantation is able to induce long-term remissions in angioimmunoblastic T-cell lymphoma: a retrospective study from the Lymphoma Working Party of the European Group for Blood and Marrow Transplantation," *Journal of Clinical Oncology*, vol. 27, no. 24, pp. 3951–3958, 2009.
 - [55] P. Corradini, A. Doderio, F. Zallio et al., "Graft-versus-lymphoma effect in relapsed peripheral T-cell non-Hodgkin's lymphomas after reduced-intensity conditioning followed by allogeneic transplantation of hematopoietic cells," *Journal of Clinical Oncology*, vol. 22, no. 11, pp. 2172–2176, 2004.
 - [56] G. G. Wulf, J. Hasenkamp, W. Jung, B. Chapuy, L. Truemper, and B. Glass, "Reduced intensity conditioning and allogeneic stem cell transplantation after salvage therapy integrating alemtuzumab for patients with relapsed peripheral T-cell non-Hodgkin's lymphoma," *Bone Marrow Transplantation*, vol. 36, no. 3, pp. 271–273, 2005.
 - [57] C. Haioun, E. Lepage, C. Gisselbrecht et al., "Survival benefit of high-dose therapy in poor-risk aggressive non-Hodgkin's lymphoma: final analysis of the prospective LNH87-2 protocol—a groupe d'étude des lymphomes de l'adulte study," *Journal of Clinical Oncology*, vol. 18, no. 16, pp. 3025–3030, 2000.
 - [58] N. Mounier, D. Simon, C. Haioun, P. Gaulard, C. Gisselbrecht, and I. Khouri, "Impact of high-dose chemotherapy on peripheral T-cell lymphomas," *Journal of Clinical Oncology*, vol. 20, no. 5, pp. 1426–1427, 2002.
 - [59] C. Gisselbrecht, E. Lepage, T. Molina et al., "Shortened first-line high-dose chemotherapy for patients with poor-prognosis aggressive lymphoma," *Journal of Clinical Oncology*, vol. 20, no. 10, pp. 2472–2479, 2002.
 - [60] N. Mounier, C. Gisselbrecht, J. Brière et al., "All aggressive lymphoma subtypes do not share similar outcome after front-line autotransplantation: a matched-control analysis by the Groupe d'Etude des Lymphomes de l'Adulte (GELA)," *Annals of Oncology*, vol. 15, no. 12, pp. 1790–1797, 2004.
 - [61] B. S. Sohn, I. Park, E. K. Kim et al., "Comparison of clinical outcome after autologous stem cell transplantation between patients with peripheral T-cell lymphomas and diffuse large B-cell lymphoma," *Bone Marrow Transplantation*, vol. 44, no. 5, pp. 287–293, 2009.

- [62] M. P. Escalón, N. S. Liu, Y. Yang et al., "Prognostic factors and treatment of patients with T-cell non-Hodgkin lymphoma: the M. D. Anderson Cancer Center experience," *Cancer*, vol. 103, no. 10, pp. 2091–2098, 2005.
- [63] N. Schmitz, L. Trümper, M. Ziepert et al., "Treatment and prognosis of mature T-cell and NK-cell lymphoma: an analysis of patients with T-cell lymphoma treated in studies of the German High-Grade Non-Hodgkin's Lymphoma Study Group (DSHNHL)," *Blood*, vol. 166, no. 18, pp. 3418–3425, 2010.
- [64] L. H. Trümper, K. Hohloch, M. Kloess et al., "CHOP/CHOEP-14 followed by consolidation with alemtuzumab in untreated aggressive T-cell lymphomas (DSHNHL 2003-1) and toxicity of a phase II trial of the German High Grade Non-Hodgkin's Lymphoma Group DSHNHL," *Journal of Clinical Oncology*, vol. 24, no. 18s, p. 431: abstract 7538, 2006.
- [65] J. G. Kim, S. K. Sohn, Y. S. Chae et al., "Alemtuzumab plus CHOP as front-line chemotherapy for patients with peripheral T-cell lymphomas: a phase II study," *Cancer Chemotherapy and Pharmacology*, vol. 60, no. 1, pp. 129–134, 2007.
- [66] A. Gallamini, F. Zaja, C. Patti et al., "Alemtuzumab (Campath-1H) and CHOP chemotherapy as first-line treatment of peripheral T-cell lymphoma: results of a GITIL (Gruppo Italiano Terapie Innovative nei Linfomi) prospective multicenter trial," *Blood*, vol. 110, no. 7, pp. 2316–2323, 2007.
- [67] E. Weidmann, G. Hess, K. U. Chow et al., "A phase II study of alemtuzumab, fludarabine, cyclophosphamide, and doxorubicin (Campath-FCD) in peripheral T-cell lymphomas," *Leukemia and Lymphoma*, vol. 51, no. 3, pp. 447–455, 2010.
- [68] K. C. Weisel, E. Weidmann, I. Anagnostopoulos, L. Kanz, A. Pezzutto, and M. Subklewe, "Epstein-Barr virus-associated B-cell lymphoma secondary to FCD-C therapy in patients with peripheral T-cell lymphoma," *International Journal of Hematology*, vol. 88, no. 4, pp. 434–440, 2008.
- [69] H. C. Kluin-Nelemans, J. L. Coenen, J. E. Boers, G. W. Van Imhoff, and S. Rosati, "EBV-positive immunodeficiency lymphoma after alemtuzumab-CHOP therapy for peripheral T-cell lymphoma," *Blood*, vol. 112, no. 4, pp. 1039–1041, 2008.