

## Research Article

# Polynomial Roots and Calabi-Yau Geometries

**Yang-Hui He<sup>1, 2, 3</sup>**

<sup>1</sup> School of Physics, Nankai University, Tianjin 300071, China

<sup>2</sup> Department of Mathematics, City University London, Northampton Square, London EC1V 0HB, UK

<sup>3</sup> Merton College, University of Oxford, Oxford OX1 4JD, UK

Correspondence should be addressed to Yang-Hui He, hey@maths.ox.ac.uk

Received 13 October 2010; Accepted 27 April 2011

Academic Editor: André Lukas


Copyright © 2011 Yang-Hui He. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The examination of roots of constrained polynomials dates back at least to Waring and to Littlewood. However, such delicate structures as fractals and holes have only recently been found. We study the space of roots to certain integer polynomials arising naturally in the context of Calabi-Yau spaces, notably Poincaré and Newton polynomials, and observe various salient features and geometrical patterns.

## 1. Introduction and Summary

The subject of roots of monovariate polynomials is, without doubt, an antique one and has germinated an abundance of fruitful research over the ages. It is, therefore, perhaps surprising that any new statements could at all be made regarding such roots. The advent of computer algebra, chaotic phenomena, and random ensembles has, however, indeed shed new light upon so ancient a metier.

Polynomials with constrained coefficients and form, though permitted to vary randomly, have constituted a vast field itself. As far back as 1782, Edward Waring, in relation to his famous problem on power summands, had shown that for cubic polynomials with random real coefficients, the ratio of the probability of finding nonreal zeros versus that of not finding non-real zeros is less than or equal to 2. Constraining the coefficients to be integers within a fixed range has, too, its own history. It was realised in [1] that a degree  $n$  random polynomial  $P(z) = 1 + \sum_{k=1}^n a_k z^k$  with  $a_k = -1, 0, 1$  distributed evenly, the expected number  $\nu_n$  of real roots is of order  $\mathcal{O}(n^{1/2})$  asymptotically in  $n$ . This was furthered by [2] to be essentially independent of the statistics, in that  $\nu_n$  has the same asymptotics (cf. also [3, 4]), as much for  $a_k$  being evenly distributed real numbers, in  $[-1, 1]$ , or as Gaussian distributed in  $(-\infty, \infty)$ .



**Figure 1:** The position, on the complex plane, of the zeros of 50000 random integer polynomials with coefficients  $-1, 0, \text{ or } 1$ , for degrees up to 4 in part (a), 6 in part (b), and 10 in part (c).

Continual development ensued (q.v. also [5]), notably by Littlewood [6], Erdős and Turán [7], Hammersley [8], and Kac [9]. Indeed, a polynomial with coefficients only taking values as  $\pm 1$  has come to be known as a *Littlewood polynomial*, and the *Littlewood Problem* asks for the precise asymptotics, in the degree, of such polynomials taking values, with complex arguments, on the unit circle. The classic work of Montgomery [10] and Odlyzko [11], constituting one of the most famous computer experiments in mathematics (q. v. Section 3.1 of [12] for some recent remarks on the distributions), empirically showed that the distribution of the (normalized) spacings between successive critical zeros of the Riemann zeta function is the same as that of a Gaussian unitary ensemble of random matrices, whereby infusing our subject with issues of uttermost importance.

Subsequently, combining the investigation of zeros and of random polynomials, Odlyzko and Poonen [13] studied the zeros of Littlewood-type polynomials by setting the coefficients to 0 and 1; they provided certain bounds as well as found interesting fractal structures. Thus, inspired and with the rapid advance of computational power, Borwein et al. constructed various plots of zeros of constrained random polynomials and many remarkable features were instantly visible [14, 15]. We can readily demonstrate this with Mathematica [16], as is shown in Figure 1. In the figure, we take a sample of 50000 random polynomials with coefficients  $-1, 0, \text{ or } 1$  up to various degrees, and plot, on the complex plane, their zeros. Not only do we see fractal behaviour (cf. discussions in [3, 17] on chaotic dualities in field theories) near the boundaries, the nature of the holes are intimately related [18] to the Lehmer-Mahler Conjecture: that the Mahler measure  $M(P) := \exp((1/2\pi) \int_0^{2\pi} \log |P(e^{i\theta})| d\theta)$  of any integral polynomial  $P(z)$  (which is not a multiple of cyclotomic polynomials) should be bounded below by that of  $z^{10} - z^9 + z^7 - z^6 + z^5 - z^4 + z^3 - z + 1$ , which is approximately 1.17.

High resolution variants of Figure 1 have been considered recently by Christensen [19], Jörgenson [20] and Derbyshire and Matson [21], inter alia, and many beautiful pictures can be found (cf. also a nice account in [22]). Particular striking are the coloured density plots in [21].

An interesting query, in somewhat reverse direction to the above line of thought, was posed in [23]: recalling that the Lee-Yang Circle Theorem placed severe constraints on the generating function of the partition function of the Ising Model, the author asked if one could statistically test whether a given Laurent polynomial could, in fact, be the Jones polynomial of a knot. Using a landscape of knots generated by the programme “knotscape” [24], the said work investigated many distribution properties of the zeros of known Jones polynomials.

Enchanted by this motif which has threaded varying developments over the decades while persistently generating new perspectives, a question immediately springs to mind. One of the central topics of both modern mathematics and theoretical physics is undoubtedly that of Calabi-Yau geometries. A key feature is the superabundance thereof. In complex dimension one, there is only the torus; in dimension two, there are the 4-torus and the K3-surface; however, for dimension three and above, no classification is known, and already a plethora has been constructed. The first database was that of the so-called CICYs, or, complete intersection Calabi-Yau threefolds in products of projective spaces [25] as well as hypersurfaces in weighted  $\mathbb{C}P^4$  [26], by Candelas et al. Then, over more than a decade, Kreuzer and Skarke formulated and compiled an impressive list of on the order of  $10^{10}$  threefolds as hypersurfaces in toric varieties [27]. Finding new patterns in this vast distribution of manifolds has seen some recent activity [28–30]. Indeed, the multitude of these geometries is at the core of the so-called vacuum degeneracy problem of superstring theory and constitutes a part of the landscape issue [31].

Along a parallel vein, the space of noncompact (singular) Calabi-Yau spaces (as affine varieties) has also been extensively explored, notably by Hanany et al. over the past decade [32–39], especially those which admit a toric description [33]; the discovery of their intimate relation to dimer models and brane tilings [34, 35, 38] has also generated some excitement. These supplant yet another corner in the landscape of geometries and associated supersymmetric vacua.

Thus motivated, many tasks lend themselves to automatic investigation; we here give a precis of some key points. In Section 2, we begin with the compact, smooth Calabi-Yau manifolds. As mentioned above, there had been much effort in classifying and constructing these, especially in complex dimensions three and four. An immediate polynomial, of constrained form and integer coefficients, and yet succinctly encoding some topological information, is the Poincaré polynomial, which can be readily written in terms of the Hodge numbers. We take the “experimental data” of all the known Hodge numbers of the threefolds and fourfolds spanning two decades of work and plot the complex roots of the associated Poincaré polynomials in Figures 4 and 8, respectively.

Much intricate structures are clearly visible. These are then contrasted with a “standard background sample”, namely, the roots of random integer sextic and octic polynomials, with unit leading coefficient and vanishing linear term, drawn in Figures 2 and 7. From such collections are extracted the subclass of those which admit  $-1$  as roots, which, by a theorem from differential geometry, correspond to spaces which have more than one isometry. Interestingly, they correspond to self-mirror threefolds and “quasi-”self-mirror fourfolds. We plot the roots for these in Figures 5 and 10, and see that they furnish certain substrata of the conglomerate plots mentioned above.

Thenceforth, we move on to noncompact Calabi-Yau geometries in Section 3. There, too, is a plenitude of examples, most notably those which are toric. We focus on toric threefolds because these have planar toric diagrams as lattice points in  $\mathbb{Z}^2$  due to the Calabi-Yau condition. Once more, a natural polynomial invites itself: the Newton polynomial. The Riemann surface corresponding to this bivariate polynomial has been a central subject to the gauge and brane theories in the context of string theory. Moreover, the two important projections thereof, namely, the amoeba and alga projections, have provided many beautiful Monte Carlo plots, illustrating deep algebraic geometry as well as gauge theory. Because we are confronted with two complex variables, we need to slightly deviate from our theme of complex roots; instead, we find it expedient to regard the variables as real and consider

the real projection of the Riemann surface. Subsequently, we can study the ensemble of real turning (critical) points of these planar curves.

Again, we resort to “actual data” and focus on the most well-known affine toric threefold geometries—as shown in Figure 11—corresponding to local Calabi-Yau singularities, including, of course, the famous conifold. To each space, we find the collective of critical points in  $\mathbb{R}^2$  as we vary the integer coefficients—commonly known as multiplicities in the dimer model interpretation—of the Newton polynomial, and plot them in Figure 13. We see a sensitive dependence of the emergent subtle structures upon the choice of toric data.

In many respects, we have taken a very pragmatic and empirical approach toward the data accumulated over many years of theoretical research, of quantity large enough to justify experimentation. To this philosophy of “experimental mathematics” we adhere throughout, observe wherever we should, and infer wherever we may. Without much to do, therefore, let us delve into the details of the issues summarized above.

## 2. Compact Calabi-Yau Manifolds, Poincaré Polynomials, and Complex Roots

An important quantifying polynomial for a smooth compact manifold  $X$  is the Poincaré polynomial, which is a generating function for topological invariants of  $X$  (say of dimension  $n$ ):

$$P(t; X) = \sum_{i=0}^n b_i t^i, \quad (2.1)$$


with the  $b_i$  being the  $i$ th Betti number. Indeed, this seems a more natural candidate for our present studies than some because other famous polynomials such as the Hilbert polynomial or the numerator of the Hilbert series (which of late have been instrumental in counting BPS operators [40–42]) are not topological invariants and depend on the specific projective embedding. Furthermore, by definition, at  $t = -1$ , the polynomial evaluates to the Euler characteristic; this will be of significance shortly.

The zeros of the Poincaré polynomial have rather remarkable properties. It was conjectured that [43] that if the rank of the manifold  $X$  is greater than 1, where rank is defined to be the maximal number of everywhere independent, mutually commuting, vector fields on  $X$ , that is, the number of isometries, then  $-1$  is a multiple root of the Poincaré polynomial of  $X$ . Unfortunately, this conjecture was shown to be false [44]. Nevertheless, it still holds that the rank of  $X$  exceeds unity if and only if  $-1$  is a multiple root of  $P(t; X)$ .

Moreover, of number theoretic and arithmo-geometric significance is the fact that certain Poincaré polynomials exhibit Riemann Hypothesis behaviour [45, 46], in analogy to the Hasse-Weil zeta local zeta functions. Recently, alignment of zeros of Hilbert polynomials have been studied by [47], in relation to zeta functions.

### 2.1. Calabi-Yau Threefolds

Our focus will be on (2.1). First, let us study the case of Calabi-Yau threefolds, which have been of the greatest interest, at least historically. Because we are dealing with complex (Kähler) manifolds, Hodge decomposition implies that  $b_i(X) = \sum_{p,q} h^{p,q}(X)$ , with  $h^{p,q}(X) = \dim H_{\bar{\partial}}^{p,q}(X)$  the dimensions of the Dolbeault cohomology groups. Indeed, for



**Figure 2:** (a) The position, on the complex plane, of the zeros of 50000 random integer degree six polynomials with coefficients between 0 and 1000. (b) The same, but with monic palindromic sextics. (c) Monic palindromic sextics, and with linear term vanishing.

(compact, smooth, connected) Calabi-Yau threefolds, the Hodge diamond, and subsequently the Betti numbers and the Poincaré polynomial, can be written as


$$\begin{array}{ccccccc}
 & & & & & & 1 \\
 & & & & & 0 & \\
 & & & 0 & & 0 & \\
 & 0 & & h^{1,1} & & 0 & \\
 1 & h^{2,1} & & h^{2,1} & & h^{2,1} & 1 \\
 & 0 & & h^{1,1} & & 0 & \\
 & & 0 & & 0 & & \\
 & & & & & & 1
 \end{array}
 \quad (b_0 = b_6, b_1 = b_5, b_2 = b_4, b_3) = (1, 0, h^{1,1}, 2 + 2h^{2,1}), \tag{2.2}$$

$$P(t; X) = 1 + h^{1,1}t^2 + (2 + 2h^{2,1})t^3 + h^{1,1}t^4 + t^6.$$

That the Poincaré polynomial is palindromic is obvious and follows from Poincaré duality. Therefore, our first constraint is palindromicity to which we will presently restrict. We recall the roots of a completely random sample of integer polynomials with coefficients in  $[-1, 1]$  up to the sextic in part (b) of Figure 1. In Figure 2, we plot, in part (a), a sample of 50000 random integer sextic polynomials with coefficients in  $[0, 1000]$  (making sure that the highest coefficient at degree 6 is not 0) as a comparative norm. Next, in part (b), we plot the same, but for monic palindromic sextics, that is,  $P(t) = 1 + b_1t + b_2t^2 + b_3t^3 + b_2t^4 + b_1t^5 + t^6$ . Then, in (c), we restrict once more, with some foresight, so that the linear term vanishes, that is,  $P(t) = 1 + b_2t^2 + b_3t^3 + b_2t^4 + t^6$ . We see that upon the condition of palindromicity, there is a marked emergence of roots on the unit circle; this of course arises from the symmetric terms of the form  $e^{it}$  combining to give (co)sines whose reality then facilitates the addition to zero. The symmetry about the  $x$ -axis is simply that all roots appear in conjugate pairs because our polynomials have real coefficients.

For our amusement, seeing the form of the semiunit-circular shape being prominent, we are reminded of the conformal map  $z \rightarrow (z/(z - 1))$  which takes the unit circle to the critical strip of the Riemann Hypothesis, as shown in detail by part (a) of Figure 3. We take the space of roots of monic palindromic sextics with vanishing linear/quintic terms from part (c) of Figure 2, apply the inverse map  $z \rightarrow (z/(z+1))$  to map to the critical strip, and redo the plot in part (b) of Figure 3. We see that the plot resembles the zero-free region of the Riemann zeta function inside the critical strip.

Founded upon these above discussions, we are ready to approach “actual data.” We now collect all known Calabi-Yau threefolds, which come from three major databases,



**Figure 3:** (a) The conformal map  $z \rightarrow (z/(z-1))$  takes the left half of the critical strip to the inside of the unit disk, with the boundaries mapped as shown by the arrows. It takes the mirror image, in the right half of the critical strip, to the complement of the unit disk. The inverse map is given by  $z \rightarrow (z/(z+1))$ . (b) The position of 50000 randomly integer monic palindromic sextic polynomials with vanishing linear/quintic terms and with coefficients ranging in  $[0, 1000]$ , applying the map  $z \rightarrow (z/(z+1))$ .

namely, the aforementioned CICYs, the hypersurfaces in toric varieties, as well as the collection of individually tailored ones of small Hodge numbers (cf. [28, 29]). These total, respectively, 30108, 266 and 54 distinct pairs of Hodge numbers  $(h^{1,1}(X), h^{2,1}(X))$ . In all, there are 30237 distinct pairs (of course, each with much degeneracy) of Hodge numbers; to our present knowledge, these are all the ones circulated in the literature.


We plot (Traditionally, the now-famous plot, first appearing in [26], is done with  $\chi = 2(h^{1,1} - h^{2,1})$  as abscissa and  $h^{1,1} + h^{2,1}$  as ordinate.) these, with  $h^{1,1}$  as the abscissa and  $h^{2,1}$ , ordinate, in part (a) of Figure 4, the largest amongst these is (491, 11). Note that because of mirror symmetry, there is a symmetry interchanging the two coordinates. It is still an open question whether there exists any Calabi-Yau threefold whose either Hodge number exceeds 491, a bound which has defied constructions so far. This is why in our random standard background sample in Figure 2, we have conveniently selected the largest integer coefficient to be  $1000 \sim 2 \cdot (491 + 1)$ . In part (b) of the said Figure, we plot, on the complex plane, the roots of the Poincaré polynomials of all these known threefolds. Because we are dealing with polynomials of nonnegative coefficients, there should be many generic roots with negative real parts. Comparing with the random sample in part (c) of Figure 2, we see a beautiful clustering of points in the first quadrant (and, by complex conjugation, the fourth).

Next, let us test for how many Poincaré polynomials  $-1$  is a root; these, as mentioned above, would correspond to manifolds which have more than one isometries. Interestingly, of the some 30000, only 148 pass the test. These turn out to be only the 148 known self-mirror manifolds; we plot their Hodge numbers in Part (a) of Figure 5 (the values of Hodge numbers range from (1, 1) to (251, 251), skipping many high values, as well as the number 13). Indeed, this is a simple consequence of palindromicity as one sees that, upon substituting  $t = -1$  into  $P(t; X)$  in (2.2), we obtain  $P(-1; X) = 2(h^{1,1} - h^{2,1}) = \chi(X)$ , the Euler characteristic. In Part (b), we plot all the other roots as well, and see that these constitute a portion of the small crescent around the origin.


## 2.2. Calabi-Yau Fourfolds

Having explored the space of Calabi-Yau threefolds, it is automatic to proceed onto the space of fourfolds, a relative *terra incognita*. We again resort to the wonderful database compiled by





**Figure 5:** (a) The Hodge number of self-mirror Calabi-Yau threefolds; these have  $-1$  as a root of the Poincaré polynomial. (b) All of the roots of the Poincaré polynomial of these self-mirrors.




**Figure 6:** (a) The Hodge numbers, with  $h^{1,1} - h^{3,1}$  as the abscissa and  $h^{1,1} + h^{3,1}$ , the ordinate, of the fourfolds from Kreuzer-Skarke's database. (b) The same, but with  $h^{1,1} - h^{2,1}$ , as the abscissa and  $h^{1,1} + h^{2,1}$  as the ordinate.


Following [49], we plot  $h^{1,1} + h^{3,1}$  as ordinate versus  $h^{1,1} - h^{3,1}$  as abscissa, which demonstrates mirror-like behaviour. (Though in *cit. ibid.*, only the hypersurfaces in weighted  $\mathbb{C}P^5$  were considered, whereas here we plot the entirety of the known fourfolds.) We also plot  $h^{1,1} + h^{2,1}$  versus  $h^{1,1} - h^{2,1}$ , showing that the behaviour in the application direction is rather trivial. These are shown in Figure 6.

We now repeat the experiment undertaken for threefolds. First, we plot the space of generic roots, and present them in Figure 7. In part (a), a sample of 50000 random integer octic polynomials with coefficients in  $[0, 2500000]$  (making sure that the highest coefficient at degree 8 is not 0) as a comparative basis: octic, since we will be contrasting with degree 8 Poincaré polynomials, upper bound of 2500000, since we can see from Figure 6 and (2.3), that the Min and Max of the Hodge numbers are, respectively,  $(h^{1,1}, h^{2,1}, h^{3,1}) \in ([1, 303148], [0, 2010], [1, 3030148])$ , so that the upper bound to the  $b_4$  term is 2425228. Next, in part (b), we plot the same, but for monic palindromic octics, that is,  $P(t) = 1 + b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_3t^5 + b_2t^6 + b_1t^7 + t^8$ . Finally, in (c), we restrict once more, so that the linear term vanishes, that is,  $P(t) = 1 + b_2t^2 + b_3t^3 + b_4t^4 + b_3t^5 + b_2t^6 + t^8$ .

In contradistinction to these generic results, we can now find all the complex roots of the Poincaré polynomials of all known Calabi-Yau fourfolds. The three million or so distinct Hodge data now presents a heavy computational challenge, on which a Quadra-core MacPro



**Figure 7:** (a) The position, on the complex plane, of the zeros of 50000 random integer polynomials with coefficients between 0 and 2500000. (b) The same, but with monic palindromic octics. (c) Monic palindromic octics, and with linear term vanishing.




**Figure 8:** (a) The position, on the complex plane, of the some 23 million zeros of the Poincaré polynomials of the approximately 1 million smooth Calabi-Yau fourfolds arising as hypersurfaces in toric fivefolds. (b) A slightly magnified area emphasizing the ordinate in the range  $[-1, 1]$ .

with 40 Gb of memory laboured for a week, to produce some 23 million complex roots. We present a scatter plot of these roots in part (a) of Figure 8. In part (b) of the same figure, we magnify it slightly to emphasize the same range as the random plots in Figure 7.


The theorem that  $-1$  being a root of the Poincaré polynomial of  $X$  implies  $X$  has rank exceeding unity is generally applicable. Hence, we can continue with this analysis. Now, (2.4) implies that (of course, the last equality follows directly for the definition of the the Euler characteristic  $\chi(X)$ )

$$P(-1, X) = 48 + 6(h^{1,1} - h^{2,1} + h^{3,1}) = \chi(X). \tag{2.5}$$

However, the relation between  $\chi$  and being self-mirror, or even the concept of the latter, is obviously not as clear in complex dimension greater than three. Be that as it may, we can still examine (2.5) in the the space of fourfolds. Of the some 3 million distinct triplets, there are only 61 with vanishing Euler number, which we demonstrate in Figure 9: in part (a),  $h^{1,1} + h^{3,1}$  against  $h^{1,1} - h^{3,1}$ , and in part (b),  $h^{1,1} + h^{2,1}$  against  $h^{1,1} - h^{2,1}$ . We see that these are all of relatively small Hodge numbers, and in part (b), we see that in spite of the general linear behaviour seen in part (b) of Figure 6, there is some substructure. In part (c), we plot the interesting shape of the roots of the Poincaré polynomials for these 61 members. In retrospect to Figure 6, we see that perhaps the closest notion to mirror symmetry in complex dimension four is the interchange of  $h^{1,1}$  and  $h^{3,1}$ . Of the circa 3 million, we find 5009 which



**Figure 9:** (a) The Hodge numbers, with  $h^{1,1} - h^{3,1}$  as the abscissa and  $h^{1,1} + h^{3,1}$ , the ordinate, of the fourfolds which have vanishing Euler number, and hence rank exceeding unity. (b) The same, but with  $h^{1,1} - h^{2,1}$ , as the abscissa and  $h^{1,1} + h^{2,1}$  as the ordinate. (c) The position of the roots, on the complex plane, of the Poincaré polynomial of these 61 spaces out of the some 3 million.



**Figure 10:** (a) The Hodge numbers, with  $h^{1,1} = h^{3,1}$  as the abscissa and  $h^{2,1}$ , the ordinate, of the “self-mirror” Calabi-Yau fourfolds. (b) The position of the roots of the Poincaré polynomials of these 5009 spaces.


have the property that  $h^{1,1} = h^{3,1}$ . We plot the pairs  $(h^{1,1} = h^{3,1}, h^{2,1})$  in part (a) of Figure 10 and the roots of their Poincaré polynomials in part (b).

### 3. Noncompact Calabi-Yau Geometries, Toric Diagrams, and Newton Polynomials

Having indulged in an excursion into the space of compact smooth Calabi-Yau threefolds and fourfolds, as well as their Poincaré polynomials, proceeding to the space of noncompact Calabi-Yau geometries is almost a perfunctory next step. These are affine varieties such as flat space  $\mathbb{C}^d$  and singularities which locally admit Gorenstein resolutions, and are central to McKay Correspondence and generalizations in mathematics as well as AdS/CFT and branes in string theory. A rich tapestry on this subject has been woven over the past few decades, whereby augmenting the relevance of our present investigation.

The most important family of non-compact Calabi-Yau geometries is indubitably those which afford toric description, as mentioned in the introduction. In complex dimension three, the Calabi-Yau condition compels the toric diagram to be coplanar, whence each is characterized by a (convex) lattice polygon  $D = \{v_i\}$ , with each  $v_i \in \mathbb{Z}^2$ . Therefore, a polynomial which instantly springs to mind is the Newton polynomial

$$D = \{(x_i, y_i)\} \implies P(z, w; X) = \sum_i a_i z^{x_i} w^{y_i} \in \mathbb{C}[z, w], \quad (3.1)$$




**Figure 11:** The most popular affine Calabi-Yau threefold toric diagrams, corresponding, respectively, to (a)  $\mathbb{C}^3$ , (b) conifold, (c) SPP, (d)  $\mathbb{F}_0$ , and (e, f, g, h)  $dP_{0,1,2,3}$ . The end points are at the standard lattice points in  $\mathbb{Z}^2$ .

where we have inserted potential coefficients  $a_i$  for generality. This is not a frivolous act; indeed, when  $a_i \in \mathbb{Z}_{\geq 0}$ , they are the so-called “multiplicities” first defined in [33] and play a vital rôle in comprehending the dimer model/brane tiling interpretation of toric gauge theories [34, 35, 50].


The most famous toric diagrams for affine Calabi-Yau threefolds are depicted in Figure 11, with the endpoints at the self-explanatory lattice points in  $\mathbb{Z}^2$ ; these include the reflexive polytopes in dimension two, and are commonly known as (a)  $\mathbb{C}^3$ , (b) the conifold, (c) the suspended pinched point (SPP), (d) affine cone over the zeroth Hirzebruch surface  $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$ , (e)  $dP_0 = \mathbb{C}^3/\mathbb{Z}_3$ , the affine cone over  $\mathbb{P}^2$ , and (f, g, h)  $dP_n$ , cones over, respectively, the first, second, and third del Pezzo surfaces which are  $\mathbb{P}^2$  blown up at  $n = 1, 2,$  and  $3$  generic points.

An immediate difficulty with (3.1) is, of course, that the polynomial is bi-variate, whereby describing, algebraically, Riemann surfaces. Even though such surfaces are crucial in the understanding of the gauge theory constructed on branes probing these affine toric Calabi-Yau spaces (cf. [50] for discussion on a web of inter-relations and various projections and spines of these Riemann surfaces), the notion of zeros is not obvious. We could, for example, set one of the coördinates to a fixed value, and consider the roots of the resulting univariate projection. This, however, does not seem particularly natural. Nevertheless, for illustrative purposes, we include a few examples in Figure 12, wherein we have set  $z$  to 1, varied the coefficients  $a_i$  randomly and integrally in  $[-5, 5]$ , and plotted the roots of the resulting polynomial in  $w$  for 5000 samples.

A much more natural and, as it turns out, interesting direction to take is to consider (3.1) not as a complex, but as a real curve. For comparative purposes, we should be mindful of the “amoebae” and the so-dubbed “algae” projections proposed in [50], which are, respectively the (natural log of the) real and imaginary projections of the Newton polynomials of the associated toric Calabi-Yau threefold.



**Figure 12:** The roots of the Newton polynomials  $P(z, w)$  at  $z = 1$ , for (a) the conifold, (b)  $\mathbb{F}_0$ , (c)  $dP_1$ , and (d)  $dP_3$ .



**Figure 13:** In reference to the toric diagrams in Figure 11, for each Calabi-Yau threefold geometry, we find the corresponding Newton polynomial as a real algebraic curve. We sample over 50000 random integer coefficients in the range  $[-10, 10]$ , and isolate the real critical points in  $\mathbb{R}^2$ , which are then plotted collectively for (b) conifold, (c) SPP, (d)  $\mathbb{F}_0$ , and (e, f, g, h)  $dP_{0,1,2,3}$ .

In this regard, perhaps the most significant quantity for our complex Newton polynomials is the set of turning points in  $\mathbb{R}^2$ , namely, the set of real critical points of  $P(z, w) \in \mathbb{R}[z, w]$ . In other words, we find the simultaneous real solutions to

$$\partial_z P(z, w) = \sum_i a_i x_i z^{x_i-1} w^{y_i} = 0, \quad \partial_w P(z, w) = \sum_i a_i y_i z^{x_i} w^{y_i-1} = 0, \quad (3.2)$$

with  $a_i$  randomly sampled and discard the imaginary solutions. This, indeed, brings us back to Waring's original considerations on nonreal roots.

In Figure 13, we take a fixed toric diagram corresponding to a given Calabi-Yau geometry, and consider the Newton Polynomial in (3.1). Then, we sample 50000 random integer coefficients  $a_i$  in an appropriate range, here taken to be  $[-10, 10]$ . For each, we find the real critical points, and collectively plot them. We note that  $\mathbb{C}^3$  does not have any real critical points and is thus left out. This is because the Newton polynomial is simply  $a + bz + cw$  for  $a, b, c \in \mathbb{Z}$ . Thus, (3.2) gives  $b = c = 0$ , independent of  $(z, w)$  coordinates. Similarly, the case of (b), the conifold, can be considered a reference point. The Newton polynomial is  $a + bz + cw + dzw$ , whence, the critical points are given by the solutions of  $b + dw = c + dz = 0$ , or,  $w_0 = -b/d, z_0 = -c/d$ . Hence, given that each of  $b, c, d$  is independently randomly evenly distributed, the turning points  $(z_0, w_0)$  are then distributed as quotients of even random samples, and whence the clustering nearer to the lower values as seen in the darker region in the figure.

## Dedication

To the occasion of the happy Installation of Professor Sir Martin Taylor as the Fiftieth Warden of Merton College, Oxford, on the Second Day of the Month of October, in the Year of Our Lord Two Thousand and Ten, and to the noble retirement of Professor Dame Jessica Rawson, this humble brief note is dedicated. Vivat Custos, vivat Collegium, & Stet Fortuna Domus!

## Acknowledgments





Ad Catharinae Sanctae Alexandriae et ad Maiorem Dei Gloriam, cum Universitate Nankai, Universitate Civitate Londiniensis, Scientiae et Technologiae Concilio Anglicae, et Collegio Mertonense Oxoniensis, atque amore Elizabethae Katherinae Hunter, Rosae Hiberniae, multas gratias YHH agit.

## References

- [1] A. Bloch and G. Pólya, "On the zeros of certain algebraic equations," *Proceedings London Mathematical Society*, vol. 33, pp. 102–114, 1932.
- [2] J. Littlewood and A. Offord, "On the number of real roots of a random algebraic equation," *Journal London Mathematical Society*, vol. 13, pp. 288–295, 1938.
- [3] J. H. Hannay, "Chaotic analytic zero points: exact statistics for those of a random spin state," *Journal of Physics. A*, vol. 29, no. 5, pp. L101–L105, 1996.
- [4] Y. Peres and B. Virag, "Zeros of the i.i.d. Gaussian power series: a conformally invariant determinantal process," *Journal of Physics A*, vol. 29, no. 5.
- [5] A. T. Bharucha-Reid and M. Sambandham, *Random polynomials*, Probability and Mathematical Statistics, Academic Press, Orlando, Fla, USA, 1986.
- [6] J Littlewood, "Some problems in real and complex analysis," in *Heath Mathematical Monographs*, Lexington, Mass, USA, 1968.
- [7] P. Erdős and P. Turán, "On the distribution of roots of polynomials," *Annals of Mathematics*, vol. 51, pp. 105–119, 1950.
- [8] J. M. Hammersley, "The zeros of a random polynomial," in *Proceedings of the 3rd Berkeley Symposium on Mathematical Statistics and Probability*, vol. 2, pp. 89–111, University of California Press, Los Angeles, Calif, USA.
- [9] M. Kac, "On the average number of real roots of a random algebraic equation," *Bulletin of the American Mathematical Society*, vol. 49, pp. 314–320, 1943.
- [10] H. Montgomery, "The pair correlation of zeros of the zeta function," in *Proceedings of Symposia in Pure Mathematics*, vol. 24, pp. 181–193, American Mathematical Society, Providence, RI, USA, 1973.
- [11] A. Odlyzko, "10<sup>20</sup>-th zero of the Riemann zeta function and 70 million of its neighbors," 1989, preprint, A.T.T.
- [12] Y. H. He, V. Jejjala, and D. Minic, "On the physics of the Riemann zeros," In press, <http://arxiv.org/abs/1004.1172>.
- [13] A. M. Odlyzko and B. Poonen, "Zeros of polynomials with 0,1 coefficients," *L'Enseignement Mathématique*, vol. 39, no. 3-4, pp. 317–348, 1993.
- [14] P. Borwein and L. Jörgenson, "Visible structures in number theory," *The American Mathematical Monthly*, vol. 108, no. 10, pp. 897–910, 2001.
- [15] D. Bailey, J. Borwein, N. Calkin, R. Girgensohn, D. Luke, and V. Moll, *Experimental Mathematics in Action*, A. K. Peters Ltd., Wellesley, Mass, USA, 2007.
- [16] "Mathematica Edition: Version 7.0," Wolfram Research.
- [17] S. Franco, Y. H. He, C. Herzog, and J. Walcher, "Chaotic duality in string theory," *Physical Review D*, vol. 70, article 046006, 2004.

- [18] F. Beaucoup, P. Borwein, D. Boyd, and C. Pinner, "Multiple roots of  $[[\text{minus}]1, 1]$  power series," *Journal of the London Mathematical Society*, vol. 57, no. 1, pp. 135–147, 1998.
- [19] D. Christensen, <http://jdc.math.uwo.ca/roots/>.
- [20] J. Jörgenson, <http://oldweb.cecm.sfu.ca/personal/loki/Projects/Roots/>.
- [21] S. Derbyshire and J. Matson, "Polynomial plot: simple math expressions yield intricate visual patterns," *Scientific American*, 2009.
- [22] J. Baez, <http://math.ucr.edu/home/baez/roots/>.
- [23] X. S. Lin, "Zeros of the Jones polynomial," <http://math.ucr.edu/~xl/abs-jk.pdf>.
- [24] J. Hoste and M. Thistlethwaite, <http://www.math.utk.edu/~morwen/knotscape.html>.
- [25] P. Candelas, A. M. Dale, C. A. Lütken, and R. Schimmrigk, "Complete intersection Calabi-Yau manifolds," *Nuclear Physics*, vol. 298, no. 3, pp. 493–525, 1988.
- [26] P. Candelas, M. Lynker, and R. Schimmrigk, "Calabi-Yau manifolds in weighted  $P(4)$ ," *Nuclear Physics*, vol. 341, no. 2, pp. 383–402, 1990.
- [27] M. Kreuzer and H. Skarke, "Complete classification of reflexive polyhedra in four dimensions," *Advances in Theoretical and Mathematical Physics*, vol. 4, pp. 1209–1230, 2002.
- [28] P. Candelas, X. de la Ossa, Y. H. He, and B. Szendroi, "Triadophilia: a special corner in the landscape," *Advances in Theoretical and Mathematical Physics*, vol. 12, p. 2, 2008.
- [29] P. Candelas and R. Davies, "New Calabi-Yau manifolds with small Hodge numbers," *Fortschritte der Physik*, vol. 58, p. 383, 2010.
- [30] V. Braun, "On free quotients of complete intersection Calabi-Yau manifolds," In press, <http://arxiv.org/abs/1003.3235>.
- [31] F. Denef and M. R. Douglas, "Computational complexity of the landscape. I," *Annals of Physics*, vol. 322, p. 1096, 2007.
- [32] A. Hanany and Y. H. He, "Non-Abelian finite gauge theories," *Journal of High Energy Physics*, vol. 2, article 013, 1999.
- [33] B. Feng, A. Hanany, and Y. H. He, "D-brane gauge theories from toric singularities and toric duality," *Nuclear Physics*, vol. B595, p. 165, 2001.
- [34] A. Hanany and K. D. Kennaway, "Dimer models and toric diagrams," In press, <http://arxiv.org/abs/hep-th/0503149>.
- [35] S. Franco, A. Hanany, D. Martelli, J. Sparks, D. Vegh, and B. Wecht, "Gauge theories from toric geometry and brane tilings," *Journal of High Energy Physics*, vol. 1, article 128, 2006.
- [36] D. Forcella, A. Hanany, Y. H. He, and A. Zaffaroni, "The master space of  $N=1$  Gauge theories," *Journal of High Energy Physics*, vol. 8, article 012, 2008.
- [37] A. Hanany and Y. H. He, "M2-branes and quiver Chern-Simons: a taxonomic study," In press, <http://arxiv.org/abs/0811.4044v2>.
- [38] J. Davey, A. Hanany, N. Mekareeya, and G. Torri, "Brane tilings, M2-branes and Chern-Simons theories," in *Proceedings of the 49th Cracow School of Theoretical Physics: Non-Perturbative Gravity and Quantum Chromodynamics*, Zakopane, Poland, May-June 2009.
- [39] J. Davey, A. Hanany, and R. K. Seong, "Counting orbifolds," *Journal of High Energy Physics*, vol. 6, no. 6, article 010, 2010.
- [40] S. Benvenuti, B. Feng, A. Hanany, and Y. H. He, "Counting BPS operators in gauge theories: quivers, syzygies and plethystics," *Journal of High Energy Physics*, vol. 11, article 050, 2007.
- [41] D. Forcella, "Operators and vacua of  $N=1$  field theories," *Nuovo Cimento*, vol. 125, pp. 905–914, 2010.
- [42] A. Hanany, N. Mekareeya, and G. Torri, "The Hilbert series of adjoint SQCD," *Nucl. Phys.*, vol. 825, no. 1-2, pp. 52–97, 2010.
- [43] T. Emery, "Vector fields on manifolds," *Bulletin of the American Mathematical Society*, vol. 75, pp. 643–683, 1969.
- [44] G. E. Bredon, "Counterexamples on the rank of a manifold," *Proceedings of the American Mathematical Society*, vol. 27, no. 3, pp. 592–594, 1971.
- [45] D. Bump, K. K. Choi, P. Kurlberg, and J. Vaaler, "A local Riemann hypothesis. I," *Mathematische Zeitschrift*, vol. 233, no. 1, pp. 1–19, 2000.
- [46] A. Postnikov and R. P. Stanley, "Deformations of Coxeter hyperplane arrangements," *Journal of Combinatorial Theory. Series A*, vol. 91, no. 1-2, pp. 544–597, 2000.
- [47] F. Rodriguez-Villegas, "On the zeros of certain polynomials," *Proceedings of the American Mathematical Society*, vol. 130, no. 8, pp. 2251–2245, 2002.

- [48] M. Kreuzer and H. Skarke, "Calabi-Yau 4-folds and toric fibrations," *Journal of Geometry and Physics*, vol. 26, no. 3-4, pp. 272–290, 1998.
- [49] M. Lynker, R. Schimmrigk, and A. Wisskirchen, "Landau-Ginzburg vacua of string, M-and F-theory at  $c = 12$ ," *Nuclear Physics*, vol. 550, p. 123, 1999.
- [50] B. Feng, Y. H. He, K. D. Kennaway, and C. Vafa, "Dimer models from mirror symmetry and quivering amoebae," *Advances in Theoretical and Mathematical Physics*, vol. 12, p. 3, 2008.

The Hindawi logo consists of two interlocking loops, one blue and one green.

**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

