Research Article

$B_c \rightarrow BP, BV$ Decays with the QCD Factorization Approach

Junfeng Sun,¹ Na Wang,^{1,2} Qin Chang,¹ and Yueling Yang¹

¹Institute of Particle and Nuclear Physics, Henan Normal University, Xinxiang 453007, China

²Institute of Particle and Key Laboratory of Quark and Lepton Physics, Central China Normal University, Wuhan 430079, China

Correspondence should be addressed to Qin Chang; changqin@htu.cn

Received 7 January 2015; Accepted 17 March 2015

Academic Editor: Michal Kreps

Copyright © 2015 Junfeng Sun et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

We studied the nonleptonic $B_c \rightarrow BP, BV$ decays with the QCD factorization approach. It is found that the Cabibbo favored processes of $B_c \rightarrow B_s\pi, B_s\rho, B_u\bar{K}$ are the promising decay channels with branching ratio larger than 1%, which should be observed earlier by the LHCb collaboration.

1. Introduction

The B_c meson is the ground pseudoscalar meson of the $\bar{b}c$ system [1]. Compared with the heavy unflavored charmonium $c\bar{c}$ and bottomonium $b\bar{b}$, the B_c meson is unique in some respects. (1) Heavy quarkonia could be created in the parton-parton process $ij \rightarrow Q\bar{Q}$ at the order of α_s^2 (where $ij = gg$ or $q\bar{q}$, $Q = b, c$), while the production probability for the B_c meson is at least at the order of α_s^4 via $ij \rightarrow B_c^{(*)+} + b\bar{c}$, where the gluon-gluon fusion mechanism is dominant at Tevatron and LHC [2]. The B_c meson is difficult to produce experimentally, but it was observed for the first time via the semileptonic decay mode $B_c \rightarrow J/\psi\ell\nu$ in $p\bar{p}$ collisions by the CDF collaboration in 1998 [3, 4], which showed the realistic possibility of experimental study of the B_c meson. One of the best measurements on the mass and lifetime of the B_c meson is reported recently by the LHCb collaboration, $m_{B_c} = 6276.28 \pm 1.44 \pm 0.36$ MeV [5] and $\tau_{B_c} = 513.4 \pm 11.0 \pm 5.7$ fs [6]. With the running of the LHC, the B_c meson has a promising prospect. It is estimated that one could expect some 10^{10} B_c events at the high-luminosity LHC experiments per year [7, 8]. The studies on the B_c meson have entered a new precision era. (2) For charmonium and bottomonium, the strong and electromagnetic interactions are mainly responsible for annihilation of the $Q\bar{Q}$ quark pair into final states. The B_c meson, carrying nonzero flavor number $B = C = \pm 1$ and lying below

the BD meson pair threshold, can decay only via the weak interaction, which offers an ideal sample to investigate the weak decay mechanism of heavy flavors that is inaccessible to both charmonium and bottomonium. The B_c weak decay provides great opportunities to investigate the perturbative and nonperturbative QCD, final state interactions, and so forth.

With respect to the heavy-light $B_{u,d,s}$ mesons, the doubly heavy B_c meson has rich decay channels because of its relatively large mass and that both b and c quarks can decay individually. The decay processes of the B_c meson can be divided into the following three classes [2, 9–11]: (1) the c quark decays with the b quark as a spectator; (2) the b quark decays with the c quark as a spectator; (3) the b and c quarks annihilate into a virtual W boson, with the ratios of $\sim 70\%$, 20% and 10% , respectively [2]. Up to now, the experimental evidences of pure annihilation decay mode [class (3)] are still nothing. The $b \rightarrow c$ transition, belonging to the class (2), offers a well-constructed experimental structure of charmonium at the Tevatron and LHC. Although the detection of the c quark decay is very challenging to experimentalists, the clear signal of the $B_c \rightarrow B_s\pi$ decay is presented by the LHCb group using the $B_s \rightarrow D_s\pi$ and $B_s \rightarrow J/\psi\phi$ channels with statistical significance of 7.7σ and 6.1σ , respectively [12].

Anticipating the forthcoming accurate measurements on the B_c meson at hadron colliders and the lion's share of the B_c

decay width from the c quark decay [31–33], many theoretical papers were devoted to the study of the $B_c \rightarrow BP, BV$ decays (where P and V denote the $SU(3)$ ground pseudoscalar and vector mesons, resp.), such as [17, 34–37] with the BSW model [38, 39] or IGSW model [40], [18, 19, 41] based on the Bethe-Salpeter (BS) equation, [20–24, 42] with potential models, [25] with constituent quark model, [26–28] with QCD sum rules, [43] with the quark diagram scheme, [29, 30] with the perturbative QCD approach (pQCD) [44–49], and so on. The previous predictions on the branching ratios for the $B_c \rightarrow BP, BV$ decays are collected in Table 3. The discrepancies of previous investigations arise mainly from the different model assumptions. Recently, several phenomenological methods have been fully developed to cope with the hadronic matrix elements and successfully applied to the nonleptonic B decay, such as the pQCD approach [44–49] based on the k_T factorization scheme, the soft-collinear effective theory [50–57] and the QCD-improved factorization (QCDF) approach [58–63] based on the collinear approximation and power counterterm rules in the heavy quark limits. In this paper, we will study the $B_c \rightarrow BP, BV$ decays with the QCDF approach to provide a ready reference to the existing and upcoming experiments.

This paper is organized as follows. In Section 2, we will present the theoretical framework and the amplitudes for the $B_c \rightarrow BP, BV$ decays within the QCDF framework. Section 3 is devoted to numerical results and discussion. Finally, Section 4 is our summation.

2. Theoretical Framework

2.1. The Effective Hamiltonian. The low energy effective Hamiltonian responsible for the nonleptonic bottom-conserving $B_c \rightarrow BP, BV$ decays constructed by means of the operator product expansion and the renormalization group (RG) method is usually written in terms of the four-quark interactions [64, 65]. Consider

$$\begin{aligned}
 H_{\text{eff}} &= \frac{G_F}{\sqrt{2}} \left\{ V_{ub} V_{cb}^* [C_1^a(\mu) Q_1^a(\mu) + C_2^a(\mu) Q_2^a(\mu)] \right. \\
 &\quad + \sum_{q_1, q_2=d,s} V_{uq_1} V_{cq_2}^* [C_1(\mu) Q_1(\mu) + C_2(\mu) Q_2(\mu)] \\
 &\quad \left. + \sum_{q_3=d,s} V_{uq_3} V_{cq_3}^* \sum_{k=3}^{10} C_k(\mu) Q_k(\mu) \right\} + \text{h.c.}, \quad (1)
 \end{aligned}$$

where the Fermi coupling constant $G_F \simeq 1.166 \times 10^{-5} \text{ GeV}^{-2}$ [1]; $Q_{1,2}$, $Q_{1,2}^a$, and $Q_{3,\dots,10}$ are the relevant local tree, annihilation, and penguin four-quark operators, respectively, which govern the decays in question. The Cabibbo-Kobayashi-Maskawa (CKM) factor $V_{uq_i} V_{cq_j}^*$ and Wilson coefficients C_i describe the coupling strength for a given operator.

Using the unitarity of the CKM matrix, there is a large cancellation of the CKM factors

$$V_{ud} V_{cd}^* + V_{us} V_{cs}^* = -V_{ub} V_{cb}^* \sim \mathcal{O}(\lambda^5), \quad (2)$$

where the Wolfenstein parameter $\lambda = \sin \theta_c = 0.225$ [61] and θ_c is the Cabibbo angle. Hence, compared with the tree contributions, the contributions of annihilation and penguin operators are strongly suppressed by the CKM factor. If the CP -violating asymmetries that are expected to be very small due to the small weak phase difference for c quark decay are prescinded from the present consideration, then the penguin and annihilation contributions could be safely neglected. The local tree operators $Q_{1,2}$ in (1) are expressed as follows:

$$\begin{aligned}
 Q_1 &= [\bar{q}_{2,\alpha} \gamma_\mu (1 - \gamma_5) c_\alpha] [\bar{u}_\beta \gamma^\mu (1 - \gamma_5) q_{1,\beta}], \\
 Q_2 &= [\bar{q}_{2,\alpha} \gamma_\mu (1 - \gamma_5) c_\beta] [\bar{u}_\beta \gamma^\mu (1 - \gamma_5) q_{1,\alpha}], \quad (3)
 \end{aligned}$$

where α and β are the $SU(3)$ color indices.

The Wilson coefficients $C_i(\mu)$ summarize the physics contributions from scales higher than μ . They are calculable with the RG improved perturbation theory and have properly been evaluated to the next-to-leading order (NLO) [64, 65]. They can be evolved from a higher scale $\mu \sim \mathcal{O}(m_W)$ down to a characteristic scale $\mu \sim \mathcal{O}(m_c)$ with the functions including the flavor thresholds [64, 65]

$$\vec{C}(\mu) = U_4(\mu, m_b) M(m_b) U_5(m_b, m_W) \vec{C}(m_W), \quad (4)$$

where $U_f(\mu_f, \mu_i)$ is the RG evolution matrix converting coefficients from the scale μ_i to μ_f , and $M(\mu)$ is the quark threshold matching matrix. The expressions of $U_f(\mu_f, \mu_i)$ and $M(\mu)$ can be found in [64, 65]. The numerical values of LO and NLO $C_{1,2}$ with the naive dimensional regularization scheme are listed in Table 1. The values of NLO Wilson coefficients in Table 1 are consistent with those given by [64, 65] where a trick with “effective” number of active flavors $f = 4.15$ rather than formula (4) is used.

To obtain the decay amplitudes, the remaining work is how to accurately evaluate the hadronic matrix elements $\langle BM | Q_i(\mu) | B_c \rangle$ which summarize the physics contributions from scales lower than μ . Since the hadronic matrix elements involve long distance contributions, one is forced to use either nonperturbative methods such as lattice calculations and QCD sum rules or phenomenological models relying on some assumptions. Consequently, it is very unfortunate that hadronic matrix elements cannot be reliably calculated at present, and that the most intricate part and the dominant theoretical uncertainties in the decay amplitudes reside in the hadronic matrix elements.

2.2. Hadronic Matrix Elements. Phenomenologically, based on the power counting rules in the heavy quark limit, Beneke et al. proposed that the hadronic matrix elements could be written as the convolution integrals of hard scattering kernels and the light cone distribution amplitudes with the QCDF master formula [58–63]. The QCDF approach is widely applied to nonleptonic B decays and it works well [66–76],

TABLE 1: The numerical values of the Wilson coefficients and the effective coefficients for $B_c \rightarrow B\pi$ decay, where $m_c = 1.275 \pm 0.025$ GeV [1].

μ	LO		NLO		QCDF			
	C_1	C_2	C_1	C_2	$\text{Re}(a_1)$	$\text{Im}(a_1)$	$\text{Re}(a_2)$	$\text{Im}(a_2)$
$0.8m_c$	1.334	-0.587	1.274	-0.503	1.270	0.096	-0.450	-0.218
m_c	1.275	-0.503	1.222	-0.424	1.216	0.068	-0.361	-0.173
$1.2m_c$	1.239	-0.449	1.189	-0.373	1.184	0.054	-0.306	-0.148

which encourage us to apply the QCDF approach to the study of $B_c \rightarrow BP, BV$ decays. Since the spectator is the heavy b quark who is almost always on shell, the virtuality of the gluon linked with the spectator should be $\sim \mathcal{O}(\Lambda_{\text{QCD}}^2)$. The dominant behavior of the $B_c \rightarrow B$ transition form factors and the contributions of hard spectator scattering interactions are governed by soft processes. According to the basic idea of the QCDF approach [69, 70], the hard and soft contributions to the form factors entangle with each other and cannot be identified reasonably, so the physical form factors are used as the inputs. The hard spectator scattering contributions are power suppressed in the heavy quark limit. Finally, the hadronic matrix elements can be written as

$$\begin{aligned} \langle BM | Q_{1,2} | B_c \rangle &= \sum_i F_i^{B_c \rightarrow B} \int dx H_i(x) \Phi_M(x) \\ &\propto \sum_i F_i^{B_c \rightarrow B} f_M \{1 + \alpha_s r_1 + \dots\}, \end{aligned} \quad (5)$$

where $F_i^{B_c \rightarrow B}$ is the transition form factor and $\Phi_M(x)$ is the light-cone distribution amplitudes of the emitted meson M with the decay constant f_M . The hard scattering kernels $H_i(x)$ are computable order by order with the perturbation theory in principle. At the leading order α_s^0 , $H_i(x) = 1$, that is, the convolution integral of (5) results in the meson decay constant. The hadronic matrix elements are parameterized by the product of form factors and decay constants, which are real and renormalization scale independent. One goes back to the simple “naive factorization” (NF) scenario. At the order α_s and higher orders, the information of strong phases and the renormalization scale dependence of hadronic matrix elements could be partly recuperated. Combined the nonfactorizable contributions with the Wilson coefficients, the scale independent effective coefficients at the order α_s can be obtained [58–63] as follows:

$$\begin{aligned} a_1 &= C_1^{\text{NLO}} + \frac{1}{N_c} C_2^{\text{NLO}} + \frac{\alpha_s}{4\pi} \frac{C_F}{N_c} C_2^{\text{LO}} V, \\ a_2 &= C_2^{\text{NLO}} + \frac{1}{N_c} C_1^{\text{NLO}} + \frac{\alpha_s}{4\pi} \frac{C_F}{N_c} C_1^{\text{LO}} V, \end{aligned} \quad (6)$$

where the expressions of vertex corrections are [58–63]

$$\begin{aligned} V &= 6 \log \left(\frac{m_c^2}{\mu^2} \right) - 18 - \left(\frac{1}{2} + i3\pi \right) a_0^M \\ &\quad + \left(\frac{11}{2} - i3\pi \right) a_1^M - \frac{21}{20} a_2^M + \dots, \end{aligned} \quad (7)$$

with the twist-2 quark-antiquark distribution amplitudes of pseudoscalar P and longitudinally polarized vector V meson in terms of Gegenbauer polynomials [14–16]. One has

$$\phi_M(x) = 6x\bar{x} \sum_{n=0}^{\infty} a_n^M C_n^{3/2}(x - \bar{x}), \quad (8)$$

where $\bar{x} = 1 - x$; a_n^M is the Gegenbauer moment and $a_0^M \equiv 1$.

From the numbers in Table 1, it is found that (1) for the coefficient a_1 the nonfactorizable contributions accompanied by the Wilson coefficient C_2 can provide $\geq 10\%$ enhancement compared with the NF's result, and a relatively small strong phase $\leq 5^\circ$; (2) for the coefficient a_2 , the nonfactorizable contributions assisted with the large Wilson coefficient C_1 are significant. In addition, a relatively large strong phase $\sim -155^\circ$ is obtained; (3) the QCDF's values of $a_{1,2}$ agree basically with the real coefficients $a_1 \simeq 1.20$ and $a_2 \simeq -0.317$ which are used by previous studies on the $B_c \rightarrow BP, BV$ decays in [17, 18, 20–28, 34–37, 42], but with more information on the strong phases.

2.3. Decay Amplitudes. Within the QCDF framework, the amplitudes for $B_c \rightarrow BM$ decays are expressed as

$$\begin{aligned} \mathcal{A}(B_c \rightarrow BM) &= \langle BM | \mathcal{H}_{\text{eff}} | B_c \rangle \\ &= \frac{G_F}{\sqrt{2}} V_{uq_1} V_{cq_2}^* a_i \langle M | J^\mu | 0 \rangle \langle B | J_\mu | B_c \rangle. \end{aligned} \quad (9)$$

The matrix elements of current operators are defined as

$$\begin{aligned} \langle P(p) | \bar{q}_1 \gamma^\mu (1 - \gamma_5) q_2 | 0 \rangle &= -if_P p^\mu, \\ \langle V(\epsilon, p) | \bar{q}_1 \gamma^\mu (1 - \gamma_5) q_2 | 0 \rangle &= f_V m_V \epsilon^\mu, \end{aligned} \quad (10)$$

where f_P and f_V are the decay constants of pseudoscalar P and vector V mesons, respectively; m_V and ϵ denote the mass and polarization of vector meson, respectively.

For the mixing of physical pseudoscalar η and η' meson, we adopt the quark-flavor basis description proposed in [13] and neglect the contributions from possible gluonium and $c\bar{c}$ compositions; that is,

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \eta_q \\ \eta_s \end{pmatrix}, \quad (11)$$

where $\eta_q = (u\bar{u} + d\bar{d})/\sqrt{2}$ and $\eta_s = s\bar{s}$; the mixing angle $\phi = (39.3 \pm 1.0)^\circ$ [13]. We assume that the vector mesons are ideally mixed; that is, $\omega = (u\bar{u} + d\bar{d})/\sqrt{2}$ and $\phi = s\bar{s}$.

TABLE 2: The numerical values of input parameters.

Wolfenstein parameters	
$\lambda = 0.22537 \pm 0.00061$ [1]	$A = 0.814^{+0.023}_{-0.024}$ [1]
$\bar{\rho} = 0.117 \pm 0.021$ [1]	$\bar{\eta} = 0.353 \pm 0.013$ [1]
Decay constant of mesons	
$f_\pi = 130.41 \pm 0.20$ MeV [1]	$f_K = 156.2 \pm 0.7$ MeV [1]
$f_{\eta_q} = (1.07 \pm 0.02)f_\pi$ [13]	$f_{\eta_s} = (1.34 \pm 0.06)f_\pi$ [13]
$f_\rho = 216 \pm 3$ MeV [14–16]	$f_\omega = 187 \pm 5$ MeV [14–16]
$f_{K^*} = 220 \pm 5$ MeV [14–16]	
Gegenbauer moments at the scale $\mu = 1$ GeV	
$a_1^\pi = a_1^{\eta_q} = a_1^{\eta_s} = 0$ [14–16]	$a_2^\pi = a_2^{\eta_q} = a_2^{\eta_s} = 0.25 \pm 0.15$ [14–16]
$a_1^{\bar{K}} = -a_1^K = 0.06 \pm 0.03$ [14–16]	$a_2^K = a_2^{\bar{K}} = 0.25 \pm 0.15$ [14–16]
$a_{1,\rho}^\parallel = a_{1,\omega}^\parallel = 0$ [14–16]	$a_{2,\rho}^\parallel = a_{2,\omega}^\parallel = 0.15 \pm 0.07$ [14–16]
$a_{1,\bar{K}^*}^\parallel = -a_{1,K^*}^\parallel = 0.03 \pm 0.02$ [14–16]	$a_{2,K^*}^\parallel = a_{2,\bar{K}^*}^\parallel = 0.11 \pm 0.09$ [14–16]

The transition form factors are defined as [38, 39]

$$\begin{aligned}
& \langle B(k) | \bar{q} \gamma^\mu (1 - \gamma_5) c | B_c(p) \rangle \\
&= \left[p + k - \frac{m_{B_c}^2 - m_B^2}{q^2} q \right]^\mu F_1^{B_c \rightarrow B}(q^2) \\
&+ \frac{m_{B_c}^2 - m_B^2}{q^2} q^\mu F_0^{B_c \rightarrow B}(q^2),
\end{aligned} \quad (12)$$

where $q = p - k$, and the condition of $F_0^{B_c \rightarrow B}(0) = F_1^{B_c \rightarrow B}(0)$ is required compulsorily to cancel the singularity at the pole $q^2 = 0$.

For the $B_c \rightarrow B$ transition form factors, since the velocity of the recoiled B meson is very low in the rest frame of the B_c meson, the wave functions of B and B_c mesons overlap strongly. It is believed that the form factors $F_{0,1}^{B_c \rightarrow B}$ should be close to the result using the nonrelativistic harmonic oscillator wave functions with the BSW model [17]. Consider

$$F_{0,1}^{B_c \rightarrow B} \simeq \left(\frac{2m_{B_c}m_B}{m_{B_c}^2 + m_B^2} \right)^{1/2} \simeq 0.99. \quad (13)$$

The flavor symmetry breaking effects on the form factors are neglectable in (13). For simplification, we take $F_{0,1}^{B_c \rightarrow B_{u,d,s}} = 1.0$ in our numerical calculation to give a rough estimation.

3. Numerical Results and Discussions

The branching ratios of nonleptonic two-body B_c decays in the rest frame of the B_c meson can be written as

$$\mathcal{Br}(B_c \rightarrow BM) = \frac{\tau_{B_c}}{8\pi} \frac{p}{m_{B_c}^2} |\mathcal{A}(B_c \rightarrow BM)|^2, \quad (14)$$

where the lifetime of the B_c meson $\tau_{B_c} = 513.4 \pm 11.0 \pm 5.7$ fs [6] and p is the common momentum of final particles. The decay amplitudes $\mathcal{A}(B_c \rightarrow BM)$ are listed in the Appendix.

The input parameters in our calculation, including the CKM Wolfenstein parameters, decay constants, and Gegenbauer moments of distribution amplitudes in (8), are collected in Table 2. If not specified explicitly, we will take their central values as the default inputs. Our numerical results on the CP -averaged branching ratios are presented in Table 3, where theoretical uncertainties of the ‘‘QCDF’’ column come from the CKM parameters, the renormalization scale $\mu = (1 \pm 0.2)m_{B_c}$, decay constants, and Gegenbauer moments, respectively. For comparison, previous results calculated with the fixed coefficients $a_1 \simeq 1.22$ and $a_2 \simeq -0.4$ are also listed. There are some comments on the branching ratios.

(1) From the numbers in Table 3, it is seen that different branching ratios for the $B_c \rightarrow BP, BV$ decays were obtained with different approaches in previous works, although the same coefficients $a_{1,2}$ are used. Much of the discrepancy comes from the different values of the transition form factors. If the same value of the form factor is used, then the disparities on branching ratios for the a_1 -dominated B_c decays will be highly alleviated. For example, all previous predictions on $\mathcal{Br}(B_c \rightarrow B_s \pi)$ will be about 10% with the same form factor $F_0^{B_c \rightarrow B_s} \simeq 1.0$, which is generally in line with the QCDF estimation within uncertainties and also agrees with the recent LHCb measurement [12].

(2) There is a hierarchical structure between the QCDF's results on branching ratios for the $B_c \rightarrow BP$ and BV decays with the same final B_q meson, for example,

$$\begin{aligned}
& \mathcal{Br}(B_c \rightarrow B_q \pi) > \mathcal{Br}(B_c \rightarrow B_q \rho), \\
& \mathcal{Br}(B_c \rightarrow B_q K) \geq 5 \mathcal{Br}(B_c \rightarrow B_q K^*),
\end{aligned} \quad (15)$$

which differs from the previous results. There are two decisive factors. One is the kinematic factor. The phase space for the $B_c \rightarrow BV$ decays is more compressed than that for the $B_c \rightarrow BP$ decays, because the mass of the light pseudoscalar meson (except for the exotic η' meson) is generally less than the mass of the corresponding vector meson with the same valence quark components. The other is the dynamical factor. The

TABLE 3: The CP-averaged branching ratios for the $B_c \rightarrow BP, BV$ decays.

Decay mode	Case	Reference [17] ^a	Reference [18] ^b	Reference [19] ^c	Reference [20] ^d	Reference [21, 22] ^e	Reference [23] ^f	Reference [24] ^g	Reference [25] ^h	Reference [26-28] ⁱ	Reference [29, 30] ^j	QCDF
$B_c \rightarrow B_s^0 \pi^+$	1-a	1.0×10^{-1}	6.8×10^{-2}	1.8×10^{-2}	2.9×10^{-2}	4.0×10^{-2}	4.3×10^{-2} (4.3×10^{-2})	1.3×10^{-1}	4.6×10^{-2}	1.9×10^{-1}	8.8×10^{-2}	$(1.13^{+0.00+0.11+0.01}_{-0.00-0.06-0.01}) \times 10^{-1}$
$B_c \rightarrow B_s^0 K^+$	1-b	7.6×10^{-3}	4.9×10^{-3}	2.0×10^{-3}	2.4×10^{-3}	3.3×10^{-3}	3.3×10^{-3} (3.3×10^{-3})	8.5×10^{-3}	3.4×10^{-3}	1.2×10^{-2}	5.2×10^{-3}	$(7.41^{+0.04+0.70+0.09}_{-0.04-0.39-0.09}) \times 10^{-3}$
$B_c \rightarrow B_s^0 \rho^+$	1-a	6.3×10^{-2}	5.2×10^{-2}	4.6×10^{-2}	1.6×10^{-2}	2.7×10^{-2}	3.0×10^{-2} (2.7×10^{-2})	1.1×10^{-1}	2.7×10^{-2}	8.4×10^{-2}	3.2×10^{-2}	$(4.44^{+0.00+0.41+0.13}_{-0.00-0.23-0.13}) \times 10^{-2}$
$B_c \rightarrow B_s^0 K^{*+}$	1-b	3.2×10^{-4}		1.2×10^{-3}	3.5×10^{-5}	1.5×10^{-4}	8.0×10^{-5} (7.1×10^{-5})	4.0×10^{-4}	1.3×10^{-4}		9.7×10^{-5}	$(1.25^{+0.01+0.12+0.06}_{-0.01-0.07-0.06}) \times 10^{-4}$
$B_c \rightarrow B_d^0 \pi^0$	1-b	7.4×10^{-3}	3.8×10^{-3}	1.2×10^{-3}	1.2×10^{-3}	1.3×10^{-3}	1.8×10^{-3} (1.5×10^{-3})	8.4×10^{-3}	2.4×10^{-3}	1.2×10^{-2}	6.9×10^{-3}	$(7.83^{+0.04+0.73+0.04}_{-0.04-0.41-0.04}) \times 10^{-3}$
$B_c \rightarrow B_d^0 K^+$	1-c		3.0×10^{-4}	1.2×10^{-4}	1.0×10^{-4}	1.1×10^{-4}	1.5×10^{-4} (1.2×10^{-4})	5.9×10^{-4}	1.8×10^{-4}	8.1×10^{-4}	4.4×10^{-4}	$(5.29^{+0.06+0.50+0.07}_{-0.06-0.28-0.07}) \times 10^{-4}$
$B_c \rightarrow B_d^0 \rho^+$	1-b	8.3×10^{-3}	6.9×10^{-3}	3.3×10^{-3}	1.5×10^{-3}	1.6×10^{-3}	2.2×10^{-3} (1.7×10^{-3})	1.4×10^{-2}	2.4×10^{-3}	1.1×10^{-2}	4.3×10^{-3}	$(5.32^{+0.03+0.49+0.15}_{-0.03-0.28-0.15}) \times 10^{-3}$
$B_c \rightarrow B_d^0 K^{*+}$	1-c		2.1×10^{-4}	1.5×10^{-4}	4.6×10^{-5}	4.4×10^{-5}	4.9×10^{-5} (3.7×10^{-5})	3.5×10^{-4}	5.7×10^{-5}	1.7×10^{-4}	8.3×10^{-5}	$(1.06^{+0.01+0.10+0.05}_{-0.01-0.06-0.05}) \times 10^{-4}$
$B_c \rightarrow B_u^+ \bar{K}^0$	2-a	2.1×10^{-2}	1.2×10^{-2}	4.9×10^{-3}	4.2×10^{-3}	4.4×10^{-3}	6.0×10^{-3} (4.9×10^{-3})	2.3×10^{-2}	6.8×10^{-3}	3.6×10^{-2}	2.2×10^{-3}	$(1.97^{+0.00+1.11+0.05}_{-0.00-0.54-0.05}) \times 10^{-2}$
$B_c \rightarrow B_u^+ K^0$	2-c						1.6×10^{-5} (1.3×10^{-5})				6.3×10^{-6}	$(5.71^{+0.06+3.20+0.15}_{-0.06-1.58-0.14}) \times 10^{-5}$
$B_c \rightarrow B_u^+ \bar{K}^{*0}$	2-a	7.8×10^{-3}	8.5×10^{-3}	5.8×10^{-3}	1.6×10^{-3}	1.6×10^{-3}	1.9×10^{-3} (1.4×10^{-3})	1.3×10^{-2}	2.1×10^{-3}	8.0×10^{-3}	2.0×10^{-4}	$(3.72^{+0.00+2.09+0.21}_{-0.00-1.03-0.20}) \times 10^{-3}$
$B_c \rightarrow B_u^+ K^{*0}$	2-c						5.0×10^{-6} (3.7×10^{-6})				5.6×10^{-7}	$(1.07^{+0.01+0.60+0.06}_{-0.01-0.30-0.06}) \times 10^{-5}$
$B_c \rightarrow B_u^+ \pi^0$	2-b	4.0×10^{-4}	2.1×10^{-4}	6.4×10^{-5}	6.2×10^{-5}	6.7×10^{-5}	9.7×10^{-5} (8.1×10^{-5})	4.5×10^{-4}	1.3×10^{-4}	6.6×10^{-4}	5.2×10^{-5}	$(4.23^{+0.02+2.37+0.07}_{-0.02-1.17-0.07}) \times 10^{-4}$
$B_c \rightarrow B_u^+ \rho^0$	2-b	4.4×10^{-4}	3.7×10^{-4}	1.7×10^{-4}	8.7×10^{-5}	8.9×10^{-5}	1.2×10^{-4} (9.4×10^{-5})	7.4×10^{-4}	1.3×10^{-4}	5.5×10^{-4}	1.8×10^{-5}	$(2.86^{+0.01+1.60+0.10}_{-0.01-0.79-0.10}) \times 10^{-4}$
$B_c \rightarrow B_u^+ \omega$	2-b	4.1×10^{-4}					9.0×10^{-5} (7.0×10^{-5})				1.3×10^{-5}	$(2.05^{+0.01+1.15+0.12}_{-0.01-0.57-0.12}) \times 10^{-4}$

TABLE 3: Continued.

Decay mode	Case	Reference [17] ^a	Reference [18] ^b	Reference [19] ^c	Reference [20] ^d	Reference [21, 22] ^e	Reference [23] ^f	Reference [24] ^g	Reference [25] ^h	Reference [26–28] ⁱ	Reference [29, 30] ^j	QCDF
$B_c \rightarrow B_u^+ \eta$							5.0×10^{-4}				1.4×10^{-4}	$(1.46^{+0.01+0.82+0.13}_{-0.01-0.40-0.12}) \times 10^{-3}$
$B_c \rightarrow B_u^+ \eta'$							(4.1×10^{-4}) 6.7×10^{-6} (5.6×10^{-6})				4.2×10^{-6}	$(7.28^{+0.04+4.09+1.66}_{-0.04-2.01-1.47}) \times 10^{-5}$

^aIt is estimated with the form factors $F_0^{B_c \rightarrow B_s} = 0.925$, $F_0^{B_c \rightarrow B} = 0.91$ and parameter $\omega = 1$ GeV [17] based on the BSW model.

^bIt is estimated with the instantaneous nonrelativistic approximation and the potential model based on the BS equation.

^cIt is estimated in a relativistic model with a one-gluon interaction plus a scalar confinement potential based on the BS equation.

^dIt is estimated with the form factors $F_0^{B_c \rightarrow B_s} = 0.5$, $F_0^{B_c \rightarrow B} = 0.39$ using a quasipotential in the relativistic quark model [20].

^eIt is estimated with the form factors $F_0^{B_c \rightarrow B_s} = 0.58$, $F_0^{B_c \rightarrow B} = 0.39$ in the nonrelativistic constituent quark model [21, 22].

^fIt is estimated with the form factors $F_0^{B_c \rightarrow B_s} = 0.573$ (0.571), $F_0^{B_c \rightarrow B} = 0.467$ (0.426) in the light-front quark model based on the Coulomb plus linear (harmonic oscillator) potential, together with the hyperfine interaction [23].

^gIt is estimated with the form factors $F_0^{B_c \rightarrow B_s} = 1.03$, $F_0^{B_c \rightarrow B} = 1.01$ in the relativistic independent quark model [24].

^hIt is estimated within a relativistic constituent quark model [25].

ⁱIt is estimated with the form factors $F_0^{B_c \rightarrow B_s} = 1.3$, $F_0^{B_c \rightarrow B} = 1.27$ in the QCD sum rules [26–28].

^jIt is estimated with the perturbative QCD approach based on the k_T factorization scheme [29, 30].

TABLE 4: Hierarchy of amplitudes among the QCDF's branching ratios for B_c decay.

Case	Coefficient	CKM factor	Branching ratio	Decay modes
1 a	a_1	$ V_{ud}V_{cs}^* \sim 1$	$\geq 10^{-2}$	$B_s\pi, B_s\rho$
1 b	a_1	$ V_{ud}V_{cd}^* , V_{us}V_{cs}^* \sim \lambda$	$\geq 10^{-3}$	$B_sK, B_d\pi, B_d\rho$
1 c	a_1	$ V_{us}V_{cd}^* \sim \lambda^2$	$\geq 10^{-5}$	B_dK, B_dK^*
2 a	a_2	$ V_{ud}V_{cs}^* \sim 1$	$\geq 10^{-3}$	$B_u^+\bar{K}^0, B_u^+\bar{K}^{*0}$
2 b	a_2	$ V_{ud}V_{cd}^* , V_{us}V_{cs}^* \sim \lambda$	$\geq 10^{-4}$	$B_u\pi, B_u\rho, B_u\omega$
2 c	a_2	$ V_{us}V_{cd}^* \sim \lambda^2$	$\geq 10^{-6}$	$B_u^+K^0, B_u^+K^{*0}$

orbital angular momentum for the BP final states is $\ell_{BP} = 0$, while the orbital angular momentum is $\ell_{BV} = 1$ for the BV final states.

(3) According to the CKM factors and the coefficients $a_{1,2}$, there is another hierarchy of amplitudes among the QCDF's branching ratios for the B_c decays, which could be subdivided into different cases (see Table 4). The CKM-favored a_1 -dominated $B_c \rightarrow B_s\pi$ decays are expected to have the largest branching ratio, $\sim 10\%$, within the QCDF framework. In addition, the branching ratios for the Cabibbo favored $B_c^+ \rightarrow B_s^0\rho^+, B_u^+\bar{K}^0$ decays are also larger than 1%, which might be promisingly detected at experiments.

(4) There are many uncertainties on the QCDF's results. The first uncertainty from the CKM factors is small due to the high precision on Wolfenstein parameter λ with only 0.3% relative errors [1]. Large uncertainty comes from the renormalization scale, especially for the a_2 dominated $B_c \rightarrow B_uP, B_uV$ decays. In principle, the second uncertainty could be reduced by the inclusion of higher order α_s corrections to hadronic matrix elements. It has been showed [77, 78] that tree amplitudes incorporating with the NNLO corrections are relatively less sensitive to the choice of scale than the NLO amplitudes. As aforementioned, large uncertainty mainly comes from hadron parameters, such as the transition form factors, which is expected to be cancelled from the rate of branching ratios. For example,

$$\frac{\mathcal{B}r(B_c \rightarrow B_sK)}{\mathcal{B}r(B_c \rightarrow B_s\pi)} \approx |V_{us}|^2 \frac{f_K^2}{f_\pi^2} \approx \frac{\mathcal{B}r(B_c \rightarrow B_dK)}{\mathcal{B}r(B_c \rightarrow B_d\pi)}, \quad (16)$$

$$\frac{\mathcal{B}r(B_c \rightarrow B_sK^*)}{\mathcal{B}r(B_c \rightarrow B_s\rho)} \approx |V_{us}|^2 \frac{f_{K^*}^2}{f_\rho^2} \approx \frac{\mathcal{B}r(B_c \rightarrow B_dK^*)}{\mathcal{B}r(B_c \rightarrow B_d\rho)}, \quad (17)$$

$$\frac{\mathcal{B}r(B_c \rightarrow B_u\pi)}{\mathcal{B}r(B_c \rightarrow B_d\pi)} \approx \frac{1}{2} \frac{|a_2|^2}{|a_1|^2} \approx \frac{\mathcal{B}r(B_c \rightarrow B_u\rho)}{\mathcal{B}r(B_c \rightarrow B_d\rho)}. \quad (18)$$

Particularly, the relation of (18) might be used to give some information on the coefficients $a_{1,2}$ and to provide an interesting feasibility research on the validity of the QCDF approach for the charm quark decay. Finally, we would like to point out that many uncertainties from other factors, such as the final state interactions, which deserve the dedicated study, are not considered here. So one should not be too serious about the numbers in Table 3. Despite this, our results will still

provide some useful information to experimental physicists; that is, the Cabibbo favored $B_c \rightarrow B_s\pi, B_s\rho, B_u\bar{K}$ decays have large branching ratios $\geq 1\%$, which could be detected earlier.

4. Summary

In prospects of the potential B_c meson at the LHCb experiments, accurate and thorough studies of the B_c decays will be accessible very soon. The carefully theoretical study on the B_c decays is urgently desiderated. In this paper, we concentrated on the nonfactorizable contributions to hadronic matrix elements within the QCDF framework, while the transition form factors are taken as nonperturbative inputs, which is different from previous studies. It is found that the branching ratios for the Cabibbo favored $B_c \rightarrow B_s\pi, B_s\rho, B_u\bar{K}$ decays are very large and could be measured earlier by the running LHCb experiment in the forthcoming years.

Appendix

Decay Amplitudes

$$\begin{aligned}
\mathcal{A}(B_c^+ \rightarrow B_s^0\pi^+) &= -i \frac{G_F}{\sqrt{2}} F_0^{B_c \rightarrow B_s} f_\pi (m_{B_c}^2 - m_{B_s}^2) V_{ud} V_{cs}^* a_1, \\
\mathcal{A}(B_c^+ \rightarrow B_s^0K^+) &= -i \frac{G_F}{\sqrt{2}} F_0^{B_c \rightarrow B_s} f_K (m_{B_c}^2 - m_{B_s}^2) V_{us} V_{cs}^* a_1, \\
\mathcal{A}(B_c^+ \rightarrow B_s^0\rho^+) &= \sqrt{2} G_F F_1^{B_c \rightarrow B_s} f_\rho m_\rho (\epsilon_\rho \cdot p_{B_c}) V_{ud} V_{cs}^* a_1, \\
\mathcal{A}(B_c^+ \rightarrow B_s^0K^{*+}) &= \sqrt{2} G_F F_1^{B_c \rightarrow B_s} f_{K^*} m_{K^*} (\epsilon_{K^*} \cdot p_{B_c}) V_{us} V_{cs}^* a_1, \\
\mathcal{A}(B_c^+ \rightarrow B_d^0\pi^+) &= -i \frac{G_F}{\sqrt{2}} F_0^{B_c \rightarrow B_d} f_\pi (m_{B_c}^2 - m_{B_d}^2) V_{ud} V_{cd}^* a_1,
\end{aligned}$$

$$\begin{aligned}
\mathcal{A}(B_c^+ \rightarrow B_d^0 K^+) &= -i \frac{G_F}{\sqrt{2}} F_0^{B_c \rightarrow B_d} f_K (m_{B_c}^2 - m_{B_d}^2) V_{us} V_{cd}^* a_1, \\
\mathcal{A}(B_c^+ \rightarrow B_d^0 \rho^+) &= \sqrt{2} G_F F_1^{B_c \rightarrow B_d} f_\rho m_\rho (\epsilon_\rho \cdot p_{B_c}) V_{ud} V_{cd}^* a_1, \\
\mathcal{A}(B_c^+ \rightarrow B_d^0 K^{*+}) &= \sqrt{2} G_F F_1^{B_c \rightarrow B_d} f_{K^*} m_{K^*} (\epsilon_{K^*} \cdot p_{B_c}) V_{us} V_{cd}^* a_1, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \bar{K}^0) &= -i \frac{G_F}{\sqrt{2}} F_0^{B_c \rightarrow B_u} f_K (m_{B_c}^2 - m_{B_u}^2) V_{ud} V_{cs}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ K^0) &= -i \frac{G_F}{\sqrt{2}} F_0^{B_c \rightarrow B_u} f_K (m_{B_c}^2 - m_{B_u}^2) V_{us} V_{cd}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \bar{K}^{*0}) &= \sqrt{2} G_F F_1^{B_c \rightarrow B_u} f_{K^*} m_{K^*} (\epsilon_{K^*} \cdot p_{B_c}) V_{ud} V_{cs}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ K^{*0}) &= \sqrt{2} G_F F_1^{B_c \rightarrow B_u} f_{K^*} m_{K^*} (\epsilon_{K^*} \cdot p_{B_c}) V_{us} V_{cd}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \pi^0) &= +i \frac{G_F}{2} F_0^{B_c \rightarrow B_u} f_\pi (m_{B_c}^2 - m_{B_u}^2) V_{ud} V_{cd}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \rho^0) &= -G_F F_1^{B_c \rightarrow B_u} f_\rho m_\rho (\epsilon_\rho \cdot p_{B_c}) V_{ud} V_{cd}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \omega) &= +G_F F_1^{B_c \rightarrow B_u} f_\omega m_\omega (\epsilon_\omega \cdot p_{B_c}) V_{ud} V_{cd}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \eta_q) &= -i \frac{G_F}{2} F_0^{B_c \rightarrow B_u} f_{\eta_q} (m_{B_c}^2 - m_{B_u}^2) V_{ud} V_{cd}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \eta_s) &= -i \frac{G_F}{\sqrt{2}} F_0^{B_c \rightarrow B_u} f_{\eta_s} (m_{B_c}^2 - m_{B_u}^2) V_{us} V_{cs}^* a_2, \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \eta) &= \cos \phi \mathcal{A}(B_c^+ \rightarrow B_u^+ \eta_q) - \sin \phi \mathcal{A}(B_c^+ \rightarrow B_u^+ \eta_s), \\
\mathcal{A}(B_c^+ \rightarrow B_u^+ \eta') &= \sin \phi \mathcal{A}(B_c^+ \rightarrow B_u^+ \eta_q) + \cos \phi \mathcal{A}(B_c^+ \rightarrow B_u^+ \eta_s).
\end{aligned} \tag{A.1}$$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The work is supported by the National Natural Science Foundation of China (Grant nos. 11475055, 11275057, and U1232101). N. Wang thanks the support from CCNU-QLPL Innovation Fund (QLPL201411). Q. Chang is also supported by the Foundation for the Author of National Excellent Doctoral Dissertation of China (Grant no. 201317) and the Program for Science and Technology Innovation Talents in Universities of Henan Province (Grant no. 14HASTIT036).

References

- [1] K. Olive, K. Agashe, C. Amsler et al., “Review of particle physics,” *Chinese Physics C*, vol. 38, no. 9, Article ID 090001, 2014.
- [2] N. Brambilla, M. Krämer, R. Mussa et al., “Heavy quarkonium physics,” <http://arxiv.org/abs/hep-ph/0412158>.
- [3] F. Abe, H. Akimoto, A. Akopian et al., “Observation of Bc mesons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV,” *Physical Review D*, vol. 58, no. 11, Article ID 112004, 29 pages, 1998.
- [4] F. Abe, H. Akimoto, A. Akopian et al., “Observation of the Bc meson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV,” *Physical Review Letters*, vol. 81, p. 2432, 1998.
- [5] R. Aaij, C. A. Beteta, B. Adeva et al., “Observation of $B_c^+ \rightarrow J/\psi D_s^+$ and $B_c^+ \rightarrow J/\psi D_s^{*+}$ decays,” *Physical Review D*, vol. 87, Article ID 112012, 2013.
- [6] R. Aaij, B. Adeva, M. Adinolfi et al., “Measurement of the lifetime of the Bc meson using the $B_c^+ \rightarrow J/\psi \pi^+$ decay mode,” *Physics Letters B*, vol. 742, pp. 29–37, 2015.
- [7] I. P. Gouz, V. V. Kiselev, A. K. Likhoded, V. I. Romanovsky, and O. P. Yushchenko, “Prospects for the Bc studies at LHCb,” *Physics of Atomic Nuclei*, vol. 67, no. 8, pp. 1559–1570, 2004.
- [8] A. Likhoded and A. Luchinsky, “Light hadron production in $B_c \rightarrow B_s^{(*)} + X$ decays,” *Physical Review D*, vol. 82, Article ID 014012, 2010.
- [9] M. Lusignoli and M. Masetti, “Bc decays,” *Zeitschrift für Physik C: Particles and Fields*, vol. 51, no. 4, pp. 549–555, 1991.
- [10] C. Chang and Y. Chen, “Decays of the Bc meson,” *Physical Review D*, vol. 49, no. 7, pp. 3399–3411, 1994.
- [11] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. V. Tkabladze, “Reviews of topical problems: physics of Bc-mesons,” *Physics-Uspekhi*, vol. 38, no. 1, pp. 1–37, 1995.
- [12] R. Aaij, B. Adeva, M. Adinolfi et al., “Observation of the decay $B_c^+ \rightarrow B_s^0 \pi^+$,” *Physical Review Letters*, vol. 111, Article ID 181801, 2013.
- [13] T. Feldmann, P. Kroll, and B. Stech, “Mixing and decay constants of pseudoscalar mesons,” *Physical Review D*, vol. 58, Article ID 114006, 1998.
- [14] P. Ball, “Theoretical update of pseudoscalar meson distribution amplitudes of higher twist: the nonsinglet case,” *Journal of High Energy Physics*, vol. 1999, no. 1, article 10, 1999.
- [15] P. Ball, V. M. Braun, and A. Lenz, “Higher-twist distribution amplitudes of the K meson in QCD,” *Journal of High Energy Physics*, vol. 2006, no. 5, article 4, 2006.

- [16] P. Ball and G. Jones, "Twist-3 distribution amplitudes of K^* and ϕ mesons," *Journal of High Energy Physics*, vol. 2007, no. 3, article 069, 2007.
- [17] D. Du and Z. Wang, "Predictions of the standard model for $B_c^\pm$ weak decays," *Physical Review D*, vol. 39, no. 5, pp. 1342–1348, 1989.
- [18] C.-H. Chang and Y.-Q. Chen, "Decays of the B_c meson," *Physical Review D*, vol. 49, no. 7, Article ID 3399, 1994.
- [19] A. El-Hady, J. Muñoz, and J. Vary, "Semileptonic and nonleptonic B_c decays," *Physical Review D*, vol. 62, Article ID 014019, 2000.
- [20] D. Ebert, R. N. Faustov, and V. O. Galkin, "Weak decays of the B_c meson to B_s and B mesons in the relativistic quark model," *European Physical Journal C*, vol. 32, no. 1, pp. 29–43, 2003.
- [21] E. Hernández, J. Nieves, and J. Verde-Velasco, "Study of exclusive semileptonic and nonleptonic decays of B_c in a nonrelativistic quark model," *Physical Review D*, vol. 74, Article ID 074008, 2006.
- [22] E. Hernández, J. Nieves, and J. M. Verde-Velasco, "Study of semileptonic and nonleptonic decays of the B_c -Meson," *European Physical Journal A*, vol. 31, no. 4, pp. 714–717, 2007.
- [23] H. Choi and C. Ji, "Nonleptonic two-body decays of the B_c meson in the light-front quark model and the QCD factorization approach," *Physical Review D*, vol. 80, Article ID 114003, 2009.
- [24] S. Naimuddin, S. Kar, M. Priyadarsini, N. Barik, and P. C. Dash, "Nonleptonic two-body B_c -meson decays," *Physical Review D*, vol. 86, Article ID 094028, 2012.
- [25] M. Ivanov, J. Körner, and P. Santorelli, "Exclusive semileptonic and nonleptonic decays of the B_c meson," *Physical Review D*, vol. 73, Article ID 054024, 2006.
- [26] V. V. Kiselev, A. E. Kovalsky, and A. K. Likhoded, " B_c decays and lifetime in QCD sum rules," *Nuclear Physics B*, vol. 585, no. 1-2, pp. 353–382, 2000.
- [27] V. V. Kiselev, A. E. Kovalsky, and A. K. Likhoded, "Bc-Meson decays and lifetime within QCD sum rules," *Physics of Atomic Nuclei*, vol. 64, no. 10, pp. 1860–1875, 2001.
- [28] I. P. Gouz, V. V. Kiselev, A. K. Likhoded, V. I. Romanovsky, and O. P. Yushchenko, "Prospects for the B_c studies at LHCb," *Physics of Atomic Nuclei*, vol. 67, no. 8, pp. 1559–1570, 2004.
- [29] J. Sun, Y. Yang, Q. Chang, and G. Lu, "Phenomenological study of the $B_c \rightarrow BP, BV$ decays with perturbative QCD approach," *Physical Review D*, vol. 89, no. 11, Article ID 114019, 17 pages, 2014.
- [30] J. Sun, Y. Yang, and G. Lu, "Study of the $B_c \rightarrow B_s \pi$ decay with the perturbative QCD approach," *Science China Physics, Mechanics & Astronomy*, vol. 57, no. 10, pp. 1891–1897, 2014.
- [31] M. Beneke and G. Buchalla, " B_c meson lifetime," *Physical Review D*, vol. 53, article 4991, 1996.
- [32] C. Chang, S. Chen, T. Feng, and X. Li, "Lifetime of the B_c meson and some relevant problems," *Physical Review D*, vol. 64, Article ID 014003, 2001.
- [33] C.-H. Chang, S.-L. Chen, T.-F. Feng, and X.-Q. Li, "Study of the B_c -Meson Lifetime," *Communications in Theoretical Physics*, vol. 35, no. 1, p. 57, 2001.
- [34] M. Lusignol and M. Masetti, " B_c decays," *Zeitschrift für Physik C: Particles and Fields*, vol. 51, no. 4, pp. 549–555, 1991.
- [35] A. Y. Anisimov, P. Y. Kulikov, I. M. Narodetskii, and K. A. Ter-Martirosyan, "Exclusive and inclusive decays of the B_c meson in the light-front ISGW model," *Physics of Atomic Nuclei*, vol. 62, no. 10, pp. 1739–1753, 1999.
- [36] R. Dhir, N. Sharma, and R. Verma, "Flavor dependence of B_c meson form factors and $B_c \rightarrow PP$ decays," *Journal of Physics G: Nuclear and Particle Physics*, vol. 35, no. 8, Article ID 085002, 2008.
- [37] R. Dhir and R. Verma, " B_c meson form factors and $B_c \rightarrow PV$ decays involving flavor dependence of transverse quark momentum," *Physical Review D*, vol. 79, Article ID 034004, 2009.
- [38] M. Wirbel, B. Stech, and M. Bauer, "Exclusive semileptonic decays of heavy mesons," *Zeitschrift für Physik C Particles and Fields*, vol. 29, no. 4, pp. 637–642, 1985.
- [39] M. Bauer, B. Stech, and M. Wirbel, "Exclusive non-leptonic decays of D^- , D_s^- and B -mesons," *Zeitschrift für Physik C Particles and Fields*, vol. 34, no. 1, pp. 103–115, 1987.
- [40] N. Isgur, D. Scora, B. Grinstein, and M. B. Wise, "Semileptonic B and D decays in the quark model," *Physical Review D*, vol. 39, no. 3, pp. 799–818, 1989.
- [41] J. Liu and K. Chao, " B_c meson weak decays and CP violation," *Physical Review D*, vol. 56, no. 7, pp. 4133–4145, 1997.
- [42] P. Colangelo and F. de Fazio, "Using heavy quark spin symmetry in semileptonic B_c decays," *Physical Review D*, vol. 61, no. 3, Article ID 034012, 11 pages, 2000.
- [43] R. C. Verma and A. Sharma, "Quark diagram analysis of weak hadronic decays of the B_c^+ meson," *Physical Review D*, vol. 65, Article ID 114007, 2002.
- [44] C. Chang and H. Li, "Three-scale factorization theorem and effective field theory: analysis of nonleptonic heavy meson decays," *Physical Review D*, vol. 55, p. 5577, 1997.
- [45] T.-W. Yeh and H.-N. Li, "Factorization theorems, effective field theory, and nonleptonic heavy meson decays," *Physical Review D*, vol. 56, pp. 1615–1631, 1997.
- [46] Y. Keum, H. Li, and A. Sanda, "Fat penguins and imaginary penguins in perturbative QCD," *Physics Letters B*, vol. 504, no. 1-2, pp. 6–14, 2001.
- [47] Y.-Y. Keum and H.-N. Li, "Nonleptonic charmless B decays: factorization versus perturbative QCD," *Physical Review D*, vol. 63, Article ID 074006, 2001.
- [48] C. Lü, K. Ukai, and M. Yang, "Branching ratio and CP violation of $\bar{B}\pi\pi$ decays in the perturbative QCD approach," *Physical Review D*, vol. 63, Article ID 074009, 2001.
- [49] C.-D. Lü and M.-Z. Yang, " $B \rightarrow \pi\rho, \pi\omega$ decays in perturbative QCD approach," *The European Physical Journal C*, vol. 23, no. 2, pp. 275–287, 2002.
- [50] C. Bauer, S. Fleming, and M. Luke, "Summing Sudakov logarithms in $B \rightarrow X_s \gamma$ in effective field theory," *Physical Review D*, vol. 63, Article ID 014006, 2001.
- [51] C. W. Bauer, S. Fleming, D. Pirjol, and I. W. Stewart, "An effective field theory for collinear and soft gluons: heavy to light decays," *Physical Review D*, vol. 63, Article ID 114020, 2001.
- [52] C. W. Bauer and I. W. Stewart, "Invariant operators in collinear effective theory," *Physics Letters B*, vol. 516, no. 1-2, pp. 134–142, 2001.
- [53] C. W. Bauer, D. Pirjol, and I. W. Stewart, "Soft-collinear factorization in effective field theory," *Physical Review D*, vol. 65, Article ID 054022, 2002.
- [54] C. Bauer, S. Fleming, D. Pirjol, I. Z. Rothstein, and I. W. Stewart, "Hard scattering factorization from effective field theory," *Physical Review D*, vol. 66, no. 1, Article ID 014017, 23 pages, 2002.
- [55] M. Beneke, A. P. Chapovsky, M. Diehl, and T. Feldmann, "Soft-collinear effective theory and heavy-to-light currents beyond

- leading power,” *Nuclear Physics B*, vol. 643, no. 1–3, pp. 431–476, 2002.
- [56] M. Beneke and T. Feldmann, “Multipole-expanded soft-collinear effective theory with non-Abelian gauge symmetry,” *Physics Letters B*, vol. 553, no. 3–4, pp. 267–276, 2003.
- [57] M. Beneke and T. Feldmann, “Factorization of heavy-to-light form factors in soft-collinear effective theory,” *Nuclear Physics B*, vol. 685, no. 1–3, pp. 249–296, 2004.
- [58] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, “QCD factorization for $B \rightarrow \pi\pi$ decays: strong phases and CP violation in the heavy quark limit,” *Physical Review Letters*, vol. 83, no. 10, pp. 1914–1917, 1999.
- [59] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, “QCD factorization for exclusive non-leptonic B -meson decays: general arguments and the case of heavy-light final states,” *Nuclear Physics B*, vol. 591, no. 1–2, pp. 313–418, 2000.
- [60] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, “QCD factorization in $B \rightarrow \pi K, \pi\pi$ decays and extraction of Wolfenstein parameters,” *Nuclear Physics B*, vol. 606, no. 1–2, pp. 245–321, 2001.
- [61] D. Du, D. Yang, and G. Zhu, “Infrared divergence and twist-3 distribution amplitudes in QCD factorization for $B \rightarrow PP$,” *Physics Letters B*, vol. 509, no. 3–4, pp. 263–272, 2001.
- [62] D. Du, D. Yang, and G. Zhu, “Analysis of the decays $B \rightarrow \pi\pi$ and πK with QCD factorization in the heavy quark limit,” *Physics Letters B*, vol. 488, no. 1, pp. 46–54, 2000.
- [63] D. Du, D. Yang, and G. Zhu, “QCD factorization for $\bar{B}PP$,” *Physical Review D*, vol. 64, Article ID 014036, 2001.
- [64] G. Buchalla, A. Buras, and M. Lautenbacher, “Weak decays beyond leading logarithms,” *Reviews of Modern Physics*, vol. 68, p. 1125, 1996.
- [65] A. J. Buras, “Weak hamiltonian, CP violation and rare decays,” <http://arxiv.org/abs/hep-ph/9806471>.
- [66] D. Du, H. Gong, J. Sun, D. Yang, and G. Zhu, “Phenomenological analysis of $\bar{B} PP$ decays with QCD factorization,” *Physical Review D*, vol. 65, Article ID 074001, 2002.
- [67] D. Du, H. Gong, J. Sun, D. Yang, and G. Zhu, “Phenomenological analysis of charmless decays $B \rightarrow PV$ with QCD factorization,” *Physical Review D*, vol. 65, Article ID 094025, 2002, Erratum in *Physical Review D*, vol. 66, Article ID 079904, 2002.
- [68] J. Sun, G. Zhu, and D. Du, “Phenomenological analysis of charmless decays $B_s \rightarrow PP, PV$ with QCD factorization,” *Physical Review D*, vol. 68, Article ID 054003, 2003.
- [69] M. Beneke and M. Neubert, “QCD factorization for $B \rightarrow PP$ and $B \rightarrow PV$ decays,” *Nuclear Physics B*, vol. 675, no. 1–2, pp. 333–415, 2003.
- [70] M. Beneke, J. Rohrer, and D. Yang, “Branching fractions, polarisation and asymmetries of $B \rightarrow VV$ decays,” *Nuclear Physics B*, vol. 774, no. 1–3, pp. 64–101, 2007.
- [71] X. Li and Y. Yang, “Reexamining charmless $B \rightarrow PV$ decays in the QCD factorization approach,” *Physical Review D*, vol. 73, no. 11, Article ID 114027, 24 pages, 2006.
- [72] H.-Y. Cheng and K.-C. Yang, “Branching ratios and polarization in $B \rightarrow VV, VA, AA$ decays,” *Physical Review D*, vol. 78, Article ID 094001, 2008, Erratum in *Physical Review D*, vol. 79, Article ID 039903, 2009.
- [73] H.-Y. Cheng and C.-K. Chua, “Resolving B -CP puzzles in QCD factorization,” *Physical Review D*, vol. 80, Article ID 074031, 2009.
- [74] H. Cheng and C. Chua, “Revisiting charmless hadronic $B_{u,d}$ decays in QCD factorization,” *Physical Review D*, vol. 80, no. 11, Article ID 114008, 33 pages, 2009.
- [75] H. Cheng and C. Chua, “QCD factorization for charmless hadronic B_s decays revisited,” *Physical Review D*, vol. 80, no. 11, Article ID 114026, 25 pages, 2009.
- [76] H. Cheng and C. Chua, “Charmless hadronic B decays into a tensor meson,” *Physical Review D*, vol. 83, Article ID 034001, 2011.
- [77] G. Bell, “NNLO vertex corrections in charmless hadronic B decays: real part,” *Nuclear Physics B*, vol. 822, no. 1–2, pp. 172–200, 2009.
- [78] M. Beneke, T. Huber, and X.-Q. Li, “NNLO vertex corrections to non-leptonic B decays: tree amplitudes,” *Nuclear Physics B*, vol. 832, no. 1–2, pp. 109–151, 2010.