Research Article

π_{e3} Form Factor f_- Near the Mass Shell

M. I. Krivoruchenko^{1,2,3}

¹*Institute for Theoretical and Experimental Physics, B. Cheremushkinskaya 25, Moscow 117218, Russia*

²*Moscow Institute of Physics and Technology, Dolgoprudny 141700, Russia*

³*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna 141980, Russia*

Correspondence should be addressed to M. I. Krivoruchenko; mikhail.krivoruchenko@itep.ru

Received 29 November 2014; Accepted 6 April 2015

Academic Editor: Enrico Lunghi

Copyright © 2015 M. I. Krivoruchenko. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

The generalized Ward-Takahashi identity (gWTI) in the pion sector for broken isotopic symmetry is derived and used for the model-independent calculation of the longitudinal form factor f_- of the π_{e3} vector vertex. The on-shell f_- is found to be proportional to the mass difference of the pions and the difference between the vector isospin $T = 1$ and scalar isospin $T = 2$ pion radii. A numerical estimate of the form factor yields a value two times higher than the previous estimate from the quark model. Off-shell form factors are known to be ambiguous because of the gauge dependence and the freedom in the parameterization of the fields. The near-mass-shell f_- appears to be an exception, allowing for experimental verification of the consequences of the gWTI. We calculate the near-mass-shell f_- using the gWTI and dispersion techniques. The results are discussed in the context of the conservation of vector current (CVC) hypothesis.

1. Introduction

The $\pi^+ \rightarrow \pi^0 e^+ \nu_e$ decay (π_{e3}) is one of the main semileptonic electroweak processes. The vector nature of the transition, its simple kinematics, and the precise measurement of the partial width make this decay particularly attractive for testing the Standard Model.

The decay amplitude is proportional to the V_{ud} element of the Cabibbo-Kobayashi-Maskawa (CKM) matrix. Radiative corrections and pion structure effects in the π_{e3} decay have been calculated with high accuracy [1–6], sufficient for verification of the unitarity of the CKM matrix. The experimental data, however, are not yet sufficiently precise for this purpose [7, 8].

Measurements of the π_{e3} decay are also motivated by the possibility of testing the conservation of vector current (CVC) in the meson sector. The CVC hypothesis [9–11] suggests that the isovector component of the electromagnetic current and the charged components of the weak vector current belong to the same isospin triplet. In the limit of exact isotopic symmetry, conservation of the electromagnetic current implies conservation of the weak vector current.

Off the mass shell, the CVC is equivalent to the Ward-Takahashi identity (WTI) for the isospin $SU(2)$ group. The WTI, however, is of greater generality and leads to useful relationships between off-shell form factors, including those that vanish when some of the external legs are on shell. The violation of isotopic symmetry, associated with the small mass difference between the up and down quarks and the electromagnetic and weak interactions, results in the nonconservation of the charge-changing components of the weak vector current. For the broken symmetry, the CVC condition and the WTI are replaced by partial CVC and the generalized WTI (gWTI), which are especially sensitive to the pattern of isotopic symmetry breaking.

Off-shell form factors enter the descriptions of nucleon knockout reactions [12, 13] and bremsstrahlung reactions [14]. As a result of shifting away from the mass shell, the electromagnetic currents of bound nucleons are different from the free currents, which also leads to observable effects [15].

The parameterization of the degrees of freedom associated with the pion field can be performed in various ways, which produces off-shell ambiguity in the amplitudes.

The on-shell form factors are related to the asymptotic states and are uniquely defined. This statement is known as the equivalence theorem (ET) [16–18]. The off-shell form factors, although they contribute to the physical amplitudes, depend on this parameterization and cannot be measured experimentally [19, 20].

We report a notable exception to this rule. The longitudinal part of the π_{e3} vertex is shown to be uniquely defined near the mass shell and fundamentally accessible to measurements, and therefore, it can be investigated for its consistency with the gWTI.

In quark models, pions are usually poorly described because of their Goldstone boson nature. As an application of the gWTI, we derive a model-independent expression and provide a numerical estimate for the longitudinal form factor f_- . The structure of this paper is as follows. Section 2 begins with a discussion of the constraints imposed by the WTI of the symmetry group $U(1)$ on the electromagnetic pion form factors. In Section 3, a generalization of the WTI for broken isotopic symmetry is derived, and in Section 4, it is used to find the relationships between weak vector form factors. The near-mass-shell form factor f_- is then expressed in terms of the pion mass difference and the pion radii in the isospin $T = 1$ and 2 channels. In Section 5, we focus on extracting the isospin $T = 2$ pion radius from the experimental data on $\pi\pi$ -scattering phase shifts and numerically evaluate f_- . Finally, we conclude in Section 6 with a summary and a discussion.

2. $U(1)$ Vector Vertex

The on-shell conserved vector current of a charged scalar particle is determined by one form factor. Off the mass shell, there are two form factors. In the most general case, the current can be written as follows:

$$\Gamma_\mu(p', p) = (p' + p)_\mu \mathcal{F}_1 + q_\mu (p'^2 - p^2) \mathcal{F}_2, \quad (1)$$

where $q = p' - p$ is the momentum transfer. The form factors $\mathcal{F}_i$ are symmetric functions of p'^2 and p^2 and arbitrary functions of q^2 and the physical mass m of the charged pion. The factor $p'^2 - p^2$ in the second term is added to ensure the negative C parity of the current. The form factor decomposition (1) arises in scalar quantum electrodynamics (QED) (see, e.g., [21]) and in chiral perturbation theory (χ PT) (see, e.g., [22]). The WTI of the $U(1)$ symmetry group establishes a relationship between $\mathcal{F}_1$ and $\mathcal{F}_2$ that is identical to that of (4).

Isotopic symmetry implies that the mass difference of nonstrange quarks can be neglected and that the electromagnetic and weak interactions are switched off beyond tree level. In this limit, the weak interaction is described by an isovector current $\Gamma_\mu^a(p', p) = T^a \Gamma_\mu(p', p)$, where the T^a are isospin generators. As a result, exact CVC occurs: a dressed vertex forms an isospin triplet, and the weak vector current is conserved.

The WTI associated with the $U(1)$ symmetry group imposes a constraint on the vertex (see, e.g., [23]):

$$\Delta^{-1}(p') - \Delta^{-1}(p) = q_\mu \Gamma_\mu(p', p), \quad (2)$$

where $\Delta(p) = p^2 - m^2 - \Sigma(p^2, m)$ is the renormalized pion propagator. The self-energy operator satisfies

$$\begin{aligned} \Sigma(m^2, m) &= 0, \\ \frac{\partial}{\partial p^2} \Sigma(p^2, m) \Big|_{p^2=m^2} &= 0. \end{aligned} \quad (3)$$

Combining (2) with (1) yields

$$\begin{aligned} q^2 \mathcal{F}_2(p'^2, p^2, q^2) &= \frac{\Delta^{-1}(p') - \Delta^{-1}(p)}{p'^2 - p^2} \\ &\quad - \mathcal{F}_1(p'^2, p^2, q^2). \end{aligned} \quad (4)$$

In the limit $p'^2 = p^2 = m^2$, we obtain

$$\mathcal{F}_2(m^2, m^2, q^2) = \frac{1 - \mathcal{F}_1(m^2, m^2, q^2)}{q^2}. \quad (5)$$

In the vicinity of $q^2 = 0$, the form factor $\mathcal{F}_1$ can be expanded to give

$$\mathcal{F}_2(m^2, m^2, 0) = -\frac{1}{6} \langle r^2 \rangle_v, \quad (6)$$

where $\langle r^2 \rangle_v$ is the vector charge radius.

The pion form factor $\mathcal{F}_1(p'^2, p^2, q^2)$, which describes strong interaction effects, is an analytic function of the complex variable p^2 at least in the circle $|p^2 - m^2| < 8m^2$. The nearest to the mass shell singularity at $p^2 = 9m^2$ is caused by the three-pion threshold. A Taylor series in powers of $p^2 - m^2$ converges within that circle. The same is true for the variable p'^2 . A Taylor series in powers of q^2 converges for $|q^2| < 4m^2$, where the radius of convergence is determined by the two-pion threshold.

The equivalence of the Coulomb and Lorentz gauges in QED was rigorously proved in [24, 25]. On shell, the amplitudes are gauge invariant, whereas the off-shell dependence on the gauge persists. $\mathcal{F}_1$ and, by virtue of (5), $\mathcal{F}_2$ are thus uniquely defined, when both legs of the charged pion are on the mass shell. To first order in the displacement from the mass shell, the longitudinal component of the vertex is also gauge invariant, as is evident from (1). Aside from these cases, $\Gamma_\mu(p', p)$ depends on the gauge and cannot be measured.

$\mathcal{F}_2$ with two on-shell legs $p'^2 = p^2 = m^2$ does not contribute to the current. Off the mass shell, however, $\mathcal{F}_2$ does contribute, and its contribution is uniquely determined by the WTI. Isotopic rotation of $\mathcal{F}_2$ is not sufficient to obtain a full weak-interaction vertex. We show that the violation of isotopic symmetry generates an isospin $T = 2$ contribution that is unrelated to isotopic rotation.

3. Generalized Ward-Takahashi Identity for Broken Isotopic Symmetry

In order to find the longitudinal form factor of the weak vector current, we derive a generalization of the WTI that is

associated with the replacement of the exact $U(1)$ symmetry by the broken $SU(2)$ symmetry.

Let us consider the variation of the pion propagator

$$i\Delta^{\alpha\beta}(x', x) = \langle 0 | T\varphi^\alpha(x')\varphi^\beta(x) | 0 \rangle \quad (7)$$

under the $SU(2)$ transformation

$$\varphi \longrightarrow \varphi' = e^{-i\chi}\varphi. \quad (8)$$

In terms of a Cartesian basis, the pion field $\varphi^\alpha(x)$ is Hermitian, and χ is an infinitesimal imaginary antisymmetric matrix that can be expanded in the group generators with real coefficients $\chi = \sum_a \chi^a T^a$. In the conventional normalization, $\text{Tr}(T^a T^b) = 2\delta^{ab}$. The variation of the pion propagator is given by

$$i\delta\Delta(x', x) = \chi(x')\Delta(x', x) - \Delta(x', x)\chi(x). \quad (9)$$

In matrix notation, $\Delta^{-1}i\delta\Delta\Delta^{-1} = -i\delta\Delta^{-1} = [\Delta^{-1}, \chi]$.

Independent calculation of $\delta\Delta$ involves the effective Lagrangian, which we consider to be of the form

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{eff}}\left((D_\mu\varphi)^\dagger, D_\mu\varphi, \varphi^\dagger, \varphi\right), \quad (10)$$

where $iD_\mu = i\partial_\mu - eA_\mu - B_\mu$, A_μ is the electromagnetic field, $e = T^3$, and $B_\mu = B_\mu^a T^a$ is the weak vector field associated with Z^0 and $W^\pm$ bosons. In a Cartesian basis, D_μ is real. The isotopic invariance in $\mathcal{L}_{\text{eff}}$ is broken by the mass term $\varphi^\dagger m^2 \varphi$ with $[m^2, T^a] \neq 0$ and by the electromagnetic and weak interactions. In other respects, $\mathcal{L}_{\text{eff}}$ is an arbitrary function, which reflects the ambiguity in the parameterization of the pion field.

The combination of $eA_\mu + B_\mu$ ensures the implementation of the CVC condition at a bare interaction vertex: the electromagnetic current coincides with the third component of the weak vector current by construction. If the isotopic symmetry in $\mathcal{L}_{\text{eff}}$ would not have been broken, the CVC condition could also hold for a dressed vertex, and the weak vector current could also be conserved. In the case of this broken symmetry, however, the dressed vertex acquires an admixture of tensor components, while the charge-changing weak vector current is no longer conserved. The WTI can be viewed as a generalization of the CVC for an off-shell vertex. It permits generalizations for broken symmetries; thus, when the isotopic symmetry is broken, the CVC becomes partial CVC, whereas the WTI of the exact isotopic symmetry turns into the WTI of the broken isotopic symmetry.

Transformation (8) is equivalent to the compensating transformation in $\mathcal{L}_{\text{eff}}$:

$$\varphi \longrightarrow \varphi' = e^{i\chi}\varphi, \quad (11)$$

which generates variations of the pion mass term and the vector fields in $\mathcal{L}_{\text{eff}}$:

$$\begin{aligned} m^2 &\longrightarrow m'^2 = m^2 + [m^2, i\chi], \\ eA_\mu &\longrightarrow eA'_\mu = eA_\mu + [e, i\chi]A_\mu, \\ B_\mu &\longrightarrow B'_\mu = B_\mu + \partial_\mu\chi + [B_\mu, i\chi]. \end{aligned} \quad (12)$$

As a result, $\mathcal{L}_{\text{eff}}$ acquires the correction

$$\begin{aligned} \delta\mathcal{L}_{\text{eff}} &= -\text{Tr}\left(\mathcal{F}_\mu(\partial^\mu\chi + [e, i\chi]A^\mu + [B^\mu, i\chi])\right) \\ &\quad - \text{Tr}\left(\mathcal{F}[m^2, i\chi]\right), \end{aligned} \quad (13)$$

where

$$\begin{aligned} \mathcal{F}_{\mu\beta}^\alpha &= -\frac{\partial\mathcal{L}_{\text{eff}}}{\partial B_{\alpha}^{\mu\beta}}, \\ \mathcal{F}_\beta^\alpha &= -\frac{\partial\mathcal{L}_{\text{eff}}}{\partial(m^2)_\alpha^\beta} \end{aligned} \quad (14)$$

are the vector and scalar currents of the pion.

The response of the pion propagator to the compensating transformation is

$$\begin{aligned} i\delta\Delta(x', x) &= \langle 0 | T\varphi(x')\tilde{\varphi}(x) i \int d^4y \delta\mathcal{L}_{\text{eff}}(y) | 0 \rangle \\ &= \int d^4y d^4z' d^4z i\Delta(x', z') \\ &\quad \cdot [-i\Gamma_\mu^a(z', z, y)\partial^\mu\chi^a(y) \\ &\quad + \Theta^a(z', z, y)\chi^a(y) + \Omega^a(z', z, y)\chi^a(y)] \\ &\quad \cdot i\Delta(z, x), \end{aligned} \quad (15)$$

with vertex functions defined by

$$\begin{aligned} \Gamma_\mu^a(z', z, y) &= -\int d^4x' d^4x \Delta^{-1}(z', x') \langle 0 | T\varphi(x') \\ &\quad \cdot \tilde{\varphi}(x) \text{Tr}(\mathcal{F}_\mu(y)T^a) | 0 \rangle \Delta^{-1}(x, z), \\ \Theta^a(z', z, y) &= -\int d^4x' d^4x \Delta^{-1}(z', x') \langle 0 | T\varphi(x') \\ &\quad \cdot \tilde{\varphi}(x) \text{Tr}(\mathcal{F}(y)[m^2, T^a]) | 0 \rangle \Delta^{-1}(x, z), \\ \Omega^a(z', z, y) &= -\int d^4x' d^4x \Delta^{-1}(z', x') \langle 0 | T\varphi(x') \\ &\quad \cdot \tilde{\varphi}(x) \text{Tr}(\mathcal{F}_\mu(y)[eA^\mu(y) + B^\mu(y), T^a]) | 0 \rangle \\ &\quad \cdot \Delta^{-1}(x, z). \end{aligned} \quad (16)$$

Now, let us consider momentum space:

$$\begin{aligned} (2\pi)^4 \delta^4(p' - p - q) \Gamma_\mu^a(p', p) \\ &= \int d^4z d^4z' d^4y e^{ip'z' - ipz - iqy} \Gamma_\mu^a(z', z, y), \\ (2\pi)^4 \delta^4(p' - p - q) \Theta^a(p', p) \\ &= \int d^4z d^4z' d^4y e^{ip'z' - ipz - iqy} \Theta^a(z', z, y), \\ (2\pi)^4 \delta^4(p' - p - q) \Omega^a(p', p) \\ &= \int d^4z d^4z' d^4y e^{ip'z' - ipz - iqy} \Omega^a(z', z, y). \end{aligned} \quad (17)$$

$$-i\text{Tr}(\mathcal{S}_\mu \partial^\mu \chi) \quad \text{Tr}(\mathcal{S}[m^2, \chi]) \quad \text{Tr}(\mathcal{S}_\mu [eA^\mu, \chi]) \quad \text{Tr}(\mathcal{S}_\mu [B^\mu, \chi])$$

FIGURE 1: Schematic representation of the gWTI for the pion propagator with broken isotopic symmetry. The blobs denote the dressed vertices. The first one is a Lorentz vector entering the pion β decay. The last three are Lorentz scalars; they are responsible for the symmetry breaking by the pion masses and the electromagnetic and weak interactions.

Combining (9) and (15) leads to

$$\begin{aligned} \Delta^{-1}(p') T^a - T^a \Delta^{-1}(p) \\ = q_\mu \Gamma_\mu^a(p', p) - \Theta^a(p', p) - \Omega^a(p', p). \end{aligned} \quad (18)$$

This equation generalizes the WTI for the broken $SU(2)$ symmetry. It is shown graphically in Figure 1.

4. $SU(2)$ Vector Vertex

In a Cartesian basis, $\Delta(x, y) = \tilde{\Delta}(y, z)$, which means that $\Delta(p) = \tilde{\Delta}(-p)$. Given that $\Delta^{-1}(p) = p^2 - m^2 - \Sigma(p^2)$, the pion propagator is symmetric under transposition of the isospin indices: $\Delta(p) = \tilde{\Delta}(p)$.

The isospin content of a 3×3 matrix $\mathcal{M}$ can be determined using projection operators:

$$\begin{aligned} (\mathcal{M})^{T=0} &= \frac{1}{2} \text{Tr}[\mathcal{M}], \\ (\mathcal{M})_a^{T=1} &= \frac{1}{2} \text{Tr}[\mathcal{M} T^a], \\ (\mathcal{M})_{ab}^{T=2} &= \frac{1}{2} \text{Tr} \left[\mathcal{M} \left(T^a T^b + T^b T^a - \frac{2}{3} \delta^{ab} T(T+1) \right) \right]. \end{aligned} \quad (19)$$

Among symmetric matrices, there are only isospin-0 or isospin-2 matrices that are proportional to the unit matrix or to quadratic forms of T^a . The allowed pion mass operator is therefore $m^2 + \Sigma(m^2) = C_0 + C_1(T^3)^2$, where the C_i are numbers. The charged pions are degenerate in mass; therefore, C invariance holds.

The vertex functions obey

$$\begin{aligned} \Gamma_\mu^a(p', p) &= \tilde{\Gamma}_\mu^a(-p, -p'), \\ \Theta^a(p', p) &= \tilde{\Theta}^a(-p, -p'), \\ \Omega^a(p', p) &= \tilde{\Omega}^a(-p, -p'). \end{aligned} \quad (20)$$

The functions $\Theta^a(p', p)$ and $\Omega^a(p', p)$ depend on the variables p'^2 , p^2 , and q^2 . Symmetry (antisymmetry) in the isospin indices implies that vertex function is symmetric (antisymmetric) under the permutation of p'^2 and p^2 . Functions that are symmetric in isospin describe states with isospin $T = 0$

and 2, whereas those that are antisymmetric describe isospin $T = 1$ states. Similar properties characterize the form factors of $\Gamma_\mu^a(p', p)$.

The vertex functions can be expanded in scalar functions $\mathcal{F}_{i\pm}^a$ that are symmetric in p'^2 and p^2 . The lower index $\pm$ indicates the symmetry with respect to permutation of the isospin indices: $\tilde{\mathcal{F}}_{i\pm}^a = \pm \mathcal{F}_{i\pm}^a$ ($i = 1, 2, 3$). The expansion takes the form

$$\begin{aligned} \Gamma_\mu^a(p', p) \\ = (p' + p)_\mu (\mathcal{F}_{1-}^a + (p'^2 - p^2) \mathcal{F}_{1+}^a) \end{aligned} \quad (21)$$

$$+ q_\mu ((p'^2 - p^2) \mathcal{F}_{2-}^a + \mathcal{F}_{2+}^a),$$

$$\Theta^a(p', p) + \Omega^a(p', p) = \mathcal{F}_{3+}^a + (p'^2 - p^2) \mathcal{F}_{3-}^a. \quad (22)$$

If there was no isospin symmetry breaking, we could have $\mathcal{F}_{i-}^a = T^a \mathcal{F}_{i-}$, as there are no other $SU(2)$ generators, and $\mathcal{F}_{i+}^a = 0$, implying that $\Gamma_\mu^a(m^2, m^2, q^2) \propto T^a$ and $\Theta^a(m^2, m^2, q^2) + \Omega^a(m^2, m^2, q^2) = 0$.

The gWTI (18) can be split into isospin-symmetric and isospin-antisymmetric parts:

$$\begin{aligned} - \left[m^2 + \frac{1}{2} (\Sigma(p') + \Sigma(p)), T^a \right] \\ = (p'^2 - p^2)^2 \mathcal{F}_{1+}^a + q^2 \mathcal{F}_{2+}^a - \mathcal{F}_{3+}^a, \end{aligned} \quad (23)$$

$$\begin{aligned} T^a - \frac{1}{2} \left\{ \frac{\Sigma(p') - \Sigma(p)}{p'^2 - p^2}, T^a \right\} \\ = \mathcal{F}_{1-}^a + q^2 \mathcal{F}_{2-}^a - \mathcal{F}_{3-}^a, \end{aligned} \quad (24)$$

where $[,]$ is a commutator and $\{, \}$ is an anticommutator.

Consider some of the consequences of (23). For $p' = p$, $q = 0$, we obtain

$$[m^2 + \Sigma(p^2), T^a] = \mathcal{F}_{3+}^a(p^2, p^2, 0). \quad (25)$$

In a cyclic basis, the $SU(2)$ generators are equal to $T_{\pm 1} = \mp(T^1 \pm iT^2)/\sqrt{2}$ and $T_0 = T^3$. The left-hand side of (25), when sandwiched between the initial and final states in the pion β decay, gives the pion mass difference:

$$\langle \pi^0 | [m^2 + \Sigma(p^2), T_{-1}] | \pi^+ \rangle = m_f^2 - m_i^2 \equiv -\Delta m_\pi^2, \quad (26)$$

where $m_{\pi^0} = m_f$ and $m_{\pi^+} = m_i$. In the more general case of $p'^2 = p^2$ and $q \neq 0$, one finds

$$\begin{aligned} [m^2 + \Sigma(p^2), T^a] &= -q^2 \mathcal{F}_{2+}^a(p^2, p^2, q^2) \\ &+ \mathcal{F}_{3+}^a(p^2, p^2, q^2). \end{aligned} \quad (27)$$

The trace of the commutator of finite-dimensional matrices vanishes, and therefore, the left-hand side of (25) and (27) describes a state of total isospin 2. The isospin-zero component of (27) satisfies the equation $q^2 (\mathcal{F}_{2+}^a)^{T=0} - (\mathcal{F}_{3+}^a)^{T=0} = 0$.

At zero momentum transfer, $(\mathcal{F}_{3+}^a)^{T=0} = 0$, and the overall normalization of the $T = 0$ form factors is not fixed. In the t channel of the pion β decay, the state $\pi^+\pi^0$ has the $T^3 = 1$ isospin projection, so the $T = 0$ channel is of no interest.

As a consequence of elastic unitarity and analyticity, the q^2 dependence of the on-mass-shell form factor is determined by the Jost function $D_J^T(q^2)$ that can be constructed in terms of the phase shift in the corresponding channel (see, e.g., [23]). We consider an S-wave because the last three vertices in $\delta\mathcal{L}_{\text{eff}}$ induced by the transformation (11) do not contain derivatives of χ^a , and thus, the S-wave $T = 2$ channel of the pion-pion scattering is relevant. We therefore write

$$(\mathcal{F}_{3+}^a(m_f^2, m_i^2, q^2))^{T=2} = \frac{\mathcal{F}_{3+}^a(m_f^2, m_i^2, 0)}{D_J^{T=2}(q^2)}, \quad (28)$$

where $D_J^{T=2}(0) = 1$. The value of $\mathcal{F}_{3+}^a(m_f^2, m_i^2, 0)$ is proportional to the mass splitting in the pion multiplet. We restrict ourselves to the first order in Δm_π^2 . Accordingly, the physical pion masses in the arguments of the vertex functions can be replaced by a mean value, for example, $\mu = (m_f + m_i)/2$, provided that these functions, such as $\mathcal{F}_{2+}^a$ and $\mathcal{F}_{3+}^a$, are already small. Using (25)–(28), we obtain

$$\begin{aligned} (\mathcal{F}_{3+}^a(\mu^2, \mu^2, q^2))^{T=2} &= \frac{[m^2 + \Sigma(\mu^2), T^a]}{D_J^{T=2}(q^2)}, \\ (\mathcal{F}_{2+}^a(\mu^2, \mu^2, q^2))^{T=2} &= [m^2 + \Sigma(\mu^2), T^a] \mathcal{F}_{2+}(\mu^2, \mu^2, q^2), \end{aligned} \quad (29)$$

where

$$\mathcal{F}_{2+}(\mu^2, \mu^2, q^2) = \frac{1/D_J^{T=2}(q^2) - 1}{q^2}. \quad (30)$$

Let us consider the consequences of (24). Equation (21) shows that $\mathcal{F}_{2-}^a$ can be evaluated with the required accuracy in the limit of exact isotopic symmetry with $[m^2 + \Sigma(p), T^a] = \mathcal{F}_{i+}^a = \mathcal{F}_{3\pm}^a = 0$ and $\mathcal{F}_{1-}^a = T^a \mathcal{F}_{1-}$. On the mass shell, moreover, $\Sigma(p) = 0$, and thus,

$$\mathcal{F}_{2-}(\mu^2, \mu^2, q^2) = \frac{1 - \mathcal{F}_{1-}(\mu^2, \mu^2, q^2)}{q^2}. \quad (31)$$

To the leading order of the expansion in Δm_π^2 , the form factor $\mathcal{F}_{2-}^a$ is determined by the isotopic rotation of the electromagnetic form factor (5).

The form factors $\mathcal{F}_{2\pm}$, as defined by (30) and (31), are not singular at $q^2 = 0$. For small q^2 , they are determined by the pion radii. The on-shell weak vector current is usually parameterized in the form

$$\begin{aligned} \langle \pi^0(p') | \bar{d} \gamma_\mu (1 - \gamma_5) u | \pi^+(p) \rangle \\ = \sqrt{2} \left((p' + p)_\mu f_+ + q_\mu f_- \right), \end{aligned} \quad (32)$$

where $q_\mu = (p' - p)_\mu$.

The exact CVC condition implies

$$f_- = 0. \quad (33)$$

Partial CVC, as follows from the comparison of (21) and (32), yields

$$\begin{aligned} f_- &= (m_{\pi^0}^2 - m_{\pi^+}^2) (\mathcal{F}_{2-}(\mu^2, \mu^2, 0) + \mathcal{F}_{2+}(\mu^2, \mu^2, 0)) \\ &= \frac{m_{\pi^+}^2 - m_{\pi^0}^2}{6} \left(\langle r^2 \rangle_v^{T=1} - \langle r^2 \rangle_s^{T=2} \right). \end{aligned} \quad (34)$$

The isovector part is recovered from the electromagnetic vertex, whereas the isotensor part is independent of it. Remarkably, for a dressed vertex, the $W^\pm$ boson, being a member of the weak isospin triplet, is coupled to both the strong isospin triplet and the strong isospin quintet.

Equation (34) can also be derived in a simpler and more direct way, analogously to the standard analysis of $K_{\ell 3}$ decays [26].

First, we introduce the form factors

$$f_v(q^2) = \mathcal{F}_{1-}(m_f^2, m_i^2, q^2), \quad (35)$$

$$\begin{aligned} f_s(q^2) &= \mathcal{F}_{1-}(m_f^2, m_i^2, q^2) \\ &\quad + q^2 (\mathcal{F}_{2-}(m_f^2, m_i^2, q^2) + \mathcal{F}_{2+}(m_f^2, m_i^2, q^2)), \end{aligned} \quad (36)$$

with $f_+ = f_v(0) = f_s(0) = 1$. In the t channel, the space-like component of the current is proportional to $f_v(q^2)$, whereas the time-like component is proportional to $f_s(q^2)$. By applying the t -channel unitarity condition for the form factors, we find that f_v is determined by the $J = T = 1$ scattering amplitude, whereas f_s is determined by the $J = 0$, $T = 2$ scattering amplitude, where J is the angular momentum of the pions. f_v and f_s can thus be identified as vector and scalar form factors, respectively, which is consistent with the gWTI.

Second, we assume an expansion in q^2 for both $f_v(q^2)$ and $f_s(q^2)$. This assumption conveys the analyticity of the form factors. Equation (36) leads to

$$\begin{aligned} 1 + \frac{1}{6} \langle r^2 \rangle_s^{T=2} q^2 &= 1 + \frac{1}{6} \langle r^2 \rangle_v^{T=1} q^2 + \frac{q^2}{m_f^2 - m_i^2} f_- \\ &\quad + O(q^4), \end{aligned} \quad (37)$$

such that we recover (34) and, in addition, observe that it is exact for the on-shell case. However, the connection to the CVC condition remains hidden in this simplified approach.

The nature of the form factor f_s can also be clarified with the aid of (32). Taking the scalar product of both sides of the equation with the vector q_μ yields

$$\begin{aligned} \langle \pi^0(p') | i \partial^\mu (\bar{d} \gamma_\mu (1 - \gamma_5) u) | \pi^+(p) \rangle \\ = -\Delta m_\pi^2 \sqrt{2} f_s(q^2). \end{aligned} \quad (38)$$

On the left-hand side, we obtain the form factor, which is a Lorentz scalar. Equation (38) is the formal statement of the nonconservation of the lowering component of the weak vector current.

Equation (34) can therefore be derived using either the gWTI or the kinematic decomposition of (35) and (36). Because recourse to the gWTI is optional, (34) is less interesting for testing the gWTI. We therefore analyze the form factors off the mass shell.

The WTI of exact symmetry implies, to first order in the displacement $p'^2 - p^2$ and for low momentum transfers,

$$f_- = -\frac{p'^2 - p^2}{6} \langle r^2 \rangle_v^{T=1}. \quad (39)$$

The WTI of broken symmetry implies, to first order in Δm_π^2 and the displacement $p'^2 - p^2$ and for low momentum transfers,

$$f_- = -\frac{p'^2 - p^2}{6} \langle r^2 \rangle_v^{T=1} + \frac{m_{\pi^0}^2 - m_{\pi^+}^2}{6} \langle r^2 \rangle_s^{T=2}. \quad (40)$$

The $\pi^+ \rightarrow \pi^0 e^+ \nu_e \gamma$ decay rate depends on the near-mass-shell behavior of the form factor f_- . Any such reaction may serve for testing the gWTI.

For large momentum transfers, the pion radii in (34), (39), and (40) should be replaced with the form factors of (30) and (31). Expressions (33), (34), (39), and (40) correspond to the various versions of the CVC condition.

The on-shell form factor $\mathcal{F}_{1-}^a$ is independent of both the gauge and the parameterization. By virtue of (30) and (31), the on-shell form factors $\mathcal{F}_{2\pm}^a$ are also uniquely defined. The longitudinal component of the vertex (21) contains the factor $p'^2 - p^2$ in $\mathcal{F}_{2-}^a$, whereas $\mathcal{F}_{2+}^a$ has smallness of $O(\Delta m_\pi^2)$. We thus conclude that the longitudinal component of $\Gamma_\mu^a(p', p)$ is uniquely defined in the neighborhood of the mass shell to first order in the pion mass splitting and the displacement $p'^2 - p^2$. In the neighborhood of the mass shell, the form factor f_- thus avoids the general rule [16–20, 24, 25] that states that off-shell amplitudes are ambiguous.

The results obtained for the near-mass-shell representation of f_- in terms of the physical masses and radii of the pions exhibit explicit independence of the gauge and the parameterization of the pion field.

From a mathematical point of view, one can speak of the uniqueness of the longitudinal component of $\Gamma_\mu^a(p', p)$ in an infinitesimal neighborhood of the mass shell or, equivalently, of the uniqueness of the longitudinal component of $\Gamma_\mu^a(p', p)$ and its first derivatives with respect to p'_ν and p_ν on the mass shell to first order in the pion mass splitting.

The transverse component of the vertex does not exhibit any specific behavior, and thus, we consider f_+ to be ambiguous off shell.

The generality of the relationships (34) and (40) is influenced by only the CVC condition at a bare interaction vertex. This condition is satisfied in the Standard Model, so any violation of these relationships can be interpreted as an indication for new physics at or above the electroweak scale.

5. Numerical Estimates

In this section, we employ the dispersion techniques for numerical estimation of the pion radius $\langle r^2 \rangle_s^{T=2}$, entering (34) and (40), and of the form factor f_- .

In the scattering of $T = 2$ pions, the inelastic channels do not become relevant until an invariant mass of 1.2 GeV [27–29]. The elastic scattering amplitude is determined by the phase shift through the Jost function:

$$D_J^T(s) = \exp \left(-\frac{s}{\pi} \int_{4\mu^2}^{+\infty} \frac{\delta^T(s') ds'}{s'(s' - s)} \right). \quad (41)$$

The unitarity relation for the form factor implies (cf. (28))

$$(\mathcal{F}_{3+}(s))^T = \frac{P(s)}{D_J^T(s)}, \quad (42)$$

where $P(s)$ is a finite-degree polynomial. The degree of $P(s)$ can be fixed by the quark-counting rules, which, in our case, take the form $\mathcal{F}_{2\pm}(s) \sim 1/s$ for $s \rightarrow -\infty$. The asymptotes of the Jost function are determined by value of the phase at infinity. The experimental phase is known for energies up to $\sqrt{s_{\max}} = 2.1$ GeV. At this energy, $\delta(s_{\max}) \approx 0$. If one takes $\delta(s) = 0$ for $\sqrt{s} > 2.1$ GeV, then the quark-counting rules are not satisfied, even for the zero-degree polynomial $P(s) = 1$. The minimal required modification is to introduce a hypothetical resonance with mass $m_X > \sqrt{s_{\max}}$. Such a resonance should have a large width because it is not observed experimentally. Note that, in the Veneziano model, there are no resonances in the $T = 2$ channel. The contribution of a large s is suppressed in the dispersion integral by the inverse of the second power of s , and one can assume that this contribution is not dominant. The contribution of the large- s region can be evaluated by comparing the contributions of the intervals $\sqrt{s} = 2\mu - 1.5$ GeV and $1.5 - 2.1$ GeV, for which experimental data are available. A polynomial $P(s)$ of degree n requires $n + 1$ resonances. If there are primitives, then the number of resonances increases. The numbers of resonances and primitives are correlated with the number of Castillejo-Dalitz-Dyson poles [30].

Expanding the Jost function in the vicinity of $s = 0$, we obtain

$$\langle r^2 \rangle_s^T = \frac{6}{\pi} \int_{4\mu^2}^{+\infty} \frac{\delta^T(s')}{s'^2} ds'. \quad (43)$$

For numerical estimates, the upper limit of the integral is replaced by $s_{\max}$; the scattering phase shift between the experimental points is interpolated linearly. The results are shown in Table 1 for the data sets from [27–29]. A further refinement of the estimate can be done using more theoretical inputs from semiphenomenological models of pion-pion scattering [31] and lattice QCD [32].

The negative “mean square radius” occurs because of the negative t -channel scattering phase, which is a signature of repulsion. In the general case, the mean square radius is defined as the first expansion coefficient of the corresponding form factor in powers of q^2 , and the sign of this expansion

TABLE 1: Lorentz scalar isospin $T = 2$ mean square radius of the pion, as determined using (43). The phase-shift analyses of [27–29] are used; (a) and (b) denote nonequivalent datasets of scattering phases. $\sqrt{s_{\max}}$ is the maximum energy up to which the phase analysis is performed.

$\langle r^2 \rangle_s^{T=2} [\text{fm}^2]$	$\sqrt{s_{\max}} [\text{GeV}]$	References
-0.09 ± 0.05	1.4	[27]
-0.10 ± 0.01	2.1	[28] (a)
-0.10 ± 0.03	2.1	[28] (b)
-0.11 ± 0.02	1.5	[29] (a)
-0.13 ± 0.01	1.5	[29] (b)

coefficient is not fixed a priori. Vector boson exchange produces repulsive forces between identical particles. The $T = 2$ channel with $T_3 = 2$ contains a pair of positively charged pions, and thus, the repulsive nature of the phase can be attributed to the dominance of the t -channel ρ -meson exchange between the pions.

Equation (43) has a variety of applications. In the zero-width approximation with $\delta^{T=1}(s) = \pi\theta(s - m_\rho^2)$, where m_ρ is the mass of the ρ meson, one arrives at the well-known expression for the pion charge radius, $\langle r^2 \rangle_v^{T=1} = 6/m_\rho^2$. In the $T = 1$ channel with an attraction, $\langle r^2 \rangle_v^{T=1}$ becomes positive and approaches the experimental value, thereby supporting the vector meson dominance model. The pion scalar radius can be determined using the data for pion-pion scattering in the $T = 0$ channel, yielding $\langle r^2 \rangle_s^{T=0} = 0.61 \pm 0.04 \text{ fm}^2$ [33]. The positive value of $\langle r^2 \rangle_s^{T=0}$ reflects the dominance of the attractive interaction, which is consistent with the existence of the σ meson.

In the dispersion theory, the contribution from the high energy part of the dispersion integrals is considered to be the important source of theoretical uncertainties. The representation (41) is justified in an approximation of neglecting all the inelastic channels. In our case, the elasticity holds for $\sqrt{s} = 2\mu - 1.2 \text{ GeV}$ [27–29], so the dispersion integral from this region accurately assesses the contribution to the pion radius. With increasing the energy above 1.2 GeV, the parameter of inelasticity increases smoothly up to 0.5 for $\sqrt{s} = 2.1 \text{ GeV}$. The part of the integral (43) corresponding to this region gives therefore an approximate estimate of the contribution. Using results of [28], we find that this contribution is numerically small and equals only -0.007 fm^2 . The uncertainty associated with higher energies $\sqrt{s} > 2.1 \text{ GeV}$ can be estimated by assuming the existence of a hypothetical resonance X with mass $m_X > 2.1 \text{ GeV}$. In the zero-width approximation, the error equals $6/m_X^2 \sim 0.03 \text{ fm}^2$ for $m_X = 3 \text{ GeV}$, which is comparable to the experimental uncertainty of the contribution from the interval $\sqrt{s} = 2\mu - 1.2 \text{ GeV}$.

Experimental value of the π^+ charge radius equals $\langle r^2 \rangle_v^{T=1} = (0.672 \pm 0.008 \text{ fm})^2$ [34]. Using the value $\langle r^2 \rangle_s^{T=2} = -0.10 \pm 0.03 \pm 0.03 \text{ fm}^2$, we obtain with the help of (34)

$$f_- = (2.97 \pm 0.25) \times 10^{-3}, \quad (44)$$

which is a factor of two greater than the light-front quark model prediction [5]. The estimated error corresponds to

the experimental errors in the $\pi\pi$ -scattering phases and the uncertainty of the high energy part of the integral representation (43).

Contribution of the longitudinal form factor to the $\pi^+ \rightarrow \pi^0 e^+ \nu_e$ decay rate can be estimated to give $\Delta B/B = -0.94 \times 10^{-3} f_-$; the additional small factor in f_- arises for kinematic reasons. The experimental error in B is 0.6% [34]. We thus reaffirm earlier conclusions that f_- is currently beyond the capabilities of the experimental study. The possibility of measuring the longitudinal weak vector current in the neutron β decay [35], muon capture [36], and in the $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$ and $\tau^- \rightarrow K^- K^0 \nu_\tau$ decays is, perhaps, more promising.

A favorable experimental situation exists in the $K_{\ell 3}$ decays. The large difference in the masses of π - and K -mesons hampers the near-mass-shell expansion, so the $K_{\ell 3}$ decays seem to be not suitable for the measuring f_- off shell and testing the gWTI. At the same time the dispersion techniques allowing determining f_- on shell have found in these decays a successful application [37, 38].

The dominant part of Δm_π^2 is electromagnetic in origin [39–41]. Small mass difference between up and down quarks leads to the $\pi^0 - \eta$ mixing, which also contributes to Δm_π^2 . Equation (34) describes therefore the electromagnetic contribution of order $O(\alpha)$ to the longitudinal weak vector current and the small $\pi^0 - \eta$ mixing effect.

χ PT has been very successful in describing properties of light mesons in the nonperturbative regime [40–43]. Predictions for the $\pi\pi$ -scattering lengths, first obtained by Weinberg [44] and improved by the development of χ PT, agree with experiment to a high degree of accuracy [45, 46]. The pion radii $\langle r^2 \rangle_s^{T=0}$ and $\langle r^2 \rangle_v^{T=1}$ are also well understood [33, 40, 47, 48], while the pion radius $\langle r^2 \rangle_s^{T=2}$ has not been discussed so far. The embedding of isotopic symmetry breaking into the χ PT framework on line with [3, 4, 41, 49] could allow also describing the near-mass-shell form factor f_- , starting from the first principles of QCD.

6. Conclusion

In this paper, the longitudinal component of the weak vector current in the π_{e3} decay was calculated. This component arises due to the violation of isotopic symmetry by the pion masses and the electromagnetic and weak interactions. A generalization of the WTI was derived in the pion sector to account for the isotopic symmetry breaking. It was shown that the isovector $T = 1$ part of the current can be reconstructed through isotopic rotation of the off-shell pion electromagnetic form factors, whereas the isotensor $T = 2$ part has no analogs but rather is uniquely determined by the gWTI combined with the elastic unitarity, analyticity, and the pion-pion scattering data. Because of the low momentum transfer, the on-shell form factor is proportional to the pion mass difference and the difference of the vector-isovector and scalar-isotensor pion radii. The relationship between the near-mass-shell form factor and the pion radii (40) is a consequence of the CVC condition extended beyond the approximation of exact isotopic symmetry.

The various versions of the CVC condition are distinguished depending on whether the isotopic symmetry is exact or broken and whether the outer legs in the vertex are on or off shell. The corresponding predictions for the longitudinal form factor f_- are given in (33), (34), (39), and (40). In the case of the exact isotopic symmetry, the bare and dressed weak vertices are pure isospin triplets; the CVC and the WTI hold on and off shell, respectively. In the case of the broken isotopic symmetry, the dressed weak vertex is no longer a pure isospin triplet; partial CVC and the gWTI hold on and off shell.

Favorable conditions for testing the gWTI by measuring the longitudinal weak vector current appear to exist in the $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau \gamma$ and $\tau^- \rightarrow K^- K^0 \nu_\tau \gamma$ decays with soft photon emission because of the involvement of the near-mass-shell form factors, the relatively large momentum transfer in the weak vertex, and the small mass differences within the pion and kaon multiplets.

A nontrivial consequence of partial CVC occurs in the off-shell case, in which the vertex in general is not gauge invariant and depends on the parameterization of the pion field. The only exception is the longitudinal component of the vertex in the neighborhood of the mass shell. To first order in the pion mass splitting and the displacement from the mass shell, the longitudinal component is independent of both the gauge and the parameterization, and therefore, the near-mass-shell form factor f_- appears to be a unique object whose properties are unambiguously determined by the partial CVC condition (the gWTI) and also fundamentally allow for experimental verification.

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This work was supported by RFBR Grant no. 13-02-01442, Grant no. HLP-2015-18 of Heisenberg-Landau Program, and Grant no. 3172.2012.2 for Leading Scientific Schools of Russian Federation.

References

- [1] A. Sirlin, "Current algebra formulation of radiative corrections in gauge theories and the universality of the weak interactions," *Reviews of Modern Physics*, vol. 50, p. 573, 1978, Erratum in: *Reviews of Modern Physics*, vol. 50, p. 905, 1978.
- [2] A. Sirlin, "Radiative corrections in the standard model," *AIP Conference Proceedings*, vol. 300, no. 1, pp. 331–341, 1994.
- [3] V. Cirigliano, M. Knecht, H. Neufeld, and H. Pichl, "The pionic beta decay in chiral perturbation theory," *The European Physical Journal C*, vol. 27, no. 2, pp. 255–262, 2003.
- [4] H. Pichl, "Electromagnetic radiative corrections to pionic beta decay $\pi^+ \rightarrow \pi^0 e^+ \nu_e$," *Nuclear Physics B—Proceedings Supplements*, vol. 121, pp. 191–194, 2003.
- [5] W. Jaus, "Covariant analysis of the light-front quark model," *Physical Review D*, vol. 60, Article ID 054026, 1999.
- [6] W. Jaus, "Radiative corrections of $O(\alpha)$ for pion beta decay in the light-front quark model," *Physical Review D*, vol. 63, Article ID 053009, 2001.
- [7] D. Počanić, E. Frlež, V. A. Baranov et al., "Precise measurement of the $\pi^+ \rightarrow \pi^0 e^+ \nu$ Branching ratio," *Physical Review Letters*, vol. 93, Article ID 181803, 2004.
- [8] D. Počanić, "Pion, muon decays and weak interaction symmetries," *Nuclear Physics A*, vol. 844, no. 1–4, pp. 26c–29c, 2010.
- [9] S. S. Gerstein and Y. B. Zeldovich, "On meson corrections in the theory of β -decay," *ZhETF*, vol. 29, p. 698, 1955 (Russian).
- [10] S. S. Gerstein and Y. B. Zeldovich, "On meson corrections in the theory of beta-decay," *Soviet Physics—JETP*, vol. 2, p. 576, 1956.
- [11] R. P. Feynman and M. Gell-Mann, "Theory of the Fermi interaction," vol. 109, pp. 193–198, 1958.
- [12] P. C. Tiemeijer and J. A. Tjon, "Electromagnetic form factors for an off-shell nucleon in a vector meson dominance model," *Physical Review C*, vol. 42, no. 2, pp. 599–609, 1990.
- [13] J. W. Bos and J. H. Koch, "The off-shell electromagnetic vertex of the nucleon in chiral perturbation theory," *Nuclear Physics A*, vol. 563, no. 4, pp. 539–548, 1993.
- [14] H. Haberzettl and K. Nakayama, "Gauge-invariant formulation of $NN \rightarrow NN\gamma$," *Physical Review C*, vol. 85, no. 6, Article ID 064001, 14 pages, 2012.
- [15] M. Schwamb and H. Arenhovel, "Off-shell effects in electromagnetic reactions on the deuteron," *Nuclear Physics A*, vol. 696, no. 3–4, pp. 556–580, 2001.
- [16] R. Haag, "Quantum field theories with composite particles and asymptotic conditions," *Physical Review*, vol. 112, pp. 669–673, 1958.
- [17] J. S. R. Chisholm, "Change of variables in quantum field theories," *Nuclear Physics*, vol. 26, pp. 469–479, 1961.
- [18] S. Kamefuchi, L. O’Raifeartaigh, and A. Salam, "Change of variables and equivalence theorems in quantum field theories," *Nuclear Physics*, vol. 28, no. 1, pp. 529–549, 1961.
- [19] H. W. Fearing, "The off-shell nucleon-nucleon amplitude: why it is unmeasurable in nucleon-nucleon bremsstrahlung," in *Few-Body Problems in Physics '99*, vol. 12 of *Few-Body Systems*, pp. 263–268, Springer, 2000.
- [20] H. W. Fearing and S. Scherer, "Field transformations and simple models illustrating the impossibility of measuring off-shell effects," *Physical Review C*, vol. 62, Article ID 034003, 2000.
- [21] J. S. Ball and T.-W. Chiu, "Analytic properties of the vertex function in gauge theories. I," *Physical Review D*, vol. 22, no. 10, pp. 2542–2549, 1980.
- [22] T. E. Rudy, H. W. Fearing, and S. Scherer, "Off-shell electromagnetic form factors of pions and kaons in chiral perturbation theory," *Physical Review C*, vol. 50, no. 1, pp. 447–459, 1994.
- [23] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Fields*, McGraw-Hill, 1965.
- [24] I. Bialynicki-Birula, "Renormalization, diagrams, and gauge invariance," *Physical Review D*, vol. 2, no. 12, pp. 2877–2886, 1970.
- [25] G. ’tHooft and M. Veltman, "Combinatorics of gauge fields," *Nuclear Physics B*, vol. 50, no. 1, pp. 318–353, 1972.
- [26] L. B. Okun, *Leptons and Quarks*, North-Holland, Amsterdam, The Netherlands, 1982.
- [27] D. Cohen, T. Ferbel, P. Slattery, and B. Werner, "Study of $\pi\pi$ scattering in the isotopic-spin-2 channel," *Physical Review D*, vol. 7, no. 3, pp. 661–668, 1973.

- [28] N. B. Durusoy, M. Baubillier, R. George, M. Goldberg, and A. M. Touchard, “Study of the $I = 2\pi\pi$ scattering from the reaction $\pi^- d \rightarrow \pi^- \pi^- p_s p$ at 9.0 GeV/c,” *Physics Letters B*, vol. 45, no. 5, pp. 517–520, 1973.
- [29] W. Hoogland, S. Peters, G. Grayer et al., “Measurement and analysis of the $\pi^+ \pi^+$ system produced at small momentum transfer in the reaction $\pi^+ p \rightarrow \pi^+ \pi^+ n$ at 12.5 GeV,” *Nuclear Physics B*, vol. 126, no. 1, pp. 109–123, 1977.
- [30] M. I. Krivoruchenko, “Remarks on the origin of Castillejo-Dalitz-Dyson poles,” *Physical Review C*, vol. 82, no. 1, Article ID 018201, 4 pages, 2010.
- [31] N. T. Hong Xiem and S. Shinmura, “Pion-pion, pion-kaon, and kaon-kaon interactions in the one-meson-exchange model,” *Progress of Theoretical and Experimental Physics*, no. 2, Article ID 023D04, 16 pages, 2014.
- [32] S. R. Beane, E. Chang, W. Detmold et al., “ $I = 2\pi\pi$ S-wave scattering phase shift from lattice QCD,” *Physical Review D*, vol. 85, no. 3, Article ID 034505, 12 pages, 2012.
- [33] G. Colangelo, J. Gasser, and H. Leutwyler, “ $\pi\pi$ scattering,” *Nuclear Physics B*, vol. 603, no. 1-2, pp. 125–179, 2001.
- [34] K. A. Olive, K. Agashe, C. Amsler et al., “Review of particle physics,” *Chinese Physics C*, vol. 38, no. 9, Article ID 090001, 2014.
- [35] H. Abele, “The neutron. Its properties and basic interactions,” *Progress in Particle and Nuclear Physics*, vol. 60, no. 1, pp. 1–81, 2008.
- [36] D. Gazit, “Muon capture on ^3He and the weak structure of the nucleon,” *Physics Letters B*, vol. 666, no. 5, pp. 472–476, 2008.
- [37] V. Bernard, M. Oertel, E. Passemar, and J. Stern, “ $K_{\mu 3}^L$ decay: a stringent test of right-handed quark currents,” *Physics Letters B*, vol. 638, no. 5-6, pp. 480–486, 2006.
- [38] V. Bernard, M. Oertel, E. Passemar, and J. Stern, “Dispersive representation and shape of the Kl3 form factors: robustness,” *Physical Review D*, vol. 80, no. 3, Article ID 034034, 2009.
- [39] T. Das, G. Guralnick, V. Mathur, F. Low, and J. Young, “Electromagnetic mass difference of pions,” *Physical Review Letters*, vol. 18, no. 18, pp. 759–761, 1967.
- [40] J. Gasser and H. Leutwyler, “Chiral perturbation theory to one loop,” *Annals of Physics*, vol. 158, no. 1, pp. 142–210, 1984.
- [41] R. Urech, “Virtual photons in chiral perturbation theory,” *Nuclear Physics B*, vol. 433, no. 1, pp. 234–254, 1995.
- [42] J. Gasser and H. Leutwyler, “Chiral perturbation theory: expansions in the mass of the strange quark,” *Nuclear Physics B*, vol. 250, no. 1-4, pp. 465–516, 1985.
- [43] S. Scherer, “Introduction to Chiral perturbation theory,” *Advances in Nuclear Physics*, vol. 27, p. 277, 2003.
- [44] S. Weinberg, “Pion scattering lengths,” *Physical Review Letters*, vol. 17, no. 11, pp. 616–621, 1966.
- [45] J. Gasser, “On the history of pion-pion scattering,” in *Proceedings of the International Workshop on Effective Field Theories: From the Pion to the Upsilon (EFT '09)*, p. 29, Valencia, Spain, February 2009.
- [46] G. Colangelo, “Precise standard model tests,” in *Proceedings of the Kaon International Conference (KAON '09)*, p. 53, Tsukuba, Japan, 2009.
- [47] J. Gasser and U. G. Meissner, “Chiral expansion of pion form factors beyond one loop,” *Nuclear Physics B*, vol. 357, no. 1, pp. 90–128, 1991.
- [48] J. Bijnens, G. Colangelo, and P. Talavera, “The vector and scalar form factors of the pion to two loops,” *Journal of High Energy Physics*, vol. 5, article 014, 1998.
- [49] V. Cirigliano and I. Rosell, “Two-loop effective theory analysis of $\pi(K) \rightarrow e\bar{\nu}_e[\gamma]$ Branching ratios,” *Physical Review Letters*, vol. 99, Article ID 231801, 2007.