

Research Article

On the Compatibility of the Diboson Excess with a gg -Initiated Composite Sector

Verónica Sanz

Department of Physics and Astronomy, University of Sussex, Brighton BN1 9QH, UK

Correspondence should be addressed to Verónica Sanz; v.sanz@sussex.ac.uk

Received 22 November 2015; Accepted 18 February 2016

Academic Editor: George Siopsis

Copyright © 2016 Verónica Sanz. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

We propose that recent results by ATLAS and CMS searching for heavy resonances decaying into bosons could be a first hint of a new sector of pure gauge confining physics, possibly linked to the origin of the Higgs as a Composite Higgs. The lightest resonances (glueballs) of this new sector would be neutral, spin-zero, and spin-two, and their behaviour would resemble that of a radion and a massive graviton of extra dimensions. We outline how 13 TeV LHC data could be used to improve sensitivity on this scenario, as well as future characterization during the 13 TeV LHC run.

1. Introduction

Search for heavy resonances decaying into a pair of bosons performed by CMS and ATLAS [1–5] shows tantalizing hints towards the existence of a new resonance at a mass of around 2 TeV, a possibility which has created quite some excitement [6–21].

In this paper we provide an alternative interpretation in terms of a new strong sector, possibly linked to the origin of the Higgs particle as a composite state. We will consider new states, singlet under the SM interactions, which can be produced and decay through their coupling to the stress-energy tensor. An example of such a theory is a new pure gauge sector which undergoes confinement at energies around the TeV scale. The spectrum of this theory contains glueballs, with spin-zero and spin-two resonances at the bottom of the spectrum [22–26]. Focusing on the low-lying, conceivably narrow states, we concentrate in the scenario where the new states couple to gluons through anomalies and decay predominantly to massive vector bosons or Higgses.

2. Glueballs: Theoretical Aspects

Consider a new nonabelian gauge sector, for example, $SU(n)$ gauge group, which undergoes confinement leading to a low energy spectrum of glueballs. Glueballs are bound states of

gauge fields and their behaviour has been studied both in the case of QCD and in the case of more general gauge theories. The ordering of states can be understood by examining the interpolating operators of minimal canonical dimension [27, 28], a prescription which lattice simulations seem to confirm [22].

With this prescription, the lightest states would then correspond to those generated by the lowest dimensional singlet operator, namely, a four-dimension operator $\text{Tr } F_{\mu\nu} F_{\alpha\beta}$, which generates glueballs with quantum numbers $J^{\text{PC}} = 0^{++}, 2^{++}, 0^{-+}$, and 2^{-+} (note that these quantum numbers could also be achieved within *oddballs* [29]).

The next level of resonances would be associated with the five-dimension operator, $\text{Tr } F_{\mu\nu} D_\rho F_{\alpha\beta}$, leading to resonances with quantum numbers $J^{\text{PC}} = 1^{++}$ and 3^{++} . One could continue this procedure to classify resonances by examining six-dimension and higher operators generated by gauge fields.

In the following we are going to focus on the lowest resonances: spin-zero and spin-two. Lattice studies on $SU(n)$ gauge theories find that the lowest resonances correspond to $\text{PC} = ++$; hence we will denote them by

$$\begin{aligned} \phi &: J^{\text{PC}} = 0^{++}, \\ G_{\mu\nu} &: J^{\text{PC}} = 2^{++}. \end{aligned} \tag{1}$$

Determining the separation between the scalar (0^{++}) and tensor states (2^{++}) is a difficult task in lattice gauge theories. Lattice resulting in a pure glueball calculation indicates that the tensor mode is about 60% heavier than the scalar, even in the large- n limit. Nevertheless, this result will likely change once the pure gauge theory is coupled to the Standard Model. In the following, I will consider these two lightest states as two distinct possibilities for the lightest state in a glueball spectrum.

The resonances ϕ and G propagate as Klein-Gordon and Fierz-Pauli [30] fields. The Fierz-Pauli Lagrangian describes a massive spin-two field, a rank-two symmetric, and traceless tensor. Additionally, a positive-energy condition must be satisfied (see [31] and references therein).

2.1. Generic Couplings of the Spin-Zero and Spin-Two States. Contrary to the case of QCD, a pure gauge theory has no global (chiral) symmetries which would be broken by the confinement dynamics. On the other hand, space-time symmetries can be broken by confinement. For example, glueballs break spontaneously scale invariance of the gauge theory. Hence, the lightest spin-zero resonance could play the role of a dilaton, the Goldstone boson of the spontaneous breaking of scale invariance. In this case, the couplings of the dilaton resonance are of the form $(\phi/\Lambda)\partial_\mu J^\mu$, where J^μ is the global current whose spontaneous breaking at the scale Λ leads to the emergence of the Goldstone boson ϕ . The global current is given by $J^\mu = T^{\mu\nu}v_\nu$, where v_μ is the generator of dilatation symmetry $v_\nu = \lambda x_\nu$, which then implies that the dilaton couples to trace of the stress tensor $T = T^\mu_\mu$; consider

$$-\frac{a_i}{\Lambda}\phi T_i, \quad (2)$$

where Λ is the symmetry breaking scale and the index i refers to species; that is, $i = \Phi, \gamma, g, \dots$. In our set-up the dimensionless parameter a_i encodes the degree of compositeness of species i , with an order one value indicating a large mixture with the composite sector. Note that this coupling vanishes for massless gauge bosons (gluon and photon) as $\text{Tr} T_{\mu\nu}^{\gamma, g} = 0$, but a coupling would be induced nevertheless at one loop through the anomaly [32–34]

$$-\frac{a_{g,\gamma}}{\Lambda}\phi F_{\mu\nu}F^{\mu\nu}, \quad (3)$$

where F here denotes the gluon or photon field strength.

We encounter a similar situation for the spin-two state $G_{\mu\nu}$ whose properties are derived from diffeomorphism invariance, broken spontaneously by the gauge dynamics. Indeed, a massless spin-two object $\theta^{\mu\nu}$ is conserved, $\partial_\mu \theta^{\mu\nu} = 0$, in the absence of breaking. As in 0^+ case, $\theta_{\mu\nu}$ couples to a conserved current $\partial_\mu J^\mu = 0$. The breaking of this diffeomorphism invariance can be parametrized by $\partial_\mu \theta^{\mu\nu} = v^\nu$, where v^μ corresponds to a massive vector field, which is *eaten* by the massless spin-two field [35–37]. When joining together, the spin-two massless field and the massive vector will lead to the massive spin-two state $G_{\mu\nu}$. As long as the composite sector preserves Lorentz, gauge, and CP invariance, the coupling of

the massive spin-two resonance to *two* SM particles will be given by [31]

$$-\frac{b_i}{\Lambda}G_{\mu\nu}T_i^{\mu\nu}, \quad (4)$$

where $T_i^{\mu\nu}$ corresponds to the stress tensor of species i . Studies of couplings to the tensor state to SM particles have been done in the context of glueballs in QCD [38]. Note that these analyses differ from ours in which the tensor two-point function is treated as containing a massless state, and the constraints from diffeomorphism invariance are not included.

2.2. A Set-Up for Glueballs and a Composite Higgs. In a scenario where EWSB is due to strong dynamics, such as Composite Higgs scenarios [39–41], the glueball sector could be involved in causing the spontaneous breaking of the global symmetry responsible for the pseudo-Goldstone Higgs. In this section we explore this possibility.

Let us imagine a sector with a large global symmetry G and a gauge symmetry G_g with no fundamental fermions. Assume then that the dynamics of G_g become strong at some scale Λ , which then triggers the spontaneous breaking of G down to a smaller subgroup H . An example would be $\text{SO}(5) \times \text{SU}(3)$ sector and $\text{SU}(3)$ gauge strong dynamics leading to the breaking of $\text{SO}(5)$ to $\text{SO}(4)$; see Figure 1. One could then proceed as usual in the minimal Composite Higgs scenarios, by partly gauging $\text{SO}(4)$ interactions. To achieve EWSB, one would likely need to use a mechanism as explored in [42], where the Higgs potential is generated via a sequential breaking which one could link to the strong sector producing both the spin-zero and spin-two resonances and breaking the global symmetry. The spectrum at low energies would then contain the Goldstone bosons (a doublet under $\text{SU}(2) \subset \text{SO}(4)$) and the glueballs.

In this case, the glueballs will exhibit a hierarchy of couplings. The SM particles with larger couplings to the glueballs would be the Higgs degrees of freedom through a mechanism such as *partial compositeness* [43–45]. As a result, the resonances of the composite sector would couple preferentially to the Higgs doublet, namely, with the Higgs particle h and the *longitudinal* W^\pm and Z bosons.

Couplings to gluons and photons would be induced through the anomaly terms, and couplings to light fermions would be suppressed by their mass. In Section 2.4 we describe how to parametrize these couplings. But before that we will discuss an alternative view in the context of extra dimensional theories, which shows the same hierarchical couplings.

2.3. Holographic Radion and Graviton as Proxies for Glueballs. Glueballs can be treated as fields arising from extra dimensions using dualities. These dualities are based on the AdS/CFT correspondence [46], which draws a duality between strongly coupled theories in D dimensions and a gravitational dual in $D + 1$ dimensions, reaching beyond supersymmetric or exactly conformal theories [47]. The holographic duality is not an exact mathematical statement but has to be taken as a qualitative statement between strongly




FIGURE 1: (a) Spectrum of glueballs in a pure gauge theory, including the prior operators. The two lightest states are singled out as ϕ (0^{++}) and $G_{\mu\nu}$ (2^{++}). (b) The new sector of strong interactions exhibits a global symmetry G broken spontaneously by G_g strong dynamics.

coupled theories (the target theory) and an *analogue computer* [48–51], a theory on higher dimensions with improved calculability.

Most of the information of the dual extra dimensional theory can be understood by means of the metric, which can be expressed as

$$ds^2 = w(z)^2 (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2) \quad (5)$$

with z being the extra dimension and $w(z)$ being the warp factor. The extra dimension is often compactified, $z \in [z_{UV}, z_{IR}]$, with $z_{UV(IR)}$ being the UV(IR) brane.

Physics at different positions inside the extra dimension (the bulk) correspond to snapshots of the 4D theory at different energy scales. In other words, the bulk of the extra dimension encodes the RG evolution of the 4D Lagrangian, with the UV brane and IR brane representing the UV and IR boundary conditions on the RGE. Propagation of extra dimensional fields from the UV to the IR branes corresponds to integrating out degrees of freedom: at a position z_* the *local* cutoff is related to the UV cutoff as $\Lambda(z_*) = w(z_*)\Lambda_{UV}$, a change in cutoff which can be understood à la Wilsonian integrating out heavy degrees of freedom [52].

The running stops at the IR brane, with the presence of the IR brane signalling that a sector of the 4D theory is undergoing confinement. 4D composite states are therefore dual to the Kaluza-Klein modes due to compactification. Localization of a field in extra dimensions has a dual meaning as well. Fields localized near or on the UV brane do not strongly participate in the strong dynamics encoded near the IR brane and are then called *elementary*. Localization towards a brane is then the equivalent to the degree of compositeness of the field. On the other hand, gauge fields in the extra dimension which are not localized anywhere specifically in the extra dimension, hence with flat profiles, represent global symmetries of the composite sector, weakly gauged by the UV dynamics [53]. Flat fields are, hence, a mixture of composite and elementary field, much the same as the ρ - γ mixing in QCD [48, 54, 55].

In the context of the extra dimensional theory the roles of the 0^{++} and 2^{++} states are clear [32, 33, 56–58],

$$\begin{aligned} \phi &\longrightarrow \text{radion}, \\ G_{\mu\nu} &\longrightarrow \text{KK-graviton}, \end{aligned} \quad (6)$$

and their behaviours match in both theories in the sense that the structure of couplings of the glueballs and their holographic duals are dictated by symmetries. This dual can be used as a framework to parametrize the properties of the glueballs. Note though that there are differences between the two pictures. The 4D theory does not contain gravity; hence the holographic graviton and radion masses are unrelated to the scale of quantum gravity, which is assumed to be much higher than the physics we focus on. Also, terms involving the radion and dilaton at quadratic order may differ [59] when dilatation symmetry is not extended to gravity [60]. An extensive literature on radion/KK-graviton properties can be found elsewhere, for example, [44, 61–70].

2.4. Glueball Parametrization. In the partial compositeness picture, larger couplings between the resonances and SM particles indicate a more direct communication between the new strong dynamics and the SM sector. The hierarchy of couplings we expect is as follows [34, 57]: (1) *Higgs degrees of freedom*: $a_H, b_H \simeq \mathcal{O}(1)$, (2) *gluon and photon couplings*: $a_{\gamma,g}, b_{\gamma,g} \simeq \alpha_{s,EM}$, and (3) *light SM fermions*: $a_f, b_f \propto (m_f/v)^\gamma$, where γ is some number, larger than one. In the dual extra dimensional picture, the same hierarchy of couplings has a geometrical meaning [34, 48]. The radion/KK-graviton are bulk states localized near the IR brane; hence, with larger overlap with states there, the gluon/photon are delocalized states, and light fermions are bound near the UV brane. Hence, the couplings would go as above (see [57]).

3. Glueballs at the LHC

In this section we discuss the signatures of glueballs and their interpretation in terms of the reported excess in the diboson channels [1–5]. This excess is not statistically significant at the moment; hence concrete examples of specific




FIGURE 2: Total cross sections at 8 TeV LHC for the spin-zero (a) and spin-two (b) hypothesis in the HV and VV channels and digluon final state. Note that cross sections do not include efficiencies to cuts.

theoretical frameworks are useful to interpret the signatures by performing combinations of different channels. As there is no sensitivity in the fully leptonic channels and at the same time there is a sizeable overlap among the boosted hadronic W , Z , and Higgs channels, it turns out that an excess on a channel, for example, WZ , could be in fact due to an excess in the neutral channels.

In the following, we will assume that the total cross section into dibosons (WW or ZZ) is of the order of 1–10 fb, a number to be taken as a ballpark figure. Increasing sensitivity on this excess using the 8 TeV LHCLHC data would require combination of ATLAS and CMS analyses, as well as tailoring to more specific scenarios such as the one presented here.

Firstly, let us comment on bounds from precision tests of the electroweak sector, which can be parametrized with the help of S and T parameters [71, 72]. These bounds on composite G and ϕ can be obtained from [73, 74] with suitable modifications. In both cases, the contribution to electroweak observables is induced at loop level. Assuming there is no Higgs- ϕ mixing term, the contribution to S parameter due to ϕ is scaled as $S_\phi \propto \alpha_{\tilde{H}}^2 v^2 / \Lambda^2$, whereas the spin-two state is $S_G \approx s_W^2 c_W^2 (b_H^2 m_G^2 / \Lambda^2)$ [73]. Both contributions also receive a logarithmic enhancement. As the coupling of $G_{\mu\nu}$ to longitudinal W and Z is through their mass, $\alpha T = 0$ at one loop.

3.1. Branching Ratios and Total Width. The partial widths of the 2^+ state to Higgs degrees of freedom take the following form: $\Gamma(G \rightarrow W^+W^-)/2 \approx \Gamma(G \rightarrow ZZ) \approx \Gamma(G \rightarrow hh) \approx (b_H^2/480\pi)(m_G^3/\Lambda^2)$, which are correct up to factors of $\mathcal{O}(m_{Z,W,h}/m_G)^2$. More accurate predictions of the branching ratios can be found in [57]. Similarly, the partial widths of

TABLE 1

	ggF 8 TeV LHC	VBF 8 TeV LHC
ϕ	$a_g^2 (3 \text{ TeV}/\Lambda)^2 (370\text{--}740) \text{ fb}$	$a_g^2 (3 \text{ TeV}/\Lambda)^2 (0.002\text{--}0.004) \text{ fb}$
G	$b_g^2 (3 \text{ TeV}/\Lambda)^2 (90\text{--}190) \text{ fb}$	$b_g^2 (3 \text{ TeV}/\Lambda)^2 (0.6\text{--}1.3) \text{ fb}$
	ggF 13 TeV LHC	VBF 13 TeV LHC
ϕ	$a_g^2 (3 \text{ TeV}/\Lambda)^2 (5\text{--}8) \text{ pb}$	$a_H^2 (3 \text{ TeV}/\Lambda)^2 (0.03\text{--}0.05) \text{ pb}$
G	$b_g^2 (3 \text{ TeV}/\Lambda)^2 (1\text{--}2) \text{ pb}$	$b_H^2 (3 \text{ TeV}/\Lambda)^2 (0.02\text{--}0.03) \text{ pb}$

the 0^+ state to electroweak bosons are as follows: $\Gamma(\phi \rightarrow W^+W^-)/2 \approx \Gamma(\phi \rightarrow ZZ) \approx \Gamma(\phi \rightarrow hh) \approx (a_H^2/16\pi)(m_\phi^3/\Lambda^2)$.

The partial width to gluons would be given by $\Gamma(\phi \rightarrow gg) = (a_g^2/2\pi\Lambda^2)m_\phi^3$ and $\Gamma(G \rightarrow gg) = (b_g^2/120\pi\Lambda^2)m_G^3$. Therefore, the resonances remain narrow as long as $\Lambda \geq m_{\phi,G}$. Assuming a_H and b_H are the largest couplings, the total width would be given by a simple expression: $\Gamma_\phi \approx (a_H^2/4\pi)(m_\phi^3/\Lambda^2)$ and $\Gamma_G \approx (b_H^2/240\pi)(m_G^3/\Lambda^2)$.

3.2. Production Cross Section. The production of the glueballs is dominated by gluon fusion (ggF) and possibly vector boson fusion (VBF), as couplings to light fermions are very suppressed. Whether ggF or VBF is the dominant production mechanism depends on the suppression of couplings to gluons with respect to W and Z couplings, that is, the value of the coefficients (a_g and b_g) and (a_H and b_H), respectively. The production cross section through gluon fusion and VBF is given by Table 1, in the range $m_{\phi,G} = 2\text{--}1.8 \text{ TeV}$. Vector boson fusion is kinematically suppressed, although it is enhanced by couplings to Higgs degrees of freedom, a_H and b_H . Note that the VBF 13 TeV LHC is more steep for the spin-two resonance.

In Figure 2 we show the total cross sections at 8 TeV LHC for the spin-zero and spin-two hypothesis in the HV and VV

channels and digluon final state. As one can see, the spin-two resonance remains narrow in a larger region of the parameter space, a behaviour which is related to the kinematic features discussed in the next section. These numbers have been obtained by implementing the glueballs in the context of `feynrules` [75], exported to `Madgraph5` [76] using the `UFO` [77] format.

The coupling of the glueball ϕ to gluons, photons, and vector bosons has the same Lorentz structure as the Higgs (see (2) and (3)), whereas the spin-two resonance has a more interesting Lorentz structure. In particular, the Feynman rule (in the unitary gauge) involving this resonance with vector bosons is given by [57]

$$\begin{aligned} & [G_{\mu\nu}, V_\alpha(k_1), V_\beta(k_2)] : \\ & -i \frac{1}{\Lambda} (b_H m_V^2 C_{\mu\nu,\alpha\beta} + b_V W_{\mu\nu,\alpha\beta}), \end{aligned} \quad (7)$$

where

$$\begin{aligned} W_{\mu\nu,\alpha\beta} & \equiv \eta_{\alpha\beta} k_{1\mu} k_{2\nu} + \eta_{\mu\alpha} (k_1 \cdot k_2 \eta_{\nu\beta} - k_{1\beta} k_{2\nu}) \\ & - \eta_{\mu\beta} k_{1\nu} k_{2\alpha} + \frac{1}{2} \eta_{\mu\nu} (k_{1\beta} k_{2\alpha} - k_1 \cdot k_2 \eta_{\alpha\beta}) \\ & + (\mu \longleftrightarrow \nu), \end{aligned} \quad (8)$$

$$C_{\mu\nu,\alpha\beta} \equiv \eta_{\mu\alpha} \eta_{\nu\beta} + \eta_{\nu\alpha} \eta_{\mu\beta} - \eta_{\mu\nu} \eta_{\alpha\beta}$$

and V here denotes either a massive vector boson ($b_H \neq 0$) or a massless gluon or photon ($b_H = 0$ and $b_V = b_{g,\gamma}$).

These differences between the structures of couplings of ϕ and $G_{\mu\nu}$ are manifested in the total cross section and the ratio of ggF versus VBF discussed in the last section. Kinematic distributions and hence acceptance to cuts and reconstruction will be modified. As an illustration, we show in Figure 3 the p_T distribution of the leading V-jet for both spin hypotheses. Note that this simulation has been done at parton level, and showering/hadronization and detector effects would distort the distribution.

The higher spin resonance exhibits more boosted spectrum due to the different Lorentz structure of the coupling; see [78–82] for a discussion in the context of the Higgs candidate. Therefore, criteria such as mass drop or mass reconstruction will depend on the spin of the resonance. The spin of the resonance determines not only the p_T spectrum but also angular distributions. We leave for a future study [83] a more realistic analysis of this scenario within a feasibility study at Les Houches.

4. Discussion of the Results

The Run1 data has been able to probe diboson production in the very boosted regime, providing a tantalizing hint of new physics at around 2 TeV. How fast and reliably this hint can be confirmed or discarded depends on contrasting the signatures with specific models. Combination of different diboson channels as well as with dijet, ditop, and Higgs-vector boson relies on benchmarks against which the analyses can be tested. Besides trivial issues of branching ratios to specific




FIGURE 3: Distribution of the transverse momentum of the leading vector boson in the decay of 0^{++} and 2^{++} resonances at 8 TeV LHC.

final states, the analysis depends on whether a resonance is gluon, quark, or vector boson initiated and its quantum numbers, as they determine how the decay products are distributed.

In this paper we have such benchmark of interpretation in the context of glueballs of a new strong sector. We have discussed how the lightest states, massive spin-zero and spin-two resonances, could inherit their couplings from the breaking of diffeomorphism invariance due to the strong dynamics. They would then behave in a similar way as expected from a radion and a massive Kaluza-Klein resonance of extra dimensions.

If the new sector dynamics are linked to EWSB, the resonances will couple preferentially to Higgs degrees of freedom, that is, the Higgs and longitudinal polarizations of W and Z bosons, whereas the best way to produce them is through gluon fusion with a possible vector boson fusion component.

Regarding the collider analysis, the results presented here are performed at parton level. A more detailed analysis, including prospects to measure properties with 13 TeV LHC, will be done in a future publication [83].

Additional Points

After this paper was written, new results from the Run2 LHC looking for the diboson resonance show no significant excess in the region around 2 TeV. More data in the summer of 2016 would be required to determine whether the Run1 excess is ruled out by a larger Run2 dataset.

Competing Interests

The author declares that there are no competing interests regarding the publication of this paper.

Acknowledgments

The author would like to thank Gustaaf Broojimans and Chris Pollard for discussions on the diboson analyses; the author is also grateful for the excellent atmosphere in Les Houches. Many thanks are due to Axel Maas and Agostino Patella for discussions on the spectrum of glueballs. This work is supported by the Science Technology and Facilities Council (STFC) under Grant no. ST/L000504/1.

References

- [1] V. Khachatryan, A. M. Sirunyan, A. Tumasyan et al., “Search for massive resonances in dijet systems containing jets tagged as W or Z boson decays in pp collisions at $\sqrt{s} = 8\text{TeV}$,” *Journal of High Energy Physics*, vol. 2014, article 173, 2014.
- [2] G. Aad, B. Abbott, J. Abdallah et al., “Search for high-mass diboson resonances with boson-tagged jets in proton-proton collisions at $\sqrt{s} = 8\text{TeV}$ with the ATLAS detector,” *Journal of High Energy Physics*, vol. 2015, article 55, 2015.
- [3] G. Aad, B. Abbott, J. Abdallah et al., “Search for production of WW/WZ resonances decaying to a lepton, neutrino and jets in pp collisions at $\sqrt{s} = 8\text{TeV}$ with the ATLAS detector,” *The European Physical Journal C*, vol. 75, article 209, 2015.
- [4] V. Khachatryan, A. M. Sirunyan, A. Tumasyan et al., “Search for massive resonances decaying into pairs of boosted bosons in semi-leptonic final states at $\sqrt{s} = 8\text{TeV}$,” *Journal of High Energy Physics*, vol. 2014, article 174, 2014.
- [5] G. Aad, B. Abbott, J. Abdallah et al., “Search for resonant diboson production in the $\ell\ell q\bar{q}$ final state in pp collisions at $\sqrt{s} = 8\text{TeV}$ with the ATLAS detector,” *The European Physical Journal C*, vol. 75, article 69, p. 69, 2015.
- [6] C. W. Chiang, H. Fukuda, K. Harigaya, M. Ibe, and T. T. Yanagida, “Diboson resonance as a portal to hidden strong dynamics,” <http://arxiv.org/abs/1507.02483>.
- [7] B. A. Dobrescu and Z. Liu, “Heavy Higgs bosons and the 2 TeV W' boson,” *Journal of High Energy Physics*, vol. 2015, no. 10, article 118, 2015.
- [8] A. Carmona, A. Delgado, M. Quiros, and J. Santiago, “Diboson resonant production in non-custodial composite Higgs models,” <http://arxiv.org/abs/1507.01914>.
- [9] G. Cacciapaglia and M. T. Frandsen, “Unitarity implications of a diboson resonance in the TeV region for Higgs physics,” *Physical Review D*, vol. 92, no. 5, Article ID 055035, 2015.
- [10] B. C. Allanach, B. Gripaios, and D. Sutherland, “Anatomy of the ATLAS diboson anomaly,” *Physical Review D*, vol. 92, no. 5, Article ID 055003, 9 pages, 2015.
- [11] Q.-H. Cao, B. Yan, and D.-M. Zhang, “Simple non-Abelian extensions of the standard model gauge group and the diboson excesses at the LHC,” *Physical Review D*, vol. 92, no. 9, Article ID 095025, 18 pages, 2015.
- [12] J. Brehmer, J. Hewett, J. Kopp, T. Rizzo, and J. Tattersall, “Symmetry restored in dibosons at the LHC?” <http://arxiv.org/abs/1507.00013>.
- [13] A. Thamm, R. Torre, and A. Wulzer, “Composite heavy vector triplet in the ATLAS diboson excess,” *Physical Review Letters*, vol. 115, no. 22, Article ID 221802, 2015.
- [14] Y. Gao, T. Ghosh, K. Sinha, and J. H. Yu, “ $SU(2) \times SU(2) \times U(1)$ interpretations of the diboson and Wh excesses,” *Physical Review D*, vol. 92, no. 5, Article ID 055030, 2015.
- [15] A. Alves, A. Berlin, S. Profumo, and F. S. Queiroz, “Dirac-fermionic dark matter in $U(1)_X$ models,” *Journal of High Energy Physics*, vol. 2015, article 76, pp. 1–33, 2015.
- [16] T. Abe, R. Nagai, S. Okawa, and M. Tanabashi, “Unitarity sum rules, three-site moose model, and the ATLAS 2 TeV diboson anomalies,” *Physical Review D*, vol. 92, no. 5, Article ID 055016, 14 pages, 2015.
- [17] T. Abe, T. Kitahara, and M. M. Nojiri, “Prospects for spin-1 resonance search at 13 TeV LHC and the ATLAS diboson excess,” *Journal of High Energy Physics*, vol. 2016, article 84, 2016.
- [18] D. B. Franzosi, M. T. Frandsen, and F. Sannino, “Diboson signals via Fermi scale spin-one states,” *Physical Review D*, vol. 92, no. 11, Article ID 115005, 5 pages, 2015.
- [19] J. A. Aguilar-Saavedra, “Triboson interpretations of the ATLAS diboson excess,” <http://arxiv.org/abs/1506.06739>.
- [20] D. B. Franzosi, M. T. Frandsen, and F. Sannino, “Diboson signals via Fermi scale spin-one states,” *Physical Review D*, vol. 92, Article ID 115005, 2015.
- [21] H. S. Fukano, M. Kurachi, S. Matsuzaki, K. Terashi, and K. Yamawaki, “2 TeV walking technirho at LHC?” *Physics Letters B*, vol. 750, pp. 259–265, 2015.
- [22] M. J. Teper, “Glueball masses and other physical properties of $SU(N)$ gauge theories in $D=3+1$: a review of lattice results for theorists,” <http://arxiv.org/abs/hep-th/9812187>.
- [23] C. J. Morningstar and M. J. Peardon, “Glueball spectrum from an anisotropic lattice study,” *Physical Review D*, vol. 60, no. 3, Article ID 034509, 1999.
- [24] Y. Chen, A. Alexandru, S. J. Dong et al., “Glueball spectrum and matrix elements on anisotropic lattices,” *Physical Review D*, vol. 73, no. 1, Article ID 014516, 21 pages, 2006.
- [25] E. Gregory, A. Irving, B. Lucini et al., “Towards the glueball spectrum from unquenched lattice QCD,” *Journal of High Energy Physics*, vol. 2012, article 170, 2012.
- [26] B. Lucini, A. Rago, and E. Rinaldi, “Glueball masses in the large N limit,” *Journal of High Energy Physics*, vol. 2010, no. 8, article 119, 2010.
- [27] R. L. Jaffe, K. Johnson, and Z. Ryzak, “Qualitative features of the glueball spectrum,” *Annals of Physics*, vol. 168, no. 2, pp. 344–367, 1986.
- [28] J. Kuti, “Exotica and the confining flux,” *Nuclear Physics B: Proceedings Supplements*, vol. 73, no. 1-3, pp. 72–85, 1986.
- [29] F. J. Llanes-Estrada, P. Bicudo, and S. R. Cotanch, “ J^- glueballs and a low odderon intercept,” *Physical Review Letters*, vol. 96, no. 8, Article ID 081601, 4 pages, 2006.
- [30] M. Fierz and W. Pauli, “On relativistic wave equations for particles of arbitrary spin in an electromagnetic field,” *Proceedings of the Royal Society of London A*, vol. 173, no. 953, article 211, 1939.
- [31] R. Fok, C. Guimarães, R. Lewis, and V. Sanz, “It is a graviton! or maybe not,” *Journal of High Energy Physics*, vol. 2012, article 62, 2012.
- [32] C. Csaki, J. Hubisz, and S. J. Lee, “Radion phenomenology in realistic warped space models,” *Physical Review D*, vol. 76, no. 12, Article ID 125015, 15 pages, 2007.

- [33] J. Fan, W. D. Goldberger, A. Ross, and W. Skiba, “Standard model couplings and collider signatures of a light scalar,” *Physical Review D*, vol. 79, no. 3, Article ID 035017, 11 pages, 2009.
- [34] M. Gouzevitch, A. Oliveira, J. Rojo, R. Rosenfeld, G. P. Salam, and V. Sanz, “Scale-invariant resonance tagging in multijet events and new physics in Higgs pair production,” *Journal of High Energy Physics*, vol. 2013, no. 7, article 148, 2013.
- [35] E. Kiritsis, “Product CFTs, gravitational cloning, massive gravitons and the space of gravitational duals,” *Journal of High Energy Physics*, no. 11, p. 49, 2006.
- [36] E. D’Hoker and D. Z. Freedman, “Supersymmetric gauge theories and the AdS/CFT correspondence,” <http://arxiv.org/abs/hep-th/0201253>.
- [37] O. Aharony, A. B. Clark, and A. Karch, “CFT/AdS correspondence, massive gravitons, and a connectivity index conjecture,” *Physical Review D*, vol. 74, no. 8, Article ID 086006, 2006.
- [38] S. Narison, “Masses, decays and mixings of gluonia in QCD,” *Nuclear Physics B*, vol. 509, no. 1-2, pp. 312–356, 1998.
- [39] R. Contino, Y. Nomura, and A. Pomarol, “Higgs as a holographic pseudo-Goldstone boson,” *Nuclear Physics B*, vol. 671, pp. 148–174, 2003.
- [40] K. Agashe, R. Contino, and A. Pomarol, “The minimal composite Higgs model,” *Nuclear Physics B*, vol. 719, no. 1-2, pp. 165–187, 2005.
- [41] G. F. Giudice, C. Grojean, A. Pomarol, and R. Rattazzi, “The strongly-interacting light Higgs,” *Journal of High Energy Physics*, vol. 2007, no. 6, article 045, 2007.
- [42] V. Sanz and J. Setford, “Composite Higgses with seesaw EWSB,” *Journal of High Energy Physics*, vol. 2015, article 154, 2015.
- [43] D. B. Kaplan, “Flavor at SSC energies: a new mechanism for dynamically generated fermion masses,” *Nuclear Physics B*, vol. 365, no. 2, pp. 259–278, 1991.
- [44] T. Gherghetta and A. Pomarol, “Bulk fields and supersymmetry in a slice of AdS,” *Nuclear Physics B*, vol. 586, no. 1-2, pp. 141–162, 2000.
- [45] R. Contino, T. Kramer, M. Son, and R. Sundrum, “Warped/composite phenomenology simplified,” *Journal of High Energy Physics*, vol. 2007, no. 5, article 074, 2007.
- [46] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri, and Y. Oz, “Large N field theories, string theory and gravity,” *Physics Reports*, vol. 323, no. 3-4, pp. 183–386, 2000.
- [47] N. Arkani-Hamed, M. Porrati, and L. Randall, “Holography and phenomenology,” *Journal of High Energy Physics*, vol. 2001, no. 8, article 017, 2001.
- [48] J. Hirn and V. Sanz, “The fifth dimension as an analogue computer for strong interactions at the LHC,” *Journal of High Energy Physics*, vol. 2007, no. 3, article 100, 2007.
- [49] J. Hirn and V. Sanz, “Negative S parameter from holographic technicolor,” *Physical Review Letters*, vol. 97, Article ID 121803, 2006.
- [50] J. Hirn, A. Martin, and V. Sanz, “Benchmarks for new strong interactions at the LHC,” *Journal of High Energy Physics*, vol. 2008, no. 5, article 084, 2008.
- [51] J. Hirn, N. Rius, and V. Sanz, “Geometric approach to condensates in holographic QCD,” *Physical Review D*, vol. 73, no. 8, Article ID 085005, 2006.
- [52] L. Randall, V. Sanz, and M. D. Schwartz, “Entropy-area relations in field theory,” *Journal of High Energy Physics*, vol. 2002, no. 6, article 008, 2002.
- [53] C. Csáki, Y. Shirman, and J. Terning, “Seiberg dual for the MSSM: partially composite W and Z ,” *Physical Review D*, vol. 84, no. 9, Article ID 095011, 10 pages, 2011.
- [54] J. Erlich, E. Katz, D. T. Son, and M. A. Stephanov, “QCD and a holographic model of hadrons,” *Physical Review Letters*, vol. 95, no. 26, Article ID 261602, 4 pages, 2005.
- [55] L. Da Rold and A. Pomarol, “Chiral symmetry breaking from five dimensional spaces,” *Nuclear Physics B*, vol. 721, no. 1-3, pp. 79–97, 2005.
- [56] E. Katz, A. Lewandowski, and M. D. Schwartz, “Tensor mesons in AdS/QCD,” *Physical Review D*, vol. 74, no. 8, Article ID 086004, 2006.
- [57] H. M. Lee, M. Park, and V. Sanz, “Gravity-mediated (or composite) dark matter,” *The European Physical Journal C*, vol. 74, article 2715, 2014.
- [58] H. M. Lee, M. Park, and V. Sanz, “Gravity-mediated (or composite) Dark Matter confronts astrophysical data,” *Journal of High Energy Physics*, vol. 2014, article 63, 2014.
- [59] C. Csáki, M. Graesser, L. Randall, and J. Terning, “Cosmology of brane models with radion stabilization,” *Physical Review D*, vol. 62, no. 4, Article ID 045015, 2000.
- [60] W. D. Goldberger, B. Grinstein, and W. Skiba, “Distinguishing the Higgs boson from the dilaton at the Large Hadron Collider,” *Physical Review Letters*, vol. 100, no. 11, Article ID 111802, 2008.
- [61] A. L. Fitzpatrick, J. Kaplan, L. Randall, and L.-T. Wang, “Searching for the Kaluza-Klein graviton in bulk RS models,” *Journal of High Energy Physics*, vol. 2007, no. 9, article 013, 2007.
- [62] T. Gherghetta and A. Pomarol, “A warped supersymmetric standard model,” *Nuclear Physics B*, vol. 602, no. 1-2, pp. 3–22, 2001.
- [63] Y. Grossman and M. Neubert, “Neutrino masses and mixings in non-factorizable geometry,” *Physics Letters B*, vol. 474, no. 3-4, pp. 361–371, 2000.
- [64] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, “Bulk gauge fields in the Randall–Sundrum model,” *Physics Letters B*, vol. 473, no. 1-2, pp. 43–49, 2000.
- [65] A. Pomarol, “Gauge bosons in a five-dimensional theory with localized gravity,” *Physics Letters B*, vol. 486, no. 1-2, pp. 153–157, 2000.
- [66] S. Chang, J. Hisano, H. Nakano, N. Okada, and M. Yamaguchi, “Bulk standard model in the Randall–Sundrum background,” *Physical Review D*, vol. 62, no. 8, Article ID 084025, 2000.
- [67] G. F. Giudice, R. Rattazzi, and J. D. Wells, “Graviscalars from higher-dimensional metrics and curvature-Higgs mixing,” *Nuclear Physics B*, vol. 595, no. 1-2, pp. 250–276, 2001.
- [68] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, “Experimental probes of localized gravity: on and off the wall,” *Physical Review D*, vol. 63, no. 7, Article ID 075004, 20 pages, 2001.
- [69] K. Agashe, H. Davoudiasl, G. Perez, and A. Soni, “Warped gravitons at the CERN LHC and beyond,” *Physical Review D*, vol. 76, no. 3, Article ID 036006, 2007.
- [70] R. Bao, M. S. Carena, J. Lykken, M. Park, and J. Santiago, “Revamped braneworld gravity,” *Physical Review D*, vol. 73, no. 6, Article ID 064026, 25 pages, 2006.
- [71] M. E. Peskin and T. Takeuchi, “New constraint on a strongly interacting Higgs sector,” *Physical Review Letters*, vol. 65, no. 8, pp. 964–967, 1990.
- [72] M. E. Peskin and T. Takeuchi, “Estimation of oblique electroweak corrections,” *Physical Review D*, vol. 46, no. 1, pp. 381–409, 1992.

- [73] C. Csáki, M. L. Graesser, and G. D. Kribs, “Radion dynamics and electroweak physics,” *Physical Review D*, vol. 63, no. 6, Article ID 065002, 2001.
- [74] A. Kobakhidze and J. Yue, “Excluding a generic spin-2 Higgs impostor,” *Physics Letters B*, vol. 727, no. 4-5, pp. 456–460, 2013.
- [75] N. D. Christensen and C. Duhr, “FeynRules—Feynman rules made easy,” *Computer Physics Communications*, vol. 180, no. 9, pp. 1614–1641, 2009.
- [76] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer, and T. Stelzer, “MadGraph 5: going beyond,” *Journal of High Energy Physics*, vol. 2011, no. 6, article 128, 2011.
- [77] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer, and T. Reiter, “UFO—the universal FeynRules output,” *Computer Physics Communications*, vol. 183, no. 6, pp. 1201–1214, 2012.
- [78] J. Ellis, D. S. Hwang, V. Sanz, and T. You, “A fast track towards the ‘Higgs’ spin and parity,” *Journal of High Energy Physics*, vol. 2012, article 134, 2012.
- [79] J. Ellis, R. Fok, D. S. Hwang, V. Sanz, and T. You, “Distinguishing ‘Higgs’ spin hypotheses using $\gamma\gamma$ and WW^* decays,” *The European Physical Journal C*, vol. 73, article 2488, 2013.
- [80] A. Djouadi, R. M. Godbole, B. Mellado, and K. Mohan, “Probing the spin-parity of the Higgs boson via jet kinematics in vector boson fusion,” *Physics Letters B*, vol. 723, no. 4-5, pp. 307–313, 2013.
- [81] J. Ellis, V. Sanz, and T. You, “Prima facie evidence against spin-two Higgs impostors,” *Physics Letters B*, vol. 726, no. 1–3, pp. 244–250, 2013.
- [82] J. Ellis, V. Sanz, and T. You, “Associated production evidence against Higgs impostors and anomalous couplings,” *The European Physical Journal C*, vol. 73, article 2507, 2013.
- [83] J. Brehmer, G. Brooijmans, G. Cacciapaglia et al., “The Diboson excess: experimental situation and classification of explanations; a Les Houches pre-proceeding,” <http://arxiv.org/abs/1512.04357>.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

