Research Article

Space-Time Defects and Group Momentum Space

Michele Arzano¹ and Tomasz Trzeńniewski²

¹*Dipartimento di Fisica and INFN, "Sapienza" University of Rome, P.le A. Moro 2, 00185 Roma, Italy*

²*Institute of Theoretical Physics, University of Wrocław, Pl. M. Borna 9, 50-204 Wrocław, Poland*

Correspondence should be addressed to Tomasz Trzeńniewski; tbwbt@ift.uni.wroc.pl

Received 11 May 2017; Accepted 16 July 2017; Published 17 August 2017

Academic Editor: Angel Ballesteros

Copyright © 2017 Michele Arzano and Tomasz Trzeńniewski. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

We study massive and massless conical defects in Minkowski and de Sitter spaces in various space-time dimensions. The energy momentum of a defect, considered as an (extended) relativistic object, is completely characterized by the holonomy of the connection associated with its space-time metric. The possible holonomies are given by Lorentz group elements, which are rotations and null rotations for massive and massless defects, respectively. In particular, if we fix the direction of propagation of a massless defect in $n + 1$ -dimensional Minkowski space, then its space of holonomies is a maximal Abelian subgroup of the $AN(n - 1)$ group, which corresponds to the well known momentum space associated with the n -dimensional κ -Minkowski noncommutative space-time and κ -deformed Poincaré algebra. We also conjecture that massless defects in n -dimensional de Sitter space can be analogously characterized by holonomies belonging to the same subgroup. This shows how group-valued momenta related to four-dimensional deformations of relativistic symmetries can arise in the description of motion of space-time defects.

1. Introduction

Conical space-time defects were first introduced by Staruszkiewicz [1] as point particles coupled to gravity in $2 + 1$ space-time dimensions. Several years later, Deser et al. [2] generalized this result to configurations of many arbitrary particles. The peculiarity of such systems is that particles are represented by topological defects, due to the lack of local degrees of freedom for gravity in three space-time dimensions, and thus strictly speaking they do not interact gravitationally. The same kind of particle solution was also shown to exist for nonzero cosmological constant, corresponding to $2 + 1$ -dimensional de Sitter or anti-de Sitter space [3].

In $3 + 1$ dimensions, a conical defect can be obtained simply by replacing the point-like singularity with a singular one-dimensional object. Such linear defects are known under the name of *cosmic strings* and first turned out to be possibly generated during a spontaneous gauge symmetry breaking in the early universe [4]. They were then studied in cosmology as a possible source of primordial density fluctuations [5], where their contribution was subsequently constrained by observations of the cosmic microwave background (see,

e.g., [6]). Cosmic “superstrings” can be also produced in string inflation models [7]. At a more abstract level, ’t Hooft has recently employed cosmic strings as ingredients of his *piecewise flat* model of gravity [8]. In full analogy with a point particle in $2 + 1$ dimensions, the strings he considers are infinitely thin and straight. The generalization of the concept of a codimension two conical defect to an arbitrary number of space-time dimensions is then completely straightforward: a straight line is replaced by a hyperplane. Such idealized defects are also the focus of our paper.

A somewhat surprising feature of the description of a point particle in $2 + 1$ gravity (with vanishing cosmological constant) is that its extended momentum space is actually a Lie group, $SO(2, 1)$, which as a manifold has the form of three-dimensional anti-de Sitter space [9]. Interestingly, a nontrivial geometry of momentum space is also one of the effects of (quantum) deformations of relativistic symmetries in four space-time dimensions, which are at the basis of the so-called *doubly special relativity* framework [10, 11] and of its most recent incarnation: the *relative locality* program [12–14]. In this context, the role of deformed symmetries within $2 + 1$ gravity has been studied in various approaches since the late 1990s [15–21].

One might wonder whether conical defects in higher dimensional (Minkowski) spaces possess properties similar to the three-dimensional case and, in particular, if their motion can be parametrized by some Lie group related to deformations of relativistic symmetries. To this end, let us recall that the best studied example of deformed relativistic symmetries in four dimensions is given by the κ -Poincaré (Hopf) algebra [22], associated with the noncommutative κ -Minkowski space [23]. The momentum space corresponding to such a deformation is given by the Lie group $\text{AN}(3)$, which is a subgroup of the five-dimensional Lorentz group $\text{SO}(4, 1)$ [24, 25]. Therefore, it is of interest to investigate whether this group plays any role in the characterization of space-time defects. We argue here that, at least to a certain extent, this is indeed the case when one considers *light-like* or *massless* defects.

A massless conical defect can be obtained by boosting a (timelike) massive defect to the speed of light. In order to achieve a nontrivial limit, one usually applies the boost following a prescription of Aichelburg and Sexl, which was first proposed to derive the gravitational field of a photon from the Schwarzschild metric [26]. This method was subsequently extended to other singular null objects, in particular massless cosmic strings [27–29]. Some of the null sources coexist with impulsive gravitational waves but this is not the case for the straight strings [29]. Moreover, since $3 + 1$ -dimensional (anti-)de Sitter space can be represented as a hyperboloid embedded in the $4 + 1$ -dimensional flat space-time, the Aichelburg-Sexl boost can also be utilized [30, 31] to obtain massless particle solutions in (anti-)de Sitter space from the Schwarzschild (anti-)de Sitter solution (with particles generating impulsive gravitational waves). The same approach was applied to derive null particle solutions in $2 + 1$ -dimensional (anti-)de Sitter space [32]. Since in $2 + 1$ dimensions there are no propagating degrees of freedom, in this case, particles are not accompanied by gravitational waves.

In this work, we analyze how momenta of massless defects in $n + 1$ -dimensional Minkowski space can be described by null rotations belonging to the $\text{AN}(n - 1)$ group, the momentum space corresponding to the n -dimensional noncommutative κ -Minkowski space. We also discuss the relation between massless defects in Minkowski space and the same defects in de Sitter space of one dimension less. The structure of the paper is as follows. In Section 2, we begin with a discussion of massive conical defects generalized to $4 + 1$ -dimensional Minkowski space. Static defects are presented first and then we turn to moving defects. In particular, we describe how such defects are characterized by holonomies of the metric connection, which encode their energy and momentum. In Section 3, we explain how massless defects in $n + 1$ dimensions are parametrized by elements of the $\text{AN}(n - 1)$ group, especially for a fixed direction of propagation. In Section 4, we discuss both massive and massless defects in de Sitter space, focusing on the $3 + 1$ -dimensional case. We conclude with a summary of our analysis.

2. Conical Defects in Minkowski Space

An intuitive picture of a conical space-time defect is obtained by considering Minkowski space with a wedge removed and the faces of the wedge “glued together” to form a cone. The gluing is realized by identifying the opposite faces via a rotation by the *deficit angle* characterizing the defect. One can generalize this construction to include identifications of the two faces (not necessarily forming a wedge) via a general Poincaré transformation, thus producing defects known as “dislocations” and “disclinations,” adopting the terminology used in the classification of defects in solid media (see [33] for an extended discussion).

As we will discuss below, conical defects are curvature singularities. Thus, a given defect carries some mass/energy on its infinitely thin hyperplane. In $2 + 1$ -dimensional gravity, when one works with units $c = 1$, Newton’s constant G has the dimension of inverse mass (it is the inverse Planck mass) and hence provides a natural mass scale for the theory. The quantity $\mu := Gm$ is the dimensionless “rest energy per point,” which is proportional to the deficit angle of a particle at rest $\alpha = 8\pi Gm$. If we generalize this picture to $n + 1$ dimensions, G will have the dimension of length to the power $n - 2$ times inverse mass. Similar to the three-dimensional case, we can then define the dimensionless energy density $\mu := G\rho$, where ρ is the mass per unit of volume of the defect’s hyperplane (e.g., length of the string in $3 + 1$ dimensions).

The metric of a conical defect can be written in terms of cylindrical coordinates, in which its geometric properties are most transparent. Here, we focus on a defect in $4 + 1$ space-time dimensions, for reasons which will become clear in the following, but an extension to any number of dimensions is straightforward. In analogy with a conical defect in $2 + 1$ [2] and $3 + 1$ dimensions [29], in the five-dimensional case, the metric has the form

$$ds^2 = -dt^2 + dz^2 + dw^2 + dr^2 + (1 - 4\mu)^2 r^2 d\phi^2. \quad (1)$$

This metric describes an infinite, flat “cosmic brane” identified with the zw -plane. The brane can be thought of as the “tip” of a conical defect whose deficit angle $\alpha = 8\pi\mu$ is determined by the (dimensionless) *rest energy density* of the brane $\mu \in (0, 1/4)$. The surrounding space-time is locally isometric to Minkowski space. Indeed, as one can verify by a straightforward calculation, the Riemann curvature vanishes everywhere outside the brane’s world volume, which is the tzw -hyperplane. Consequently, the result of the parallel transport of a vector along an arbitrary loop encircling the defect depends only on the conical curvature singularity and the loop’s winding number. The parallel transport around a loop $\gamma(\lambda)$ is measured by the *holonomy* of the Levi-Civita connection given by

$$(h_\gamma)^\mu{}_\nu = \mathcal{P} \exp \left(- \int_\gamma \Gamma^\mu{}_{\sigma\nu} \frac{d\gamma^\sigma}{d\lambda} d\lambda \right), \quad (2)$$

where $\mathcal{P}$ denotes the path ordering and $\Gamma^\mu{}_{\sigma\nu}$ are Christoffel symbols associated with metric (1). One may notice that in four space dimensions it is possible to take a loop around a two-dimensional plane due to the fact that closing a path

around a given hyperplane requires at least two directions orthogonal to it. Holonomy (2) of a loop containing the defect is a nontrivial Lorentz transformation, which can be understood as a result of the presence of a delta-peaked curvature on the defect's world volume [34]. Not surprisingly, this Lorentz transformation is also gluing the faces of the defect's wedge [9].

A generalization of holonomy (2) can be used to characterize both position and momentum of a (moving) defect. This is achieved by considering the parallel transport of a coordinate frame instead of just a vector. If the defect is displaced from the frame's origin, the resulting *Poincaré holonomy* in addition to a Lorentz transformation contains also a translation, carrying information about the defect's position [8, 35]. However, here we will restrict our considerations to Lorentz holonomies (2), which completely describe the energy-momentum density of a moving defect, as it was first shown for $2 + 1$ dimensions [9]. Let us start by explaining how the rest energy density μ appearing in the conical metric is encoded in the Lorentz holonomy of a static defect. We do this by deriving the explicit form of holonomy (2) corresponding to metric (1). The calculation is most easily carried out by picking a simple loop, for example, a circle parametrized by $r = (1 - 4\mu)^{-1}$, $\phi = 2\pi\lambda$, and $t, z, w = 0$. The result written in terms of Cartesian coordinates $x = r \cos \phi$ and $y = r \sin \phi$ is given by

$$h_R(\alpha) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & \cos(\alpha) & -\sin(\alpha) & 0 & 0 \\ 0 & \sin(\alpha) & \cos(\alpha) & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (3)$$

Such a holonomy is simply a rotation (an *elliptic* Lorentz transformation) by the deficit angle $\alpha = 8\pi\mu$ around the origin in the xy -plane.

The metric of a moving defect can be obtained [29] by “boosting” the static metric (1) in Cartesian coordinates using an ordinary boost in a direction which lies in the xy -plane. Indeed, metric (1) is obviously invariant under boosts in the

directions z and w and therefore the defect can only move in the plane perpendicular to itself, like a point particle in $2 + 1$ dimensions. For simplicity, we may take a boost in the x direction:

$$\begin{aligned} t &\mapsto t \cosh \chi + x \sinh \chi, \\ x &\mapsto x \cosh \chi + t \sinh \chi, \end{aligned} \quad (4)$$

where χ is the rapidity parameter. Introducing light-cone coordinates u, v via $t = (v - u)/\sqrt{2}$, $x = (v + u)/\sqrt{2}$, we obtain the metric

$$ds^2 = 2dudv + dy^2 + dz^2 + dw^2 - \left(1 - (1 - 4\mu)^2\right) \cdot \frac{(e^\chi(udy - ydu) + e^{-\chi}(vdy - ydv))^2}{e^{2\chi}u^2 + e^{-2\chi}v^2 + 2(uv + y^2)}, \quad (5)$$

describing a massive brane traveling with velocity $V = \tanh \chi$ in the direction x . It can be shown after a simple calculation [35] that the deficit angle α' of a moving defect becomes wider than the deficit angle α at rest and that they are related by the formula $\tan(\alpha'/2) = \tan(\alpha/2) \cosh \chi$. Here, we observe that, in the limit of small α and α' , the latter simplifies to

$$\alpha' = \alpha \cosh \chi, \quad (6)$$

which can be interpreted as the familiar expression for the relativistic energy density. Namely, as we already explained, the rest energy density $\mathcal{E}_0 = \mu \sim \alpha$, while the factor $\cosh \chi = (1 - V^2)^{-1/2}$ and hence the total energy density $\mathcal{E}_0(1 - V^2)^{-1/2} = \mathcal{E} \sim \alpha'$.

One can also notice that, due to the nontrivial parallel transport of frames, the presence of a conical singularity in the otherwise flat space-time introduces an ambiguity in the direction of a boost as perceived by different observers. To resolve this issue, we have to consistently choose one frame for defining boosts (e.g., such one that the defect's wedge lies symmetrically behind the boost's direction). On the other hand, the whole problem can be avoided by directly boosting holonomy (3). This is achieved by acting on the rotation $h_R(\alpha)$ by a boost $b(\chi)$ via the conjugation $b^{-1}(\chi)h_R(\alpha)b(\chi)$. The resulting Lorentz holonomy

$$h_R(\alpha, \chi) = \begin{pmatrix} 1 + 2 \sinh^2(\chi) \sin^2(\alpha/2) & -\sinh(2\chi) \sin^2(\alpha/2) & -\sinh(\chi) \sin(\alpha) & 0 & 0 \\ \sinh(2\chi) \sin^2(\alpha/2) & 1 - 2 \cosh^2(\chi) \sin^2(\alpha/2) & -\cosh(\chi) \sin(\alpha) & 0 & 0 \\ -\sinh(\chi) \sin(\alpha) & \cosh(\chi) \sin(\alpha) & \cos(\alpha) & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad (7)$$

is thus an element of the *conjugacy class* of rotations by the angle α , which fully characterizes the motion of a defect described at rest by metric (1).

2.1. Massless Defects. So far, we have dealt with massive defects. Not surprisingly, it is also possible to consider

massless (light-like) defects, moving with the speed of light. This can be achieved by deriving the theoretical limiting case of the calculation described above, which consists in performing an “infinite” boost via the prescription first introduced by Aichelburg and Sexl [26] in the case of the Schwarzschild metric. The trick is to keep the quantity

$\varrho \equiv 8\pi\mu \cosh \chi$ fixed (which is the laboratory energy density, as can be shown by a simple calculation [29]) while taking the limit of the rapidity $\chi \rightarrow \infty$. If we use the well known distributional identity

$$\lim_{b \rightarrow 0} \frac{b}{a^2 + b^2} = \pi \delta(a), \quad (8)$$

metric (5) in the Aichelburg-Sexl limit becomes

$$ds^2 = 2dudv + dy^2 + dz^2 + dw^2 - \sqrt{2}\varrho |y| \delta(u) du^2 \quad (9)$$

and it describes the geometry of a massless cosmic brane. One might be worried about the distributional nature of this metric. However, solutions to Einstein equations involving distributions have a long history and have been extensively studied (see, e.g., [36]). Metric (9) corresponds to a zw -plane moving along the null direction v . Space-time is still flat outside the defect's world volume and hence the holonomy of a loop around the defect reflects the presence of a curvature singularity at the hyperplane $u, y = 0$. In the considered coordinates, it is convenient to take a square loop with $u, y \in [-1, 1]$ and transform the associated holonomy to Cartesian coordinates, which gives

$$h_P(\varrho) = \begin{pmatrix} 1 + \frac{\varrho^2}{2} & -\frac{\varrho^2}{2} & -\varrho & 0 & 0 \\ \frac{\varrho^2}{2} & 1 - \frac{\varrho^2}{2} & -\varrho & 0 & 0 \\ -\varrho & \varrho & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (10)$$

This holonomy is a *null rotation* by the deficit angle ϱ with respect to the null axis v in the uy -plane. Such Lorentz transformations are called *parabolic* since their orbits are parabolas on a given light-like plane (i.e., open curves) in contrast to the circular orbits of usual rotations. Null rotations can be expressed by the homogeneous combinations of boosts and usual rotations, as we will discuss in detail in the next section.

An alternative construction of the metric of a massless defect, free from the “ambiguity” associated with performing a boost in the conical space-time, was given in [29]. As mentioned above, one can perform a boost $b(\chi)$ of the holonomy describing a static defect (3) via the conjugation $b^{-1}(\chi)h_R(\alpha)b(\chi)$. If we take the Aichelburg-Sexl limit, the resulting Lorentz group element will be a null rotation (10). The metric leading to such a holonomy can be then reconstructed following the example of a massive defect (1). The geometries will be similar but now the deficit angle has to be cut out from a null plane instead of a spacelike one. To this end, we need cylindrical-like light-cone coordinates, which can be derived starting from usual light-cone coordinates and making a transformation to $q = v + (1/2)y^2u^{-1}$, $\varphi = yu^{-1}$. However, the “angular” coordinate φ has an infinite range and therefore the cut cannot be introduced by a simple rescaling

of it, like it is done for ϕ in (1). One rescales φ only in the region $u > 0$, obtaining the metric of the form

$$ds^2 = 2dudq + dz^2 + dw^2 + (1 - f(\varphi)\Theta(u))^2 u^2 d\varphi^2, \quad (11)$$

$$\int_{-\infty}^{\infty} f(\varphi) d\varphi = \sqrt{2}\varrho,$$

where a smooth function $f(\varphi) < 1$ with the compact support is the analog of the rescaling factor 4μ in (1). As one can verify, this metric has the same holonomy as (9) and thus they are equivalent. Furthermore, (11) has the advantage of being smooth away from the $u = 0$ hyperplane and it shows that a massless conical defect is not accompanied by gravitational waves (see [29] for details).

In the next section, we will focus on the group theoretic structure needed to characterize the motion of a massless defect. This will lead us to a suggestive connection with the momentum space of certain widely studied models of *deformed relativistic symmetries*.

3. Momentum Space of Massless Defects and the $AN(n)$ Lie Group

As discussed in the previous section, the motion of a conical defect in Minkowski space (with a given number of dimensions) is completely characterized by the Lorentz holonomy associated with a loop encircling the defect. Thus, the space of all possible holonomies can be thought of as energy-momentum space of the defect. In order to describe this momentum space in more detail in the case of massless defects, we will now parametrize such a defect using space-time vectors, as it is customarily done for a moving point particle. To get an intuitive picture, we start from familiar defects in $3+1$ dimensions and then generalize the discussion to higher dimensional cases and in particular to $4+1$ dimensions, which we are especially interested in.

Let us first consider a massless cosmic string in four-dimensional Minkowski space that is oriented along the spacelike direction given by the vector $\mathbf{d} = (0, 0, 0, 1)$ and moving in the null direction $\mathbf{n} = (1, 1, 0, 0)$ (in Cartesian coordinates t, x, y, z). Its motion is completely captured by the holonomy (cf. (10))

$$h_P(\varrho) = \begin{pmatrix} 1 + \frac{\varrho^2}{2} & -\frac{\varrho^2}{2} & -\varrho & 0 \\ \frac{\varrho^2}{2} & 1 - \frac{\varrho^2}{2} & -\varrho & 0 \\ -\varrho & \varrho & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (12)$$

The Lorentz group element (12) is a null rotation by the angle ϱ with respect to the null axis $\mathbf{n}$, in the plane spanned by the light-like vector $(1, -1, 0, 0)$ and spacelike vector $(0, 0, 1, 0)$. In general, a massless string in $3+1$ dimensions can be completely characterized by 4 parameters [35]. Indeed, one needs to specify a parabolic angle ϱ , carrying the defect's

energy density, a light-like vector $\mathbf{n}$, along which the defect propagates, and a spacelike vector $\mathbf{d}$, which is the direction of the spatial extension of the defect, with the orthogonality condition $\mathbf{n} \cdot \mathbf{d} = 0$ (a defect is invariant under boosts acting along $\mathbf{d}$). The overall scaling of $\mathbf{n}$ and $\mathbf{d}$ is irrelevant. In the end, one has $2 + 2 - 1 + 1 = 4$ independent coefficients, which parametrize all possible holonomies. Thus, the space of holonomies/momenta of a massless string is bigger than momentum space of a massless particle in $3 + 1$ dimensions, which is only three-dimensional. The reason is that conical defects in space-time with more than three dimensions are extended objects and the holonomies must account for their nontrivial orientation in space.

The full space of momenta described above can be restricted in two simple ways. Firstly, we may fix the spatial orientation of a string $\mathbf{d}$. Then, however, the defect effectively behaves like a point particle in $2 + 1$ dimensions, since it can only move in two directions perpendicular to itself. Secondly, we may choose to fix the direction of motion $\mathbf{n}$. It turns out that this case is more interesting. To be specific, let us introduce a complete orthogonal set of null vectors $\mathbf{n}_{(x)} = (1, 1, 0, 0)$, $\mathbf{n}_{(y)} = (1, 0, 1, 0)$, and $\mathbf{n}_{(z)} = (1, 0, 0, 1)$, as well as a set of orthonormal spacelike vectors $\mathbf{d}_{(x)} = (0, 1, 0, 0)$, $\mathbf{d}_{(y)} = (0, 0, 1, 0)$, and $\mathbf{d}_{(z)} = (0, 0, 0, 1)$. Suppose that we fix the direction of defect's velocity as $\mathbf{n} = \mathbf{n}_{(a)}$, $a \in \{x, y, z\}$. Then, the spatial orientation of a defect is orthogonal to $\mathbf{n}_{(a)}$ and is a linear combination of two vectors $\mathbf{d}_{(j)}$, $j = 1, 2$, which can be chosen to be $\mathbf{d}_{(b)}$, $b \neq a$. For example, for $\mathbf{n}_{(x)}$, we may take $\mathbf{d}_{(1)} \equiv \mathbf{d}_{(z)}$ and $\mathbf{d}_{(2)} \equiv \mathbf{d}_{(y)}$. We observe that, for a given $\mathbf{n}_{(a)}$, the space of holonomies of a massless defect is a subgroup of the Lorentz group $\text{SO}(3, 1)$ determined by the generators of null rotations $X_j^{(a)} := J_{bt} + J_{ba}$, with each j corresponding to one of the spatial indices $b \neq a$ (see also (20)). The holonomy of a defect characterized by $\mathbf{n}_{(a)}$ and $\mathbf{d}_{(j)}$ is obtained by exponentiating such a generator to

$$h_j^{(a)}(k_j) = e^{ik_j X_j^{(a)}}, \quad a = x, y, z, \quad j = 1, 2 \quad (13)$$

(no summation in the exponent), while the parabolic angle $\varrho \equiv k_j$ parametrizes the family of holonomies. For example, if the defect moves along $\mathbf{n}_{(x)}$, we have two generators $X_1^{(x)} = J_{yt} - J_{yx}$ and $X_2^{(x)} = J_{zt} + J_{zx}$, which can be written in a four-dimensional matrix representation as

$$\begin{aligned} X_1^{(x)} &= i \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ X_2^{(x)} &= i \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (14)$$

Holonomy (12) in this picture is given by the group element $h_1^{(x)} = e^{ik_1 X_1^{(x)}}$, where $\varrho \equiv k_1$, as can be verified by a direct calculation.

To obtain the holonomy of a defect with a fixed $\mathbf{n}_{(a)}$ but an arbitrary $\mathbf{d}$, we just take the product

$$h_1^{(a)}(k_1) h_2^{(a)}(k_2) = e^{ik^j X_j^{(a)}} \quad (15)$$

and then the (normalized) spacelike vector is a linear combination $\mathbf{d} = \sqrt{k_1^2 + k_2^2}^{-1} (-k_2 \mathbf{d}_{(1)} + k_1 \mathbf{d}_{(2)})$, while the parabolic angle is given by $\varrho \equiv \sqrt{k_1^2 + k_2^2}$, as one can straightforwardly calculate. Therefore, the defect's energy density is naturally expressed in terms of group coordinates k_1 and k_2 , which also specify the defect's orientation. This agrees with the counting of degrees of freedom discussed above since the choice of $\mathbf{n}$ fixes two parameters and another two remain free.

Let us now notice that the generators $X_1^{(a)}$ and $X_2^{(a)}$ commute and thus span a *maximal Abelian subalgebra* of the Lorentz algebra $\mathfrak{so}(3, 1)$. To complete the picture, we may additionally consider the boost generator $X_0^{(a)} := J_{0a}$, which in our matrix representation for $a = x$ is given by

$$X_0^{(x)} = -i \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (16)$$

(The representations of generators for other choices of $\mathbf{n}_{(a)}$ are collected in Appendix A.) Group elements generated by a given $X_0^{(a)}$, namely,

$$g_0^{(a)} = e^{ik_0 X_0^{(a)}}, \quad a = x, y, z, \quad (17)$$

form a one-dimensional subgroup of hyperbolic Lorentz transformations in the direction of the spatial component of $\mathbf{n}_{(a)}$, which can be seen as complementary with respect to null rotations generated by $X_1^{(a)}$ and $X_2^{(a)}$. Acting on a holonomy $h_j^{(a)}(k_j)$ via the conjugation

$$g_0^{(a)} h_j^{(a)}(k_j) (g_0^{(a)})^{-1} = h_j^{(a)}(e^{k_0} k_j), \quad (18)$$

one rescales the parabolic angle $\varrho \equiv k_j$ by the factor of e^{k_0} , while preserving the orientation of the defect. Thus, the effect of a boost is to red- or blue-shift the defect's energy density. We should stress that the boost, analogously to the case of massive defects, has to act via conjugation since such an action preserves the trace of a Lorentz group element and consequently the property that the holonomy belongs to the parabolic conjugacy class.

Finally, we notice that the generator $X_0^{(a)}$ does not commute with $X_1^{(a)}$ and $X_2^{(a)}$, and the respective commutators are given by

$$[X_0^{(a)}, X_j^{(a)}] = -i X_j^{(a)}, \quad j = 1, 2. \quad (19)$$

The Lie algebra generated by $X_0^{(a)}$, $X_1^{(a)}$, and $X_2^{(a)}$ (with a trivial commutator between $X_1^{(a)}$ and $X_2^{(a)}$) is usually denoted by

$\mathfrak{an}(2)$ and called the (three-dimensional) *Abelian nilpotent algebra*, since it possesses an Abelian subalgebra spanned by the generators $X_1^{(a)}$ and $X_2^{(a)}$, which are also nilpotent (e.g., $(X_j^{(a)})^3 = 0$) in a four-dimensional matrix representation.

The generalization of the above picture to any number of space-time dimensions is conceptually straightforward. In $n + 1$ dimensions, we have n independent null directions, corresponding to n families of generators of null rotations and complementary boosts, labeled by $a = 1, \dots, n$:

$$\begin{aligned} X_0^{(a)} &= J_{0a}, \\ X_j^{(a)} &= J_{j0} + J_{ja}, \\ j \neq a &= 1, \dots, n, \end{aligned} \quad (20)$$

where $J_{\mu\nu}$ are generators of the Lorentz group $\text{SO}(n, 1)$. More precisely, J_{0a} generates boosts in the x_a -direction, J_{j0} boosts in one of the directions x_j orthogonal to x_a , and J_{ja} rotations in the plane spanned by x_a and x_j . Similar to the generators appearing in (19), families (20) are actually representations of the generators X_0 and X_j (with nilpotent generators satisfying $(X_j)^3 = 0$ in an $(n + 1) \times (n + 1)$ matrix representation) of the $\mathfrak{an}(n - 1)$ algebra, a subalgebra of the Lorentz algebra $\mathfrak{so}(n, 1)$.

Quite interestingly, the Lie algebra $\mathfrak{an}(n - 1)$ is very popular in the noncommutative geometry community, where it is known as the κ -Minkowski space-time (first introduced in the four-dimensional version in [23]) and the generators $X_0^{(a)}$ and $X_j^{(a)}$, after the appropriate rescaling by a dimensionful constant, are identified with time and space coordinates, respectively. Furthermore, the group $\text{AN}(n - 1)$ generated by $\mathfrak{an}(n - 1)$ can be obtained from the (local) Iwasawa decomposition of the Lorentz group [37] and, in the field theoretic models on κ -Minkowski space, coordinates on this group are *momentum variables* determined by a *quantum group Fourier transform* [20, 38, 39] of functions of noncommuting space-time coordinates. These momentum coordinates transform under the nonlinear boosts and correspond to translation generators of a quantum deformation of the Poincaré algebra known as the κ -Poincaré Hopf algebra [22, 40]. From the geometric point of view, the Lie group manifold $\text{AN}(n - 1)$ is given by half of the n -dimensional de Sitter space [24, 25] and thus the momentum space of such a model is curved, which is a characteristic feature of the recently proposed “relative locality” approach to modeling phenomenological aspects of quantum gravity [12, 13].

In more than $3 + 1$ dimensions, to completely characterize a given massless defect, it is enough to take a parabolic deficit angle, carrying its energy density, and two vectors that determine a two-form normal to the defect’s world volume (which is of codimension 2) [35]. The vectors (one light-like and one spacelike) are mutually orthogonal, and their scaling is irrelevant, and thus in $n + 1$ dimensions we deal in total with $2n - 2$ parameters, which should be encoded in the defect’s holonomy. If we now consider the restriction imposed on a defect by fixing its direction of motion, as we did in the

$3 + 1$ -dimensional case, then the holonomy will have $n - 1$ parameters less and hence only $n - 1$ parameters will remain. Therefore, momentum space of a restricted defect is always given by the maximal Abelian subgroup of $\text{AN}(n - 1)$, formed by null rotations.

The most interesting is the $4 + 1$ -dimensional case, corresponding to (a subgroup of) the $\text{AN}(3)$ group, which is the momentum space associated with the κ -Poincaré algebra in $3 + 1$ space-time dimensions. The holonomy of an arbitrary massless cosmic brane with the fixed velocity vector $\mathbf{n}$ (we drop here the superscript (a)) can be written as

$$h(k_1, k_2, k_3) = h_1(k_1) h_2(k_2) h_3(k_3), \quad (21)$$

where h_j , $j = 1, 2, 3$, are defined analogously to their four-dimensional counterparts (13) and (17). As can be shown after some calculations, the energy density of a brane characterized by this holonomy is given by $\varrho = \sqrt{k_1^2 + k_2^2 + k_3^2}$. The brane’s spatial plane is spanned by two perpendicular vectors, which can be put in the form $\mathbf{d}_I = n\mathbf{d}_{(1)} + n'\mathbf{d}_{(2)} + \mathbf{d}_{(3)}$, $\mathbf{d}_{II} = n''\mathbf{d}_{(1)} - (1 + nn'')(n')^{-1}\mathbf{d}_{(2)} + \mathbf{d}_{(3)}$, where the coefficients $n = (k_2 - k_3)/(k_1 - k_2)$, $n' = (k_3 - k_1)/(k_1 - k_2)$, and $n'' = (k_2(k_2 - k_1) + k_3(k_3 - k_1))/(k_1(k_1 - k_3) + k_2(k_2 - k_3))$ and the vectors $\mathbf{d}_{(j)}$, $j = 1, 2, 3$, form an orthonormal set orthogonal to $\mathbf{n}_{(a)}$. Thus, there are three parameters n, n', n'' specifying the brane as an extended physical object but one of them is irrelevant since $\mathbf{d}_I$ and $\mathbf{d}_{II}$ give a purely geometrical internal construction, while it is enough to know a single spacelike vector that is orthogonal to the defect’s world volume.

In the section below, we extend our discussion to (massless) defects in $3 + 1$ -dimensional de Sitter space, with cosmological constant $\Lambda > 0$, showing how their space-time metric is strictly related to the metric of defects in $4 + 1$ -dimensional Minkowski space.

4. Conical Defects in de Sitter Space

We look here at a generalization of the derivation of a massless conical defect as it was done in $2 + 1$ -dimensional de Sitter space [32]. (In Appendix B, we have included a similar discussion for defects in anti-de Sitter space.) We begin with a static massive conical defect in $3 + 1$ dimensions [41, 42], analogous to a point particle in $2 + 1$ dimensions [3], whose metric in static de Sitter coordinates has the form

$$\begin{aligned} ds^2 &= -\left(1 - \lambda r^2\right) d\tau^2 + \left(1 - \lambda r^2\right)^{-1} dr^2 \\ &\quad + r^2 \left(d\theta^2 + (1 - 4\mu)^2 \sin^2\theta d\phi^2\right), \end{aligned} \quad (22)$$

(where we denote $\lambda \equiv \Lambda/3$) and describes a cosmic string of rest energy density μ , corresponding to the deficit angle $8\pi\mu$. Let us observe that the above metric can be written in terms

of embedding Cartesian coordinates of the 4 + 1-dimensional Minkowski space

$$\begin{aligned} t &= \sqrt{\lambda^{-1} - r^2} \sinh(\sqrt{\lambda} \tau), \\ w &= \pm \sqrt{\lambda^{-1} - r^2} \cosh(\sqrt{\lambda} \tau), \\ x &= r \cos \phi \sin \theta, \\ y &= r \sin \phi \sin \theta, \\ z &= r \cos \theta, \end{aligned} \quad (23)$$

subject to the (4,1)-hyperboloid condition $\lambda^{-1} = -t^2 + x^2 + y^2 + z^2 + w^2$. Transformed (22) turns out to have an identical form to metric (1) of a defect in 4 + 1-dimensional Minkowski space. There are only two obvious subtleties in the de Sitter case: space-time coordinates are constrained by the hyperboloid condition and the interpretation of μ has to be adjusted according to the number of dimensions, since it now represents the *linear* energy density rather than energy per surface. A consequence of this situation is that we could indirectly apply the derivation of light-like defects from Section 2 to the de Sitter case. Nevertheless, it may be more illuminating to stick to the approach of [32] and see whether we will obtain in this way the same metric as (9).

Our goal is to derive a massless defect. Therefore, for convenience, we first rescale the radial coordinate r to $r/(1 - 4\mu)$ and expand the metric to the first order in μ , which gives

$$\begin{aligned} ds^2 &\approx ds_{ds}^2 \\ &+ 8\mu \left(\lambda r^2 d\tau^2 + (1 - \lambda r^2)^{-2} dr^2 + r^2 d\theta^2 \right), \end{aligned} \quad (24)$$

where ds_{ds}^2 is the pure de Sitter metric $ds_{ds}^2 = -(1 - \lambda r^2) d\tau^2 + (1 - \lambda r^2)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)$. We subsequently transform (24) to embedding coordinates (23) and perform a boost $t \mapsto t \cosh \chi + x \sinh \chi$, $x \mapsto x \cosh \chi + t \sinh \chi$, with the rapidity parameter χ , obtaining

$$\begin{aligned} ds^2 &= ds_{ds}^2 + \frac{8\mu}{(T^2 - w^2)^2} \left((\lambda^{-1} + T^2 - w^2) (wdT - Tdw)^2 \right. \\ &\quad \left. + \frac{\lambda^{-2} (TdT - wdw)^2}{\lambda^{-1} + T^2 - w^2} \right) + 8\mu \\ &\quad \cdot \frac{(z(-TdT + zdz + wdw) + (\lambda^{-1} + T^2 - z^2 - w^2) dz)^2}{(\lambda^{-1} + T^2 - w^2)(\lambda^{-1} + T^2 - z^2 - w^2)}, \end{aligned} \quad (25)$$

where $T \equiv t \cosh \chi - x \sinh \chi$ (and ds_{ds}^2 is now given in embedding coordinates). The metric ds_{ds}^2 is preserved by the boost since the latter belongs to the isometry group of de Sitter space. To proceed further, we introduce light-cone coordinates $u = (x - t)/\sqrt{2}$ and $v = (x + t)/\sqrt{2}$ and employ the Aichelburg-Sexl boost prescription, taking $\chi \rightarrow \infty$ and keeping the laboratory energy density $\varrho := 8\pi\mu \cosh \chi$

constant. Using the distributional identity [30]

$$\lim_{\chi \rightarrow \infty} f(T^2) \cosh \chi = \frac{1}{\sqrt{2}} \delta(u) \int_{-\infty}^{+\infty} f(T^2) dT, \quad (26)$$

we finally obtain

$$ds^2 = 2dudv + dy^2 + dz^2 + dw^2 - \sqrt{2}\varrho |y| \delta(u) du^2. \quad (27)$$

Metric (27) corresponds to a conical defect at the hypersurface $u, y = 0, z^2 + w^2 = \lambda^{-1}$, that is, a light-like, circular string on a meridian of the cosmological horizon of de Sitter space (for an observer on the worldline $y, z, w = 0, x = x(t) > \lambda^{-1}$, the future/past horizon is the sphere $y^2 + z^2 + w^2 = \lambda^{-1}$, $x \pm t = 0$). Such a result is qualitatively different from 2 + 1 dimensions, where there is a pair of massless particles (similar to the massive solution [3]) on the opposite points of the horizon [32]. However, it can simply be seen as the dimensional reduction of a circle to the degenerate case of a pair of points.

The string can be better visualized in a different coordinate system, defined in analogy with [31]. To this end, we first make the following transformation of coordinates:

$$\begin{aligned} t &= \frac{1}{2\eta} \left(\lambda^{-1} - \eta^2 + (X + \lambda^{-1/2})^2 + Y^2 + Z^2 \right), \\ x &= \frac{1}{\sqrt{\lambda}\eta} (X + \lambda^{-1/2}), \\ y &= \frac{1}{\sqrt{\lambda}\eta} Y, \\ z &= \frac{1}{\sqrt{\lambda}\eta} Z, \\ w &= \frac{1}{2\eta} \left(\lambda^{-1} + \eta^2 - (X + \lambda^{-1/2})^2 - Y^2 - Z^2 \right), \end{aligned} \quad (28)$$

and then we set $X = \rho \cos \phi \sin \theta$, $Y = \rho \sin \phi \sin \theta$, and $Z = \rho \cos \theta$, so that metric (27) takes the form

$$\begin{aligned} ds^2 &= \frac{1}{\lambda\eta^2} (-d\eta^2 + d\rho^2 + \rho^2 (d\theta^2 + \sin^2 \theta d\phi^2)) \\ &\quad - \frac{\varrho}{\sqrt{\lambda}} |\sin \phi \sin \theta| \\ &\quad \cdot (\delta(\eta - \rho) (d\eta - d\rho)^2 + \delta(\eta + \rho) (d\eta + d\rho)^2). \end{aligned} \quad (29)$$

As one can observe, this metric describes a light-like circular string lying at the great circle $\phi = 0, \pi, \theta \in [0, \pi]$ on the cosmological horizon $\rho = |\eta|$ of de Sitter space, whose metric is in the first line. The first term containing the Dirac delta is responsible for the string at times $\eta \geq 0$ and the other at times $\eta \leq 0$.

The form of (27) is exactly the same as metric (9) of a light-like cosmic brane in 4 + 1-dimensional Minkowski space, as could be expected from the analogous situation for (22). This result easily generalizes to an arbitrary number of dimensions and thus we might say that a conical defect

in the $n + 1$ -dimensional de Sitter space is a “projection” of the corresponding defect in the embedding $(n + 1) + 1$ -dimensional Minkowski space. The projection is understood in simple geometric terms: the $n - 1$ -dimensional hyperplane of a Minkowski defect is cutting through the n -dimensional sphere of a de Sitter spatial slice, determining the $n - 2$ -dimensional sphere of the projected defect.

5. Summary

In this work, we have provided an exploration of the relation between the holonomies of conical defects in more than three space-time dimensions and group-valued momenta, which appear in scenarios of deformed relativistic symmetries. Our motivation was the well established fact that momentum space of point particles in $2 + 1$ gravity, treated as conical defects, is a Lie group. We started by recalling how the motion of massive conical defects can be characterized by their holonomies, belonging to the Lorentz group. Generalizing the discussion to massless defects, we observed that holonomies describing such defects in $n + 1$ -dimensional Minkowski space, given by null rotations, are elements of the maximal Abelian subgroup of the $AN(n - 1)$ group. In particular, for a fixed direction of propagation, the space of all possible holonomies of a massless defect is exactly the above subgroup. Since the full $AN(n - 1)$ group plays the role of momentum space associated with n -dimensional κ -Minkowski space, this fortunate coincidence provides a partial “physical” rendition of the group-valued momenta emerging in the context of deformed relativistic symmetries, which were originally introduced as rather formal mathematical structures. At the same time, group momentum space (or, more broadly, momentum space with a nontrivial geometry) is a crucial ingredient of doubly special relativity and relative locality approaches to the problem of quantum gravity. Both of these frameworks are based on heuristic assumptions and often use classical and quantum $2 + 1$ gravity as a toy model for their concepts. Thus, our analysis of massless defects offers a practical setting to explore such ideas in a higher number of dimensions and in the context of the well studied κ -Minkowski space, though only to a certain extent (a subgroup of $AN(n - 1)$ will correspond to a subset of κ -Minkowski space). Furthermore, similar group momentum spaces are most likely associated with defects coupled to the BF theory [43], a topological field analog of gravity in more than three dimensions. It remains to be investigated how the space of holonomies generalizes when one considers spinning defects, which is a somewhat problematic case, leading to the occurrence of closed timelike curves [35]. In this paper, we also discussed the case of massless conical defects in $3 + 1$ -dimensional de Sitter space, showing how they can be seen as closed light-like strings and how their metric is strictly related to the one of massless defects in $4 + 1$ -dimensional Minkowski space. This led us to conjecture that the same holonomies, belonging to the $AN(3)$ group, which characterize the latter, could completely capture the motion of massless defects in de Sitter space and that such a statement could be generalized to more space-time dimensions. A proof of this conjecture is postponed to future studies.

Appendix

A. Representations of κ -Minkowski Generators in $3 + 1$ Dimensions

For the purpose of illustration, we write down here the three independent 4×4 matrix representations of the $\mathfrak{an}(2)$ generators that can be used to reconstruct holonomies of massless defects for our choice of vectors $\mathbf{n}_{(a)}$ and $\mathbf{d}_{(b)}$, $a, b = x, y, z$, which we introduced in Section 3. Defects with the vectors $\mathbf{n}_{(x)}$ and $\mathbf{d}_{(1)} \equiv \mathbf{d}_{(z)}$, $\mathbf{d}_{(2)} \equiv \mathbf{d}_{(y)}$ correspond to the representation $X_0^{(x)} = K_x$, $X_1^{(x)} = K_y - J_z$, $X_2^{(x)} = K_z + J_y$, where

$$\begin{aligned} X_0^{(x)} &= -i \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ X_1^{(x)} &= i \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ X_2^{(x)} &= i \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \end{pmatrix}; \end{aligned} \quad (\text{A.1})$$

the vectors $\mathbf{n}_{(y)}$ and $\mathbf{d}_{(1)} \equiv \mathbf{d}_{(z)}$, $\mathbf{d}_{(2)} \equiv \mathbf{d}_{(x)}$ correspond to $X_0^{(y)} = K_y$, $X_1^{(y)} = K_x + J_z$, $X_2^{(y)} = K_z - J_x$, where

$$\begin{aligned} X_0^{(y)} &= -i \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ X_1^{(y)} &= i \begin{pmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ X_2^{(y)} &= i \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 1 & 0 \end{pmatrix}; \end{aligned} \quad (\text{A.2})$$

the vectors $\mathbf{n}_{(z)}$ and $\mathbf{d}_{(1)} \equiv \mathbf{d}_{(y)}$, $\mathbf{d}_{(2)} \equiv \mathbf{d}_{(x)}$ correspond to $X_0^{(z)} = K_z$, $X_1^{(z)} = K_x - J_y$, $X_2^{(z)} = K_y + J_x$, where

$$\begin{aligned} X_0^{(z)} &= K_z = -i \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \\ X_1^{(z)} &= i \begin{pmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \\ X_2^{(z)} &= i \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}. \end{aligned} \quad (\text{A.3})$$

J_a and K_a , $a = x, y, z$, respectively, denote the generators of rotations and boosts.

B. Conical Defects in Anti-de Sitter Space

For completeness, let us also discuss conical defects in anti-de Sitter space, with $\Lambda < 0$. In this case, the situation is very similar to the de Sitter case and therefore we will be very brief. We again study the generalization of a massless defect from 2 + 1 dimensions [32] to 3 + 1 dimensions.

For brevity, we set $\Lambda = -3$. Then, the metric of a static massive defect in 3 + 1 dimensions [44, 45], analogous to a particle in 2 + 1 dimensions [3], in static anti-de Sitter coordinates has the form

$$\begin{aligned} ds^2 &= -(1 + r^2) d\tau^2 + (1 + r^2)^{-1} dr^2 \\ &\quad + r^2 (d\theta^2 + (1 - 4\mu)^2 \sin^2 \theta d\phi^2) \end{aligned} \quad (\text{B.1})$$

and describes a cosmic string of rest energy density μ and the deficit angle $8\pi\mu$. One may expand it to the first power in μ and transform to embedding coordinates

$$\begin{aligned} t &= \sqrt{1 + r^2} \sin \tau, \\ w &= \pm \sqrt{1 + r^2} \cos \tau, \\ x &= r \cos \phi \sin \theta, \\ y &= r \sin \phi \sin \theta, \\ z &= r \cos \theta, \end{aligned} \quad (\text{B.2})$$

which satisfy the (2, 3)-hyperboloid condition $-1 = -t^2 + x^2 + y^2 + z^2 - w^2$. We subsequently perform the Aichelburg-Sexl boost, obtaining eventually (there is no difference for $\Lambda \neq -3$)

$$\begin{aligned} ds^2 &= 2dudv + dy^2 + dz^2 - dw^2 \\ &\quad - \sqrt{2}\ell |y| \delta(u) du^2. \end{aligned} \quad (\text{B.3})$$

The above metric corresponds to a curvature singularity at the hypersurface $u, y = 0, -z^2 + w^2 = 1$, that is, a light-like hyperbolic string. The hyperbola has two branches but these actually belong to the same worldsheet, analogously to 2 + 1 dimensions [32] (in contrast to the de Sitter case), and therefore it is a single defect. Moreover, the form of (B.3) is different from metric (9) of a massless defect in 4 + 1-dimensional Minkowski space. For an additional insight, we change the coordinates to

$$\begin{aligned} t &= \frac{1}{2Z} (1 - \eta^2 + (X + 1)^2 + Y^2 + Z^2), \\ x &= \frac{1}{Z} (X + 1), \\ y &= \frac{1}{2Z} (1 + \eta^2 - (X + 1)^2 - Y^2 - Z^2), \\ z &= \frac{1}{Z} Y, \\ w &= \frac{1}{\eta} Z \end{aligned} \quad (\text{B.4})$$

and once more to $X = \rho \cos \phi \sin \theta$, $Y = \rho \sin \phi \sin \theta$, and $Z = \rho \cos \theta$. Restoring $\Lambda \neq -3$, we find that

$$\begin{aligned} ds^2 &= \frac{3}{|\Lambda| \rho^2 \cos^2 \theta} (-d\eta^2 + d\rho^2 \\ &\quad + \rho^2 (d\theta^2 + \sin^2 \theta d\phi^2)) - \ell \sqrt{\frac{3}{|\Lambda|}} \\ &\quad \cdot \frac{|\sin \phi \sin \theta|}{\cos^2 \theta} (\delta(\eta - \rho) (d\eta - d\rho)^2 \\ &\quad + \delta(\eta + \rho) (d\eta + d\rho)^2). \end{aligned} \quad (\text{B.5})$$

The string lies on the surface $\rho = |\eta|$, which is a hyperboloid in anti-de Sitter space. Again, one term with the Dirac delta is responsible for the defect at times $\eta \geq 0$ and the other at times $\eta \leq 0$.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Acknowledgments

The authors would like to thank M. van de Meent for the very useful correspondence and J. Kowalski-Glikman for comments on the manuscript. The work of Michele Arzano is supported by a Marie Curie Career Integration Grant within the 7th European Community Framework Programme and in part by a grant from the John Templeton Foundation. Tomasz Trzeźniewski acknowledges the support by the Foundation for Polish Science International PhD Projects Programme cofinanced by the EU European Regional Development Fund and the additional funds provided by the National Science

Center under Agreements nos. DEC-2011/02/A/ST2/00294 and 2014/13/B/ST2/04043.

References

- [1] A. Staruszkiewicz, "Gravitation theory in three-dimensional space," *Acta Physica Polonica*, vol. 24, pp. 734–740, 1963.
- [2] S. Deser, R. Jackiw, and G. 't Hooft, "Three-dimensional Einstein gravity: dynamics of flat space," *Annals of Physics*, vol. 152, no. 1, pp. 220–235, 1984.
- [3] S. Deser and R. Jackiw, "Three-dimensional cosmological gravity: dynamics of constant curvature," *Annals of Physics*, vol. 153, no. 2, pp. 405–416, 1984.
- [4] T. W. B. Kibble, "Topology of cosmic domains and strings," *Journal of Physics A*, vol. 9, pp. 1387–1398, 1976.
- [5] M. B. Hindmarsh and T. W. Kibble, "Cosmic strings," *Reports on Progress in Physics*, vol. 58, no. 5, pp. 477–562, 1995.
- [6] N. Bevis, M. Hindmarsh, M. Kunz, and J. Urrestilla, "CMB power spectra from cosmic strings: predictions for the Planck satellite and beyond," *Physical Review D*, vol. 82, Article ID 065004, 18 pages, 2010.
- [7] M. Sakellariadou, "Cosmic strings and cosmic superstrings," *Nuclear Physics B. Proceedings Supplement*, vol. 192/193, pp. 68–90, 2009.
- [8] G. 't Hooft, "A locally finite model for gravity," *Foundations of Physics*, vol. 38, no. 8, pp. 733–757, 2008.
- [9] H.-J. Matschull and M. Welling, "Quantum mechanics of a point particle in (2+1)-dimensional gravity," *Classical and Quantum Gravity*, vol. 15, pp. 2981–3030, 1998.
- [10] G. Amelino-Camelia, "Relativity in spacetimes with short-distance structure governed by an observer-independent (planckian) length scale," *International Journal of Modern Physics D*, vol. 11, pp. 35–59, 2002.
- [11] G. Amelino-Camelia, "Relativity: special treatment," *Nature*, vol. 418, no. 6893, pp. 34–35, 2002.
- [12] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman, and L. Smolin, "Principle of relative locality," *Physical Review D*, vol. 84, no. 8, Article ID 084010, 2011.
- [13] G. Gubitosi and F. Mercati, "Relative locality in κ -Poincaré," *Classical and Quantum Gravity*, vol. 30, Article ID 145002, 21 pages, 2013.
- [14] L. Freidel, R. G. Leigh, and D. Minic, "Quantum gravity, dynamical phase-space and string theory," *International Journal of Modern Physics D*, vol. 23, Article ID 1442006, 9 pages, 2014.
- [15] F. A. Bais and N. M. Muller, "Topological field theory and the quantum double of $SU(2)$," *Nuclear Physics B*, vol. 530, no. 1-2, pp. 349–400, 1998.
- [16] F. A. Bais, N. M. Muller, and B. J. Schroers, "Quantum group symmetry and particle scattering in (2+1)-dimensional quantum gravity," *Nuclear Physics B*, vol. 640, no. 1-2, pp. 3–45, 2002.
- [17] L. Freidel and E. R. Livine, "3D quantum gravity and effective noncommutative quantum field theory," *Physical Review Letters*, vol. 96, no. 22, Article ID 221301, 4 pages, 2006.
- [18] K. Noui, "Three-dimensional loop quantum gravity: towards a self-gravitating quantum field theory," *Classical and Quantum Gravity*, vol. 24, no. 2, pp. 329–360, 2007.
- [19] M. Arzano, J. Kowalski-Glikman, and T. Trzeźniewski, "Beyond Fock space in three-dimensional semiclassical gravity," *Classical and Quantum Gravity*, vol. 31, no. 3, Article ID 035013, 13 pages, 2014.
- [20] M. Arzano, D. Latini, and M. Lotito, "Group momentum space and Hopf algebra symmetries of point particles coupled to 2+1 gravity," *Symmetry, Integrability and Geometry: Methods and Applications (SIGMA)*, vol. 10, no. 079, 23 pages, 2014.
- [21] J. Kowalski-Glikman and T. Trzeźniewski, "Deformed Carroll particle from 2+1 gravity," *Physics Letters B*, vol. 737, pp. 267–271, 2014.
- [22] J. Lukierski, A. Nowicki, and H. Ruegg, "New quantum Poincaré algebra and κ -deformed field theory," *Physics Letters B*, vol. 293, no. 3-4, pp. 344–352, 1992.
- [23] S. Majid and H. Ruegg, "Bicrossproduct structure of κ -Poincaré group and non-commutative geometry," *Physics Letters B*, vol. 334, no. 3-4, pp. 348–354, 1994.
- [24] J. Kowalski-Glikman, "De Sitter space as an arena for doubly special relativity," *Physics Letters B*, vol. 547, no. 3-4, pp. 291–296, 2002.
- [25] J. Kowalski-Glikman and S. Nowak, "Doubly special relativity and de Sitter space," *Classical and Quantum Gravity*, vol. 20, no. 22, pp. 4799–4816, 2003.
- [26] P. C. Aichelburg and R. U. Sexl, "On the gravitational field of a massless particle," *General Relativity and Gravitation*, vol. 2, no. 4, pp. 303–312, 1971.
- [27] C. Lousto and N. G. Sanchez, "Gravitational shock waves generated by extended sources: ultrarelativistic cosmic strings, monopoles and domain walls," *Nuclear Physics B*, vol. 355, pp. 231–249, 1991.
- [28] C. Barrabès, P. A. Hogan, and W. Israel, "Aichelburg-Sexl boost of domain walls and cosmic strings," *Physical Review D*, vol. 66, no. 2, Article ID 025032, 6 pages, 2002.
- [29] M. van de Meent, "Geometry of massless cosmic strings," *Physical Review D*, vol. 87, Article ID 025020, 8 pages, 2013.
- [30] M. Hotta and M. Tanaka, "Shock-wave geometry with nonvanishing cosmological constant," *Classical and Quantum Gravity*, vol. 10, no. 2, pp. 307–314, 1993.
- [31] J. Podolský and J. B. Griffiths, "Impulsive gravitational waves generated by null particles in de Sitter and anti-de Sitter backgrounds," *Physical Review D*, vol. 56, pp. 4756–4767, 1997.
- [32] R.-G. Cai and J. B. Griffiths, "Null particle solutions in three-dimensional (anti-) de Sitter spaces," *Journal of Mathematical Physics*, vol. 40, no. 7, pp. 3465–3475, 1999.
- [33] R. A. Puntigam and H. H. Soleng, "Volterra distortions, spinning strings, and cosmic defects," *Classical and Quantum Gravity*, vol. 14, no. 5, pp. 1129–1149, 1997.
- [34] T. Regge, "General relativity without coordinates," *Il Nuovo Cimento*, vol. 19, no. 3, pp. 558–571, 1961.
- [35] M. van de Meent, "Piecewise flat gravity in 3+1 dimensions," <https://arxiv.org/abs/1111.6468>.
- [36] R. Steinbauer and J. A. Vickers, "The use of generalized functions and distributions in general relativity," *Classical and Quantum Gravity*, vol. 23, no. 10, pp. R91–R114, 2006.
- [37] N. Ya. Vilenkin and A. U. Klimyk, "Representations of Lie Groups and Special Functions," *Kluwer*, 1992.
- [38] C. Guedes, D. Oriti, and M. Raasakka, "Quantization maps, algebra representation, and non-commutative Fourier transform for Lie groups," *Journal of Mathematical Physics*, vol. 54, no. 8, Article ID 083508, 31 pages, 2013.
- [39] L. Freidel, J. Kowalski-Glikman, and S. Nowak, "From non-commutative κ -Minkowski to Minkowski space-time," *Physics Letters B*, vol. 648, no. 1, pp. 70–75, 2007.
- [40] J. Lukierski and H. Ruegg, "Quantum κ -Poincaré in any dimension," *Physics Letters B*, vol. 329, no. 2-3, pp. 189–194, 1994.

- [41] A. M. Ghezelbash and R. B. Mann, "Vortices in de Sitter spacetimes," *Physics Letters B*, vol. 537, no. 3-4, pp. 329–339, 2002.
- [42] E. R. Bezerra de Mello and A. A. Saharian, "Vacuum polarization by a cosmic string in de Sitter spacetime," *Journal of High Energy Physics*, vol. JHEP04, no. 46, 21 pages, 2009.
- [43] J. C. Baez, D. K. Wise, and A. S. Crans, "Exotic statistics for strings in 4D BF theory," *Advances in Theoretical and Mathematical Physics*, vol. 11, no. 5, pp. 707–749, 2007.
- [44] M. H. Dehghani, A. M. Ghezelbash, and R. B. Mann, "Vortex holography," *Nuclear Physics B*, vol. 625, no. 1-2, pp. 389–406, 2002.
- [45] E. R. Bezerra de Mello and A. A. Saharian, "Vacuum polarization induced by a cosmic string in anti-de Sitter spacetime," *Journal of Physics A: Mathematical and Theoretical*, vol. 45, no. 11, Article ID 115402, 18 pages, 2012.