

Research Article

Fermions Tunneling and Quantum Corrections for Quintessential Kerr-Newman-AdS Black Hole

Wajiha Javed  and Rimsha Babar

Department of Mathematics, University of Education, Township, Lahore-54590, Pakistan

Correspondence should be addressed to Wajiha Javed; wajiha.javed@ue.edu.pk

Received 20 October 2018; Revised 9 December 2018; Accepted 16 December 2018; Published 14 January 2019

Academic Editor: Elias C. Vagenas

Copyright © 2019 Wajiha Javed and Rimsha Babar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

This paper is devoted to study charged fermion particles tunneling through the horizon of Kerr-Newman-AdS black hole surrounded by quintessence by using Hamilton-Jacobi ansatz. In our analysis, we investigate Hawking temperature as well as quantum corrected Hawking temperature on account of generalized uncertainty principle. Moreover, we discuss the effects of correction parameter β on the corrected Hawking temperature T_{e-H} , graphically. We conclude that the temperature T_{e-H} vanishes when $\beta = 100$, whereas for $\beta < 100$ and $\beta > 100$, the temperature turns out to be positive and negative, respectively. We observe that the graphs of T_{e-H} w.r.t. quintessence parameter α exhibit behavior only for the particular ranges, i.e., $0 < \alpha < 1/6$, charge $0 < Q \leq 1$, and rotation parameter $0 < a \leq 1$. For smaller and larger values of negative Λ , as horizon increases, the temperature decreases and increases, respectively.

1. Introduction

Hawking radiation is the black-body radiation emitted by a black hole (BH) due to quantum effects near the horizon of a BH [1, 2]. It is named after the physicist Stephen Hawking (1979) who derived a theoretical argument for its existence [3]. Hawking with his collaborators [4] studied quantum mechanical uncertainty principle and observed that the rotating BH should create and emit quantum particles, while the base of BH thermodynamics is propounded by Bekenstein [5], who predicted that the BH must have a finite entropy.

There are various methods to investigate the imaginary part of outward radiated particles action. One of the tunneling methods is named as null geodesic method utilized by Parik and Wilczek [6–8], which is the extension of the analysis of Kraus and Wilczek [9–11]. They studied the tunneling of massless scalar particles. For an outgoing massive particle, the equation of motion is different from that of massless particle. The trajectory of a massless particle is a null geodesic. Another technique to investigate BH's tunneling process is Hamilton-Jacobi ansatz introduced by Agheben et al. [12], which is the extension of complex route analysis of Padmanabhan et al. [13–16].

Later, Kerner and Mann [17, 18] investigated tunneling process of fermion particles by using Hamilton-Jacobi technique and obtained the corresponding Hawking temperature. This phenomenon is based on the calculations of imaginary part of action at horizon which in turn is associated with the Boltzmann factor of emission at Hawking temperature. Also, they [19] investigated fermions tunneling from Rindler and typical nonrotating BH horizons. The WKB approximation is used to investigate the tunneling probability for classically forbidden trajectory from interior to exterior region through horizon. The expression of the tunneling probability Γ is given by

$$\Gamma = \exp \left[-\frac{2}{\hbar} \text{Im } I \right], \quad (1)$$

where I is the semiclassical action of the outgoing particle and \hbar is Planck's constant.

Black holes are effective modes to explore the effects of quantum gravity by investigating their thermodynamical properties. Using generalized uncertainty principle (GUP) for BH physics, some thrilling implications and consequences have been performed in literature [20–22]. Nozari and Saghafi [23] discussed Hawking radiation for massless scalar

particles in the background geometry of Schwarzschild BH by following the Parikh-Wilczek tunneling technique and recovered the tunneling rate as well as corrected Hawking temperature by considering GUP. Kerner et al. [24], Li et al. [25], and Jian et al. [26] investigated the tunneling phenomenon of fermions from the Kerr and Kerr-Newman BHs by applying the WKB approximation to the Dirac equation. Using Hamilton-Jacobi ansatz, the GUP-deformed corrected Hawking temperatures for fermions are derived for various curved spacetimes [27–35].

By taking into account the quantum corrections and back-reaction effects, Singleton et al. [36, 37] studied the information loss paradox and conservation of energy and entropy. Banerjee and Majhi [38–42] studied the Hawking radiation process by using quantum tunneling phenomenon at horizons and investigated the back-reaction effects. The quantum tunneling spectrum for scalar and fermion particles has also been discussed in literature [43–49].

Using Kerner and Mann's formulation, Sharif and Javed [50–55] studied the fermions tunneling phenomenon to investigate Hawking temperatures for charged anti-de Sitter BHs, charged torus-like BHs, *Plebański–Demiański* family of BHs, regular BHs, and traversable wormholes. They [56] also discussed Hawking radiation for a pair of charged accelerating and rotating BHs involving NUT parameter. Moreover, they [57] investigated quantum corrections for regular BHs, i.e., Bardeen and ABGB BHs. Recently, Javed et al. [58] discussed charged vector particles tunneling for a pair of accelerating and rotating BHs as well as for 5D gauged super-gravity BHs.

During 1990's, it is observed that the universe is filled with matter and the appealing power of gravity, which pulls all matter together. Later, the Hubble space telescope perceptions for extremely far off supernovae demonstrated that quite a while prior, the universe was really expanding more gradually than it is today. So, the expansion of universe has not been slow due to gravity, but it is accelerating. Eventually scholars thought that possibly it was a consequence of Einstein's hypothesis of gravity, known as cosmological constant (Λ). Initially, Λ is added by Einstein as a steady term in his field equations of General Relativity in order to study the static universe, later this effort was unsuccessful due to Hubble's observations, which confirmed that our universe is expanding [59]. Recently, Λ is considered as a source term in the field equations, which can be considered as mass of empty space or vacuum energy, adequately utilizing dark energy to adjust gravity [60].

The value of Λ could be either positive or negative as per background geometry. Dark energy is gravitationally repulsive, not attractive. The nature of dark energy is still not well understood but detected by its effects on the rate at which universe expand. The idea of dark energy is more theoretical and numerous things are still remain as matter of consideration [61]. Dark energy as a transient vacuum energy resulting from the potential energy of a dynamical field, is known as *quintessence*. This form of dark energy varies in space and time and distinguished from Λ . It plays a vital role in the expansion theory of big bang [62].

The description of dark energy in terms of negative cosmological constant (AdS space) has already been discussed in literature. Xu and Wang [63] investigated Kerr-Newman BH solution in the background of quintessential field by utilizing the Newman-Janis algorithm. It is well known fact that the Newman-Janis algorithm does not deal with the cosmological constant, so they extended the solution of Kerr-Newman metric to the Kerr-Newman-AdS solution surrounded by the quintessential dark energy through direct computations satisfying Einstein Maxwell field equations in the background of quintessential matter with negative cosmological constant. Also, they analyzed the singularity of Kerr-Newman AdS BH in the presence of quintessential field which is similar as the case of Kerr BH.

In our analysis, the quintessential field has an equation of state

$$p = \omega\rho, \quad (2)$$

where ρ and p denote the energy density and pressure, respectively. While ω represents the state parameter having range, $-1 < \omega < -1/3$ [63]. For the given AdS BH which is surrounded by quintessential field, there exist two cosmological horizons [64]. As $\omega \rightarrow -1$, the effects of quintessence and cosmological constant will be similar [65]. Moreover, the PV criticality and thermodynamical properties (pressure, volume and Hawking temperature) of quintessential RN-AdS BH are discussed by Li [66].

In our analysis, we are going to extend the work of Xu and Wang in order to investigate the thermodynamical properties of Kerr-Newman-AdS BH in the presence of quintessential field for fermion particles. For this purpose, we have considered quantum tunneling phenomenon of Hawking radiation to investigate Hawking temperature at BH's horizon as well as its modified quantum corrected form by utilizing Hamilton-Jacobi ansatz. Moreover, we investigate the effects of quintessential field on BH's thermodynamics. The main purpose of this paper is to investigate Hawking temperature from different aspects for AdS BH in the background of quintessential field incorporating quantum effects. Moreover, we observe the graphical behavior of corrected Hawking temperature with respect to horizon (r_+) and quintessence parameter for correction parameter (β) and cosmological constant.

The paper is outlined as follows. In Section 2, we introduce metric of the Kerr-Newman-AdS BH surrounded by the quintessence. In Section 3, we investigate the Hawking radiation phenomenon for charged fermion particles for the above mentioned BH and recoup the tunneling probability and Hawking temperature. By utilizing the modified Dirac equation incorporating GUP, the corrected Hawking temperature is determined in Section 4, while Section 5 provides entropy corrections. Section 6 consists of the graphical analysis of quantum corrected Hawking temperature, where we study the effects of β and Λ in detail. The last section contains concluding remarks.

2. Kerr-Newman-AdS BH with Quintessence

The accelerating expansion of the universe implies the crucial contribution of matter with negative pressure in the evolution of universe. This expansion could also be the result of cosmological constant or quintessence matter. If quintessence matter exists all over the universe, it can also be around a BH. Kerr BH has many interesting properties distinct from its nonspinning counterpart; i.e., Schwarzschild BH. Newman and Janis [67–70] analyzed that the Kerr metric [71] could be obtained from the Schwarzschild metric using a complex transformation within the framework of the Newman-Penrose formalism [72]. A similar procedure was applied to the Reissner-Nordstrom metric to generate Kerr-Newman metric [68]. The Newman-Janis algorithm proved to be prosperous in generating new stationary solutions of the Einstein field equations [73–76]. Zhaoyi and Wang [63] derived the Kerr-Newman-AdS solution in the presence of quintessence by using Newman-Janis algorithm and complex computations.

The line-element can be written as [63]

$$ds^2 = \frac{\Sigma^2}{\Delta_r} dr^2 + \frac{\Sigma^2}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta \sin^2 \theta}{\Sigma^2} \left(a \frac{dt}{\Xi} - (r^2 + a^2) \frac{d\phi}{\Xi} \right)^2 - \frac{\Delta_r}{\Sigma^2} \left(\frac{dt}{\Xi} - a \sin^2 \theta \frac{d\phi}{\Xi} \right)^2, \quad (3)$$

where

$$\begin{aligned} \Delta_r &= R^2 \left(1 - \frac{r^2}{\ell^2} \right) + Q^2 - 2Mr - \alpha r^{1-3\omega}, \\ \Delta_\theta &= 1 + \frac{a^2 \cos^2 \theta}{\ell^2}, \\ R^2 &= r^2 + a^2, \\ \ell^2 &= \frac{3}{\Lambda}, \\ \Xi &= 1 + \frac{a^2}{\ell^2}, \\ \Sigma^2 &= r^2 + a^2 \cos^2 \theta. \end{aligned} \quad (4)$$

In above expressions, M is BH mass and Q is BH charge defined as $Q^2 = q_e^2 + q_m^2$, while q_e^2 and q_m^2 are being electric and magnetic charge parameters, respectively, a is the rotation parameter, and ℓ is the curvature radius represented by cosmological constant $\Lambda < 0$, whereas α is the quintessence parameter and the state parameter ω ranges from $-1 < \omega < -1/3$.

For $\Lambda = 0$, the relationship between α and ω is defined as follows [63]:

$$\alpha \leq \frac{2}{(1-3\omega)} 8^\omega. \quad (5)$$

It is important to note that only for the fixed value of the state parameter $\omega = -2/3$, one can obtain four horizons; these are, inner, outer, and two cosmological horizons r_q (determined by the quintessence) and r_c (influenced by the cosmological constant). For $\omega = -2/3$, the above expression (5) implies $\alpha \leq 1/6$. For $\Lambda \neq 0$, four roots can be obtained by taking $\Delta_r = 0$. For $\omega = -2/3$, the horizon equation will become

$$r^4 + \frac{3\alpha}{\Lambda} r^3 + \left(a^2 - \frac{3}{\Lambda} \right) r^2 + \frac{6M}{\Lambda} r - \frac{3}{\Lambda} (a^2 + Q^2) = 0. \quad (6)$$

The above fourth-order algebraic equation can be expressed in terms of its roots, i.e.,

$$(r - r_{in})(r - r_{out})(r - r_q)(r - r_c) = 0, \quad (7)$$

where r_- is a cauchy (inner) horizon, r_+ is the event (outer) horizon, r_q and r_c are two cosmological horizons.

The line-element (3) can be expressed as

$$ds^2 = -f(r, \theta) dt^2 + \frac{1}{g(r, \theta)} dr^2 + \frac{1}{\rho(r, \theta)} d\theta^2 + K(r, \theta) d\phi^2 - 2H(r, \theta) dt d\phi, \quad (8)$$

where the metric functions are defined by

$$\begin{aligned} f(r, \theta) &= \frac{-a^2 \sin^2 \theta \Delta_\theta + \Delta_r}{\Sigma^2 \Xi^2}, \\ H(r, \theta) &= \frac{a \sin^2 \theta (-\Delta_r + R^2 \Delta_\theta)}{\Sigma^2 \Xi^2}, \\ K(r, \theta) &= \frac{\sin^2 \theta (\Delta_\theta R^4 - a^2 \sin^2 \theta \Delta_r)}{\Sigma^2 \Xi^2}, \end{aligned} \quad (9)$$

$$g(r, \theta) = \frac{\Delta_r}{\Sigma^2},$$

$$\rho(r, \theta) = \frac{\Delta_\theta}{\Sigma^2}.$$

The electromagnetic vector potential A_μ is given by [77]

$$A_\mu = \frac{1}{\Sigma^2 \Xi} \left[-(q_e r + a q_m \cos \theta) dt + (a \sin^2 \theta q_e r + q_m R^2 \cos \theta) d\phi \right]. \quad (10)$$

The angular velocity is defined as [78]

$$\Omega = -\frac{g_{t\phi}}{g_{\phi\phi}} = \frac{a(R^2 \Delta_\theta - \Delta_r)}{R^4 \Delta_\theta - a^2 \sin^2 \theta \Delta_r}; \quad (11)$$

at horizon it can be expressed as [56]

$$\Omega_H = \frac{H(r_+, \theta)}{K(r_+, \theta)} = \frac{a}{r_+^2 + a^2}. \quad (12)$$

The inverse of $f(r, \theta)$ can be written as [56]

$$\begin{aligned} F(r, \theta) &= f(r, \theta) + \frac{H^2(r, \theta)}{K(r, \theta)} \\ &= \frac{\Delta_r \Delta_\theta \Sigma^2(r, \theta)}{\Xi^2 (R^4 \Delta_\theta - a^2 \sin^2 \theta \Delta_r)}. \end{aligned} \quad (13)$$

3. Charged Fermions Tunneling

This section is devoted to investigate massive charged fermions tunneling phenomenon for Kerr-Newman-AdS BH surrounded by quintessence having electric and magnetic charges. For this purpose, we will consider the covariant Dirac equation, given as [56]

$$\iota\gamma^\mu \left(D_\mu - \frac{\iota q}{\hbar} A_\mu \right) \Psi + \frac{m}{\hbar} \Psi = 0, \quad \mu = 0, 1, 2, 3 \quad (14)$$

where q is electric charge and Ψ is wave function, while

$$D_\mu = \partial_\mu + \Omega_\mu, \quad \Omega_\mu = \frac{1}{2} \iota \Gamma_\mu^{\alpha\beta} \Sigma_{\alpha\beta}, \quad \Sigma_{\alpha\beta} = \frac{1}{4} \iota [\gamma^\alpha, \gamma^\beta]. \quad (15)$$

Here, γ^μ satisfies the following identities $[\gamma^\alpha, \gamma^\beta] = -[\gamma^\beta, \gamma^\alpha]$ for $\alpha \neq \beta$ and $[\gamma^\alpha, \gamma^\beta] = 0$ for $\alpha = \beta$. Using these relationships as well as the symmetric property of connection symbol $\Gamma_\mu^{\alpha\beta}$, we can obtain $\Omega_\mu = 0$, which yields $D_\mu = \partial_\mu$. Thus, the given (14) reduces to the following form [56, 79]:

$$\iota\gamma^\mu \left(\partial_\mu - \frac{\iota q}{\hbar} A_\mu \right) \Psi + \frac{m}{\hbar} \Psi = 0, \quad (16)$$

where Dirac matrices γ^μ are

$$\begin{aligned} \gamma^t &= \frac{1}{\sqrt{F(r, \theta)}} \gamma^0, \\ \gamma^r &= \sqrt{g(r, \theta)} \gamma^3, \\ \gamma^\theta &= \sqrt{\rho(r, \theta)} \gamma^1, \\ \gamma^\phi &= \frac{1}{\sqrt{K(r, \theta)}} \left(\gamma^2 + \frac{H(r, \theta)}{\sqrt{F(r, \theta) K(r, \theta)}} \gamma^0 \right), \end{aligned} \quad (17)$$

here γ^c 's are for chiral and γ^s are for Minkowski space, defined as [24]

$$\begin{aligned} \gamma^0 &= \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}, \\ \gamma^1 &= \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}, \\ \gamma^2 &= \begin{pmatrix} 0 & \sigma^2 \\ \sigma^2 & 0 \end{pmatrix}, \\ \gamma^3 &= \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix}. \end{aligned} \quad (18)$$

Here, Pauli sigma matrices σ^s are defined as

$$\begin{aligned} \sigma^1 &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ \sigma^2 &= \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \end{aligned}$$

$$\sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (19)$$

Spin particles have two types of spin states, spin-up and spin-down, and the corresponding wave functions are defined as, respectively,

$$\Psi_\uparrow(t, r, \theta, \phi) = \begin{bmatrix} A(t, r, \theta, \phi) \\ 0 \\ B(t, r, \theta, \phi) \\ 0 \end{bmatrix} \exp \left[\frac{i}{\hbar} I_\uparrow(t, r, \theta, \phi) \right], \quad (20)$$

$$\begin{aligned} \Psi_\downarrow(t, r, \theta, \phi) \\ = \begin{bmatrix} 0 \\ C(t, r, \theta, \phi) \\ 0 \\ D(t, r, \theta, \phi) \end{bmatrix} \exp \left[\frac{i}{\hbar} I_\downarrow(t, r, \theta, \phi) \right], \end{aligned} \quad (21)$$

where I_\uparrow (spin-up) and I_\downarrow (spin-down) represent particles action. Here, we discuss only the spin-up case; the spin-down case is similar. Moreover, the particles motion is considered in positive radial direction and the corresponding action can be considered as [56]

$$I_\uparrow = -Et + J\phi + R(r, \theta), \quad (22)$$

where E and J represent particles energy and angular momentum, respectively, while R is arbitrary function of r and θ . Using WKB approximation and the ansatz (22) for spin-up particles into the Dirac equation with $\iota A = B$ and $\iota B = A$, by applying the Taylor's expansion near the event horizon, we can obtain the set of equations, i.e.,

$$\begin{aligned} -B \left[\frac{-E + \Omega_H J - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)^2}{\sqrt{(r - r_+) F_r(r_+, \theta)}} \right. \\ \left. + \sqrt{(r - r_+) g_r(r_+, \theta)} R_r(r_+, \theta) \right] + mA = 0, \end{aligned} \quad (23)$$

$$\begin{aligned} -B \left[\frac{\iota}{\sqrt{K(r_+, \theta)}} \left(J + q \frac{a \sin^2 \theta q_e r_+ + R_+^2 q_m \cos \theta}{\Sigma^2(r_+, \theta)} \right) \right. \\ \left. + \sqrt{\rho(r_+, \theta)} R_\theta(r, \theta) \right] = 0, \end{aligned} \quad (24)$$

$$+ A \left[\frac{-E + \Omega_H J - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)}{\sqrt{(r - r_+) F_r(r_+, \theta)}} \right. \\ \left. + \sqrt{(r - r_+) g_r(r_+, \theta)} R_r(r_+, \theta) \right] + mB = 0, \quad (25)$$

$$- A \left[\frac{i}{\sqrt{K(r_+, \theta)}} \left(J + q \frac{a \sin^2 \theta q_e r_+ + R_+^2 q_m \cos \theta}{\Sigma^2(r_+, \theta)} \right) \right. \\ \left. + \sqrt{\rho(r_+, \theta)} R_\theta(r, \theta) \right] = 0. \quad (26)$$

In the above set of (23)-(26), the expressions can be defined as

$$F_r(r_+, \theta) = \frac{2 \left[r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2 \right] (r_+^2 + a^2 \cos^2 \theta)}{(r_+^2 + a^2)^2}, \quad (27)$$

$$g_r(r_+, \theta) = \frac{\Delta_r(r_+)}{\Sigma^2(r_+, \theta)} = \frac{2 \left[r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2 \right]}{(r_+^2 + a^2 \cos^2 \theta)}.$$

For $m = 0$, it is feasible to extract $1/\sqrt{\Sigma(r_+, \theta)}$ from (23) and (25), to make these equations independent of θ . Moreover, (24) and (26) have no definite r dependence. It is possible to separate the function R . So, near the horizon, the arbitrary function $R(r, \theta)$ can be separated as follows:

$$R(r, \theta) = R(r) + \Theta(\theta). \quad (28)$$

Equations (24) and (26) provide the same equation for Θ disregarding of A or B . For $m = 0$, (23) and (25) have two feasible solutions, given below:

$$R'_+(r_+) = R'_+(r_+) \\ = \left[\frac{\{E - \Omega_H J - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)\} (r_+^2 + a^2)}{\Delta_r(r_+) (r - r_+)} \right], \quad (29)$$

$$R'_-(r_+) = R'_-(r_+) \\ = - \left[\frac{\{E + \Omega_H J - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)\} (r_+^2 + a^2)}{\Delta_r(r_+) (r - r_+)} \right], \quad (30)$$

where

$$\Delta_r(r_+) \\ = 2 \left[r_+ \left(1 - \frac{a^2}{\ell^2} \right) - \frac{2r_+^3}{\ell^2} - M - \frac{\alpha (1 - 3\omega) r_+^{-3\omega}}{2} \right], \quad (31)$$

R_+ and R_- represent the radial solution of outgoing and incoming particles action, respectively. Solving for $R_+(r_+)$, (29) implies

$$R_+(r_+) \\ = \int \left[\frac{\{E - \Omega_H J - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)\} (r_+^2 + a^2)}{\Delta_r(r_+) (r - r_+)} \right] dr. \quad (32)$$

After integrating the above expression at the pole, we get

$$R_+(r_+) = \pm i\pi \\ \cdot \frac{\{E - \Omega_H J - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)\} (r_+^2 + a^2)}{2 \left[r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2 \right]}. \quad (33)$$

The above equation implies that

$$\text{Im } R_+ = \pm \pi \\ \cdot \frac{\{E - \Omega_H J - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)\} (r_+^2 + a^2)}{2 \left[r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2 \right]}. \quad (34)$$

The tunneling probability in terms of spatial ($\oint p_r dr$) and temporal contribution ($\check{E}\Delta t^{out, in}$) is given as follows [80–86]:

$$\Gamma = \exp \left[\frac{1}{\hbar} \left(\text{Im} (\check{E}\Delta t^{out}) + \text{Im} (\check{E}\Delta t^{in}) \right) \right. \\ \left. - \text{Im} \oint p_r dr \right], \quad (35)$$

where $\oint p_r dr = \int p_r^+ dr - \int p_r^- dr$, whereas $p_r^\pm = \pm \partial_r R$. The spatial contribution can be calculated as

$$\Gamma_{spatial} \propto \exp \left[-\frac{1}{\hbar} \text{Im} \oint p_r dr \right] = \exp \left[-\frac{1}{\hbar} \text{Im} \left(\int p_r^+ dr - \int p_r^- dr \right) \right] \\ = \exp \left[-\frac{\pi}{\hbar} \left(\frac{\check{E} (r_+^2 + a^2)}{r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2} \right) \right]. \quad (36)$$

The connection between interior and exterior regions of a BH defines the temporal part. For $t \rightarrow t - i\pi/(2\kappa)$, we can define the temporal contribution as

$$\text{Im}(\check{E}\Delta t^{out,in}) = \frac{-\check{E}\pi}{(2\kappa)}, \quad (37)$$

where

$$\check{E} = \left[E - \Omega_H j - q \left(\frac{q_e r + a q_m}{(r_+^2 + a^2)} \right) \right] \quad (38)$$

and

$$\kappa = \left[\frac{r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha(1 - 3\omega) r_+^{-3\omega}/2}{(r_+^2 + a^2)} \right]. \quad (39)$$

Thus, the total temporal rate can be obtained as

$$\begin{aligned} \Gamma_{temp.} &\propto \exp \left[\frac{1}{\hbar} (\text{Im}(\check{E}\Delta t^{out}) + \text{Im}(\check{E}\Delta t^{in})) \right], \\ &= \exp \left[-\frac{\pi}{\hbar} \left(\frac{\check{E} (r_+^2 + a^2)}{r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha(1 - 3\omega) r_+^{-3\omega}/2} \right) \right]. \end{aligned} \quad (40)$$

Using (35), the total tunneling rate at horizon $r = r_+$ for $\hbar = 1$ is derived as

$$\Gamma = \exp \left[\frac{-2\pi \left[E - \Omega_H J - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right) \right] (r_+^2 + a^2)}{r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha(1 - 3\omega) r_+^{-3\omega}/2} \right]. \quad (41)$$

Thus, by utilizing Boltzmann equation

$$\Gamma_B = \exp \left[\frac{E - j\Omega_H - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)}{T_H} \right] \quad (42)$$

the Hawking temperature T_H at the horizon r_+ can be obtained as

$$\begin{aligned} T_H &= \left[\frac{r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha(1 - 3\omega) r_+^{-3\omega}/2}{2\pi (r_+^2 + a^2)} \right]. \end{aligned} \quad (43)$$

The above Hawking temperature for the massive particles case is similar as for massless case for Kerr-Newman-AdS BH with quintessence. Moreover, the results for spin-down particles are similar as for spin-up case with the change of sign. The Hawking temperature is acquired for both cases and we conclude that both spin-up and spin-down particles are radiate with alike rate.

4. Quantum Corrections of T_H

In this section, we analyze Hawking temperature for massive charged fermions by considering tunneling procedure

incorporating quantum gravitational effects. The modified form of Dirac equation (14) is given as follows [87]:

$$\begin{aligned} &-\gamma^0 \left(i\partial_0 + \frac{q}{\hbar} A_0 \right) \Psi \\ &= \left(i\gamma^i \partial_i + q \frac{\gamma^i}{\hbar} A_i + \frac{m}{\hbar} \right) (1 + \beta \hbar^2 \partial_j \partial^j - \beta m^2) \Psi, \end{aligned} \quad (44)$$

where $i = 1, 2, 3$ signifies the spatial coordinates. Moreover, the correction parameter β for the minimal length M_f is defined as $\beta = \beta_0/M_f^2$ and m is the mass of fermion particles. Equation (44) can be rewritten as

$$\begin{aligned} &\left[i\gamma^0 \partial_0 + q \frac{\gamma^0}{\hbar} A_0 + i\gamma^i \partial_i (1 - \beta m^2) + i\gamma^i \beta \hbar^2 (\partial_j \partial^j) \partial_i \right. \\ &+ q \frac{\gamma^i}{\hbar} A_i (1 + \beta \hbar^2 \partial_j \partial^j - \beta m^2) \\ &\left. + \frac{m}{\hbar} (1 + \beta \hbar^2 \partial_j \partial^j - \beta m^2) \right] \Psi = 0. \end{aligned} \quad (45)$$

Using coordinate transformation, $\varphi = \phi - \Omega t$, where

$$\Omega = \frac{a (\Delta_\theta (r^2 + a^2) - \Delta_r)}{(r^2 + a^2)^2 \Delta_\theta - \Delta_r a^2 \sin^2 \theta}, \quad (46)$$

the line-element (3) reduces to the following form:

$$ds^2 = -\frac{\Delta_r \Sigma^2}{(r^2 + a^2)^2 \Delta_\theta - \Delta_r a^2 \sin^2 \theta} dt^2 + \frac{\Sigma^2}{\Delta_r} dr^2 + \frac{\Sigma^2}{\Delta_\theta} d\theta^2 + \left(\frac{(r^2 + a^2)^2 \Delta_\theta - \Delta_r a^2 \sin^2 \theta}{\Sigma^2} \right) \sin^2 \theta d\varphi^2, \quad (47)$$

which can also be expressed as

$$ds^2 = -F(r, \theta) dt^2 + \frac{1}{G(r, \theta)} dr^2 + \frac{1}{K(r, \theta)} d\theta^2 + H(r, \theta) d\varphi^2. \quad (48)$$

We can choose γ matrices in the following form:

$$\begin{aligned} \gamma^t &= \frac{1}{\sqrt{F(r, \theta)}} \begin{pmatrix} \iota & 0 \\ 0 & -\iota \end{pmatrix}, \\ \gamma^r &= \sqrt{g(r, \theta)} \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix}, \\ \gamma^\theta &= \sqrt{K(r, \theta)} \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}, \\ \gamma^\varphi &= \frac{1}{\sqrt{H(r, \theta)}} \begin{pmatrix} 0 & \sigma^2 \\ \sigma^2 & 0 \end{pmatrix}, \end{aligned} \quad (49)$$

where σ 's are defined by (19). For sake of simplicity, here we discuss the case of spin-up, following the outward radial trajectory. The wave function of spin-up particles is defined as

$$\Psi_\uparrow(t, r, \theta, \varphi) = \begin{bmatrix} A(t, r, \theta, \varphi) \\ 0 \\ B(t, r, \theta, \varphi) \\ 0 \end{bmatrix} \exp \left[\frac{i}{\hbar} I_\uparrow(t, r, \theta, \varphi) \right]. \quad (50)$$

The terms \hbar , β , ∂A , and ∂B are considered only for the first order; i.e., the higher order contributions are ignored. Using WKB approximation and (50) and (49) in (45), we can obtain the following set of equations:

$$\begin{aligned} & -\frac{\iota A}{\sqrt{F(r, \theta)}} (\partial_t I) + q \frac{\iota A}{\sqrt{F(r, \theta)}} A_t - \sqrt{G(r, \theta)} (1 - \beta m^2) (\partial_r I) + (1 - \beta m^2) A + B\beta \sqrt{G(r, \theta)} (\partial_r I) \\ & \cdot \left[G(r, \theta) (\partial_r I)^2 + K(r, \theta) (\partial_\theta I)^2 + \frac{1}{H(r, \theta)} (\partial_\varphi I)^2 \right] - Am\beta \left[G(r, \theta) (\partial_r I)^2 + K(r, \theta) (\partial_\theta I)^2 + \frac{1}{H(r, \theta)} (\partial_\varphi I)^2 \right] = 0, \end{aligned} \quad (51)$$

$$\begin{aligned} & \frac{\iota B}{\sqrt{F(r, \theta)}} (\partial_t I) - \frac{\iota B}{\sqrt{F(r, \theta)}} A_t - A \sqrt{G(r, \theta)} (1 - \beta m^2) (\partial_r I) + A\beta \sqrt{G(r, \theta)} (\partial_r I) \left[G(r, \theta) (\partial_r I)^2 + K(r, \theta) (\partial_\theta I)^2 + \frac{1}{H(r, \theta)} (\partial_\varphi I)^2 \right] + Bm (1 - \beta m^2) - mB\beta \left[G(r, \theta) (\partial_r I)^2 + K(r, \theta) (\partial_\theta I)^2 + \frac{1}{H(r, \theta)} (\partial_\varphi I)^2 \right] = 0, \end{aligned} \quad (52)$$

$$\begin{aligned} & B \left[-\sqrt{K(r, \theta) (1 - \beta m^2)} (\partial_\theta I) - \frac{\iota (1 - \beta m^2)}{\sqrt{H(r, \theta)}} (\partial_\varphi I) + \beta (\partial_\theta I) \sqrt{K(r, \theta)} \left\{ G(r, \theta) (\partial_r I)^2 + K(r, \theta) (\partial_\theta I)^2 + \frac{1}{H(r, \theta)} (\partial_\varphi I)^2 \right\} + \frac{\iota \beta}{\sqrt{H(r, \theta)}} (\partial_\varphi I) \cdot \left\{ G(r, \theta) (\partial_r I)^2 + K(r, \theta) (\partial_\theta I)^2 + \frac{1}{H(r, \theta)} (\partial_\varphi I)^2 \right\} \right] = 0, \end{aligned} \quad (53)$$

$$\begin{aligned} & A \left[-\sqrt{K(r, \theta) (1 - \beta m^2)} (\partial_\theta I) - \frac{\iota (1 - \beta m^2)}{\sqrt{H(r, \theta)}} (\partial_\varphi I) + \beta (\partial_\theta I) \sqrt{K(r, \theta)} \left\{ G(r, \theta) (\partial_r I)^2 + K(r, \theta) (\partial_\theta I)^2 + \frac{1}{H(r, \theta)} (\partial_\varphi I)^2 \right\} + \frac{\iota B}{\sqrt{H(r, \theta)}} (\partial_\varphi I) \cdot \left\{ G(r, \theta) (\partial_r I)^2 + K(r, \theta) (\partial_\theta I)^2 + \frac{1}{H(r, \theta)} (\partial_\varphi I)^2 \right\} \right] = 0. \end{aligned} \quad (54)$$

The particle's action is given by

$$I = -(E - \Omega j) t + R(r) + \Theta(\theta, \varphi). \quad (55)$$

Using (55) in (51)-(54) and by focusing on (53) and (54), we observe that they are similar after dividing by A and B and can be expressed as

$$\begin{aligned} & \left\{ \beta G_r(r_+, \theta) R'^2 + K(r_+, \theta) \beta J_\theta^2 + \beta H(r_+, \theta) J_\varphi^2 - (1 - \beta m^2) \right\} \times \left[\sqrt{K(r_+, \theta)} J_\theta + \iota \frac{1}{\sqrt{H(r_+, \theta)}} J_\varphi \right] = 0, \end{aligned} \quad (56)$$

where $R' = \partial_r R$, $J_\theta = \partial_\theta \Theta$, and $\partial_\varphi = \partial_\varphi \Theta$. In (56), β indicates the quantum gravity effects so it cannot be considered as zero;

thus the term in large bracket equals zero and provides the solution for Θ . Thus, we can write

$$\left[\sqrt{K(r_+, \theta)} J_\theta + i \frac{1}{\sqrt{H(r_+, \theta)}} J_\varphi \right] = 0. \quad (57)$$

After removing A and B from (51) and (52), we get the following expression:

$$P_6 (\partial_r R)^6 + P_4 (\partial_r R)^4 + P_2 (\partial_r R)^2 + P_0 = 0, \quad (58)$$

where

$$\begin{aligned} P_6 &= \beta^2 G^3 F, \\ P_4 &= \beta G^2 F (m^2 \beta + 2\beta C - 2), \\ P_2 &= GF \left[(1 - \beta m^2)^2 + \beta (2m^2 - 2m^4 \beta - 2C + \beta C^2) \right], \\ P_0 &= m^2 (1 - \beta m^2 - \beta C)^2 F - (E - \Omega j + q A_t)^2, \\ C &= K(r_+, \theta) J_\theta^2 + \frac{1}{H(r_+, \theta)} J_\varphi^2. \end{aligned} \quad (59)$$

From (57), we note that $C = 0$. Considering β only for the first order, the solution of (58) at horizon provides [80, 88, 89]

$$\begin{aligned} R(r) &= \pm \int \frac{1}{\sqrt{GF}} \sqrt{m^2 (1 - 2\beta m^2) F + (E - \Omega j - q A_t)^2} \\ &\times \left[1 + \beta \left(m^2 + \frac{(E - \Omega j - q A_t)^2}{F} \right) \right] dr. \end{aligned} \quad (60)$$

The above equation implies

$$R(r) = \pm i\pi \frac{(r_+^2 + a^2) (E - \Omega_H j - q A_t)}{\Delta_r(r_+)} (1 + \beta \Xi), \quad (61)$$

where

$$\Xi = 6m^2 + \frac{6}{r^2 + a^2 \cos^2 \theta} (J_\theta^2 + J_\varphi^2 \csc^2 \theta). \quad (62)$$

The positive/negative signs indicate outgoing/incoming particles. Hence, the tunneling rate [90] of charged fermions across the horizon is calculated as

$$\begin{aligned} \Gamma &= \frac{\Gamma_{(emission)}}{\Gamma_{(absorption)}} = \frac{\exp[-(2/\hbar)(\text{Im } R_+ + \text{Im } \Theta)]}{\exp[-(2/\hbar)(\text{Im } R_- + \text{Im } \Theta)]} = \exp\left[-\frac{4}{\hbar} \text{Im } R_+\right], \\ &= \exp\left[\frac{-2\pi \left\{ E - \Omega_H j - q \left((q_e r + a q_m) / (r_+^2 + a^2) \right)^2 \right\} (r_+^2 + a^2)}{r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2} \right] (1 + \beta \Xi). \end{aligned} \quad (63)$$

For $\hbar = 1$, the corrected Hawking temperature T_{e-H} is

$$T_{e-H} = \left[\frac{r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2}{2\pi (r_+^2 + a^2)} \right] \quad (64)$$

$$\cdot (1 + \beta \Xi)^{-1},$$

$$= T_0 [1 - \beta \Xi + (\beta \Xi)^2 + \dots], \quad (65)$$

$$\approx T_0 (1 - \beta \Xi). \quad (66)$$

where the semiclassical Hawking temperature is

$$T_0 = \left[\frac{r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2}{2\pi (r_+^2 + a^2)} \right]. \quad (67)$$

The corrected temperature is based on quantum numbers, i.e., mass, energy, and angular momentum.

5. Logarithmic Entropy Corrections

This section is devoted to calculate the entropy corrections for Kerr-Newman-AdS BH with quintessence. Using null geodesic technique, Banerjee and Majhi [38–42] investigated the corrected Hawking temperature and corrected entropy by taking into account the back-reaction effects. Majhi [44, 45] also analyzed the corrected temperature and entropy by using the first and second laws of thermodynamics. In our investigation, we calculate the entropy corrections for Kerr-Newman-AdS BH involving quintessence by considering the generic formula for leading order corrections to Bekenstein-Hawking formula [91]. It is worth mentioning here that one could calculate the logarithmic corrections to the BH entropy at inner/outer horizon (r_\pm) without knowing the values of any specific heat of the BH but only knowing the values of Hawking temperature ($T_{e-H,\pm}$) and entropy ($S_{0,\pm}$), for the given BH. The BH entropy corrections (S_\pm) can be defined as

$$S_\pm = S_{0,\pm} - \frac{1}{2} \ln |T_{e-H,\pm}^2 S_{0,\pm}| + \dots \quad (68)$$




FIGURE 1: For $\alpha = 0.01$, Hawking temperature T_{e-H} with respect to horizon r_+ .

The classical entropy for given BH at $r = r_+$ can be calculated as follows:

$$S_{0,+} = \frac{\mathcal{A}_+}{4} = \frac{\pi (r_+^2 + a^2)}{(1 - a^2/\ell^2)}, \quad (69)$$

where

$$\mathcal{A}_+ = \int_0^{2\pi} \int_0^\pi \sqrt{g_{\theta\theta} g_{\phi\phi}} d\theta d\phi. \quad (70)$$

After substituting the corrected Hawking temperature T_{e-H} (given by (66)) in the above (68), we can obtain the logarithmic corrections of entropy in the following form:

$$S = S_{0,+} - \frac{1}{2} \cdot \ln \left| \frac{(r_+ (1 - a^2/\ell^2) - 2r_+^3/\ell^2 - M - \alpha (1 - 3\omega) r_+^{-3\omega}/2)^2}{4\pi (r_+^2 + a^2) (1 - a^2/\ell^2)} \right| - \beta \Xi \Big| + \dots \quad (71)$$

This expression represents the corrected entropy for Kerr-Newman-AdS BH involving quintessence parameter.

6. Effects of β on T_{e-H}

In this section, we analyze graphically the effects of β on T_{e-H} with respect to various parameters.

6.1. Temperature T_{e-H} with Horizon r_+ . This subsection is devoted to discuss the behavior of corrected Hawking temperature in different domains of event Horizon r_+ with fixed BH mass $M = 1$, state parameter $\omega = -2/3$, and $\Xi = 0.01$.



(i) Figure 1 indicates the behavior of Hawking temperature with $r_+ > 0$, for different values of cosmological

constant $\Lambda < 0$ with fixed values of rotation parameter $a = 0.3$ and correction parameter $\beta = 1$. The cosmic repulsion indicates that the recent value of the cosmological constant is $\Lambda \approx -1.3 \times 10^{-56} \text{cm}^{-2}$ [63]. In our analysis, we consider this fixed value of $\Lambda = -1.3 \times 10^{-56} \text{cm}^{-2}$, and we consider Λ greater and lesser than this fixed value. (a) For different values of Λ near to the fixed value, we can observe that the temperature has its maximum value and its behavior is linear. As horizon increases the temperature also increases. The change in Λ defines the diverging temperature T_{e-H} as r_+ increases. (b) It indicates the behavior of temperature for fixed values of $a = 0.3$ and $\Lambda = -1.3 \times 10^{-56}$, while the value of β varies. We can observe that, for $\beta < 100$, the temperature is going to increase as β going lesser from 100, while for $\beta = 100$, the temperature will be zero. The temperature has linear behavior, it increases as horizon increases.

(ii) Figure 2 indicates the behavior of T_{e-H} for fixed values of a and Λ , while for varying β . (a): We can observe that for fixed value of $a = 0.5$, $\Lambda = -1$ and for varying $\beta < 100$, the behavior of temperature is positively increasing $T_{e-H} > 0$, while for $\beta = 100$, the graph shows zero temperature, i.e., $T_{e-H} = 0$. (b): for fixed $a = 0.1$ and $\Lambda = -1$ and for varying $\beta > 100$, we observe the negatively divergent behavior of temperature; i.e., the temperature is decreasing as horizon increases in the negative range. This negative and divergent behavior of temperature reflects the nonphysical unstable state of BH.

It is also worth mentioning here that, for the values of correction parameter $1 \leq \beta < 100$, we observe positive values of temperature and for $\beta = 100$, the temperature vanishes, while for $\beta > 100$, we observe nonphysical behavior with negative temperature.



(iii) Figure 3 shows the behavior of temperature for fixed values of $a = 0.3$ and varying Λ and β . (a):

FIGURE 2: For $\alpha = 1$, relation between the temperature and r_+ .FIGURE 3: For $\alpha = 1$, relation between the temperature and r_+ .

We can observe that for $a = 0.3$ and $\beta = 1$, as Λ decreases, the temperature will be small and after attaining its maximum value, the temperature decreases. For $\Lambda \ll 0$, the temperature decreases as horizon increases. In all curves, the temperature has its maximum value at small horizon. The horizon of BH can never be zero, the nonzero value of horizon leads to a BH remnant with maximum temperature. (b): We observe that for $a = 0.3$, $\Lambda = -200$ and for varying $\beta < 100$, the temperature attains positive values, Initially, the temperature will be maximum at nonzero horizon and later decreases exponentially as horizon increases, which indicates a physical stable state of BH.

6.2. Temperature T_{e-H} with Quintessence α . This subsection gives the analysis of corrected Hawking temperature T_{e-H} with quintessence parameter α for different values of rotation parameters a , BH charge, Q and cosmological constant Λ for fixed $M = 1$, $\omega = -2/3$, and $\Xi = 0.01$.

- (i) Figure 4 shows the behavior of temperature for fixed a , Q , and Λ and for varying β . (a) We can observe that, for fixed values of $a = 0.1$, $Q = 0.3$, and $\Lambda = -10^{-5}$ and for different values of β , the temperature gradually increases as β decreases. It is to be noted that, for these parameters, the temperature increases with increase in α as $\beta \rightarrow 1$. (b) This graph indicates the behavior of temperature for fixed values of $a = 1$, $Q = 0.1$, and $\Lambda = -10^{-7}$ and for varying β . It is to be noted that, for $\beta < 100$, the behavior of temperature is from negative to positive, while it attains maximum values and temperature increases with increase in α . While, for $\beta = 100$, the temperature vanishes, i.e., $T_{e-H} = 0$. It is important to note that the both negative and positive behaviors of temperature shows that the initial unstable behaviors of temperature shows that the initial unstable state of BH, which turns out to be stable with time. This negative temperature is the effect of rotation parameter $a = 1$ (maximum value).
- (ii) Figure 5 shows the behavior of temperature for fixed Λ and β , while for varying a (in Figure (a)) and varying




 FIGURE 4: Relation between temperature and α in terms of β .

 FIGURE 5: Relation between temperature and α in terms of a and Q .

Q (in Figure (b)). (a) We can observe that, for fixed values of $Q = 0.3, \beta = 1$, and $\Lambda = -10^{-5}$ and for varying a , the temperature gradually increases for increasing α . It is to be noted that as $a \rightarrow 1$, the temperature will go on from negative to positive. (b) We can observe that, for fixed values of $a = 0.3, \beta = 1$, and $\Lambda = -10^{-7}$ and for varying Q , the temperature will be high enough and it will gradually increase with increase in α .

6.3. 3D Plots: T_{e-H} and r_+ with Λ, α , and β . This section is based on the analysis of Hawking temperature with quintessence parameter α , horizon r_+ , and cosmological constant Λ for fixed $M = 1, \omega = -2/3$, and $\Xi = 0.01$. This section consists of 3D graphs.


(i) Figure 6 shows the behavior of temperature for fixed a, Λ and α , while for $\beta < 100$. (a) Here, for fixed

$a = 0.3, \Lambda = -100$ and $\alpha = 0.16$, the behavior of T_{e-H} indicates that the temperature attains negative values and its behavior is constant w.r.t. β . After attaining its maximum value at nonzero horizon, the Hawking temperature decreases as horizon increases. This indicates that there exists a BH remnant during the evaporation process. The BH temperature attains its maximum value as horizon shrinks. For $\Lambda = -100$, the temperature is negative for $\beta < 100$ and invert for other β (already mentioned in 2D graphs). (b) Graph indicates the behavior of temperature for fixed $a = 0.3, \Lambda = -26$, and $\alpha = 0.16$. We can observe that the behavior of temperature is same as in (a). For $\Lambda = -100$, the temperature is negative but as we increase the value of Λ till $\Lambda = -26$, the graph will exhibit the behavior of temperature from negative to positive and, for $\Lambda = -25$, the graph will exhibit the positive temperature.

FIGURE 6: Relation between temperature with r_+ and β .FIGURE 7: Relation between temperature with r_+ and α .

(ii) Figure 7 shows the behavior of T_{e-H} with r_+ and α for fixed a , β , and Λ . (a) Graph indicates the behavior of T_{e-H} for fixed $a = 0.1$, $\beta = 1$, and $\Lambda = -10^{-7}$. We can observe that, for larger value of Λ , the behavior of temperature is linear w.r.t. r_+ . The temperature attains high values and increases with increase in α . (b) This figure indicates the behavior of temperature for fixed $a = 0.5$, $\beta = 200$, and $\Lambda = -1$. We can observe that as horizon increases the temperature decreases, while for α , the temperature increases as α increases. In Figure 7, the behavior of temperature is same as proved in 2D graphs.

(iii) Figure 8 shows the behavior of temperature with horizon r_+ and cosmological constant Λ for fixed $a = 0.5$, $\beta = 100$, and $\alpha = 0.16$. We can observe that, for $\beta = 100$, the effect of temperature will be zero, as concluded in 2D graphs.

FIGURE 8: Relation between temperature with r_+ and Λ .

behavior is linear w.r.t. horizon. Moreover, for large Λ , the temperature increases with increasing α .

7. Conclusion

In this paper, we have computed radiation spectrum by analyzing Hawking temperature for Kerr-Newman-AdS BH surrounded by quintessence. For this purpose, we have

It is worth mentioning here that, from 2D and 3D graphs, we conclude that the temperature increases with decreasing horizon and increasing α . For small values of $\Lambda \neq -100$ and $\beta > 100$, the temperature shows negative values. For larger values of Λ , the temperature will be maximum and its

utilized the WKB approximation and the Hamilton-Jacobi ansatz for massive charged spin-1/2 particles (fermions). The investigation yields the corrected Hawking temperature T_{e-H} , reliable with BH universality. In our analysis, we have altered the Dirac equation in curved spacetime by incorporating quantum gravity effects through GUP. We have evaluated the tunneling rate at horizon as well as the corresponding Hawking temperature, quantum corrected Hawking temperature, and quantum corrected entropy. We have analyzed in detail quantum corrected Hawking temperature graphically.

We have summarized the detailed graphical analysis of this paper in the following points:

- (i) The derived Hawking temperature and its modified form T_{e-H} depend on BH's mass, charge, and rotation parameters, as well as on the mass and angular momentum of the emitted fermion particles, quintessence parameter α , and state parameter ω .
- (ii) When the quantum gravity effects are neglected ($\beta = 0$), we have recovered the Hawking temperature of Kerr Newman AdS BH with quintessence. For $\alpha = 0$ and $\omega = 0$, we have obtained the Hawking temperature of Kerr-Newman AdS BH. Moreover, when $\Lambda = 0$, the temperature of Kerr Newman BH has obtained. In addition, for charge-free ($Q = 0$) and nonrotating ($a = 0$) case, the temperature and its correction reduce to the case of Schwarzschild BH.
- (iii) In our analysis, we have considered $\Xi = 0.01$; then the condition of GUP must be satisfied for $0 < \beta < 100$. We have substituted the above mentioned values in (65) for positive values of T_e ; the correction terms became smaller than the previous terms given in series. When we consider the first order quantum corrections, the correction term is smaller than T_e . For $\beta = 100$, the first-order correction term is same as semiclassical term T_e , showing invalidity of GUP. When $\beta > 100$, the correction term became greater than the preceding term and the condition of GUP is not satisfied.
- (iv) Graphical analysis showed that the behavior of T_{e-H} is positive and negative when $\beta < 100$ and $\beta > 100$, respectively, while for $\beta = 100$, the temperature vanishes.
- (v) For fixed $\Lambda \approx -1.3 \times 10^{-56}$ [63], as well as greater and lesser than this fixed value of Λ , we have observed that the temperature increases as horizon increases, which is nonphysical. The negative as well as divergent behavior of Hawking temperature indicates the behavior reverse to Hawking's phenomenon, representing unstable state of BH, while for smaller $\Lambda \ll 0$, the temperature decreases with increasing horizon, which is physical.
- (vi) For smaller Λ , we have obtained physical (stable, +ive T_{e-H}) and nonphysical (-ive T_{e-H}) behavior of temperature for $\beta < 100$ and $\beta > 100$, respectively.
- (vii) We have observed the behavior of T_{e-H} w.r.t. α only for the particular ranges, i.e., $0 < \alpha < 1/6$, $0 < Q \leq 1$, $\Lambda \leq -1$, and $0 < a \leq 1$.

- (viii) For T_{e-H} w.r.t. α , at maximum value of the rotation parameter (i.e., $a = 1$), we have observed nonphysical and unstable state of BH, which is due to the instability in temperature.
- (ix) The results obtained from 3D graphs are similar to the results obtained from 2D graphs.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest




The authors declare that they have no conflicts of interest.

References

- [1] S. W. Hawking, "Particle creation by black holes," *Communications in Mathematical Physics*, vol. 43, no. 3, pp. 199–220, 1975.
- [2] S. W. Hawking, "Erratum: Particle creation by black holes," *Communications in Mathematical Physics*, vol. 46, p. 206, 1976.
- [3] R. Charlie, *A conservation with Dr. Stephen Hawking and Lucy Hawking*, Archived from the original on March 29, 2013.
- [4] S. Hawking, *A Brief History of Time*, Bantam Dell, UK, 1988.
- [5] J. D. Bekenstein, "Black holes and entropy," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 7, pp. 2333–2346, 1973.
- [6] M. K. Parikh and F. Wilczek, "Hawking radiation as tunneling," *Physical Review Letters*, vol. 85, no. 24, pp. 5042–5045, 2000.
- [7] M. K. Parikh, "New coordinates for de Sitter space and de Sitter radiation," *Physics Letters B*, vol. B546, p. 189, 2002.
- [8] M. K. Parikh, "A secret tunnel through the horizon," *International Journal of Modern Physics*, vol. D13, p. 2351, 2004.
- [9] P. Kraus and F. Wilczek, "Self-interaction correction to black hole radiance," *Nuclear Physics B*, vol. 433, no. 2, pp. 403–420, 1995.
- [10] P. Kraus and F. Wilczek, "Effect of self-interaction on charged black hole radiance," *Nuclear Physics B*, vol. B437, p. 231, 1995.
- [11] P. Kraus and K. E. Vakkuri, "Microcanonical D-branes and back reaction," *Nuclear Physics B*, vol. 491, p. 249, 1997.
- [12] M. Angheben, M. Nadalini, L. Vanzo, and S. Zerbini, "Hawking radiation as tunneling for extremal and rotating black holes," *Journal of High Energy Physics*, vol. 2005, no. 5, article 014, 2005.
- [13] K. Srinivasan and T. Padmanabhan, "Particle production and complex path analysis," *Physical Review D*, vol. 60, p. 24007, 1999.
- [14] S. Shankaranarayanan, K. Srinivasan, and T. Padmanabhan, "Method of complex paths and general covariance of Hawking radiation," *Modern Physics Letters A*, vol. 16, no. 9, pp. 571–578, 2001.
- [15] S. Shankaranarayanan, T. Padmanabhan, and K. Srinivasan, "Hawking radiation in different coordinate settings: complex paths approach," *Classical and Quantum Gravity*, vol. 19, no. 10, pp. 2671–2687, 2002.
- [16] S. Shankaranarayanan, "Temperature and entropy of Schwarzschild–de Sitter space-time," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, p. 084026, 2003.

- [17] R. Kerner and R. B. Mann, "Fermions tunnelling from black holes," *Classical and Quantum Gravity*, vol. 25, no. 9, Article ID 095014, 2008.
- [18] R. Kerner and R. B. Mann, "Charged Fermions Tunnelling from Kerr-Newman Black Holes," *Physics Letters B*, p. 277, 2008.
- [19] R. Kerner and R. B. Mann, "Fermions Tunnelling from Black Holes," <https://arxiv.org/abs/0710.0612v1>.
- [20] K. Nozari and S. H. Mehdipour, "Parikh-Wilczek Tunneling from Noncommutative Higher Dimensional Black Holes," *Journal of High Energy Physics*, vol. 0903, p. 061, 2009.
- [21] A. F. Ali, "No existence of black holes at LHC due to minimal length in quantum gravity," *Journal of High Energy Physics*, p. 067, 1209.
- [22] B. Majumder, "Quantum black hole and the modified uncertainty principle," *Physics Letters B*, vol. B701, p. 384, 2011.
- [23] K. Nozari and S. Saghafi, "Natural cutoffs and quantum tunneling from black hole horizon," *Journal of High Energy Physics*, vol. 11, p. 005, 2005.
- [24] R. Kerner and R. B. Mann, "Charged fermions tunnelling from Kerr-Newman black holes," *Physics Letters B*, vol. B665, p. 277, 2008.
- [25] R. Li, J.-R. Ren, and S.-W. Wei, "Hawking radiation of Dirac particles via tunneling from the Kerr black hole," *Classical and Quantum Gravity*, vol. 25, no. 12, Article ID 125016, 2008.
- [26] T. Jian and B. B. Chen, "Fermions tunneling from Kerr and Kerr-Newman black holes," *Acta Physica Polonica*, vol. B40, p. 241, 2009.
- [27] B. Majumder, "Black hole entropy with minimal length in tunneling formalism," *General Relativity and Gravitation*, vol. 45, no. 11, pp. 2403-2414, 2013.
- [28] D. Y. Chen, H. W. Wu, and H. T. Yang, "Fermion's Tunnelling with Effects of Quantum Gravity," *Advances in High Energy Physics*, vol. 2013, Article ID 432412, 6 pages, 2013.
- [29] D. Y. Chen, Q. Q. Jiang, P. Wang, and H. T. Yang, "Remnants, Fermions' Tunnelling and Effects of Quantum Gravity," *Journal of High Energy Physics*, vol. 11, p. 176, 2013.
- [30] D. Y. Chen, H. W. Wu, and H. T. Yang, "Observing remnants by fermions' tunneling," *Journal of Cosmology and Astroparticle Physics*, vol. 03, p. 036, 2014.
- [31] P. Bargueno and E. C. Vagenas, "Semiclassical corrections to black hole entropy and the generalized uncertainty principle," *Physics Letters B*, vol. B742, p. 15, 2015.
- [32] B. R. Mu, P. Wang, and H. T. Yang, "Minimal Length Effects on Tunnelling from Spherically Symmetric Black Holes," *Advances in High Energy Physics*, vol. 2015, Article ID 898916, 8 pages, 2015.
- [33] P. Wang, H. Yang, and S. Ying, "Quantum gravity corrections to the tunneling radiation of scalar particles," *International Journal of Theoretical Physics*, vol. 55, no. 5, pp. 2633-2642, 2016.
- [34] M. A. Anacleto, F. A. Brito, and E. Passos, "Quantum-corrected self-dual black hole entropy in tunneling formalism with GUP," *Physics Letters B*, vol. 749, pp. 181-186, 2015.
- [35] Z. W. Feng, H. L. Li, X. T. Zu, and S. Z. Yang, "Quantum corrections to the thermodynamics of Schwarzschild-Tangherlini black hole and the generalized uncertainty principle," *The European Physical Journal C*, vol. 76, p. 212, 2016.
- [36] D. Singleton, E. C. Vagenas, T. Zhu, and J. R. Ren, "Insights and possible resolution to the information loss paradox via the tunneling picture," *Journal of High Energy Physics*, vol. 1008, p. 089, 2010.
- [37] D. Singleton, E. C. Vagenas, and T. Zhu, "Self-similarity, conservation of entropy/bits and the black hole information puzzle," *Journal of High Energy Physics*, vol. 1405, p. 074, 2014.
- [38] R. Banerjee and B. R. Majhi, "Quantum tunneling and back reaction," *Physics Letters B*, vol. 662, no. 1, pp. 62-65, 2008.
- [39] R. Banerjee and B. R. Majhi, "Quantum tunneling beyond semiclassical approximation," *Journal of High Energy Physics*, vol. 2008, article 095, 2008.
- [40] R. Banerjee and B. R. Majhi, "Quantum tunneling and trace anomaly," *Physics Letters B*, vol. 674, p. 218, 2009.
- [41] R. Banerjee and B. R. Majhi, "Connecting anomaly and tunneling methods for the Hawking effect through chirality," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 79, no. 6, Article ID 064024, 2009.
- [42] R. Banerjee and B. R. Majhi, "Hawking black body spectrum from tunneling mechanism," *Physics Letters B*, vol. B675, p. 243, 2009.
- [43] R. Banerjee, B. R. Majhi, and S. Samanta, "Noncommutative black hole thermodynamics," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 77, no. 12, Article ID 124035, 2008.
- [44] B. R. Majhi, "Fermion tunneling beyond semiclassical approximation," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 79, Article ID 044005, 2009.
- [45] B. R. Majhi, "Hawking radiation and black hole spectroscopy in Horava-Lifshitz gravity," *Physics Letters B*, vol. B686, p. 49, 2010.
- [46] B. R. Majhi and S. Samanta, "Hawking radiation due to photon and gravitino tunneling," *Annals of Physics*, vol. 325, no. 11, pp. 2410-2424, 2010.
- [47] R. Banerjee, B. R. Majhi, and E. C. Vagenas, "Quantum tunneling and black hole spectroscopy," *Physics Letters B*, vol. 686, no. 4-5, pp. 279-282, 2010.
- [48] R. Banerjee, B. R. Majhi, and E. C. Vagenas, "A note on the lower bound of black-hole area change in the tunneling formalism," *Europhysics Letters*, vol. 92, p. 20001, 2010.
- [49] K. Bhattacharya and B. R. Majhi, "Temperature and thermodynamic structure of Einstein's equations for a cosmological black hole," *Physical Review*, vol. D94, p. 024033, 2016.
- [50] M. Sharif and W. Javed, "Fermions tunneling from charged anti-de Sitter black holes," *Canadian Journal of Physics*, vol. 90, p. 903, 2012.
- [51] M. Sharif and W. Javed, "Fermions tunneling from Plebański-Demiański black holes," *General Relativity and Gravitation*, vol. 45, no. 5, pp. 1051-1068, 2013.
- [52] M. Sharif and W. Javed, "Thermodynamics of charged Newman-Unti-Tamburino accelerating rotating black holes," *Canadian Journal of Physics*, vol. 91, p. 43, 2013.
- [53] M. Sharif and W. J. Javed, "Charged fermions tunneling from regular black holes," *Journal of Experimental and Theoretical Physics*, vol. 115, p. 782, 2012.
- [54] M. Sharif and W. Javed, "Tunneling from reissner-nordstrom-de sitter black hole with a global monopole," in *Proceedings of the 3rd GalileoXu Guangqi Meeting, International Journal of Modern Physics: Conference Series*, vol. 23, p. 271, 2013.
- [55] M. Sharif and W. Javed, "Some interesting aspects of Hawking radiation," in *Proceedings of the 13th Marcel Grossmann Meeting*, vol. 3, pp. 1950-1952, Stockholm, 2015.
- [56] M. Sharif and W. Javed, "Fermions tunneling from charged accelerating and rotating black holes with NUT parameter," *The European Physical Journal C*, vol. 72, p. 1997, 2012.

- [57] M. Sharif and W. Javed, "Quantum Corrections for a Bardeen Regular Black Hole," *Journal of the Korean Physical Society*, vol. 57, p. 217, 2010.
- [58] W. Javed, G. Abbas, and R. Ali, "Charged Vector Particles Tunneling From 5D Black Hole and Black Ring," <https://arxiv.org/abs/1705.05702>.
- [59] M. S. Longair, *Confrontation of Cosmological Theories with Observational Data*, Springer Science and Business Media, 1974.
- [60] H. Alex, "How Einstein Discovered Dark Energy," <https://arxiv.org/abs/1211.6338>.
- [61] D. Overbye, *Astronomers Report Evidence of Dark Energy Splitting the Universe*, The New York Times, 2015.
- [62] S. M. Carroll, "Quintessence and the Rest of the World: Suppressing Long-Range Interactions," *Physical Review Letters*, vol. 81, p. 3067, 1998.
- [63] Z. Xu and J. Wang, "Kerr-Newman-AdS black hole in quintessential dark energy," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. D95, p. 064015, 2017.
- [64] V. V. Kiselev, "Quintessence and black holes," *Classical and Quantum Gravity*, vol. 20, no. 6, pp. 1187–1197, 2003.
- [65] R. R. Caldwell, "An Introduction to Quintessence," *Brazilian Journal of Physics*, vol. 30, p. 2, 2000.
- [66] G. Q. Li, "Effects of dark energy on P–V criticality of charged AdS black holes," *Physics Letters B*, vol. B735, p. 256, 2014.
- [67] E. T. Newman and A. I. Janis, "Note on the Kerr spinning-particle metric," *Journal of Mathematical Physics*, vol. 6, pp. 915–917, 1965.
- [68] E. T. Newman, E. Couch, K. Chinnapared, A. Exton, A. Prakash, and R. Torrence, "Metric of a rotating, charged mass," *Journal of Mathematical Physics*, vol. 6, pp. 918–919, 1965.
- [69] E. T. Newman, "Complex coordinate transformations and the Schwarzschild-Kerr metrics," *Journal of Mathematical Physics*, vol. 14, pp. 774–776, 1973.
- [70] E. T. Newman, "The Remarkable Efficacy of Complex Methods in General Relativity," in *Highlights in Gravitation and Cosmology. Proceedings of the International Conference on Gravitation and Cosmology (Goa, 1987)*, B. R. Iyer, A. Kembhavi, J. V. Narlikar et al., Eds., Cambridge University Press, Cambridge, UK, 1988.
- [71] R. P. Kerr, "Gravitational field of a spinning mass as an example of algebraically special metrics," *Physical Review Letters*, vol. 11, pp. 237–238, 1963.
- [72] E. Newman and R. Penrose, "An approach to gravitational radiation by a method of spin coefficients," *Journal of Mathematical Physics*, vol. 3, pp. 566–578, 1962.
- [73] C. Bambi and L. Modesto, "Rotating regular black holes," *Physics Letters B*, vol. 721, no. 4–5, pp. 329–334, 2013.
- [74] B. Toshmatov, B. Ahmedov, A. Abdujabbarov, and Z. Stuchlík, "Rotating regular black hole solution," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 89, no. 10, Article ID 104017, 2014.
- [75] A. Larranaga, A. A. Cardenas, and D. A. Torres, "RETRACTED: On a general class of regular rotating black holes based on a smeared mass distribution," *Physics Letters B*, vol. 743, p. 492, 2015.
- [76] S. G. Ghosh and S. D. Maharaj, "Radiating Kerr-like regular black hole," *The European Physical Journal C*, vol. 75, p. 7, 2015.
- [77] M. M. Caldarelli, G. Cognola, and D. Klemm, "Thermodynamics of Kerr-Newman-AdS black holes and conformal field theories," *Classical and Quantum Gravity*, vol. 17, p. 399, 2000.
- [78] A. N. Aliev, "Electromagnetic properties of Kerr–anti-de Sitter black holes," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 75, p. 084041, 2007.
- [79] J. Ahmed and K. Saifullah, "Hawking Radiation of Dirac Particles from Black Strings," *Journal of Cosmology and Astroparticle Physics*, vol. 08, p. 011, 2011.
- [80] E. T. Akhmedov, V. Akhmedova, and D. Singleton, "Hawking temperature in the tunneling picture," *Physics Letters B*, vol. 642, no. 1–2, pp. 124–128, 2006.
- [81] E. T. Akhmedov, V. Akhmedova, T. Pilling, and D. Singleton, "Thermal radiation of various gravitational backgrounds," *International Journal of Modern Physics A*, vol. A22, p. 1705, 2007.
- [82] V. Akhmedova, T. Pilling, A. de Gill, and D. Singleton, "Temporal contribution to gravitational WKB-like calculations," *Physics Letters B*, vol. 666, no. 3, pp. 269–271, 2008.
- [83] E. T. Akhmedov, T. Pilling, and D. Singleton, "Subtleties in the quasi-classical calculation of hawking radiation," *International Journal of Modern Physics D*, vol. D17, p. 2453, 2008.
- [84] V. Akhmedova, T. Pilling, A. de Gill et al., "Temporal contribution to gravitational WKB-like calculations," *Physics Letters B*, vol. 666, p. 269, 2008.
- [85] V. Akhmedova, T. Pilling, A. de Gill et al., "Comments on anomaly versus WKB/tunneling methods for calculating Unruh radiation," *Physics Letters B*, vol. B673, p. 277, 2009.
- [86] d. Gill, D. Singleton, V. Akhmedova, and T. Pilling, "A WKB-like approach to Unruh radiation," *American Journal of Physics*, vol. 78, p. 685, 2010.
- [87] D. Chen, H. Wu, H. Yang, and S. Yang, "Effects of quantum gravity on black holes," *International Journal of Modern Physics A*, vol. 29, p. 1430054, 2014.
- [88] E. T. Akhmedov, V. Akhmedova, D. Singleton, and T. Pilling, "Thermal radiation of various gravitational backgrounds," *International Journal of Modern Physics A*, vol. 22, no. 8–9, pp. 1705–1715, 2007.
- [89] B. D. Chowdhury, "Problems with Tunneling of Thin Shells from Black Holes," *Pramana*, vol. 70, p. 593, 2008.
- [90] P. Mitra, "Hawking temperature from tunnelling formalism," *Physics Letters B*, vol. 648, p. 240, 2007.
- [91] P. Pradhan, "CFT and Logarithmic Corrections to the Black Hole Entropy Product Formula," *Advances in High Energy Physics*, vol. 2017, Article ID 2367387, 8 pages, 2017.



Hindawi

Submit your manuscripts at
www.hindawi.com

