Research Article

Links between (γ_n, σ_k) -KP Hierarchy, (γ_n, σ_k) -mKP Hierarchy, and $(2+1)$ -(γ_n, σ_k)-Harry Dym Hierarchy

Yehui Huang,¹ Yuqin Yao,² and Yunbo Zeng³

¹School of Mathematics and Physics, North China Electric Power University, Beijing 102206, China

²Department of Applied Mathematics, China Agricultural University, Beijing 100083, China

³Department of Mathematical Science, Tsinghua University, Beijing 100084, China

Correspondence should be addressed to Yuqin Yao; yyqinw@126.com

Received 12 July 2015; Accepted 16 November 2015

Academic Editor: Boris G. Konopelchenko

Copyright © 2015 Yehui Huang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The new $(2+1)$ -(γ_n, σ_k)-Harry Dym hierarchy and (γ_n, σ_k) -mKP hierarchy with two new time series γ_n and σ_k , which consist of γ_n -flow, σ_k -flow, and mixed γ_n and σ_k evolution equations of eigenfunctions, are proposed. Gauge transformations and reciprocal transformations between (γ_n, σ_k) -KP hierarchy, (γ_n, σ_k) -mKP hierarchy, and $(2+1)$ -(γ_n, σ_k)-Harry Dym hierarchy are studied. Their soliton solutions are presented.

1. Introduction

Generalizations of soliton hierarchies are important topics since last century. In 2008, KdV6 equation is studied as a nonholonomic deformation of KdV equation. Kupershmidt developed a deformation from the bi-Hamiltonian structure of the soliton equations and KdV6 equation could be seen as the deformation of the KdV equation [1]. We generalized this Kupershmidt deformed system and find that KdV6 equation could be seen as the Rosochatius deformation of the KdV equation with self-consistent sources and firstly found the bi-Hamiltonian structure for the KdV6 equation [2]. The Kadomtsev-Petviashvili (KP) hierarchy is an important $2+1$ dimensional integrable system and the generalizations of KP hierarchy (KPH) attract a lot of interests from both physical and mathematical points of view [3–11]. One kind of generalization is the multicomponent KP hierarchy [3], which contains many physical relevant nonlinear integrable systems such as Davey-Stewartson equation, two-dimensional Toda lattice, and three-wave resonant integrable equations. Another kind of generalization of KP equation is the so called KP equation with self-consistent sources (KPSCS) [10, 11]. The third kind of generalization is the extended KP hierarchy (exKPH), which is constructed by introducing another time

series $\{\tau_k\}$ [12–14]. The exKPH consists of t_n -flow of KP hierarchy, τ_k -flow, and t_n -evolution equations of eigenfunctions. To make difference, we may call the exKPH as (t_n, τ_k) -KPH.

Recently, we generalize the (t_n, τ_k) -KPH to (γ_n, σ_k) -KPH by introducing two new time series γ_n and σ_k with two parameters α_n and β_k [15]:

$$L_{\gamma_n} = \left[B_n + \alpha_n \sum_{i=1}^N q_i \partial^{-1} r_i, L \right], \quad B_n = L_{\geq 0}^n, \quad (1a)$$

$$L_{\sigma_k} = \left[B_k + \beta_k \sum_{i=1}^N q_i \partial^{-1} r_i, L \right], \quad (1b)$$

$$\begin{aligned} \alpha_n (q_{i, \sigma_k} - B_k(q_i)) - \beta_k (q_{i, \gamma_n} - B_n(q_i)) &= 0, \\ \alpha_n (r_{i, \sigma_k} + B_k^*(r_i)) - \beta_k (r_{i, \gamma_n} + B_n^*(r_i)) &= 0, \end{aligned} \quad (1c)$$

$$i = 1, 2, \dots, N,$$

where the pseudodifferential operator L with potential functions u_i is defined as

$$L = \partial + u_1 \partial^{-1} + u_2 \partial^{-2} + \dots \quad (2)$$

and $*$ denotes the adjoint operator. The compatibility of γ_n -flow (1a) and σ_k -flow (1b) under (1c) gives rise to the zero-curvature representation for (1a), (1b), and (1c),

$$B_{n,\sigma_k} - B_{k,\gamma_n} + [B_n, B_k] + \beta_k \left[B_n, \sum_{i=1}^N q_i \partial^{-1} r_i \right]_+ + \alpha_n \left[\sum_{i=1}^N q_i \partial^{-1} r_i, B_k \right]_+ = 0, \quad (3a)$$

$$\begin{aligned} \alpha_n (q_{i,\sigma_k} - B_k(q_i)) - \beta_k (q_{i,\gamma_n} - B_n(q_i)) &= 0, \\ \alpha_n (r_{i,\sigma_k} + B_k^*(r_i)) - \beta_k (r_{i,\gamma_n} + B_n^*(r_i)) &= 0, \end{aligned} \quad (3b)$$

$i = 1, 2, \dots, N,$

with the Lax representation

$$\begin{aligned} \psi_{\gamma_n} &= \left(B_n + \alpha_n \sum_{i=1}^N q_i \partial^{-1} r_i \right) (\psi), \\ \psi_{\sigma_k} &= \left(B_k + \beta_k \sum_{i=1}^N q_i \partial^{-1} r_i \right) (\psi). \end{aligned} \quad (4)$$

The (γ_n, σ_k) -KPH consists of γ_n -flow, σ_k -flow, and one mixed γ_n and σ_k evolution equation of eigenfunctions. The (γ_n, σ_k) -KPH can be reduced to the KPH and (t_n, τ_k) -KPH and contains first type and second type as well as mixed type of KPESCS as special cases. We also develop the dressing method to solve the (γ_n, σ_k) -KPH [15].

The (2+1)-Harry Dym equation has been firstly defined by Konopelchenko and Dubrovsky in 1984 [16]. The Harry Dym hierarchy is the third hierarchy of soliton hierarchies presented by the pseudodifferential operator technique after the KP hierarchy and mKP hierarchy [17]. In this paper, we first constructed the (2+1)- (γ_n, σ_k) -Harry Dym hierarchy ((2+1)- (γ_n, σ_k) -HDH), which consists of γ_n -flow, σ_k -flow, and one mixed γ_n and σ_k evolution equation of eigenfunctions. The (2+1)- (γ_n, σ_k) -HDH can be reduced to the (2+1)-Harry Dym hierarchy ((2+1)-HDH) and (t_n, τ_k) -HDH [18] and contains first type and second type as well as mixed type of Harry Dym equation with self-consistent sources (HDESCS) as special cases. Then the (γ_n, σ_k) -mKPH is proposed. The generalized gauge transformations and generalized reciprocal links between these three kinds of (γ_n, σ_k) hierarchies are studied. Further, based on these transformations and the solutions of the (γ_n, σ_k) -KPH [15], the solutions of (γ_n, σ_k) -mKPH and (2+1)- (γ_n, σ_k) -HDH are obtained, respectively.

The paper is organized as follows. In Section 2, we propose a new (γ_n, σ_k) -HDH and present its reduction. A new (γ_n, σ_k) -mKPH is proposed in Section 3. Section 4 is devoted to studying the links between the three kinds of (γ_n, σ_k) hierarchies. Section 5 presents the soliton solutions and a conclusion is given in the last section.

2. A New (2+1)- (γ_n, σ_k) -Harry Dym Hierarchy

2.1. A New (2+1)- (γ_n, σ_k) -Harry Dym Hierarchy. It is well known that the pseudodifferential operator $\bar{L}$ for (2+1)-HDH with potential functions w_i is defined as

$$\bar{L} = w\partial + w_0 + w_1\partial^{-1} + w_2\partial^{-2} + \dots \quad (5)$$

The (2+1)-HDH is given by [17]

$$\bar{L}_{t_n} = [\bar{B}_n, \bar{L}], \quad n \geq 1, \quad (6)$$

where $\bar{B}_n = \bar{L}_+^n = (\bar{L}^n)_{\geq 2}$, $n \geq 1$.

The compatibility of the t_n -flow and t_k -flow of (6) leads to the zero-curvature representation of (2+1)-HDH:

$$\bar{B}_{n,t_k} - \bar{B}_{k,t_n} + [\bar{B}_n, \bar{B}_k] = 0. \quad (7)$$

In particular, when taking $\bar{B}_2 = w^2\partial^2$ and $\bar{B}_3 = w^3\partial^3 + 3w^2(w_x + w_0)\partial^2$ and setting $t_2 = y$, $t_3 = t$, and $w_0 = (1/2)\partial^{-1}(w_y/w^2) - (1/2)w_x$, (7) yields the (2+1)-HD equation

$$4w_t = w^3 w_{xxx} - 3 \frac{1}{w} \left[w^2 \partial^{-1} \left(\frac{w_y}{w^2} \right) \right]_y, \quad (8)$$

which is reduced to the Harry Dym equation. Consider

$$4w_t = w^3 w_{xxx} \quad (9)$$

by dropping the y -dependence.

Based on the squared eigenfunction symmetry constraint

$$\begin{aligned} \bar{L}^k &= \bar{B}_k + \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \\ \bar{q}_{i,t_n} &= \bar{B}_n(\bar{q}_i), \end{aligned} \quad (10)$$

$$\bar{r}_{i,t_n} = -\partial^{-2} \bar{B}_n^* \partial^2(\bar{r}_i),$$

which is compatible with (2+1)-HDH [17], we propose the following generalized (2+1)-HDH with two generalized time series γ_n and σ_k :

$$\bar{L}_{\gamma_n} = \left[\bar{B}_n + \alpha_n \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \bar{L} \right], \quad (11a)$$

$$\bar{L}_{\sigma_k} = \left[\bar{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \bar{L} \right], \quad (11b)$$

$$\alpha_n (\bar{q}_{i,\sigma_k} - \bar{B}_k(\bar{q}_i)) - \beta_k (\bar{q}_{i,\gamma_n} - \bar{B}_n(\bar{q}_i)) = 0,$$

$$\begin{aligned} \alpha_n (\bar{r}_{i,\sigma_k} + \partial^{-2} \bar{B}_k^* \partial^2(\bar{r}_i)) - \beta_k (\bar{r}_{i,\gamma_n} + \partial^{-2} \bar{B}_n^* \partial^2(\bar{r}_i)) \\ = 0, \quad i = 1, 2, \dots, N. \end{aligned} \quad (11c)$$

We will prove the compatibility of (11a) and (11b) under (11c) in the following theorem.

Theorem 1. *The γ_n -flow (11a) and σ_k -flow (11b) under (11c) are compatible.*

Proof. Denote

$$\hat{B}_n = \bar{B}_n + \alpha_n \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \quad (12)$$

$$\hat{B}_k = \bar{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2.$$

In order to prove $\bar{L}_{\gamma_n, \sigma_k} = \bar{L}_{\sigma_k, \gamma_n}$, that is,

$$[\hat{B}_n, \hat{B}_k + [\hat{B}_n, \hat{B}_k], \bar{L}] = 0, \quad (13)$$

we only need to prove

$$\hat{B}_{n, \sigma_k} - \hat{B}_{k, \gamma_n} + [\hat{B}_n, \hat{B}_k] = 0. \quad (14)$$

For convenience, we omit $\sum$. We can find that

$$\begin{aligned} \hat{B}_{n, \sigma_k} &= \bar{B}_{n, \sigma_k} + \alpha_n (\bar{q} \partial^{-1} \bar{r} \partial^2)_{\sigma_k} \\ &= [\bar{B}_k + \beta_k (\bar{q} \partial^{-1} \bar{r} \partial^2), \bar{L}^n]_+ + \alpha_n (\bar{q} \partial^{-1} \bar{r} \partial^2)_{\sigma_k} \\ &= [\bar{B}_k, \bar{L}^n]_+ + \beta_k [\bar{q} \partial^{-1} \bar{r} \partial^2, \bar{L}^n]_+ + \alpha_n \bar{q}_{\sigma_k} \partial^{-1} \bar{r} \partial^2 \\ &\quad + \alpha_n \bar{q} \partial^{-1} \bar{r}_{\sigma_k} \partial^2, \end{aligned} \quad (15)$$

and, similarly,

$$\begin{aligned} \hat{B}_{k, \gamma_n} &= [\bar{B}_n, \bar{L}^k]_+ + \alpha_n [\bar{q} \partial^{-1} \bar{r} \partial^2, \bar{L}^k]_+ + \beta_k \bar{q}_{\gamma_n} \partial^{-1} \bar{r} \partial^2 \\ &\quad + \beta_k \bar{q} \partial^{-1} \bar{r}_{\gamma_n} \partial^2. \end{aligned} \quad (16)$$

Moreover, on the other hand, under the formula [18]

$$\begin{aligned} \left[\bar{B}_n, \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right]_- &= \sum_{i=1}^N \bar{B}_n (\bar{q}_i) \partial^{-1} \bar{r}_i \partial^2 \\ &\quad - \sum_{i=1}^N \bar{q}_i \partial^{-3} \bar{B}_n^* (\bar{r}_i) \partial^4, \end{aligned} \quad (17)$$

we have

$$\begin{aligned} [\hat{B}_n, \hat{B}_k] &= [\bar{B}_n, \bar{B}_k] + [\bar{B}_n, \beta_k \bar{q} \partial^{-1} \bar{r} \partial^2] \\ &\quad + [\alpha_n \bar{q} \partial^{-1} \bar{r} \partial^2, \bar{B}_k] \\ &= [\bar{L}^n - (\bar{L}^n)_-, \bar{L}^k - (\bar{L}^k)_-]_+ \\ &\quad + \beta_k [\bar{B}_n, \bar{q} \partial^{-1} \bar{r} \partial^2]_+ + \alpha_n [\bar{q} \partial^{-1} \bar{r} \partial^2, \bar{B}_k]_+ \\ &\quad + \beta_k [\bar{B}_n, \bar{q} \partial^{-1} \bar{r} \partial^2]_- + \alpha_n [\bar{q} \partial^{-1} \bar{r} \partial^2, \bar{B}_k]_- \\ &= [\bar{B}_n, \bar{L}^k]_+ + [\bar{L}^n, \bar{B}_k]_+ - [(\bar{L}^n)_-, (\bar{L}^k)_-]_+ \\ &\quad + \beta_k [\bar{B}_n, \bar{q} \partial^{-1} \bar{r} \partial^2]_+ + \alpha_n [\bar{q} \partial^{-1} \bar{r} \partial^2, \bar{B}_k]_+ \\ &\quad + \beta_k \bar{B}_n (\bar{q}) \partial^{-1} \bar{r} \partial^2 - \beta_k \bar{q} \partial^{-3} \bar{B}_n^* (\bar{r}) \partial^4 \\ &\quad - \alpha_n \bar{B}_k (\bar{q}) \partial^{-1} \bar{r} \partial^2 + \alpha_n \bar{q} \partial^{-3} \bar{B}_k^* (\bar{r}) \partial^4. \end{aligned} \quad (18)$$

Then (15), (16), and (18) under (11c) yield

$$\begin{aligned} \hat{B}_{n, \sigma_k} - \hat{B}_{k, \gamma_n} + [\hat{B}_n, \hat{B}_k] &= [\alpha_n (\bar{q}_{\sigma_k} - \bar{B}_k (q)) \\ &\quad - \beta_k (\bar{q}_{\gamma_n} - \bar{B}_n (q))] \partial^{-1} \bar{r} \partial^2 \\ &\quad + \bar{q} \partial^{-1} [\alpha_n (\bar{r}_{\sigma_k} + \partial^{-2} \bar{B}_k^* \partial^2 (\bar{r})) \\ &\quad - \beta_k (\bar{r}_{\gamma_n} + \partial^{-2} \bar{B}_n^* \partial^2 (\bar{r}))] \partial^2 = 0. \end{aligned} \quad (19)$$

□

The compatibility of γ_n -flow (11a) and σ_k -flow (11b) under (11c) gives rise to the zero-curvature representation for (11a), (11b), and (11c):

$$\begin{aligned} \left(\bar{B}_n + \alpha_n \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right)_{\sigma_k} &- \left(\bar{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right)_{\gamma_n} \\ &+ \left[\bar{B}_n + \alpha_n \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \bar{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right] \\ &= 0 \end{aligned} \quad (20)$$

which under (11c) can be simplified as follows. Then we have the following.

Theorem 2. *The commutativity of (11a) and (11b) under (11c) gives rise to the zero-curvature equation for the generalized (2+1)-HDH with two generalized time series:*

$$\begin{aligned} \bar{B}_{n, \sigma_k} - \bar{B}_{k, \gamma_n} + [\bar{B}_n, \bar{B}_k] &+ \beta_k \left[\bar{B}_n, \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right]_+ \\ &+ \alpha_n \left[\sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \bar{B}_k \right]_+ = 0, \end{aligned} \quad (21a)$$

$$\begin{aligned} \alpha_n (\bar{q}_{i, \sigma_k} - \bar{B}_k (\bar{q}_i)) &- \beta_k (\bar{q}_{i, \gamma_n} - \bar{B}_n (\bar{q}_i)) = 0, \\ \alpha_n (\bar{r}_{i, \sigma_k} + \partial^{-2} \bar{B}_k^* \partial^2 (\bar{r}_i)) & \\ \alpha_n (\bar{r}_{i, \gamma_n} + \partial^{-2} \bar{B}_n^* \partial^2 (\bar{r}_i)) &= 0, \quad i = 1, 2, \dots, N, \end{aligned} \quad (21b)$$

with the Lax representation

$$\begin{aligned} \bar{\psi}_{\gamma_n} &= \left(\bar{B}_n + \alpha_n \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right) (\bar{\psi}), \\ \bar{\psi}_{\sigma_k} &= \left(\bar{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right) (\bar{\psi}). \end{aligned} \quad (22)$$

We briefly call (11a), (11b), and (11c) and (21a) and (21b) as (2+1)- (γ_n, σ_k) -HDH. It is easy to see that (2+1)- (γ_n, σ_k) -HDH ((11a), (11b), and (11c) and (21a) and (21b)) for $\alpha_n = \beta_k = 0$ reduces to (2+1)-HDH ((6) and (7)) and (2+1)- (γ_n, σ_k) -HDH for $\alpha_n = 0, \beta_k = 1$ reduces to (t_n, τ_k) -HDH [18]. So (2+1)- (γ_n, σ_k) -HDH ((11a), (11b), and (11c) and (21a) and (21b)) presents a more generalized (2+1)-HDH which contains the (2+1)-HDH and (2+1)- (t_n, τ_k) -HDH as the special cases.

Example 3. Let us take $n = 2$ and $k = 3$, and set $\gamma_2 = y$, $\sigma_3 = t$. Then (21a) and (21b) become

$$\begin{aligned} \bar{B}_{2,t} - \bar{B}_{3,y} + [\bar{B}_2, \bar{B}_3] + \beta_3 \left[\bar{B}_2, \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right]_+ \\ + \alpha_2 \left[\sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \bar{B}_3 \right]_+ = 0, \end{aligned} \quad (23a)$$

$$\begin{aligned} \alpha_2 (\bar{q}_{i,t} - \bar{B}_3 (\bar{q}_i)) - \beta_3 (\bar{q}_{i,y} - \bar{B}_2 (\bar{q}_i)) = 0, \\ \alpha_2 (\bar{r}_{i,t} + \partial^{-2} \bar{B}_3^* \partial^2 (\bar{r}_i)) \\ - \beta_3 (\bar{r}_{i,y} + \partial^{-2} \bar{B}_2^* \partial^2 (\bar{r}_i)) = 0, \quad i = 1, 2, \dots, N, \end{aligned} \quad (23b)$$

which gives the following nonlinear equation:

$$\begin{aligned} 2ww_t - 3[w^2(w_x + w_0)]_y \\ + w^3(ww_{xxx} + 6w_x w_{xx} + 3ww_{0xx} + 12w_x w_{0x}) \\ + 2\beta_3 w^2 \sum_{i=1}^N (\bar{q}_i \bar{r}_i)_x - 2\beta_3 w w_x \sum_{i=1}^N \bar{q}_i \bar{r}_i \\ - 3\alpha_2 w^3 \sum_{i=1}^N (\bar{q}_{i,xx} \bar{r}_i + \bar{q}_{i,x} \bar{r}_{ix}) \\ - 6\alpha_2 w^2 (w_x + w_0) \sum_{i=1}^N (\bar{q}_i \bar{r}_i)_x \\ + 3\alpha_2 [w^2(w_x + w_0)]_x \sum_{i=1}^N \bar{q}_i \bar{r}_i = 0, \end{aligned} \quad (24a)$$

$$\begin{aligned} w_y - w^2(w_{xx} + 2w_{0x}) + \alpha_2 w \sum_{i=1}^N (\bar{q}_i \bar{r}_i)_x \\ - \alpha_2 w_x \sum_{i=1}^N \bar{q}_i \bar{r}_i = 0, \\ \alpha_2 [\bar{q}_{i,t} - w^3 \bar{q}_{i,xxx} - 3w^2(w_x + w_0) \bar{q}_{i,xx}] \\ - \beta_3 (\bar{q}_{i,y} - w^2 \bar{q}_{i,xx}) = 0, \\ \alpha_2 (\bar{r}_{i,t} - w^3 \bar{r}_{i,xxx} - 3w^2 w_0 \bar{r}_{i,xx}) \\ - \beta_3 (\bar{r}_{i,y} + w^2 \bar{r}_{i,xx}) = 0, \quad i = 1, 2, \dots, N, \end{aligned} \quad (24b)$$

$$\begin{aligned} \alpha_2 [\bar{q}_{i,t} - w^3 \bar{q}_{i,xxx} - 3w^2(w_x + w_0) \bar{q}_{i,xx}] \\ - \beta_3 (\bar{q}_{i,y} - w^2 \bar{q}_{i,xx}) = 0, \\ \alpha_2 (\bar{r}_{i,t} - w^3 \bar{r}_{i,xxx} - 3w^2 w_0 \bar{r}_{i,xx}) \\ - \beta_3 (\bar{r}_{i,y} + w^2 \bar{r}_{i,xx}) = 0, \quad i = 1, 2, \dots, N, \end{aligned} \quad (24c)$$

with the Lax representation as follows:

$$\bar{\psi}_y = \left(w^2 \partial^2 + \alpha_2 \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right) (\bar{\psi}), \quad (25)$$

$$\begin{aligned} \bar{\psi}_t = \left(w^3 \partial^3 + 3w^2(w_x + w_0) \partial^2 + \beta_3 \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right) \\ \cdot (\bar{\psi}). \end{aligned} \quad (26)$$

Particularly, when taking $\alpha_2 = \beta_3 = 0$; $\alpha_2 = 0, \beta_3 = 1$; $\alpha_2 = 1, \beta_3 = 0$; and $\alpha_2 = 1, \beta_3 = 1$, respectively, (24a), (24b), and (24c) and (26) reduce to the (2+1)-HD equation, the first type of (2+1)-HD equation with self-consistent sources ((2+1)-HDESCS), the second type of (2+1)-HDESCS, and the mixed type of (2+1)-HDESCS and their Lax representations, respectively.

2.2. Reduction. Consider the constraint given by

$$\bar{L}^k = \bar{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2. \quad (27)$$

Then (11b) yields

$$(\bar{L}^k)_{\sigma_k} = \left[\bar{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \bar{L}^k \right] = 0, \quad (28)$$

$$\bar{B}_{k,\sigma_k} = (\bar{L}^k_{\sigma_k})_+ = 0, \quad (29)$$

$$\left(\sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right)_{\sigma_k} = (\bar{L}^k_{\sigma_k})_- = 0,$$

which imply that $\bar{L}$, $\bar{B}_k$, $\bar{q}_i$, and $\bar{r}_i$ under (27) are independent of σ_k . Subsequently, $\bar{q}_{i,\sigma_k}$ and $\bar{r}_{i,\sigma_k}$ in (11c) should be replaced by $\lambda_i \bar{q}_i$ and $-\lambda_i \bar{r}_i$ as in the case of constrained flow of KP [19, 20]; namely, (11c) under the constraint (27) should be replaced by

$$\alpha_n (\lambda_i \bar{q}_i - \bar{B}_k (\bar{q}_i)) - \beta_k (\bar{q}_{i,\gamma_n} - \bar{B}_n (\bar{q}_i)) = 0, \quad (30)$$

$$\begin{aligned} \alpha_n (-\lambda_i \bar{r}_i + \partial^{-2} \bar{B}_k^* \partial^2 (\bar{r}_i)) - \beta_k (\bar{r}_{i,\gamma_n} + \partial^{-2} \bar{B}_n^* \partial^2 (\bar{r}_i)) \\ = 0. \end{aligned} \quad (31)$$

We will show that constraint (27) is invariant under γ_n -flow (11a) and (31). In fact, making use of (11a), (17), and (31), we have

$$\begin{aligned} (\bar{L}^k - \bar{B}_k)_{\gamma_n} = (\bar{L}^k_{\gamma_n})_- = [\bar{B}_n, \bar{L}^k]_- \\ = \left(\beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right)_{\gamma_n} = \beta_k \sum_{i=1}^N (\bar{q}_{i,\gamma_n} \partial^{-1} \bar{r}_i \partial^2 \\ + \bar{q}_i \partial^{-1} \bar{r}_{i,\gamma_n} \partial^2) = \sum_{i=1}^N [\beta_k \bar{B}_n (\bar{q}_i) \partial^{-1} \bar{r}_i \partial^2 \\ + \alpha_n (\lambda_i \bar{q}_i - \bar{B}_k (\bar{q}_i)) \partial^{-1} \bar{r}_i \partial^2 - \beta_k \bar{q}_i \partial^{-1} \bar{B}_n (\bar{r}_i) \partial^2 \\ + \alpha_n \bar{q}_i \partial^{-1} (-\lambda_i \bar{r}_i + \partial^{-2} \bar{B}_k^* \partial^2 (\bar{r}_i)) \partial^2] = [\bar{B}_n, \end{aligned}$$

$$\begin{aligned}
& \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \Big]_- - \left[\bar{B}_k, \alpha_n \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right]_- = \left[\hat{B}_n, \right. \\
& \left. \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right]_- - \left[\bar{B}_k, \alpha_n \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right]_- = \left[\hat{B}_n, \right. \\
& \left. \bar{L}^k \right]_- - \left[\hat{B}_n, \bar{B}_k \right]_- - \left[\bar{B}_k, \hat{B}_n \right]_- + \left[\bar{B}_k, \bar{B}_n \right]_- = \left[\hat{B}_n, \right. \\
& \left. \bar{L}^k \right]_-.
\end{aligned} \tag{32}$$

Then

$$\left(\bar{L}^k - \bar{B}_k - \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right)_{\gamma_n} = 0. \tag{33}$$

This means that the submanifold determined by k -constraint (27) is invariant under the γ_n -flow (11a) and (31).

Therefore, the constrained flow of (2+1)- (γ_n, σ_k) -HDH ((11a), (11b), and (11c) and (21a) and (21b)) under (27) reads

$$\begin{aligned}
& \bar{B}_{k,\gamma_n} + \left[\bar{B}_k, \bar{B}_n \right] + \beta_k \left[\sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2, \bar{B}_n \right]_+ \\
& + \alpha_n \left[\bar{B}_k, \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right]_+ = 0,
\end{aligned} \tag{34a}$$

$$\begin{aligned}
& \alpha_n (\lambda_i \bar{q}_i - \bar{B}_k (\bar{q}_i)) - \beta_k (\bar{q}_{i,\gamma_n} - \bar{B}_n (\bar{q}_i)) = 0, \\
& \alpha_n (-\lambda_i \bar{r}_i + \partial^{-2} \bar{B}_k^* \partial^2 (\bar{r}_i)) \\
& - \beta_k (\bar{r}_{i,\gamma_n} + \partial^{-2} \bar{B}_n^* \partial^2 (\bar{r}_i)) = 0, \quad i = 1, 2, \dots, N.
\end{aligned} \tag{34b}$$

with

$$\bar{B}_n = \left(\bar{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial^2 \right)_+^{n/k}. \tag{34c}$$

Example 4. When $n = 2, k = 3, \gamma_2 = y$, (34a), (34b), and (34c) give

$$\begin{aligned}
& 3 \left[w^2 (w_x + w_0) \right]_y \\
& - w^3 (w w_{xxx} + 6 w_x w_{xx} + 3 w w_{0xx} + 12 w_x w_{0x}) \\
& - 2 \beta_3 w^2 \sum_{i=1}^N (\bar{q}_i \bar{r}_i)_x + 2 \beta_3 w w_x \sum_{i=1}^N \bar{q}_i \bar{r}_i \\
& + 3 \alpha_2 w^3 \sum_{i=1}^N (\bar{q}_{i,xx} \bar{r}_i + \bar{q}_{i,x} \bar{r}_{ix}) \\
& + 6 \alpha_2 w^2 (w_x + w_0) \sum_{i=1}^N (\bar{q}_i \bar{r}_i)_x \\
& - 3 \alpha_2 \left[w^2 (w_x + w_0) \right]_x \sum_{i=1}^N \bar{q}_i \bar{r}_i = 0,
\end{aligned} \tag{35a}$$

$$\begin{aligned}
& w_y - w^2 (w_{xx} + 2 w_{0x}) + \alpha_2 w \sum_{i=1}^N (\bar{q}_i \bar{r}_i)_x \\
& - \alpha_2 w_x \sum_{i=1}^N \bar{q}_i \bar{r}_i = 0,
\end{aligned} \tag{35b}$$

$$\begin{aligned}
& \alpha_2 \left[\lambda_i \bar{q}_i - w^3 \bar{q}_{i,xxx} - 3 w^2 (w_x + w_0) \bar{q}_{i,xx} \right] \\
& - \beta_3 (\bar{q}_{i,y} - w^2 \bar{q}_{i,xx}) = 0, \\
& \alpha_2 (-\lambda_i \bar{r}_i - w^3 \bar{r}_{i,xxx} - 3 w^2 w_0 \bar{r}_{i,xx}) \\
& - \beta_3 (\bar{r}_{i,y} + w^2 \bar{r}_{i,xx}) = 0, \quad i = 1, 2, \dots, N,
\end{aligned} \tag{35c}$$

which is the k -constraint (2+1)- (γ_n, σ_k) -Harry Dym equation.

3. The (γ_n, σ_k) -mKPH

In the same way, the (γ_n, σ_k) -mKPH can be formulated. The $\tilde{L}$ operator of mKP hierarchy is defined by

$$\tilde{L} = \partial + v_0 + v_1 \partial^{-1} + v_2 \partial^{-2} + \dots \tag{36}$$

The Lax equation of mKP hierarchy is given by

$$\begin{aligned}
& \tilde{L}_{t_n} = [\tilde{B}_n, \tilde{L}], \\
& \tilde{B}_n = \tilde{L}_{\geq 1}^n,
\end{aligned} \tag{37}$$

$n \geq 1$.

The commutativity of ∂_{t_n} and ∂_{t_k} flows gives the zero-curvature equation (7).

Since the squared eigenfunction symmetry constraint, given by

$$\begin{aligned}
& \tilde{L}^k = \tilde{B}_k + \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial, \\
& \tilde{q}_{i,t_n} = \tilde{B}_n (\tilde{q}_i), \\
& \tilde{r}_{i,t_n} = -\partial^{-1} \tilde{B}_n^* \partial (\tilde{r}_i),
\end{aligned} \tag{38}$$

is compatible with mKP hierarchy [21, 22], we have the following.

Definition 5. The (γ_n, σ_k) -mKPH is defined by

$$\tilde{L}_{\gamma_n} = \left[\tilde{B}_n + \alpha_n \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial, \tilde{L} \right], \tag{39a}$$

$$\tilde{L}_{\sigma_k} = \left[\tilde{B}_k + \beta_k \sum_{i=1}^N \bar{q}_i \partial^{-1} \bar{r}_i \partial, \tilde{L} \right], \tag{39b}$$

$$\begin{aligned}
& \alpha_n (\bar{q}_{i,\sigma_k} - \tilde{B}_k (\bar{q}_i)) - \beta_k (\bar{q}_{i,\gamma_n} - \tilde{B}_n (\bar{q}_i)) = 0, \\
& \alpha_n (\bar{r}_{i,\sigma_k} + \partial^{-1} \tilde{B}_k^* \partial (\bar{r}_i)) - \beta_k (\bar{r}_{i,\gamma_n} + \partial^{-1} \tilde{B}_n^* \partial (\bar{r}_i)) \\
& = 0, \quad i = 1, 2, \dots, N.
\end{aligned} \tag{39c}$$

(39a), (39b), and (39c) have the Lax representation

$$\begin{aligned}\tilde{\psi}_{\gamma_n} &= \left(\tilde{B}_n + \alpha_n \sum_{i=1}^N \tilde{q}_i \partial^{-1} \tilde{r}_i \partial \right) (\tilde{\psi}), \\ \tilde{\psi}_{\sigma_k} &= \left(\tilde{B}_k + \beta_k \sum_{i=1}^N \tilde{q}_i \partial^{-1} \tilde{r}_i \partial \right) (\psi).\end{aligned}\quad (40)$$

In the same way as in Section 2, we can verify the compatibility of (39a) and (39b) under (39c).

By using the formula

$$\begin{aligned}\left[\tilde{B}_n, \sum_{i=1}^N \tilde{q}_i \partial^{-1} \tilde{r}_i \partial \right]_- &= \sum_{i=1}^N \tilde{B}_n (\tilde{q}_i) \partial^{-1} \tilde{r}_i \partial \\ &\quad - \sum_{i=1}^N \tilde{q}_i \partial^{-2} \tilde{B}_n^* \partial (\tilde{r}_i) \partial,\end{aligned}\quad (41)$$

the zero-curvature equation for (γ_n, σ_k) -mKPH can be written as

$$\begin{aligned}\tilde{B}_{n,\sigma_k} - \tilde{B}_{k,\gamma_n} + [\tilde{B}_n, \tilde{B}_k] + \beta_k \left[\tilde{B}_n, \sum_{i=1}^N \tilde{q}_i \partial^{-1} \tilde{r}_i \partial \right]_+ \\ + \alpha_n \left[\sum_{i=1}^N \tilde{q}_i \partial^{-1} \tilde{r}_i \partial, \tilde{B}_k \right]_+ = 0,\end{aligned}\quad (42a)$$

$$\begin{aligned}\alpha_n (\tilde{q}_{i,\sigma_k} - \tilde{B}_k (\tilde{q}_i)) - \beta_k (\tilde{q}_{i,\gamma_n} - \tilde{B}_n (\tilde{q}_i)) &= 0, \\ \alpha_n (\tilde{r}_{i,\sigma_k} + \partial^{-1} \tilde{B}_k^* \partial (\tilde{r}_i)) \\ - \beta_k (\tilde{r}_{i,\gamma_n} + \partial^{-1} \tilde{B}_n^* \partial (\tilde{r}_i)) &= 0, \quad i = 1, 2, \dots, N.\end{aligned}\quad (42b)$$

It is easy to see that (γ_n, σ_k) -mKPH ((39a), (39b), and (39c) and (42a) and (42b)) for $\alpha_n = \beta_k = 0$ reduces to mKPH (37) and (γ_n, σ_k) -mKPH for $\alpha_n = 0, \beta_k = 1$ reduces to exmKPH [23].

Example 6. Let us take $n = 2$ and $k = 3$, and set $\gamma_2 = y, \sigma_3 = t$, and $v_0 = v$. Then (42a) and (42b) become

$$\begin{aligned}\tilde{B}_{2,t} - \tilde{B}_{3,y} + [\tilde{B}_2, \tilde{B}_3] + \beta_3 \left[\tilde{B}_2, \sum_{i=1}^N \tilde{q}_i \partial^{-1} \tilde{r}_i \partial \right]_+ \\ + \alpha_2 \left[\sum_{i=1}^N \tilde{q}_i \partial^{-1} \tilde{r}_i \partial, \tilde{B}_3 \right]_+ = 0,\end{aligned}\quad (43a)$$

$$\begin{aligned}\alpha_2 (\tilde{q}_{i,t} - \tilde{B}_3 (\tilde{q}_i)) - \beta_3 (\tilde{q}_{i,y} - \tilde{B}_2 (\tilde{q}_i)) &= 0, \\ \alpha_2 (\tilde{r}_{i,t} + \partial^{-1} \tilde{B}_3^* \partial (\tilde{r}_i)) - \beta_3 (\tilde{r}_{i,y} + \partial^{-1} \tilde{B}_2^* \partial (\tilde{r}_i)) &= 0, \quad (43b) \\ i &= 1, 2, \dots, N,\end{aligned}$$

which gives the following nonlinear equation:

$$\begin{aligned}4v_t - v_{xxx} + 6v^2 v_x - 3\partial^{-1} v_{yy} - 6v_x \partial^{-1} v_y \\ + 4\beta_3 \sum_{i=1}^N (\tilde{q}_i \tilde{r}_i)_x + \alpha_2 \sum_{i=1}^N \left[3 (\tilde{q}_i \tilde{r}_{i,xx} - \tilde{q}_{i,x} \tilde{r}_i) \right.\end{aligned}\quad (44a)$$

$$\left. - 3 (\tilde{q}_i \tilde{r}_i)_y - 6 (v \tilde{q}_i \tilde{r}_i)_x \right] = 0,$$

$$\begin{aligned}\alpha_2 \left[\tilde{q}_{i,t} - \tilde{q}_{i,xxx} - 3v \tilde{q}_{i,xx} - \frac{3}{2} (\partial^{-1} v_y) \tilde{q}_{i,x} - \frac{3}{2} v_x \tilde{q}_{i,x} \right. \\ \left. - \frac{3}{2} v^2 \tilde{q}_{i,x} - \frac{3}{2} \sum_{j=1}^N \tilde{q}_j \tilde{r}_j \tilde{q}_{ix} \right] - \beta_3 (\tilde{q}_{i,y} - \tilde{q}_{i,xx} \\ - 2v \tilde{q}_{ix}) = 0,\end{aligned}\quad (44b)$$

$$\begin{aligned}\alpha_2 \left[\tilde{r}_{i,t} - \tilde{r}_{i,xxx} - 3v \tilde{r}_{i,xx} - \frac{3}{2} (\partial^{-1} v_y) \tilde{r}_{i,x} - \frac{3}{2} v_x \tilde{r}_{i,x} \right. \\ \left. - \frac{3}{2} v^2 \tilde{r}_{i,x} - \frac{3}{2} \sum_{j=1}^N \tilde{q}_j \tilde{r}_j \tilde{r}_{ix} \right] - \beta_3 (\tilde{r}_{i,y} + \tilde{r}_{i,xx} \\ - 2v \tilde{r}_{ix}) = 0, \quad i = 1, 2, \dots, N.\end{aligned}$$

Particularly, when taking $\alpha_2 = \beta_3 = 0; \alpha_2 = 0, \beta_3 = 1; \alpha_2 = 1, \beta_3 = 0$; and $\alpha_2 = 1, \beta_3 = 1$, respectively, (44a) and (44b) reduce to the mKP equation, the first type of mKP equation with self-consistent sources, the second type of mKP equation with self-consistent sources, and the mixed type of mKP equation with self-consistent sources, respectively.

4. Links between (γ_n, σ_k) -KPH and (γ_n, σ_k) -mKPH and $(2+1)$ -(γ_n, σ_k)-HDH

In [17] the gauge transformations between KP and mKP hierarchies and the reciprocal transformation between mKP and HD hierarchies are proposed. The KP, mKP, and HD hierarchy are intimately related under these gauge transformations and reciprocal links. In [22], the constrained KP hierarchy and the constrained modified KP hierarchy are studied. And in [24], the gauge transformation between KP and mKP hierarchies with self-consistent sources is given. It is natural to think whether these transformations can be generalized to the (γ_n, σ_k) -KPH, (γ_n, σ_k) -mKPH, and $(2+1)$ -(γ_n, σ_k)-HDH. The answer is positive and the main results are as follows.

Theorem 7. (a) Suppose L, q_i, r_i satisfy (γ_n, σ_k) -KPH (1a), (1b), and (1c) and q and f are independent eigenfunctions for Lax pair (4); then

$$\tilde{L} = q^{-1} L q,$$

$$\tilde{q}_i = q^{-1} q_i,$$

$$\begin{aligned}\bar{r}_i &= -\partial^{-1}(qr_i), \\ \bar{q} &= q^{-1}f\end{aligned}\quad (45)$$

satisfy the (γ_n, σ_k) -mKPH ((39a), (39b), and (39c)) and its Lax pair (40).

(b) Suppose $\bar{L}$, $\bar{q}_i$, $\bar{r}_i$ satisfy the (γ_n, σ_k) -mKPH ((39a), (39b), and (39c)); $\bar{q}$ is eigenfunction for Lax pair (40). After the transformation $\bar{x} = \bar{q}(x, \gamma_n, \sigma_k)$, $\bar{\gamma}_n = \gamma_n$, $\bar{\sigma}_k = \sigma_k$, then

$$\begin{aligned}\bar{L} &= \bar{L}, \\ \bar{q}_i &= \bar{q}_i, \\ \bar{r}_i &= \partial^{-1}(\bar{q}\bar{r}_{ix}) - \bar{r}_i\bar{q}\end{aligned}\quad (46)$$

satisfy the (2+1)- (γ_n, σ_k) -HDH ((11a), (11b), and (11c)).

Proof. For convenience, we omit Σ .

(a) Consider

$$\begin{aligned}\bar{L}_{\gamma_n} &= -q^{-2}q_{\gamma_n}Lq + q^{-1}L_{\gamma_n}q + q^{-1}Lq_{\gamma_n} = -q^{-2}(L_{\geq 0}^n \\ &\quad + \alpha_n q_i \partial^{-1} r_i)(q) Lq + q^{-1}[L_{\geq 0}^n + \alpha_n q_i \partial^{-1} r_i, L]q \\ &\quad + q^{-1}L(L_{\geq 0}^n + \alpha_n q_i \partial^{-1} r_i)(q) \\ &= [q^{-1}Lq, q^{-1}(L_{\geq 0}^n + \alpha_n q_i \partial^{-1} r_i)(q)] + q^{-1}[L_{\geq 0}^n \\ &\quad + \alpha_n q_i \partial^{-1} r_i, L]q = [q^{-1}(L_{\geq 0}^n + \alpha_n q_i \partial^{-1} r_i)q \\ &\quad - q^{-1}(L_{\geq 0}^n + \alpha_n q_i \partial^{-1} r_i)(q), \bar{L}] = [q^{-1}L_{\geq 0}^n q \\ &\quad - q^{-1}L_{\geq 0}^n(q), \bar{L}] + \alpha_n [q^{-1}q_i \partial^{-1} r_i q \\ &\quad - q^{-1}q_i \partial^{-1}(r_i q), \bar{L}] = [\bar{L}_{\geq 1}^n, \bar{L}] + \alpha_n [\bar{q}_i \partial^{-1} \bar{r}_i \partial, \bar{L}] \\ &= [\bar{L}_{\geq 1}^n + \alpha_n \bar{q}_i \partial^{-1} \bar{r}_i \partial, \bar{L}].\end{aligned}\quad (47)$$

In the same way, (39b) can be proved. The two equations in (39c) can be proved similarly. So we only prove the first one. Consider

$$\begin{aligned}\alpha_n(\bar{q}_{i, \sigma_k} - \bar{L}_{\geq 1}^k(\bar{q}_i)) \\ &= -\alpha_n q^{-2} q_{\sigma_k} q_i + \alpha_n q^{-1} q_{i, \sigma_k} - \alpha_n \bar{L}_{\geq 1}^k(\bar{q}_i) \\ &= -\alpha_n q^{-2} (L_{\geq 0}^k + \beta_k q_i \partial^{-1} r_i)(q) q_i \\ &\quad + q^{-1} [\beta_k (q_{i, \gamma_n} - L_{\geq 0}^n(q_i)) + \alpha_n L_{\geq 0}^k(q_i)] \\ &\quad - \alpha_n \bar{L}_{\geq 1}^k(\bar{q}_i)\end{aligned}$$

$$\begin{aligned}&= -\alpha_n q^{-1} \bar{L}_{\geq 0}^k q^{-1}(q) q_i - \alpha_n \beta_k q^{-2} q_i \partial^{-1}(r_i q) q_i \\ &\quad + \beta_k q^{-1} (q_{i, \gamma_n} - L_{\geq 0}^n(q_i)) + \alpha_n \bar{L}_{\geq 0}^k(q_i q^{-1}) \\ &\quad - \alpha_n \bar{L}_{\geq 1}^k(\bar{q}_i) \\ &= -\alpha_n q^{-1} \bar{L}_{\geq 0}^k(1) q_i + \alpha_n \bar{L}_{\geq 0}^k(q_i q^{-1}) \\ &\quad - \alpha_n \beta_k q^{-2} q_i \partial^{-1}(r_i q) q_i \\ &\quad + \beta_k q^{-1} (q_{i, \gamma_n} - L_{\geq 0}^n(q_i)) - \alpha_n \bar{L}_{\geq 1}^k(\bar{q}_i) \\ &= \alpha_n \bar{L}_{\geq 1}^k(\bar{q}_i) - \alpha_n \bar{L}_{\geq 1}^k(\bar{q}_i) - \alpha_n \beta_k q^{-2} q_i \partial^{-1}(r_i q) q_i \\ &\quad + \beta_k q^{-1} (q_{i, \gamma_n} - L_{\geq 0}^n(q_i)) \\ &= -\alpha_n \beta_k q^{-2} q_i \partial^{-1}(r_i q) q_i \\ &\quad + \beta_k q^{-1} (q_{i, \gamma_n} - L_{\geq 0}^n(q_i)).\end{aligned}\quad (48)$$

Similarly, we have

$$\begin{aligned}\beta_k(\bar{q}_{i, \gamma_n} - \bar{L}_{\geq 1}^n(\bar{q}_i)) &= -\alpha_n \beta_k q^{-2} q_i \partial^{-1}(r_i q) q_i \\ &\quad + \alpha_n q^{-1} (q_{i, \sigma_k} - L_{\geq 0}^k(q_i)).\end{aligned}\quad (49)$$

Thus, we have

$$\begin{aligned}\alpha_n(\bar{q}_{i, \sigma_k} - \bar{L}_{\geq 1}^k(\bar{q}_i)) &- \beta_k(\bar{q}_{i, \gamma_n} - \bar{L}_{\geq 1}^n(\bar{q}_i)) \\ &= q^{-1} [\beta_k (q_{i, \gamma_n} - L_{\geq 0}^n(q_i)) - \alpha_n (q_{i, \sigma_k} - L_{\geq 0}^k(q_i))] \\ &= 0.\end{aligned}\quad (50)$$

In the following, we will prove that $\bar{q} = q^{-1}f$ satisfies Lax pair (40). Consider

$$\begin{aligned}\bar{q}_{\gamma_n} &= -q^{-2}q_{\gamma_n}f + q^{-1}f_{\gamma_n} \\ &= -q^{-1}(L_{\geq 0}^n + \alpha_n q_i \partial^{-1} r_i)(q) q^{-1}f \\ &\quad + q^{-1}(L_{\geq 0}^n + \alpha_n q_i \partial^{-1} r_i)(f) \\ &= -q^{-1}q \bar{L}_{\geq 0}^n q^{-1}(q) q^{-1}f + q^{-1}q \bar{L}_{\geq 0}^n q^{-1}(f) \\ &\quad - \alpha_n q^{-1} q_i \partial^{-1} r_i(q) q^{-1}f + \alpha_n q^{-1} q_i \partial^{-1} r_i(f) \\ &= -\bar{L}_{\geq 0}^n(1) \bar{q} + \bar{L}_{\geq 0}^n(\bar{q}) + \alpha_n \bar{q} \bar{q}_i \bar{r}_i + \alpha_n \bar{q}_i \bar{r}_i \partial^{-1}(r_i f) \\ &= (\bar{L}_{\geq 1}^n + \alpha_n \bar{q}_i \partial^{-1} \bar{r}_i \partial)(\bar{q}).\end{aligned}\quad (51)$$

Similarly, we can prove that $\bar{q} = q^{-1}f$ satisfies the second equation in (40).

(b) Since the first order coefficient of $\bar{L}_{\geq 1}^n = \bar{L}_{\geq 1}^n$ is given by $[\bar{L}_{\geq 1}^n \bar{x}]$, we have

$$\bar{L}_{\geq 2}^n = \bar{L}_{\geq 1}^n - \bar{L}_{\geq 1}^n(\bar{q}) \partial_{\bar{x}}. \quad (52)$$

$\tilde{L}_{\gamma_n} = [\partial_{\gamma_n}, \tilde{L}] = [\partial_{\gamma_n} + \tilde{q}_{\gamma_n} \partial_{\tilde{x}}, \tilde{L}] = \tilde{L}_{\gamma_n} + [\tilde{q}_{\gamma_n} \tilde{q}_x^{-1} \partial_x, \tilde{L}]$
yields

$$\begin{aligned}
\tilde{L}_{\gamma_n} &= \tilde{L}_{\gamma_n} - [\tilde{q}_{\gamma_n} \tilde{q}_x^{-1} \partial_x, \tilde{L}] \\
&= [\tilde{L}_{\geq 1}^n + \alpha_n \tilde{q}_i \partial^{-1} \tilde{r}_i \partial, \tilde{L}] \\
&\quad - [(\tilde{L}_{\geq 1}^n + \alpha_n \tilde{q}_i \partial^{-1} \tilde{r}_i \partial) (\tilde{q}) \tilde{q}_x^{-1} \partial_x, \tilde{L}] \\
&= [\tilde{L}_{\geq 1}^n - \tilde{L}_{\geq 1}^n (\tilde{q}) \partial_{\tilde{x}}, \tilde{L}] \\
&\quad + [\alpha_n \tilde{q}_i \partial^{-1} \tilde{r}_i \partial - \alpha_n \tilde{q}_i \partial^{-1} (\tilde{r}_i \tilde{q}_x) \partial_{\tilde{x}}, \tilde{L}] \\
&= [\tilde{L}_{\geq 2}^n, \tilde{L}] \\
&\quad + [\alpha_n \tilde{q}_i \partial^{-1} \tilde{r}_i \partial - \alpha_n \tilde{q}_i (\tilde{r}_i \tilde{q} - \partial^{-1} (\tilde{r}_{ix} \tilde{q})) \partial_{\tilde{x}}, \tilde{L}] \\
&= [\tilde{L}_{\geq 2}^n, \tilde{L}] + [-\alpha_n \tilde{q}_i \partial_{\tilde{x}}^{-1} \tilde{r}_{ix} \partial_{\tilde{x}} + \alpha_n \tilde{q}_i \tilde{r}_i \partial_{\tilde{x}}, \tilde{L}] \\
&= [\tilde{L}_{\geq 2}^n, \tilde{L}] + [\alpha_n \tilde{q}_i \partial_{\tilde{x}}^{-1} \tilde{r}_i \partial_{\tilde{x}}^2, \tilde{L}],
\end{aligned} \tag{53}$$

which implies that (11a) holds. In the same way, we can prove (11b). Consider

$$\begin{aligned}
&\alpha_n (\tilde{q}_{i, \sigma_k} - \tilde{L}_{\geq 2}^k (\tilde{q}_i)) \\
&= \alpha_n \tilde{q}_{i, \sigma_k} - \alpha_n \tilde{q}_{\sigma_k} \tilde{q}_{i, \tilde{x}} - \alpha_n \tilde{L}_{\geq 2}^k (\tilde{q}_i) \\
&= \beta_k (\tilde{q}_{i, \gamma_n} - \tilde{L}_{\geq 1}^n (\tilde{q}_i)) + \alpha_n \tilde{L}_{\geq 1}^k (\tilde{q}_i) \\
&\quad - \alpha_n (\tilde{L}_{\geq 1}^k + \beta_k \tilde{q}_i \partial^{-1} \tilde{r}_i \partial_x) (\tilde{q}) \tilde{q}_{i\tilde{x}} - \alpha_n \tilde{L}_{\geq 2}^k (\tilde{q}_i) \\
&= \alpha_n \tilde{L}_{\geq 1}^k (\tilde{q}_i) - \alpha_n \tilde{L}_{\geq 1}^k (\tilde{q}) \tilde{q}_{i\tilde{x}} \\
&\quad - \alpha_n \beta_k \tilde{q}_i \partial^{-1} \tilde{r}_i \partial_x (\tilde{q}) \tilde{q}_{i\tilde{x}} + \beta_k (\tilde{q}_{i, \gamma_n} - \tilde{L}_{\geq 1}^n (\tilde{q}_i)) \\
&\quad - \alpha_n \tilde{L}_{\geq 2}^k (\tilde{q}_i) \\
&= -\alpha_n \beta_k \tilde{q}_i \partial^{-1} \tilde{r}_i \partial_x (\tilde{q}) \tilde{q}_{i\tilde{x}} + \beta_k (\tilde{q}_{i, \gamma_n} - \tilde{L}_{\geq 1}^n (\tilde{q}_i)).
\end{aligned} \tag{54}$$

Similarly

$$\begin{aligned}
\beta_k (\tilde{q}_{i, \gamma_n} - \tilde{L}_{\geq 2}^n (\tilde{q}_i)) &= -\alpha_n \beta_k \tilde{q}_i \partial^{-1} \tilde{r}_i \partial_x (\tilde{q}) \tilde{q}_{i\tilde{x}} \\
&\quad + \alpha_n (\tilde{q}_{i, \sigma_k} - \tilde{L}_{\geq 1}^k (\tilde{q}_i)).
\end{aligned} \tag{55}$$

So

$$\begin{aligned}
&\alpha_n (\tilde{q}_{i, \sigma_k} - \tilde{L}_{\geq 2}^k (\tilde{q}_i)) - \beta_k (\tilde{q}_{i, \gamma_n} - \tilde{L}_{\geq 2}^n (\tilde{q}_i)) \\
&= \beta_k (\tilde{q}_{i, \gamma_n} - \tilde{L}_{\geq 1}^n (\tilde{q}_i)) - \alpha_n (\tilde{q}_{i, \sigma_k} - \tilde{L}_{\geq 1}^k (\tilde{q}_i)) = 0.
\end{aligned} \tag{56}$$

□

In this way, we present the connection between the solutions of (γ_n, σ_k) -KPH, (γ_n, σ_k) -mKPH, and (γ_n, σ_k) -HDH under the gauge transformations and reciprocal transformations. In [17], the original three hierarchies are intimately

summarized in a diagram. In our generalized system, we can find that similar results remain correct after we add some constraints on the hierarchies. These provide us with a convenient way to obtain the solutions of (γ_n, σ_k) -mKPH and (γ_n, σ_k) -HDH from the solutions of (γ_n, σ_k) -KPH.

5. Solutions for (γ_n, σ_k) -mKPH and $(2+1)$ -(γ_n, σ_k)-HDH

We first briefly recall the generalized dressing method for the (γ_n, σ_k) -KPH proposed in [15]. Let f_i, g_i satisfy

$$f_{i, \gamma_n} = \partial^n (f_i), \tag{57a}$$

$$f_{i, \sigma_k} = \partial^k (f_i),$$

$$g_{i, \gamma_n} = \partial^n (g_i),$$

$$g_{i, \sigma_k} = \partial^k (g_i), \tag{57b}$$

$$i = 1, \dots, N,$$

and let h_i be the linear combination of f_i and g_i :

$$h_i = f_i + F_i (\alpha_n \gamma_n + \beta_k \sigma_k) g_i \quad i = 1, \dots, N, \tag{58}$$

with $F_i(X)$ being a differentiable function of X , $X = \alpha_n \gamma_n + \beta_k \sigma_k$. The dressing operator W is defined by

$$W = \frac{Wr(h_1, \dots, h_N, \partial)}{Wr(h_1, \dots, h_N)}, \tag{59}$$

where

$$Wr(h_1, \dots, h_N) = \begin{vmatrix} h_1 & h_2 & \cdots & h_N \\ h'_1 & h'_2 & \cdots & h'_N \\ \vdots & \vdots & \ddots & \vdots \\ h_1^{(N-1)} & h_2^{(N-1)} & \cdots & h_N^{(N-1)} \end{vmatrix}, \tag{60}$$

$$Wr(h_1, \dots, h_N, \partial) = \begin{vmatrix} h_1 & h_2 & \cdots & h_N & 1 \\ h'_1 & h'_2 & \cdots & h'_N & \partial \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ h_1^{(N)} & h_2^{(N)} & \cdots & h_N^{(N)} & \partial^N \end{vmatrix}$$

is the Wronskian determinant. Define

$$q_i = -\dot{F}_i W(g_i),$$

$$r_i = (-1)^{N-i} \frac{Wr(h_1, \dots, \hat{h}_i, \dots, h_N)}{Wr(h_1, \dots, h_N)}, \tag{61}$$

$$i = 1, \dots, N,$$

where the hat $\hat{}$ means ruling out this term from the Wronskian determinant, $\dot{F}_i = dF_i/dX$. Then we have the following.

Theorem 8 (see [15]). Let W be defined by (59) and (58), let $L = W\partial W^{-1}$, and let q_i and r_i be given by (61); then W , L , q_i , r_i satisfy (γ_n, σ_k) -KPH ((1a), (1b), and (1c) and (3a) and (3b)).

Choose

$$q = W(1) = (-1)^N \frac{Wr(h'_1, h'_2, \dots, h'_N)}{Wr(h_1, h_2, \dots, h_N)} \quad (62)$$

as the particular eigenfunction for (4); based on Theorem 7, we have the Wronskian solution for (γ_n, σ_k) -mKPH:

$$\tilde{L} = \frac{Wr(h_1, \dots, h_N, \partial)}{Wr(h'_1, \dots, h'_N)} \partial \left[\frac{Wr(h_1, \dots, h_N, \partial)}{Wr(h'_1, \dots, h'_N)} \right]^{-1}, \quad (63a)$$

$$\tilde{q}_i = -\dot{F}_i \frac{Wr(h_1, \dots, h_N, g_i)}{Wr(h'_1, \dots, h'_N)}, \quad (63b)$$

$$\tilde{r}_i = \frac{Wr(h'_1, \dots, \hat{h}'_i, \dots, h'_N)}{Wr(h_1, \dots, h_N)}. \quad (63c)$$

Choose

$$f = W(e^{\eta_N})$$

$$= \frac{e^{\eta_N}}{Wr(h_1, \dots, h_N)} \begin{vmatrix} h_1 & h_2 & \dots & h_N & 1 \\ h'_1 & h'_2 & \dots & h'_N & \mu_N \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ h_1^{(N)} & h_2^{(N)} & \dots & h_N^{(N)} & \mu_N^N \end{vmatrix} \quad (64)$$

as another eigenfunction for (4). Based on Theorem 7, we have

$$\tilde{q} = \frac{(-1)^N e^{\eta_N}}{Wr(h'_1, \dots, h'_N)} \begin{vmatrix} h_1 & h_2 & \dots & h_N & 1 \\ h'_1 & h'_2 & \dots & h'_N & \mu_N \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ h_1^{(N)} & h_2^{(N)} & \dots & h_N^{(N)} & \mu_N^N \end{vmatrix}. \quad (65)$$

Using $\bar{x} = \tilde{q}$, $\partial_x = \tilde{q}_x \partial_{\bar{x}}$, based on Theorem 7, we have

$$\bar{L} = \tilde{L}(\bar{x}) = \frac{\begin{vmatrix} \bar{h}_1 & \bar{h}_2 & \dots & \bar{h}_N & 1 \\ \bar{q}_x \partial_{\bar{x}} \bar{h}_1 & \bar{q}_x \partial_{\bar{x}} \bar{h}_2 & \dots & \bar{q}_x \partial_{\bar{x}} \bar{h}_N & \bar{q}_x \partial_{\bar{x}} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_1 & \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_2 & \dots & \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_N & \bar{q}_x^N \partial_{\bar{x}}^N \end{vmatrix}}{\begin{vmatrix} \bar{q}_x \partial_{\bar{x}} \bar{h}_1 & \bar{q}_x \partial_{\bar{x}} \bar{h}_2 & \dots & \bar{q}_x \partial_{\bar{x}} \bar{h}_N \\ \bar{q}_x^2 \partial_{\bar{x}}^2 \bar{h}_1 & \bar{q}_x^2 \partial_{\bar{x}}^2 \bar{h}_2 & \dots & \bar{q}_x^2 \partial_{\bar{x}}^2 \bar{h}_N \\ \vdots & \vdots & \vdots & \vdots \\ \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_1 & \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_2 & \dots & \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_N \end{vmatrix}} \tilde{q}_x \partial_{\bar{x}} \quad (66)$$

$$\left[\frac{\begin{vmatrix} \bar{h}_1 & \bar{h}_2 & \dots & \bar{h}_N & 1 \\ \bar{q}_x \partial_{\bar{x}} \bar{h}_1 & \bar{q}_x \partial_{\bar{x}} \bar{h}_2 & \dots & \bar{q}_x \partial_{\bar{x}} \bar{h}_N & \bar{q}_x \partial_{\bar{x}} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_1 & \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_2 & \dots & \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_N & \bar{q}_x^N \partial_{\bar{x}}^N \end{vmatrix}}{\begin{vmatrix} \bar{q}_x \partial_{\bar{x}} \bar{h}_1 & \bar{q}_x \partial_{\bar{x}} \bar{h}_2 & \dots & \bar{q}_x \partial_{\bar{x}} \bar{h}_N \\ \bar{q}_x^2 \partial_{\bar{x}}^2 \bar{h}_1 & \bar{q}_x^2 \partial_{\bar{x}}^2 \bar{h}_2 & \dots & \bar{q}_x^2 \partial_{\bar{x}}^2 \bar{h}_N \\ \vdots & \vdots & \vdots & \vdots \\ \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_1 & \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_2 & \dots & \bar{q}_x^N \partial_{\bar{x}}^N \bar{h}_N \end{vmatrix}} \right]^{-1}$$

$$= \frac{\tilde{q}_x^{N(N+1)/2} \begin{vmatrix} \bar{h}_1 & \bar{h}_2 & \dots & \bar{h}_N & 1 \\ \bar{h}'_1 & \bar{h}'_2 & \dots & \bar{h}'_N & \partial_{\bar{x}} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \bar{h}_1^{(N)} & \bar{h}_2^{(N)} & \dots & \bar{h}_N^{(N)} & \partial_{\bar{x}}^N \end{vmatrix}}{\tilde{q}_x^{N(N+1)/2} \begin{vmatrix} \bar{h}'_1 & \bar{h}'_2 & \dots & \bar{h}'_N \\ \bar{h}_1 & \bar{h}_2 & \dots & \bar{h}_N \\ \vdots & \vdots & \vdots & \vdots \\ \bar{h}_1^{(N)} & \bar{h}_2^{(N)} & \dots & \bar{h}_N^{(N)} \end{vmatrix}} \cdot \tilde{q}_x \partial_{\bar{x}} \left[\frac{\begin{vmatrix} \bar{h}_1 & \bar{h}_2 & \dots & \bar{h}_N & 1 \\ \bar{h}'_1 & \bar{h}'_2 & \dots & \bar{h}'_N & \partial_{\bar{x}} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \bar{h}_1^{(N)} & \bar{h}_2^{(N)} & \dots & \bar{h}_N^{(N)} & \partial_{\bar{x}}^N \end{vmatrix}}{\begin{vmatrix} \bar{h}'_1 & \bar{h}'_2 & \dots & \bar{h}'_N \\ \bar{h}_1 & \bar{h}_2 & \dots & \bar{h}_N \\ \vdots & \vdots & \vdots & \vdots \\ \bar{h}_1^{(N)} & \bar{h}_2^{(N)} & \dots & \bar{h}_N^{(N)} \end{vmatrix}} \right]^{-1}$$

$$= \frac{Wr(\bar{h}_1, \dots, \bar{h}_N, \partial_{\bar{x}})}{Wr(\bar{h}'_1, \dots, \bar{h}'_N)} \cdot \tilde{q}_x \partial_{\bar{x}} \left[\frac{Wr(\bar{h}_1, \dots, \bar{h}_N, \partial_{\bar{x}})}{Wr(\bar{h}'_1, \dots, \bar{h}'_N)} \right]^{-1}, \quad (67)$$

where the elements $\bar{h}_i = h_i(\bar{x})$ and $\bar{h}'_i$ denotes $\partial \bar{h}_i / \partial \bar{x}$. Similarly, we have

$$\bar{q}_i = -\dot{F}_i \frac{Wr(\bar{h}_1, \dots, \bar{h}_N, \bar{g}_i)}{Wr(\bar{h}'_1, \dots, \bar{h}'_N)}, \quad (68a)$$

$$\bar{r}_i = -\partial_x^{-1} (\bar{q}_x \bar{r}_i) = -\partial_{\bar{x}}^{-1} (\bar{r}_i(\bar{x}))$$

$$= -\partial_{\bar{x}}^{-1} \frac{Wr(\bar{h}'_1, \dots, \bar{h}'_i, \dots, \bar{h}'_N)}{Wr(\bar{h}_1, \dots, \bar{h}_N)}, \quad (68b)$$

where $\bar{g}_i = g_i(\bar{x})$. Equations (67) and (68a) and (68b) give the Wronskian solution for (2+1)- (γ_n, σ_k) -HDH.

Let us illustrate it by solving (24a), (24b), and (24c) and (44a) and (44b). We take the solution of (57a) and (57b) as follows:

$$f_i := \exp(\lambda_i x + \lambda_i^2 y + \lambda_i^3 t) = e^{\xi_i}, \quad (69)$$

$$g_i := \exp(\mu_i x + \mu_i^2 y + \mu_i^3 t) = e^{\eta_i},$$

$$h_i := f_i + F_i(\alpha_2 y + \beta_3 t) g_i$$

$$= 2\sqrt{F_i} \exp\left(\frac{\xi_i + \eta_i}{2}\right) \cosh(\Omega_i), \quad (70)$$

$$\Omega_i = \frac{1}{2}(\xi_i - \eta_i - \ln F_i).$$

Example 9 (solutions for the mixed type of mKPESCS ((44a) and (44b))). Since $\tilde{L} = q^{-1}Lq$, then it is easy to find

$$v_0 = \frac{q_x}{q}. \quad (71)$$

For $N = 1$, we have

$$q = -\frac{\lambda_1 + \mu_1}{2} + \frac{\mu_1 - \lambda_1}{2} \tanh(\Omega_1). \quad (72)$$

The one-soliton solution for (44a) and (44b) with $N = 1$ is as follows:

$$\begin{aligned} v &= v_0 = \frac{(\lambda_1 - \mu_1)}{2} [\tanh(\Omega_1 + \theta_1) - \tanh(\Omega_1)], \\ \tilde{q}_1 &= \frac{\dot{F}_1}{2\sqrt{F_1\lambda_1\mu_1}} (\mu_1 - \lambda_1) e^{\xi_1 + \eta_1} \operatorname{sech}(\Omega_1 + \theta_1), \\ \tilde{r}_1 &= -\frac{1}{2\sqrt{F_1}} e^{-(\xi_1 + \eta_1)} \operatorname{sech} \Omega_1, \end{aligned} \quad (73)$$

where $\theta_1 = \ln \sqrt{\lambda_1/\mu_1}$.

Example 10 (solutions for the mixed type of HDESCS ((24a), (24b), and (24c))). Based on Theorem 7, we have that

$$\tilde{q} = \frac{(-1)^N e^{\eta_N}}{Wr(h'_1, \dots, h'_N)} \begin{vmatrix} h_1 & h_2 & \cdots & h_N & 1 \\ h'_1 & h'_2 & \cdots & h'_N & \mu_N \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ h_1^{(N)} & h_2^{(N)} & \cdots & h_N^{(N)} & \mu_N^{(N)} \end{vmatrix} \quad (74)$$

is the eigenfunction of the mixed type of mKPESCS and we have $\tilde{L} = \tilde{L}$; that is,

$$\begin{aligned} w\partial_{\tilde{x}} + w_0 + w_1\partial_{\tilde{x}}^{-1} + \cdots &= \partial_x + v_0 + v_1\partial_x^{-1} + \cdots \\ &= \tilde{q}_x\partial_{\tilde{x}} + v_0 + \cdots. \end{aligned} \quad (75)$$

From the above equation, we obtain

$$\begin{aligned} w &= \tilde{q}_x, \\ w_0 &= v_0. \end{aligned} \quad (76)$$

The implicit solutions for the mixed type of HDESCS ((24a), (24b), and (24c)) with $N = 1$ are

$$\tilde{x} = \tilde{q} = \frac{\lambda_1 - \mu_1}{2\sqrt{\lambda_1\mu_1}} e^{\eta_1 + \Omega_1} \operatorname{sech}(\Omega_1 + \theta_1), \quad (77a)$$

$$w = \frac{(\lambda_1 - \mu_1)}{2} e^{\eta_1 + \Omega_1} \operatorname{sech}^2(\Omega_1 + \theta_1) \cosh(\Omega_1), \quad (77b)$$

$$w_0 = \frac{(\lambda_1 - \mu_1)}{2} [\tanh(\Omega_1 + \theta_1) - \tanh(\Omega_1)], \quad (77c)$$

$$\tilde{q}_1 = \frac{\dot{F}_1}{2\sqrt{F_1\lambda_1\mu_1}} (\mu_1 - \lambda_1) e^{\xi_1 + \eta_1} \operatorname{sech}(\Omega_1 + \theta_1), \quad (77d)$$

$$\tilde{r}_1 = \frac{1}{2\sqrt{F_1}} \partial_{\tilde{x}}^{-1} e^{-(\xi_1 + \eta_1)} \operatorname{sech}(\Omega_1), \quad (77e)$$

where $\theta_1 = \ln \sqrt{\lambda_1/\mu_1}$, $\xi_1 = \lambda_1 x + \lambda_1^2 y + \lambda_1^3 t$, $\eta_1 = \mu_1 x + \mu_1^2 y + \mu_1^3 t$, $\Omega_1 = (1/2)(\xi_1 - \eta_1 - \ln F_1)$, and the relation between x and $\tilde{x}$ is given by (77a).

6. Conclusion

In this paper, a new (2+1)- (γ_n, σ_k) -HDH is proposed by introducing new time series γ_n and σ_k and adding eigenfunctions as components. The (2+1)- (γ_n, σ_k) -HDH includes (2+1)-HD hierarchy and extended HD hierarchy and contains first type and second type as well as mixed type of (2+1)-HD equation with self-consistent sources as special cases. The reduction of the (2+1)- (γ_n, σ_k) -HDH is studied and the k -constrained (2+1)- (γ_n, σ_k) -HDH is given. In the same way as (2+1)- (γ_n, σ_k) -HDH, the (γ_n, σ_k) -mKPH is formulated and its zero-curvature equation and Lax representation are presented.

The gauge transformations and reciprocal transformations between (γ_n, σ_k) -KPH, (γ_n, σ_k) -mKPH, and (2+1)- (γ_n, σ_k) -HDH are given. They help us to obtain solutions of (γ_n, σ_k) -mKPH and (2+1)- (γ_n, σ_k) -HDH. By making use of the transformations, we find the soliton solutions of the (γ_n, σ_k) -mKPH and (2+1)- (γ_n, σ_k) -HDH. Particularly, the soliton solutions for the mixed type of mKPESCS and HDESCS are obtained.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This work is supported by National Natural Science Foundation of China (11171175, 11201477, and 11301179) and the Fundamental Research Funds for the Central Universities (2014ZZD10).

References

- [1] B. A. Kupershmidt, "KdV6: an integrable system," *Physics Letters A*, vol. 372, no. 15, pp. 2634–2639, 2008.
- [2] Y. Q. Yao and Y. B. Zeng, "The bi-Hamiltonian structure and new solutions of KdV6 equation," *Letters in Mathematical Physics*, vol. 86, no. 2-3, pp. 193–208, 2008.
- [3] E. Date, M. Jimbo, M. Kashiwara, and T. Miwa, "Operator approach to the Kadomtsev-Petviashvili equation—transformation groups for soliton equations III," *Journal of the Physical Society of Japan*, vol. 50, no. 11, pp. 3806–3812, 1981.
- [4] M. Jimbo and T. Miwa, "Solitons and infinite dimensional Lie algebras," *Publications of the Research Institute for Mathematical Sciences*, vol. 19, no. 3, pp. 943–1001, 1983.
- [5] M. Sato and Y. Sato, "Soliton equations as dynamical systems on infinite dimensional Grassmann manifold," in *Nonlinear Partial Differential Equations in Applied Science*, Tokyo, 1982, pp. 259–271, North-Holland Publishing, Amsterdam, The Netherlands, 1983.
- [6] V. G. Kac and J. W. van de Leur, "The n -component KP hierarchy and representation theory," *Journal of Mathematical Physics*, vol. 44, no. 8, article 3245, 2003.

- [7] J. van de Leur, “Schlesinger-Bäcklund transformations for the N -component KP,” *Journal of Mathematical Physics*, vol. 39, no. 5, pp. 2833–2847, 1998.
- [8] H. Aratyn, E. Nissimov, and S. Pacheva, “A new ‘dual’ symmetry structure of the KP hierarchy,” *Physics Letters A*, vol. 244, no. 4, pp. 245–255, 1998.
- [9] L. A. Dickey, *Soliton Equations and Hamiltonian Systems*, World Scientific Publishing, River Edge, NJ, USA, 2nd edition, 2003.
- [10] V. K. Mel’nikov, “On equations for wave interactions,” *Letters in Mathematical Physics*, vol. 7, no. 2, pp. 129–136, 1983.
- [11] V. K. Mel’nikov, “A direct method for deriving a multisoliton solution for the problem of interaction of waves on the x, y plane,” *Communications in Mathematical Physics*, vol. 112, no. 4, pp. 639–652, 1987.
- [12] X. J. Liu, Y. B. Zeng, and R. L. Lin, “A new extended q -deformed KP hierarchy,” *Journal of Nonlinear Mathematical Physics*, vol. 15, no. 3, pp. 333–347, 2008.
- [13] R. L. Lin, X. J. Liu, and Y. B. Zeng, “A new extended q -deformed KP hierarchy,” *Journal of Nonlinear Mathematical Physics*, vol. 15, no. 3, pp. 333–347, 2008.
- [14] Y. Yao, X. Liu, and Y. Zeng, “A new extended discrete KP hierarchy and a generalized dressing method,” *Journal of Physics A: Mathematical and Theoretical*, vol. 42, no. 45, Article ID 454026, 10 pages, 2009.
- [15] Y. Q. Yao, Y. H. Huang, and Y. B. Zeng, “A new (γ_n, σ_k) -KP hierarchy and generalized dressing method,” *Journal of Nonlinear Mathematical Physics*, vol. 19, no. 4, Article ID 1250027, 13 pages, 2012.
- [16] B. G. Konopelchenko and V. G. Dubrovsky, “Some new integrable nonlinear evolution equations in 2+1 dimensions,” *Physics Letters A*, vol. 102, no. 1-2, pp. 15–17, 1984.
- [17] B. Konopelchenko and W. Oevel, “An r -Matrix approach to nonstandard classes of integrable equations,” *Publications of the Research Institute for Mathematical Sciences*, vol. 29, no. 4, pp. 581–666, 1993.
- [18] W.-X. Ma, “An extended Harry Dym hierarchy,” *Journal of Physics. A. Mathematical and Theoretical*, vol. 43, no. 16, Article ID 165202, 2010.
- [19] B. Konopelchenko, J. Sidorenko, and W. Strampp, “(1+1)-Dimensional integrable systems as symmetry constraints of (2+1)-dimensional systems,” *Physics Letters A*, vol. 157, no. 1, pp. 17–21, 1991.
- [20] Y. Cheng, “Constraints of the Kadomtsev–Petviashvili hierarchy,” *Journal of Mathematical Physics*, vol. 33, no. 11, article 3774, 1992.
- [21] W. Oevel and S. Carillo, “Squared eigenfunction symmetries for soliton equations: part I,” *Journal of Mathematical Analysis and Applications*, vol. 217, no. 1, pp. 161–178, 1998.
- [22] W. Oevel and W. Strampp, “Constrained KP hierarchy and bi-Hamiltonian structures,” *Communications in Mathematical Physics*, vol. 157, no. 1, pp. 51–81, 1993.
- [23] X. Liu, R. Lin, B. Jin, and Y. Zeng, “A generalized dressing approach for solving the extended KP and the extended mKP hierarchy,” *Journal of Mathematical Physics*, vol. 50, no. 5, Article ID 053506, 14 pages, 2009.
- [24] T. Xiao and Y. B. Zeng, “A new constrained mKP hierarchy and the generalized Darboux transformation for the mKP equation with self-consistent sources,” *Physica A*, vol. 353, no. 1–4, pp. 38–60, 2005.