Research Article

Orthogonal Projections Based on Hyperbolic and Spherical n -Simplex

Murat Savas,¹ Baki Karliga,¹ and Atakan T. Yakut²

¹Department of Mathematics, Faculty of Science, Gazi University, 06500 Ankara, Turkey

²Department of Mathematics, Art and Sciences Faculty, Nigde University, 51100 Nigde, Turkey

Correspondence should be addressed to Murat Savas; murat@math.columbia.edu

Received 4 December 2014; Accepted 23 February 2015

Academic Editor: Giorgio Kaniadakis

Copyright © 2015 Murat Savas et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Orthogonal projection along a geodesic to the chosen k -plane is introduced using edge and Gram matrix of an n -simplex in hyperbolic or spherical n -space. The distance from a point to k -plane is obtained by the orthogonal projection. It is also given the perpendicular foot from a point to k -plane of hyperbolic and spherical n -space.

1. Introduction

One of the fundamental notions in geometry is orthogonal projection and also studied extensively through the long history of mathematics and physics. There are many applications of orthogonal projection. The concept of orthogonal projection plays an important role in the scattering theory, the theory of many-body resonance, and different branches of theoretical and mathematical physics.

Let R_1^{n+1} be $(n+1)$ -dimensional vector space equipped with the scalar product $\langle, \rangle$ which is defined by

$$\langle x, y \rangle = -x_1 y_1 + \sum_{i=2}^{n+1} x_i y_i. \quad (1)$$

If the restriction of scalar product on a subspace W of R_1^{n+1} is positive definite, then the subspace W is called *space-like*; if it is positive semi-definite and degenerate, then W is called *light-like*; if W contains a time-like vector of R_1^{n+1} , then W is called *time-like*.

$S_1^n = \{x \in R_1^{n+1} \mid \langle x, x \rangle = 1\}$ is called *de Sitter n -space*. The n -dimensional unit pseudohyperbolic space is defined as

$$H_0^n = \{x \in R_1^{n+1} \mid \langle x, x \rangle = -1\}, \quad (2)$$

which has two connected components, each of which can be considered as a model for the n -dimensional hyperbolic space H^n . Throughout this paper we consider *hyperbolic n -space*

$$H^n = \{x \in H_0^n \mid x_1 > 0\}. \quad (3)$$

Hence, each pair of points p_i, p_j in H^n satisfy $\langle p_i, p_j \rangle < 0$. The *hyperbolic distance* for $p, q \in H^n$ is defined by $\text{arccosh}(-\langle p, q \rangle)$. Since each $e \in S_1^n$ determines a time-like hyperplane of R_1^{n+1} , we have hyperplane $e^\perp \cap H^n$ of H^n .

Let R^{n+1} be $(n+1)$ -dimensional vector space equipped with the scalar product $\langle, \rangle_E$ which is defined by

$$\langle x, y \rangle_E = \sum_{i=1}^{n+1} x_i y_i. \quad (4)$$

The n -dimensional unit spherical space S^n is given by

$$S^n = \{x \in R^{n+1} \mid \langle x, x \rangle_E = 1\}. \quad (5)$$

The spherical distance $d_s(p, q)$ between p and q is given by $\text{arccos}(\langle p, q \rangle_E)$.

We consider that W is a vector subspace spanned by the vectors $e_1, e_2, \dots, e_{n-k}$ in S_1^n . By using Lemma 27 of [1], one can easily see that W is $(n-k)$ -dimensional time-like subspace

and $V = e_1^\perp \cap e_2^\perp \cap \dots \cap e_{n-k}^\perp$ is $(k+1)$ -dimensional time-like subspace of R_1^{n+1} . Consequently, for $i = 1, 2, \dots, n-k$, the hyperplane $e_i^\perp \cap H^n$ intersects at the time-like k -plane $V \cap H^n$ of H^n . One can define the same tools for spherical n -space.

Let Δ be a hyperbolic or spherical n -simplex with vertices $p_1, \dots, p_{n+1}$ and let Δ_i be the face opposite to vertex p_i . Then, according to the first section of [2], we have the edge matrix M and Gram matrix G of Δ . Let $|M|$ and M_{ij} be the determinant and ij th-minor of M ; then the unit outer normal vector of Δ_i is given by

$$e_i = \frac{-\epsilon \sum_{j=1}^{n+1} M_{ij} p_j}{\sqrt{M_{ii} |M|}}, \quad i = 1, \dots, n+1, \quad (6)$$

where ϵ is the curvature of space.

The intersection of H^n with $(k+1)$ -dimensional time-like subspace is called k -dimensional plane of H^n [3]. Similarly, a k -plane of spherical space is given by the same way.

When a geodesic is drawn orthogonally from a point to a k -plane; its intersection with the k -plane is known as *perpendicular foot* on that k -plane in H^n or S^n . The length of a geodesic segment bounded by a point and its perpendicular foot is called *the distance between that point and k -plane*. The distance between a vertex and its any opposite k -face is called *k -face altitude* of an n -simplex.

The orthogonal projection to 2-plane in Euclidean space is well-known (see [4–7]). The orthogonal projection to k -plane in Euclidean space is given in [8]. The orthogonal projection taking a point in H^n and mapping it to its perpendicular foot on a hyperplane are studied in [3, 9], respectively. The distance between a point and a hyperbolic (spherical) hyperplane is introduced in [10]. The altitude of $(n-1)$ -face of hyperbolic n -simplex is given in [11].

The orthogonal projection taking a point along a geodesic and mapping to its perpendicular foot, where geodesic meets

orthogonally the chosen k -plane of projection, has not been studied. The aim of this paper is to study such orthogonal projections according to the edge matrix of a simplex in H^n or S^n .

Let m^{k+1} be the determinant of submatrix $M(k+1, \dots, n+1)$ of M and let g^{k+1} be the determinant of submatrix $G(k+1, \dots, n+1)$ of G . Suppose that m_i^j and g_i^j are the determinant of submatrices $M \begin{pmatrix} 1 & \dots & k+1 & i \\ 1 & \dots & k+1 & j \end{pmatrix}_{i,j=k+2, \dots, n+1}$ and $G \begin{pmatrix} 1 & \dots & k+1 & i \\ 1 & \dots & k+1 & j \end{pmatrix}_{i,j=k+2, \dots, n+1}$, respectively.

Lemma 1. *Let Δ be a hyperbolic or spherical n -simplex with the edge matrix M and Gram matrix G . Let M_{ii} and G_{ii} be i th minor of M and G , respectively. Then $M^{-1} = TGT$ and $G^{-1} = TMT$, where $T = [\sqrt{G_{ii}/\epsilon|G|}\delta_{ij}] = [\sqrt{M_{ii}/\epsilon|M|}\delta_{ij}]$.*

Proof. It can be seen from [12]. $\square$

Let M^{11}, M^{12}, M^{22} and G^{11}, G^{12}, G^{22} be $(k+1) \times (k+1)$, $(k+1) \times (n-k)$, $(n-k) \times (n-k)$ types submatrices of M and G , respectively. Suppose that M, G , and diagonal matrix T are partitioned as $\begin{bmatrix} M^{11} & M^{12} \\ M^{12} & M^{22} \end{bmatrix}$, $\begin{bmatrix} G^{11} & G^{12} \\ G^{12} & G^{22} \end{bmatrix}$, and $\begin{bmatrix} T^{11} & 0 \\ 0 & T^{22} \end{bmatrix}$, respectively.

Concerning Lemma 1 along with Schur complement of a symmetric matrix, we have the following lemma.

Lemma 2. *Let $S_{M^{ii}}$ and $S_{G^{ii}}$ be Schur complements of the submatrices M^{ii} and G^{ii} . Then*

$$(M^{ii})^{-1} = T^{ii} S_{G^{jj}} T^{ii}, \quad (G^{ii})^{-1} = T^{ii} S_{M^{jj}} T^{ii} \quad (7)$$

$i \neq j; \quad i, j = 1, 2.$

Proof. It is obvious that M, G are symmetric and M^{ii}, G^{ii} are invertible. Since the inverse of Schur complement of M^{11} in M is the submatrix of M^{-1} , we have

$$M^{-1} = \begin{bmatrix} (M^{11})^{-1} + (M^{11})^{-1} M^{12} (S_{M^{11}})^{-1} M^{21} (M^{11})^{-1} & -(M^{11})^{-1} M^{12} (S_{M^{11}})^{-1} \\ -(S_{M^{11}})^{-1} M^{21} (M^{11})^{-1} & (S_{M^{11}})^{-1} \end{bmatrix}. \quad (8)$$

Similarly, for the Schur complement of M^{22} , we obtain

$$M^{-1} = \begin{bmatrix} (S_{M^{22}})^{-1} & -(S_{M^{22}})^{-1} M^{12} (M^{22})^{-1} \\ -(M^{22})^{-1} M^{21} (S_{M^{22}})^{-1} & (M^{22})^{-1} + (M^{22})^{-1} M^{21} (S_{M^{22}})^{-1} M^{12} (M^{22})^{-1} \end{bmatrix}. \quad (9)$$

Then we have

$$M^{-1} = \begin{bmatrix} (S_{M^{22}})^{-1} & -(M^{11})^{-1} M^{12} (S_{M^{11}})^{-1} \\ -(M^{22})^{-1} M^{21} (S_{M^{22}})^{-1} & (S_{M^{11}})^{-1} \end{bmatrix}, \quad (10)$$

and, by the same way, we get

$$G^{-1} = \begin{bmatrix} (S_{G^{22}})^{-1} & -(G^{11})^{-1} G^{12} (S_{G^{11}})^{-1} \\ -(G^{22})^{-1} G^{21} (S_{G^{22}})^{-1} & (S_{G^{11}})^{-1} \end{bmatrix}. \quad (11)$$

Thus, we obtain the desired results using Lemma 1. $\square$

2. Orthogonal Projection to k -Plane of H^n Based on a Hyperbolic n -Simplex

If $p_1, p_2, \dots, p_{n+1}$ are vertices of any hyperbolic n -simplex Δ , then $\{p_1, p_2, \dots, p_{n+1}\}$ is a basis of R_1^{n+1} . Let W_j be a subspace spanned by $\{p_1, p_2, \dots, \hat{p}_j, \dots, p_{n+1}\}$, and let e_j be the unit outer normal to W_j , $j = 1, \dots, n+1$. Hence $\{e_1, e_2, \dots, e_{n+1}\}$ is another basis of R_1^{n+1} .

Let W^h be a k -plane which contains a k -face with vertices $p_1, p_2, \dots, p_{k+1}$ of Δ . Then the set $\{p_1, p_2, \dots, p_{k+1}\}$ is a basis of the $(k+1)$ -dimensional subspace of W in R_1^{n+1} . Since $p_1, p_2, \dots, p_{k+1}$ are vertices of Δ , the subset $\{p_1, p_2, \dots, p_{k+1}\}$ of R_1^{n+1} can be extended bases $\{p_1, p_2, \dots, p_{n+1}\}$ and $\{p_1, \dots, p_{k+1}, e_{k+2}, \dots, e_{n+1}\}$ of R_1^{n+1} . As a consequence, we see that $\{e_{k+2}, \dots, e_{n+1}\}$ is a basis of $(n-k)$ -dimensional subspace $W^\perp$.

Theorem 3. Let p be a point and let W^h be a k -plane in H^n . Then the orthogonal projection $\sigma(p)$ of p to W^h is given by

$$\sigma(p) = \left(p + \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle e_s}{|M|m^{k+1}} \right) \cdot \left(\sqrt{1 - \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle \langle p, e_s \rangle}{|M|m^{k+1}}} \right)^{-1}. \quad (12)$$

Proof. For a point $p \in H^n$, by [10, Theorem 3.11], there is a point $\vec{p} \in W$ such that $\vec{p}\vec{p} \in W^\perp$. Therefore, we can write

$$\vec{p}\vec{p} = \sum_{s=k+2}^{n+1} \lambda_s e_s. \quad (13)$$

Then, we have $-\langle p, e_t \rangle = \sum_{s=k+2}^{n+1} \lambda_s \langle e_s, e_t \rangle$, $t = k+2, \dots, n+1$. Taking

$$G^{22} = G \begin{pmatrix} k+2 & \cdots & n+1 \\ k+2 & \cdots & n+1 \end{pmatrix} = [\langle e_s, e_t \rangle]_{s,t=k+2, \dots, n+1}, \quad (14)$$

$$L = [\lambda_{k+2} \cdots \lambda_{n+1}],$$

we obtain

$$L = -(G^{22})^{-1} [\langle p, e_t \rangle]. \quad (15)$$

By Lemma 2 and (5) of [13], we see that

$$(G^{22})^{-1} = \left[\frac{\sqrt{M_{ii}M_{jj}}m_j^i}{-|M|m^{k+1}} \right]_{i,j=k+2, \dots, n+1}, \quad (16)$$

and this implies that

$$\lambda_s = \sum_{t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle}{|M|m^{k+1}}, \quad s = k+2, \dots, n+1. \quad (17)$$

Substituting (17) into (13), we obtain

$$\dot{p} = p + \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle e_s}{|M|m^{k+1}}. \quad (18)$$

By [10], there exists a unique $\sigma(p) \in W^h$ such that $\sigma(p) = c\dot{p}$. Since $\dot{p}$ is the orthogonal projection of p to W , we have

$$c = \left(\sqrt{1 - \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle \langle p, e_s \rangle}{|M|m^{k+1}}} \right)^{-1} \quad (19)$$

which completes the proof. $\square$

In case of the orthogonal projection to a hyperplane, we obtain $m_j^j = |M|$ and $m^{k+1} = M_{jj}$. Substituting these equalities into the statement of Theorem 3, we reach the result of [3, Theorem 4.1] and [14, Proposition 2.2], as follows:

$$\sigma(p) = \frac{p - \langle p, e_j \rangle e_j}{\sqrt{1 + \langle p, e_j \rangle^2}}, \quad j = 1, \dots, n+1, \quad (20)$$

where e_j is the unit normal of W_j in R_1^{n+1} . This result is also a generalization of Theorem 3 [3, 14].

Theorem 4. Let p be a point and let W^h be a k -plane in H^n . Then,

$$\cosh \xi(p, W^h) = \sqrt{1 - \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle \langle p, e_s \rangle}{|M|m^{k+1}}}, \quad (21)$$

where $\xi(p, W^h)$ is the distance between p and W^h .

Proof. Since $\langle p, \sigma(p) \rangle = -\cosh \xi(p, W^h)$, the result follows Theorem 3. $\square$

As an immediate consequence of Theorem 4, we obtain the following known result [10, Section 4].

Corollary 5. Let p be a point and let W_j^h be a hyperplane of H^n determined by e_j . Then the distance $\xi(p, W_j^h)$ between p and W_j^h is given by

$$\cosh \xi(p, W_j^h) = \sqrt{1 + \langle p, e_j \rangle^2}. \quad (22)$$

By taking p_j instead of p in (18) and using $\langle p_j, e_t \rangle = -\sqrt{|M|/M_{tt}}\delta_{jt}$, we obtain

$$\dot{p}_j = p_j + \sum_{s=k+2}^{n+1} \sqrt{\frac{M_{ss}}{|M|}} \frac{m_j^s}{m^{k+1}} e_s, \quad (23)$$

$$\langle \dot{p}_j, \dot{p}_j \rangle = -1 - \frac{m_j^j}{m^{k+1}},$$

where p_j is a vertex of Δ . The proof of the following corollary is obvious from Theorem 3.

Corollary 6. Let Δ be a hyperbolic simplex with vertices $p_1, \dots, p_{n+1}$. Then the perpendicular foot from p_j to k -face W^h is given by

$$\sigma(p_j) = \frac{p_j + \sum_{s=k+2}^{n+1} \left(\sqrt{M_{ss}/|M|} (m_j^s/m^{k+1}) \right) e_s}{\sqrt{1 + (m_j^j/m^{k+1})}}, \quad (24)$$

$$j = k+2, \dots, n+1,$$

where $p_1, \dots, p_{k+1}$ are vertices of k -face W^h .

If we replace p by p_j and use $\langle p_j, e_t \rangle = -\sqrt{|M|/M_{tt}} \delta_{jt}$, we see that

$$\langle \sigma(p_j), p_j \rangle = -\sqrt{1 + \frac{m_j^j}{m^{k+1}}}. \quad (25)$$

If we consider the last equation in the proof of Theorem 3, we see that $\cosh \xi(p_j, W^h) = \sqrt{1 + (m_j^j/m^{k+1})}$; that result is a generalization of [11, Proposition 4] to the k -face W^h of a hyperbolic n -simplex. Since m_j^j/m^{k+1} is the diagonal j th-entry of $S_{M^{11}} = [a_{ij}]$, the altitude from p_j to k -face W^h with vertices $p_1, \dots, p_{k+1}$ is given by

$$\cosh \xi(p_j, W^h) = \sqrt{1 + a_{jj}}, \quad (26)$$

where $\xi(p_j, W^h)$ is the distance between p_j and k -face W^h .

By $\hat{p}_j = p_j + \sqrt{|M|/M_{jj}} e_j$, for $(n-1)$ -face W_j^h , we have the following corollary.

Corollary 7. Let Δ be a hyperbolic simplex with vertices $p_1, \dots, p_{n+1}$. Then the perpendicular foot from p_j to $(n-1)$ -face W_j^h is given by

$$\sigma(p_j) = \frac{p_j + \sqrt{|M|/M_{jj}} e_j}{\sqrt{1 + (|M|/M_{jj})}}, \quad (27)$$

$$j = 1, \dots, n+1,$$

where $p_1, \dots, \hat{p}_j, \dots, p_{n+1}$ are vertices of W_j^h .

Using $G_{jj} = -(|G|M_{jj})/|M|$ for $j = 1, \dots, n+1$, we obtain the following known result [11, Proposition 4].

Corollary 8. Let Δ be a hyperbolic simplex with vertices $p_1, \dots, p_{n+1}$. Then the altitude $\xi(p_j, W_j^h)$ from p_j to $(n-1)$ -face W_j^h is given by

$$\cosh \xi(p_j, W_j^h) = \sqrt{1 + \frac{|M|}{M_{jj}}}, \quad j = 1, \dots, n+1, \quad (28)$$

where $p_1, \dots, \hat{p}_j, \dots, p_{n+1}$ are vertices of W_j^h .

3. Orthogonal Projections to a k -Plane of S^n Based on a Spherical n -Simplex

Let Δ be with vertices $p_1, \dots, p_{n+1}$. Then $\{p_1, \dots, p_{n+1}\}$ is a basis of R^{n+1} . If W_j is the subspace spanned by $\{p_1, \dots, \hat{p}_j, \dots, p_{n+1}\}$, then $\{e_1, \dots, e_{n+1}\}$ is another basis of R^{n+1} where e_j is the unit outer normal to W_j for $j = 1, \dots, n+1$.

Let W^s be a k -plane which contains a k -face with vertices $p_1, p_2, \dots, p_{k+1}$. Then the set $\{p_1, p_2, \dots, p_{k+1}\}$ is a basis of the $(k+1)$ -dimensional subspace W in R^{n+1} . As a consequence, we have a basis $\{e_{k+2}, \dots, e_{n+1}\}$ of $(n-k)$ -dimensional subspace $W^\perp$.

Theorem 9. Let p be a point and let W^s be a k -plane in S^n . Then the orthogonal projection $\sigma(p)$ of p to W^s is given by

$$\sigma(p) = \left(p - \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}} m_t^s \langle p, e_t \rangle_E e_s}{|M| m^{k+1}} \right) \cdot \left(\sqrt{1 - \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}} m_t^s \langle p, e_t \rangle_E \langle p, e_s \rangle_E}{|M| m^{k+1}}} \right)^{-1}. \quad (29)$$

Proof. By [10, Theorem 3.11], for $p \in S^n$, there is a $\hat{p} \in W$ such that $\vec{p\hat{p}} \in W^\perp$. Therefore, we can write

$$\vec{p\hat{p}} = \sum_{s=k+2}^{n+1} \lambda_s e_s. \quad (30)$$

Then, we have

$$\langle p, \vec{p\hat{p}} \rangle_E = \sum_{s=k+2}^{n+1} \lambda_s \langle e_s, e_t \rangle_E. \quad (31)$$

Taking

$$G^{22} = G \begin{pmatrix} k+2 & \dots & n+1 \\ k+2 & \dots & n+1 \end{pmatrix} = [\langle e_s, e_t \rangle_E]_{s,t=k+2, \dots, n+1},$$

$$L = [\lambda_{k+2} \dots \lambda_{n+1}], \quad (32)$$

we find

$$L = -\left(G^{22}\right)^{-1} [\langle p, e_t \rangle_E]. \quad (33)$$

By Lemma 2 and (5) of [13], we see that

$$\left(G^{22}\right)^{-1} = \left[\frac{\sqrt{M_{ii}M_{jj}} m_j^i}{|M| m^{k+1}} \right]_{i,j=k+2, \dots, n+1}. \quad (34)$$

This implies that

$$\lambda_s = - \sum_{t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}} m_t^s \langle p, e_t \rangle_E}{|M| m^{k+1}}, \quad (35)$$

$$s = k+2, \dots, n+1.$$

Substituting (35) into (30), we obtain

$$\dot{p} = p - \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle_E e_s}{|M| m^{k+1}}. \quad (36)$$

By [10], there exists a unique $\sigma(p) \in W^s$ such that $\sigma(p) = c\dot{p}$. Since $\dot{p}$ is the orthogonal projection of p to W , we have

$$c = \left(\sqrt{1 - \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle_E \langle p, e_s \rangle_E}{|M| m^{k+1}}} \right)^{-1} \quad (37)$$

which completes the proof. $\square$

By Theorem 9, we have

$$\sigma(p) = \frac{p - \langle p, e_j \rangle_E e_j}{\sqrt{1 - \langle p, e_j \rangle_E^2}}, \quad (38)$$

where e_j is the unit normal of the W_j in R^{n+1} .

Theorem 10. Let p be a point and let W^s be a k -plane in S^n . Then

$$\cos \theta(p, W^s) = \sqrt{1 - \sum_{s,t=k+2}^{n+1} \frac{\sqrt{M_{ss}M_{tt}}m_t^s \langle p, e_t \rangle_E \langle p, e_s \rangle_E}{|M| m^{k+1}}}, \quad (39)$$

where $\theta(p, W^s)$ is the distance between p and W^s .

By taking p_j instead of p and using $\langle p_j, e_j \rangle_E = -\sqrt{|M|/M_{jj}}$ in (36), we obtain

$$\dot{p}_j = p_j + \sum_{s=k+2}^{n+1} \sqrt{\frac{M_{ss}}{|M|}} \frac{m_j^s}{m^{k+1}} e_s, \quad (40)$$

$$\langle \dot{p}_j, \dot{p}_j \rangle_E = 1 - \frac{m_j^j}{m^{k+1}}, \quad j = k+2, \dots, n+1,$$

where p_j is a vertex of Δ . Hence, we have the following corollary.

Corollary 11. Let Δ be a spherical n -simplex with vertices $p_1, \dots, p_{n+1}$; then the perpendicular foot from p_j to k -face W^s is given by

$$\sigma(p_j) = \left(p_j + \sum_{s=k+2}^{n+1} \sqrt{\frac{M_{ss}}{|M|}} \frac{m_j^s}{m^{k+1}} e_s \right) \cdot \left(\sqrt{1 - \frac{m_j^j}{m^{k+1}}} \right)^{-1}, \quad j = k+2, \dots, n+1, \quad (41)$$

where $p_1, \dots, p_{k+1}$ are vertices of W^s .

Let $\theta(p_j, W^s)$ be the altitude from the vertex p_j to the k -face W^s with vertices $p_1, \dots, p_{k+1}$ for $j = k+2, \dots, n+1$. Then $\theta(p_j, W^s)$ is given by

$$\cos \theta(p_j, W^s) = \sqrt{1 - \frac{m_j^j}{m^{k+1}}}. \quad (42)$$

By equality (5) in [13], the jj -th-entry of the Schur complement $S_{M^{11}} = [b_{ij}]$ satisfies $b_{jj} = m_j^j/m^{k+1}$.

Let W_j^s be the $(n-1)$ -face with vertices $p_1, \dots, \hat{p}_j, \dots, p_{n+1}$ of Δ . Then, we have

$$\dot{p}_j = p_j + \sqrt{\frac{|M|}{M_{jj}}} e_j, \quad (43)$$

$$\langle p_j, e_j \rangle_E = -\sqrt{\frac{|M|}{M_{jj}}}, \quad j = 1, \dots, n+1.$$

The proof of the following corollary is obtained by using the above equations.

Corollary 12. Let Δ be a spherical simplex with vertices $p_1, \dots, p_{n+1}$. Then the perpendicular foot from p_j to $(n-1)$ -face W_j^s is given by

$$\sigma(p_j) = \frac{p_j + \sqrt{|M|/M_{jj}} e_j}{\sqrt{1 - (|M|/M_{jj})}}, \quad j = 1, \dots, n+1, \quad (44)$$

where $p_1, \dots, \hat{p}_j, \dots, p_{n+1}$ are vertices of W_j^s .

Corollary 13. Let Δ be a spherical simplex with vertices $p_1, \dots, p_{n+1}$. Then the altitude $\theta(p_j, W_j^s)$ from p_j to $(n-1)$ -face W_j^s is given by

$$\cos \theta(p_j, W_j^s) = \sqrt{1 - \frac{|M|}{M_{jj}}}, \quad j = 1, \dots, n+1, \quad (45)$$

where $p_1, \dots, \hat{p}_j, \dots, p_{n+1}$ are vertices of W_j^s .

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] B. O'Neill, *Semi-Riemannian Geometry*, Academic Press, New York, NY, USA, 1983.
- [2] B. Karlıga and A. T. Yakut, "Vertex angles of a simplex in hyperbolic space H^n ," *Geometriae Dedicata*, vol. 120, pp. 49–58, 2006.
- [3] S. Izumiya and F. Tari, "Projections of hypersurfaces in the hyperbolic space to hyperhorospheres and hyperplanes," *Revista Matemática Iberoamericana*, vol. 24, no. 3, pp. 895–920, 2008.
- [4] J. H. Rieger, "Families of maps from the plane to the plane," *Journal of the London Mathematical Society. Second Series*, vol. 36, no. 2, pp. 351–369, 1987.

- [5] V. I. Arnol'd, "Singularities of systems of rays," *Russian Mathematical Surveys*, vol. 38, no. 2, pp. 87–176, 1983.
- [6] J. W. Bruce, "Generic geometry, transversality and projections," *Journal of the London Mathematical Society*, vol. 49, no. 1, pp. 183–194, 1994.
- [7] J. W. Bruce, P. J. Giblin, and F. Tari, "Families of surfaces: height functions and projections to planes," *Mathematica Scandinavica*, vol. 82, no. 2, pp. 165–185, 1998.
- [8] B. Karol, *Multidimensional Analytic Geometry*, vol. 50, Instytut Matematyczny Polakiej Akad. Nauk. To., Warsaw, Poland, 1969, translation from the Polish by Halina Spalinska.
- [9] L. J. Alías, T. Kurose, and G. Solanes, "Hadamard-type theorems for hypersurfaces in hyperbolic spaces," *Differential Geometry and its Applications*, vol. 24, no. 5, pp. 492–502, 2006.
- [10] E. B. Vinberg, *Geometry II*, vol. 29 of *Encyclopaedia of Mathematical Sciences*, Springer, New York, NY, USA, 1993.
- [11] A. D. Mednykh and M. G. Pashkevich, "Elementary formulas for a hyperbolic tetrahedron," *Siberian Mathematical Journal*, vol. 47, no. 4, pp. 687–695, 2006.
- [12] B. Karliĝa, "Edge matrix of hyperbolic simplices," *Geometriae Dedicata*, vol. 109, pp. 1–6, 2004.
- [13] R. A. Brualdi and H. Schneider, "Determinantal identities: gauss, schur, cauchy, sylvestre, kronecker, jacobi, binet, laplace, muir, and cayley," *Linear Algebra and its Applications*, vol. 52-53, pp. 769–791, 1983.
- [14] A. Ushijima, "The tilt formula for generalized simplices in hyperbolic space," *Discrete & Computational Geometry*, vol. 28, no. 1, pp. 19–27, 2002.