Research Article

On a System of Generalized Mixed Equilibrium Problems Involving Variational-Like Inequalities in Banach Spaces: Existence and Algorithmic Aspects

H. Mahdioui¹ and O. Chadli²

¹ Department of Mathematics, Ibnou Zohr University, Agadir 80000, Morocco

² Department of Economics, Ibnou Zohr University, B.P. 8658 Poste Dakhla, Agadir 80000, Morocco

Correspondence should be addressed to O. Chadli, ochadli@gmail.com

Received 28 January 2012; Accepted 15 June 2012

Academic Editor: Abdellah Bnouhachem

Copyright © 2012 H. Mahdioui and O. Chadli. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the existence and the algorithmic aspect of a System of Generalized Mixed Equilibrium Problems involving variational-like inequalities (SGMEPs) in the setting of Banach spaces. The approach adopted is based on the auxiliary principle technique and arguments from generalized convexity. A new existence theorem for the auxiliary problem is established; this leads us to generate an algorithm which converges strongly to a solution of (SGMEP) under weaker assumptions. When the study is reduced to the setting of reflexive Banach spaces, then it can be more relaxed by dropping the coercivity condition. The results obtained in this paper are new and improve some recent studies in this field.

1. Introduction and the Problem Statement

Let K be a nonempty closed convex subset of a Banach space X , and let $F : K \times K \rightarrow \mathbb{R}$ be a real-valued bifunction. By equilibrium problem, in short (EP), we mean the following problem:

$$(EP) \text{ Find } \bar{u} \in K \text{ such that } F(\bar{u}, v) \geq 0 \quad \forall v \in K. \quad (1.1)$$

Equilibrium problems are suitable and common format for investigation of various applied problems arising in economics, mathematical physics, transportation, communication systems, engineering, and other fields. Moreover, equilibrium problems are closely related with other general problems in nonlinear analysis, such as fixed points, game theory, variational inequality, and optimization problems, see [1–4]. Various kinds of iterative

algorithms to solve equilibrium problems and variational inequalities have been developed by many authors. There exists a vast literature on the approximation solvability of equilibrium problems and nonlinear variational inequalities using projection type methods, proximal-type methods, or resolvent operator type methods, see [5–11]. We observe that the projection method and its variant forms cannot be applied for constructing iterative algorithms of mixed variational-like inequalities or mixed equilibrium problems involving variational-like inequalities. This fact motivated many authors to develop the auxiliary principle technique to study the existence and algorithm of solutions for variational-like inequalities and its extensions to mixed equilibrium problems, see [12–15]. Kazmi and Khan [16] studied a system of generalized variational-like inequality problems in Hilbert spaces by using the auxiliary principle technique. Recently, Ding and Wang [17] and Ding [18] introduced new iterative algorithms for solving some class of system of generalized mixed variational-like inequalities and system of mixed equilibrium problems involving variational-like inequalities.

In this paper, we study the existence and the algorithmic aspect of a System of Generalized Mixed Equilibrium Problems involving variational-like inequalities (SGMEPs) in the setting of Banach spaces. A new existence theorem for the auxiliary problem is established, this leads us to generate an algorithm which converges strongly to a solution of (SGMEP) under weaker assumptions. When the study is reduced to the setting of reflexive Banach spaces, then it can be more relaxed by dropping the coercivity condition. The results obtained in this paper are new and improve some recent studies in this field.

Throughout this paper, let $I = \{1, 2\}$ be an index set. For each $i \in I$, let X_i be a Banach space, X_i^* its dual, $\langle \cdot, \cdot \rangle_i$ is the duality pairing between X_i^* and X_i , $\|\cdot\|_i$ denotes the norm in X_i and X_i^* , and let K_i be a nonempty closed convex subset of X_i . We will denote by $\mathcal{CB}(X_i^*)$ the family of all nonempty closed and bounded subsets of X_i^* .

For each $i \in I$, let $R_i : K_1 \rightrightarrows \mathcal{CB}(X_1^*)$ and $A_i : K_2 \rightrightarrows \mathcal{CB}(X_2^*)$ be multivalued mappings, $N_i : X_1^* \times X_2^* \rightarrow X_i^*$ and $\eta_i : X_i \times X_i \rightarrow X_i$ be single-valued mappings, $F_i, \varphi_i : K_i \times K_i \rightarrow \mathbb{R}$ be two real-valued functions, and let $w_i \in X_i^*$. We consider the following System of Generalized Mixed Equilibrium Problem involving variational-like inequalities (SGMEP).

Find $(\bar{x}_1, \bar{x}_2) \in K_1 \times K_2$, $(\bar{u}_1, \bar{v}_1) \in R_1(\bar{x}_1) \times A_1(\bar{x}_2)$, and $(\bar{u}_2, \bar{v}_2) \in R_2(\bar{x}_1) \times A_2(\bar{x}_2)$ such that

$$\begin{aligned} F_1(\bar{x}_1, y_1) + \langle N_1(\bar{u}_1, \bar{v}_1) - w_1, \eta_1(y_1, \bar{x}_1) \rangle_1 + \varphi_1(\bar{x}_1, y_1) - \varphi_1(\bar{x}_1, \bar{x}_1) &\geq 0 \quad \forall y_1 \in K_1; \\ F_2(\bar{x}_2, y_2) + \langle N_2(\bar{u}_2, \bar{v}_2) - w_2, \eta_2(y_2, \bar{x}_2) \rangle_2 + \varphi_2(\bar{x}_2, y_2) - \varphi_2(\bar{x}_2, \bar{x}_2) &\geq 0 \quad \forall y_2 \in K_2. \end{aligned} \quad (1.2)$$

1.1. Some Special Cases

- (1) If for each $i \in I$, $X_i = X_i^* = H_i$ is a Hilbert space, $K_i = H_i$, $F_i \equiv 0$, $w_i = 0$ and for each $(x, y) \in H_1 \times H_2$, $R_i(x) = x$ and $A_i(y) = y$, then problem (1.2) reduces to the problem of finding $(\bar{x}_1, \bar{x}_2) \in H_1 \times H_2$ such that

$$\begin{aligned} \langle N_1(\bar{x}_1, \bar{x}_2), \eta_1(y_1, \bar{x}_1) \rangle_1 + \varphi_1(\bar{x}_1, y_1) - \varphi_1(\bar{x}_1, \bar{x}_1) &\geq 0 \quad \forall y_1 \in H_1; \\ \langle N_2(\bar{x}_1, \bar{x}_2), \eta_2(y_2, \bar{x}_2) \rangle_2 + \varphi_2(\bar{x}_2, y_2) - \varphi_2(\bar{x}_2, \bar{x}_2) &\geq 0 \quad \forall y_2 \in H_2. \end{aligned} \quad (1.3)$$

Problem (1.3) has been introduced and studied by Kazmi and Khan in [16].

- (2) If for each $i \in I$, X_i is a reflexive Banach space, $K_i = X_i$ and $F_i \equiv 0$, then problem (1.2) reduces to the problem of finding $(\bar{x}_1, \bar{x}_2) \in X_1 \times X_2$, $(\bar{u}_1, \bar{v}_1) \in R_1(\bar{x}_1) \times A_1(\bar{x}_2)$ and $(\bar{u}_2, \bar{v}_2) \in R_2(\bar{x}_1) \times A_2(\bar{x}_2)$ such that

$$\begin{aligned} \langle N_1(\bar{u}_1, \bar{v}_1) - w_1, \eta_1(y_1, \bar{x}_1) \rangle_1 + \varphi_1(\bar{x}_1, y_1) - \varphi_1(\bar{x}_1, \bar{x}_1) &\geq 0 \quad \forall y_1 \in X_1; \\ \langle N_2(\bar{u}_2, \bar{v}_2) - w_2, \eta_2(y_2, \bar{x}_2) \rangle_2 + \varphi_2(\bar{x}_2, y_2) - \varphi_2(\bar{x}_2, \bar{x}_2) &\geq 0 \quad \forall y_2 \in X_2. \end{aligned} \quad (1.4)$$

Problem (1.4) has been considered recently by Ding and Wang in [17].

- (3) In finite dimensional spaces a particular problem of (1.2) has been considered by Mordukhovich et al. in [19] for finding a common solution of a variational inequality problem and an equilibrium problem by an approach based on an hybrid proximal point algorithm.

The organization of the paper is as follows. In Section 2, we give some definitions and preliminary results that we will need in the sequel. We introduce in Section 3 an auxiliary principle, for the problem studied and we show that under some suitable conditions this problem has a unique solution. Further, by using the auxiliary principle we consider an algorithm to approach the solution of the main problem studied in this paper and discuss its convergence. Finally, we end the paper by some commentaries on the approach used and we give some comparisons with some known results in this direction.

2. Preliminaries

In this section, we present some basic concepts, properties, and notations that we will consider in the development of our work. Let X be a real Banach space with norm $\|\cdot\|$, X^* its dual space, and $\langle \cdot, \cdot \rangle$ denote the duality pairing between X^* and X , and let K be a nonempty closed convex subset of X . For a finite subset A of K , we denote by $\text{co}(A)$ the convex hull of A . Let $\mathcal{CB}(X^*)$ be the family of all nonempty closed and bounded subsets of X^* , and let $\mathcal{H}(\cdot, \cdot)$ be the Hausdorff metric on $\mathcal{CB}(X^*)$ defined for all $A, D \in \mathcal{CB}(X^*)$ by

$$\mathcal{H}(A, D) = \max \left\{ \sup_{a \in A} d(a, D), \sup_{b \in D} d(b, A) \right\}, \quad (2.1)$$

where $d(a, D) = \inf_{d \in D} \|a - d\|$.

Definition 2.1. A mapping $T : X \rightarrow X^*$ is said to be

- (i) monotone if and only if $\langle T(x) - T(y), x - y \rangle \geq 0$ for all $x, y \in X$;
- (ii) α -strongly monotone with $\alpha > 0$ if and only if

$$\langle T(x) - T(y), x - y \rangle \geq \alpha \|x - y\|^2 \quad \forall x, y \in X; \quad (2.2)$$

- (iii) L -Lipschitz continuous if there exists $L > 0$ such that

$$\|T(x) - T(y)\| \leq L \|x - y\| \quad \forall x, y \in X. \quad (2.3)$$

Definition 2.2. Let $T : X \rightarrow X^*$ be a bounded linear operator. T is said to be δ -strongly positive if there exists $\delta > 0$ such that

$$\langle T(x), x \rangle \geq \delta \|x\|^2, \quad \forall x \in X. \quad (2.4)$$

Remark 2.3. One can easily see that if a bounded linear operator $T : X \rightarrow X^*$ is δ -strongly positive, then it is δ -strongly monotone and $\|T\|$ -Lipschitz continuous, where $\|T\|$ is the operator norm of T .

Definition 2.4. Let X_1 and X_2 be two Banach spaces with respective norm $\|\cdot\|_1$ and $\|\cdot\|_2$. A mapping $A_1 : X_1 \times X_2 \rightarrow X_1$ is said to be (β_1, ξ_2) -Lipschitz continuous if there exist constants $\beta_1, \xi_2 > 0$ such that

$$\|A_1(x_1, y_1) - A_1(x_2, y_2)\|_1 \leq \beta_1 \|x_1 - x_2\|_1 + \xi_2 \|y_1 - y_2\|_2 \quad \forall (x_1, y_1), (x_2, y_2) \in X_1 \times X_2. \quad (2.5)$$

Definition 2.5. A mapping $\eta : X \times X \rightarrow X$ is said to be

(i) affine in the second argument if

$$\eta(y, tx + (1-t)z) = t\eta(y, x) + (1-t)\eta(y, z), \quad \forall t \in [0, 1], \quad \forall x, y, z \in K; \quad (2.6)$$

(ii) τ -Lipschitz continuous if there exists a constant $\tau > 0$ such that

$$\|\eta(x, y)\| \leq \tau \|x - y\|, \quad \forall x, y \in X. \quad (2.7)$$

Definition 2.6. A real-valued function $\varphi : K \times K \rightarrow \mathbb{R}$ is said to be skew-symmetric if for all $x, y \in K$,

$$\varphi(x, x) - \varphi(x, y) - \varphi(y, x) + \varphi(y, y) \geq 0. \quad (2.8)$$

The skew-symmetric functions have certain properties, see [20], which can be regarded as analogs of the conditions governing the gradient monotonicity and nonnegativity of the second derivative of convex functions.

Definition 2.7. Let $F : K \times K \rightarrow \mathbb{R}$ be a real-valued bifunction. Then,

(i) F is said to be monotone if

$$F(x, y) + F(y, x) \leq 0, \quad \forall x, y \in K; \quad (2.9)$$

(ii) F is said to be α -strongly monotone if there exists $\alpha > 0$ such that

$$F(x, y) + F(y, x) \leq -\alpha \|x - y\|^2, \quad \forall x, y \in K; \quad (2.10)$$

- (iii) F is said to be upper-hemicontinuous if, for all $x, y, z \in K$, the mapping $g : [0, 1] \rightarrow \mathbb{R}$ defined by $g(t) = F(tz + (1-t)x, y)$ is upper-semicontinuous.

Remark 2.8. Clearly, strong monotonicity of F implies monotonicity of F .

Now, let S_F denote the solution set of the equilibrium problem (EP) associated to a bifunction F , that is, $S_F = \{\bar{x} \in K : F(\bar{x}, y) \geq 0 \ \forall y \in K\}$. In many situations, we usually need some more information on the structure of the solution set S_F when it is nonempty. As a preliminary result that we will need in the sequel, the following Lemma gives some sufficient conditions which insure that S_F is convex and closed.

Lemma 2.9. *Let X be a Banach space, and let K be a closed convex subset of X . Let $F : K \times K \rightarrow \mathbb{R}$ be a real-valued bifunction such that $F(x, x) \geq 0$ for all $x \in K$. Assume that*

- (i) *F is monotone and upper-hemicontinuous;*
- (ii) *for each $x \in K$ fixed, the function $y \mapsto F(x, y)$ is convex and lower semicontinuous. Then, the solution set S_F is convex and closed whenever it is nonempty.*

Proof. Assume that $S_F \neq \emptyset$. Let $\bar{u}, \bar{v} \in S_F$ and for $\lambda \in]0, 1[$ set $x_\lambda = \lambda\bar{u} + (1-\lambda)\bar{v} \in K$. Let us show that $x_\lambda \in S_F$. Since $\bar{u}, \bar{v} \in S_F$, it follows

$$F(\bar{u}, y) \geq 0, \quad F(\bar{v}, y) \geq 0 \quad \forall y \in K. \quad (2.11)$$

From the monotonicity of F , one deduces

$$F(y, \bar{u}) \leq 0, \quad F(y, \bar{v}) \leq 0 \quad \forall y \in K. \quad (2.12)$$

Therefore, from the convexity of F with respect to its second argument, it follows

$$F(y, x_\lambda) \leq 0, \quad \forall y \in K. \quad (2.13)$$

Now for $t \in]0, 1[$ and $y \in K$, set $x_t = ty + (1-t)x_\lambda \in K$. Since F is an equilibrium bifunction and F is convex with respect to its second argument, it follows that

$$0 \leq F(x_t, x_t) \leq tF(x_t, y) + (1-t)F(x_t, x_\lambda). \quad (2.14)$$

Taking account of relation (2.13) with $y = x_t$, it follows that $tF(x_t, y) \geq 0$ for all $y \in K$, and therefore since $t > 0$

$$F(x_t, y) \geq 0, \quad \forall y \in K. \quad (2.15)$$

Since F is upper-hemicontinuous, it follows that

$$F(x_\lambda, y) \geq \limsup_{t \rightarrow 0} F(x_t, y) \geq 0. \quad (2.16)$$

Therefore, $x_\lambda \in S_F$, and hence S_F is convex.

Now, let us show that S_F is closed. To this aim, let $\{x_n\} \subset S_F$ such that $x_n \rightarrow \bar{x}$. Let us show that $\bar{x} \in S_F$. One has $F(x_n, y) \geq 0$ for all $y \in K$, since F is monotone, it follows that

$$F(y, x_n) \leq 0, \quad \forall y \in K. \quad (2.17)$$

From the lower semicontinuity of F with respect to the second argument, it follows that

$$F(y, \bar{x}) \leq \liminf_{n \rightarrow +\infty} F(y, x_n) \leq 0. \quad (2.18)$$

By a similar argument as above, one can easily show that $F(\bar{x}, y) \geq 0$ for all $y \in K$. Therefore, $\bar{x} \in S_F$, which completes the proof. $\square$

In the sequel, we will need the following result that we present in a more general setting and for which we refer to [21].

Lemma 2.10. *Let X be a Hausdorff topological vector space and K a closed convex subset of X . Consider two real bifunctions $\Phi, \Psi : K \times K \rightarrow \mathbb{R}$ such that*

- (i) *for each $x, y \in K$, if $\Psi(x, y) \leq 0$ then $\Phi(x, y) \leq 0$;*
- (ii) *for each fixed $x \in K$, the function $y \in K \mapsto \Phi(x, y)$ is lower semicontinuous on every compact subset of K ;*
- (iii) *for each finite subset A of K , one has*

$$\sup_{y \in \text{co}(A)} \min_{x \in A} \Psi(x, y) \leq 0; \quad (2.19)$$

- (iv) *coercivity: there exists a nonempty compact convex subset C of X such that either (a) or (b) in the following holds*

- (a) *for all $y \in K \setminus C$, there exists $x \in C$ such that $\Phi(x, y) > 0$;*
- (b) *there exists $x_0 \in C$ such that $\forall y \in K \setminus C$ $\Psi(x_0, y) > 0$.*

Then, there exists $\bar{y} \in C$ such that $\Phi(x, \bar{y}) \leq 0$ for all $x \in K$. Furthermore, the set of solution is compact.

Remark 2.11. (1) Condition (iii) in Lemma 2.10 is much more related to convexity assumptions of the bifunction Ψ , see Proposition 2.12 in the following.

(2) If K is compact, then condition (iv) in Lemma 2.10 can be dropped.

The following proposition gives some sufficient conditions which insure condition (iii) in Lemma 2.10.

Proposition 2.12. *Suppose that*

- (i) $\Psi(x, x) \leq 0$ for each $x \in K$;
- (ii) for each $y \in K$ fixed, the set $\{x \in K : \Psi(x, y) > 0\}$ is convex.

Then, condition (iii) of Lemma 2.10 is satisfied.

Proof. Suppose by contradiction that condition (iii) of Lemma 2.10 is not satisfied. Then, there exist $x_1, \dots, x_n \in K$ and $\lambda_1, \dots, \lambda_n \geq 0$, with $\sum_{i=1}^n \lambda_i = 1$, such that $\Psi(x_i, \sum_{j=1}^n \lambda_j x_j) > 0$.

Therefore, by setting $y = \sum_{j=1}^n \lambda_j x_j$, it follows from (ii) that $\Psi(y, y) > 0$, which contradicts assumption (i). $\square$

As a consequence of Lemma 2.10, we obtain the following result on existence of mixed equilibrium problem that we will need in the sequel. We will include its proof for completeness.

Lemma 2.13. *Let X be a Banach space and K a closed convex subset of X . Let $f, g : K \times K \rightarrow \mathbb{R}$ be two real bifunctions such that*

- (i) $f(x, x) \geq 0$ for all $x \in K$; f is monotone and upper-hemicontinuous; for each $x \in K$ fixed, the function $y \mapsto f(x, y)$ is convex and lower semicontinuous;
- (ii) $g(x, x) \geq 0$ for all $x \in K$; for each $y \in K$ fixed, the function $x \mapsto g(x, y)$ is upper-semicontinuous; for each $x \in K$ fixed, the function $y \mapsto g(x, y)$ is convex and lower semicontinuous;
- (iii) *coercivity: there exists a nonempty compact convex subset C of X and $y_0 \in C \cap K$ such that*

$$f(x, y_0) + g(x, y_0) < 0 \quad \text{for each } x \in K \setminus C. \quad (2.20)$$

Then, there exists $\bar{x} \in C \cap K$ such that

$$f(\bar{x}, y) + g(\bar{x}, y) \geq 0 \quad \forall y \in K. \quad (2.21)$$

Furthermore, the solution set $S_{f,g}$ of the mixed equilibrium problem (2.21) is compact and convex.

Proof. The proof is a direct consequence of Lemma 2.10 by setting

$$\Phi(x, y) = f(x, y) - g(y, x), \quad \Psi(x, y) = -g(y, x) - f(y, x). \quad (2.22)$$

$\square$

Remark 2.14. (1) Lemma 2.13 is in fact a slight extension of Theorem 1 in [2] and Theorem 4.5 in [21] where the equilibrium condition $f(x, x) = g(x, x) = 0$ has been relaxed by assuming $f(x, x) \geq 0$ and $g(x, x) \geq 0$ for all $x \in K$.

(2) If X is a reflexive Banach space endowed with its weak topology $\sigma(X, X^*)$, then the coercivity condition (iii) in Lemma 2.13 can be replaced by the following condition:

$$(iii)' \text{ there exists } y_0 \in K \text{ such that } \lim_{\|x - y_0\| \rightarrow +\infty} g(x, y_0) / \|x - y_0\| = -\infty.$$

Ended, let $r_1 > 0$, and set $B(y_0, r_1) = \{x \in X : \|x - y_0\| \leq r_1\}$. One has $B(y_0, r_1)$ a convex and $\sigma(X, X^*)$ -compact subset of X . Since X is a reflexive Banach space, $f(y_0, \cdot)$ is lower semicontinuous and $B(y_0, r)$ is weakly compact, it follows that there exists $\alpha_0 \in \mathbb{R}$ such that $f(y_0, y) > \alpha_0$ for all $y \in B(y_0, r_1)$. Let $x \in K \setminus B(y_0, r_1)$, and set

$$y = \frac{r_1}{\|x - y_0\|} x + \left(1 - \frac{r_1}{\|x - y_0\|}\right) y_0. \quad (2.23)$$

Since $f(y_0, \cdot)$ is convex, $y \in B(y_0, r_1)$ and $f(y_0, y_0) \geq 0$, one deduces

$$\alpha_0 \leq \frac{r_1}{\|x - y_0\|} f(y_0, x) + \left(1 - \frac{r_1}{\|x - y_0\|}\right) f(y_0, y_0) \quad \forall x \in K \setminus B(y_0, r_1). \quad (2.24)$$

It follows that

$$f(y_0, x) \geq \frac{\|x - y_0\|}{r_1} (\alpha_0 - f(y_0, y_0)) + f(y_0, y_0). \quad (2.25)$$

Thus,

$$f(y_0, x) \geq \frac{\|x - y_0\|}{r_1} (\alpha_0 - f(y_0, y_0)), \quad \forall x \in K \setminus B(y_0, r_1). \quad (2.26)$$

Since f is monotone, it follows from relation (2.26) that for all

$$f(x, y_0) + g(x, y_0) \leq g(x, y_0) - \frac{\alpha_0 - f(y_0, y_0)}{r_1} \|x - y_0\|. \quad (2.27)$$

Since $g(x, y_0)/\|x - y_0\| \rightarrow -\infty$ when $\|x - y_0\| \rightarrow +\infty$, then there exists $r_2 > 0$ such that for $x \in K$ with $\|x - y_0\| > r_2$ one has

$$g(x, y_0) - \frac{\alpha_0 - f(y_0, y_0)}{r_1} \|x - y_0\| < 0. \quad (2.28)$$

Take $r = \max\{r_1, r_2\}$ and set $C = \{x \in X : \|x - y_0\| \leq r\}$, then from relations (2.27) and (2.28), one deduces that for each $x \in K \setminus C$ one has

$$f(x, y_0) + g(x, y_0) < 0. \quad (2.29)$$

Hence, condition (iii) in Lemma 2.13 is satisfied.

We end this section by the following result related to the Hausdorff metric that we will need in the sequel and for which we refer to [22].

Lemma 2.15. *Let E be a complete metric space and $R : E \rightrightarrows \mathcal{CB}(E)$ a set-valued mapping. Then, for any given $\varepsilon > 0$ and any given $x, y \in E$ and $u \in R(x)$, there exists $v \in R(y)$ such that*

$$d(u, v) \leq (1 + \varepsilon) \mathcal{H}(R(x), R(y)). \quad (2.30)$$

3. Approximation by an Auxiliary Principle

In order to get approximate solutions for the system (1.2) of generalized mixed equilibrium problem involving generalized mixed variational-like inequality problems (SGMEP), we

consider the following *auxiliary problem*: for $i \in I = \{1, 2\}$ and for given mappings $T_i : X_i \rightarrow X_i^*$, $\rho_i > 0$, $(x_1, x_2) \in K_1 \times K_2$, $(u_1, v_1) \in R_1(x_1) \times A_1(x_2)$, and $(u_2, v_2) \in R_2(x_1) \times A_2(x_2)$,

$$(AP) \begin{cases} \text{find } (z_1, z_2) \in K_1 \times K_2 \text{ such that for all } y_i \in K_i \\ \rho_i [F_i(z_i, y_i) + \langle N_i(u_i, v_i) - w_i, \eta_i(y_i, z_i) \rangle_i + \varphi_i(z_i, y_i) - \varphi_i(z_i, z_i)] \\ + \langle T_i(y_i - z_i), z_i - x_i \rangle_i \geq 0. \end{cases} \quad (3.1)$$

In this section, we give some existence results of solutions for the auxiliary problem (AP). The results obtained will be needed in the sequel to generate a unified algorithm to approach solutions of the system (1.2) under some weaker assumptions in comparison with some known results in literature.

Theorem 3.1. *For each $i \in I = \{1, 2\}$, let X_i be a Banach space and K_i a nonempty closed convex subset of X_i , $F_i, \varphi_i : K_i \times K_i \rightarrow \mathbb{R}$ two real-valued bifunctions, $T_i : X_i \rightarrow X_i^*$ a bounded linear operator, $\rho_i > 0$ and $\eta_i : X_i \times X_i \rightarrow X_i$ single-valued mappings such that*

- (i) $F_i(x, x) \geq 0$ for all $x \in K_i$; F_i is monotone and upper-hemicontinuous; for each $x \in K_i$ fixed, the function $y \mapsto F_i(x, y)$ is convex and lower semicontinuous;
- (ii) η_i is affine in the first argument and continuous in the second argument such that

$$\eta_i(x, y) + \eta_i(y, x) = 0, \quad \forall x, y \in K_i; \quad (3.2)$$

- (iii) φ_i is skew symmetric and continuous; for each $y \in K_i$ fixed, the function $x \mapsto \varphi_i(x, y)$ is convex;
- (iv) T_i is δ_i -strongly positive;
- (v) *coercivity*: for each $x_i \in K_i$, $u_i \in X_1^*$ and $v_i \in X_2^*$, there exists a nonempty compact convex subset C_{x_i, u_i, v_i}^i of X_i and $y_i^0 \in C_{x_i, u_i, v_i}^i \cap K_i$ such that $\forall z_i \in K_i \setminus C_{x_i, u_i, v_i}^i$

$$\rho_i [F_i(z_i, y_i^0) + \langle N_i(u_i, v_i) - w_i, \eta_i(y_i^0, z_i) \rangle_i + \varphi_i(z_i, y_i^0) - \varphi_i(z_i, z_i)] + \langle T_i(y_i^0 - z_i), z_i - x_i \rangle_i < 0. \quad (3.3)$$

Then, the auxiliary problem (AP) has a unique solution.

Proof. The proof of this lemma is a direct application of Lemma 2.13 by considering, for each $i \in I = \{1, 2\}$, the bifunctions $f_i, g_i : K_i \times K_i \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} f_i(z_i, y_i) &= \rho_i [F_i(z_i, y_i) + \langle N_i(u_i, v_i) - w_i, \eta_i(y_i, z_i) \rangle_i + \varphi_i(z_i, y_i) - \varphi_i(z_i, z_i)] \\ g_i(z_i, y_i) &= \langle T_i(y_i) - T_i(z_i), z_i - x_i \rangle_i, \end{aligned} \quad (3.4)$$

where $(x_1, x_2) \in K_1 \times K_2$, $(u_1, v_1) \in R_1(x_1) \times A_1(x_2)$, and $(u_2, v_2) \in R_2(x_1) \times A_2(x_2)$ are given. We need only to show that the solution is unique. To this aim, suppose that problem (AP) has two solutions z_i^1 and z_i^2 , then for $i \in I$ and for all $z \in K_i$, we have

$$\rho_i \left[F_i(z_i^1, z) + \langle N_i(u_i, v_i) - w_i, \eta_i(z, z_i^1) \rangle_i + \varphi_i(z_i^1, z) - \varphi_i(z_i^1, z_i^1) \right] + \langle T_i(z) - T_i(z_i^1), z_i^1 - x_i \rangle_i \geq 0, \quad (3.5)$$

$$\rho_i \left[F_i(z_i^2, z) + \langle N_i(u_i, v_i) - w_i, \eta_i(z, z_i^2) \rangle_i + \varphi_i(z_i^2, z) - \varphi_i(z_i^2, z_i^2) \right] + \langle T_i(z) - T_i(z_i^2), z_i^2 - x_i \rangle_i \geq 0. \quad (3.6)$$

Take $z = z_i^2$ in relation (3.5) and $z = z_i^1$ in relation (3.6) and adding the two inequalities, one obtain

$$\begin{aligned} & \rho_i \left[F_i(z_i^1) + F_i(z_i^2, z_i^1) + \langle N_i(u_i, v_i) - w_i, \eta_i(z_i^1, z_i^2) + \eta_i(z_i^2, z_i^1) \rangle_i \varphi_i(z_i^1, z_i^2) \right. \\ & \left. + \varphi_i(z_i^2, z_i^1) - \varphi_i(z_i^1, z_i^1) - \varphi_i(z_i^2, z_i^2) \right] \geq \langle T_i(z_i^2 - z_i^1), z_i^2 - z_i^1 \rangle_i. \end{aligned} \quad (3.7)$$

Since for each $i \in I$, F_i is monotone, φ_i is skew symmetric, and T_i is δ_i -strongly positive, it follows that

$$0 \geq \langle T_i(z_i^2 - z_i^1), z_i^2 - z_i^1 \rangle_i \geq \delta_i \|z_i^1 - z_i^2\|_i^2. \quad (3.8)$$

Therefore, $z_i^1 = z_i^2$ for $i \in I = \{1, 2\}$, which completes the proof. $\square$

Theorem 3.2. For each $i \in I = \{1, 2\}$, let X_i be a reflexive Banach space and K_i a nonempty closed convex subset of X_i , $F_i, \varphi_i : K_i \times K_i \rightarrow \mathbb{R}$ two real-valued bifunctions, $T_i : X_i \rightarrow X_i^*$ a bounded linear operator, $\rho_i > 0$ and $\eta_i : X_i \times X_i \rightarrow X_i$ single-valued mappings such that

- (i) $F_i(x, x) \geq 0$ for all $x \in K_i$; F_i is monotone and upper-hemicontinuous; for each $x \in K_i$ fixed, the function $y \mapsto F_i(x, y)$ is convex and lower semicontinuous;
- (ii) η_i is affine in the first argument and continuous in the second argument such that

$$\eta_i(x, y) + \eta_i(y, x) = 0, \quad \forall x, y \in K_i; \quad (3.9)$$

- (iii) φ_i is skew symmetric and continuous; for each $y \in K_i$ fixed, the function $x \mapsto \varphi_i(x, y)$ is convex;
- (iv) T_i is δ_i -strongly positive linear operator.

Then, the auxiliary problem (AP) has a unique solution.

Proof. For each $i \in I$ and given $(x_1, x_2) \in K_1 \times K_2$, $(u_1, v_1) \in R_1(x_1) \times A_1(x_2)$, and $(u_2, v_2) \in R_2(x_1) \times A_2(x_2)$, define the following bifunctions $f_i, g_i : K_i \times K_i \rightarrow \mathbb{R}$ by

$$\begin{aligned} f_i(z_i, y_i) &= \rho_i \left[F_i(z_i, y_i) + \langle N_i(u_i, v_i) - w_i, \eta_i(y_i, z_i) \rangle_i + \varphi_i(y_i, z_i) - \varphi_i(z_i, z_i) \right], \\ g_i(z_i, y_i) &= \langle T_i(y_i) - T_i(z_i), z_i - x_i \rangle_i. \end{aligned} \quad (3.10)$$

One can easily see that conditions (i)–(iii) above imply conditions (i) and (ii) of Lemma 2.13. In order to get the conclusion, we need only to show that the coercivity condition (iii) of Lemma 2.13 is satisfied. To this aim, taking into account Remark 2.14 (2), we need only to show that for some $v_i^0 \in K_i$ one has $g_i(u_i, v_i^0)/\|v_i^0 - u_i\|_i \rightarrow -\infty$ when $\|v_i^0 - u_i\|_i \rightarrow +\infty$. Let $v_i^0 \in K_i$. Then,

$$\begin{aligned}
 g_i(u_i, v_i^0) &= \langle T_i(v_i^0) - T_i(u_i), u_i - x_i \rangle_i \\
 &= \langle T_i(v_i^0 - u_i), u_i - v_i^0 \rangle_i + \langle T_i(v_i^0 - u_i), v_i^0 - x_i \rangle_i \\
 &= -\langle T_i(v_i^0 - u_i), v_i^0 - u_i \rangle_i + \langle T_i(v_i^0 - u_i), v_i^0 - x_i \rangle_i \\
 &\leq -\delta \|v_i^0 - u_i\|_i^2 + \|T_i\| \|v_i^0 - u_i\|_i \|v_i^0 - x_i\|_i.
 \end{aligned} \tag{3.11}$$

Therefore,

$$\frac{g_i(u_i, v_i^0)}{\|v_i^0 - u_i\|_i} \leq -\delta \|v_i^0 - u_i\|_i + \|T_i\| \|v_i^0 - x_i\|_i. \tag{3.12}$$

It follows that $g_i(u_i, v_i^0)/\|v_i^0 - u_i\|_i \rightarrow -\infty$ when $\|v_i^0 - u_i\|_i \rightarrow +\infty$ which completes the proof. $\square$

Remark 3.3. Theorem 3.2 improves recent results given by Ding [18, Theorem 3.1] since the bifunction F_i is not needed to be δ_i -Lipschitz continuous and weakly upper semicontinuous with respect to the first argument. We mention also that all the results obtained in [18] are under the assumption $\text{int}\{y_i \in K_i, \varphi_i(y_i, y_i) < \infty\} = \text{int}K_i \neq \emptyset$; in our approach, this assumption is not needed.

Theorem 3.2 shows that the auxiliary problem (AP) has a unique solution; we can define the following general iterative method to approach the solution of system (1.2) of generalized mixed equilibrium problem involving variational-like inequalities (SGMEP).

Algorithm 3.4. For a given $(x_1^0, x_2^0) \in K_1 \times K_2$, $(u_1^0, v_1^0) \in R_1(x_1^0) \times A_1(x_2^0)$, and $(u_2^0, v_2^0) \in R_2(x_1^0) \times A_2(x_2^0)$. By Theorem 3.1, the auxiliary problem (AP) has a unique solution $(x_1^1, x_2^1) \in K_1 \times K_2$, that is, for each $i \in I$, we have

$$\begin{aligned}
 &\rho_i^1 \left[F_i(x_i^1, y_i) + \langle N_i(u_i^0, v_i^0) - w_i, \eta_i(y_i, x_i^1) \rangle_i + \varphi_i(x_i^1, y_i) - \varphi_i(x_i^1, x_i^1) \right] \\
 &\quad + \langle T_i(y_i) - T_i(x_i^1), x_i^1 - x_i^0 \rangle_i \geq 0 \quad \forall y_i \in K_i.
 \end{aligned} \tag{3.13}$$

Since for each $i \in I$, $u_i^0 \in R_i(x_i^0) \in \mathcal{CB}(X_1^*)$, and $v_i^0 \in A_i(x_i^0) \in \mathcal{CB}(X_2^*)$, by Lemma 2.15, there exist $u_i^1 \in R_i(x_i^1)$ and $v_i^1 \in A_i(x_i^1)$ such that

$$\begin{aligned}
 \|u_i^1 - u_i^0\|_1 &\leq (1 + 1)\mathcal{H}_1(R_i(x_i^1), R_i(x_i^0)) \\
 \|v_i^1 - v_i^0\|_2 &\leq (1 + 1)\mathcal{H}_2(A_i(x_i^1), A_i(x_i^0)),
 \end{aligned} \tag{3.14}$$

where $\mathcal{H}_1$ and $\mathcal{H}_2$ are the Hausdorff metrics on $\mathcal{CB}(X_1^*)$ and $\mathcal{CB}(X_2^*)$, respectively.

By using Theorem 3.1 again, the auxiliary problem (AP) has a unique solution $(x_1^2, x_2^2) \in K_1 \times K_2$ such that

$$\begin{aligned} & \rho_i^2 \left[F_i(x_i^2, y_i) + \left\langle N_i(u_i^1, v_i^1) - w_i, \eta_i(y_i, x_i^2) \right\rangle_i + \varphi_i(x_i^2, y_i) - \varphi_i(x_i^2, x_i^2) \right] \\ & + \left\langle T_i(y_i) - T_i(x_i^2), x_i^2 - x_i^1 \right\rangle_i \geq 0, \quad \forall y_i \in K_i. \end{aligned} \quad (3.15)$$

By induction, we can construct an iterative algorithm to compute the approximate solution for the system (1.2) as follows: for given $(x_1^0, x_2^0) \in K_1 \times K_2$, $(u_1^0, v_1^0) \in R_1(x_1^0) \times A_1(x_2^0)$, and $(u_2^0, v_2^0) \in R_2(x_1^0) \times A_2(x_2^0)$, there exist sequences $\{x_1^n\}$, $\{x_2^n\}$, $\{u_1^n\}$, $\{u_2^n\}$, $\{v_1^n\}$, and $\{v_2^n\}$ such that for each $i \in I$

$$\begin{aligned} & \|u_i^{n+1} - u_i^n\|_1 \leq \left(1 + \frac{1}{n+1}\right) \mathcal{L}_1(R_i(x_1^{n+1}), R_i(x_1^n)), \quad u_i^n \in R_i(x_1^n), \\ & \|v_i^{n+1} - v_i^n\|_2 \leq \left(1 + \frac{1}{n+1}\right) \mathcal{L}_2(A_i(x_2^{n+1}), A_i(x_2^n)), \quad v_i^n \in A_i(x_2^n), \\ & \rho_i^{n+1} \left[F_i(x_i^{n+1}, y_i) + \left\langle N_i(u_i^n, v_i^n) - w_i, \eta_i(y_i, x_i^{n+1}) \right\rangle_i + \varphi_i(x_i^{n+1}, y_i) - \varphi_i(x_i^{n+1}, x_i^{n+1}) \right] \\ & + \left\langle T_i(y_i) - T_i(x_i^{n+1}), x_i^{n+1} - x_i^n \right\rangle_i \geq 0 \quad \forall y_i \in K_i, \quad n = 0, 1, 2, \dots \end{aligned} \quad (3.16)$$

The following convergence analysis is presented for the algorithm above.

Theorem 3.5. *Under the hypotheses of Theorem 3.2, further assume that for each $i \in I$,*

- (i) F_i is σ_i -strongly monotone and upper-hemicontinuous; N_i is (β_i, ξ_i) -mixed Lipschitz continuity; R_i is k_i - $\mathcal{L}_1$ -Lipschitz continuous, and A_i is μ_i - $\mathcal{L}_2$ -Lipschitz continuous;
- (ii) η_i is τ_i -Lipschitz continuous;
- (iii) the sequence $\{\rho^n\}_{n \in \mathbb{N}}$ of positive real numbers is increasing and $\lim_{n \rightarrow +\infty} \rho_n = +\infty$.

Furthermore, assume that the following condition holds:

$$(C_1) \begin{cases} \Lambda = \max\{\theta_1 + \theta_2, \vartheta_1 + \vartheta_2\} < 1, \text{ where} \\ \theta_1 = \frac{\tau_1 \beta_1 k_1}{\sigma_1}, \quad \theta_2 = \frac{\tau_2 \beta_2 k_2}{\sigma_2}, \\ \vartheta_1 = \frac{\tau_1 \xi_1 \mu_1}{\sigma_1}, \quad \vartheta_2 = \frac{\tau_2 \xi_2 \mu_2}{\sigma_2}. \end{cases} \quad (3.17)$$

Then, the sequences $\{x_1^n\}$, $\{x_2^n\}$, $\{u_1^n\}$, $\{u_2^n\}$, $\{v_1^n\}$, and $\{v_2^n\}$ generated by Algorithm 3.4 converge strongly to $\bar{x}_1$, $\bar{x}_2$, $\bar{u}_1$, $\bar{v}_1$, $\bar{u}_2$, and $\bar{v}_2$, respectively, where $(\bar{u}_1, \bar{v}_1) \in T_1(\bar{x}_1) \times A_1(\bar{x}_2)$, $(\bar{u}_2, \bar{v}_2) \in T_2(\bar{x}_1) \times A_2(\bar{x}_2)$, and $(\bar{x}_1, \bar{x}_2, \bar{u}_1, \bar{v}_1, \bar{u}_2, \bar{v}_2)$ is a solution of the System of Generalized Mixed Equilibrium Problem involving variational-like in equalities (1.2).

Proof. By the definition of Algorithm 3.4, we have for $i \in I$,

$$\begin{aligned} & \rho_i^n \left[F_i(x_i^n, y_i) + \left\langle N_i(u_i^{n-1}, v_i^{n-1}) - w_i, \eta_i(y_i, x_i^n) \right\rangle_i + \varphi_i(x_i^n, y_i) - \varphi_i(x_i^n, x_i^n) \right] \\ & + \left\langle T_i(y_i) - T_i(x_i^n), x_i^n - x_i^{n-1} \right\rangle_i \geq 0 \quad \forall y_i \in K_i, \end{aligned} \quad (3.18)$$

$$\begin{aligned} & \rho_i^{n+1} \left[F_i(x_i^{n+1}, y_i) + \left\langle N_i(u_i^n, v_i^n) - w_i, \eta_i(y_i, x_i^{n+1}) \right\rangle_i + \varphi_i(x_i^{n+1}, y_i) - \varphi_i(x_i^{n+1}, x_i^{n+1}) \right] \\ & + \left\langle T_i(y_i) - T_i(x_i^{n+1}), x_i^{n+1} - x_i^n \right\rangle_i \geq 0 \quad \forall y_i \in K_i. \end{aligned} \quad (3.19)$$

By considering $y_i = x_i^{n+1}$ in relation (3.18) and $y_i = x_i^n$ in relation (3.19), and taking into account $\eta_i(x_i^{n+1}, x_i^n) = -\eta_i(x_i^n, x_i^{n+1})$, one obtains by adding the two inequalities

$$\begin{aligned} & F_i(x_i^{n+1}, x_i^n) + F_i(x_i^n, x_i^{n+1}) + \left\langle N_i(u_i^{n-1}, v_i^{n-1}) - N_i(u_i^n, v_i^n), \eta_i(x_i^{n+1}, x_i^n) \right\rangle_i \\ & + \varphi_i(x_i^n, x_i^{n+1}) + \varphi_i(x_i^{n+1}, x_i^n) - \varphi_i(x_i^n, x_i^n) - \varphi_i(x_i^{n+1}, x_i^{n+1}) \\ & + \frac{1}{\rho_i^n} \left\langle T_i(x_i^{n+1} - x_i^n), x_i^n - x_i^{n-1} \right\rangle_i \\ & - \frac{1}{\rho_i^{n+1}} \left\langle T_i(x_i^n - x_i^{n+1}), x_i^n - x_i^{n+1} \right\rangle_i \geq 0. \end{aligned} \quad (3.20)$$

Since F_i is σ_i -monotone, φ_i is skew-symmetric, and T_i is δ_i -strongly positive and $\|T_i\|$ -Lipschitz continuous, it follows from relation (3.20) that

$$\begin{aligned} & \left\| N_i(u_i^{n-1}, v_i^{n-1}) - N_i(u_i^n, v_i^n) \right\|_i \left\| \eta_i(x_i^{n+1}, x_i^n) \right\| + \frac{\|T_i\|}{\rho_i^n} \left\| x_i^{n+1} - x_i^n \right\|_i \left\| x_i^n - x_i^{n-1} \right\|_i \\ & \geq \left[\sigma_i + \frac{\delta_i}{\rho_i^{n+1}} \right] \left\| x_i^n - x_i^{n+1} \right\|_i^2. \end{aligned} \quad (3.21)$$

Note that N_1 is (β_1, ξ_1) -mixed Lipschitz continuous, R_1 is k_1 - $\mathcal{L}_1$ -Lipschitz continuous, and A_1 is μ_1 - $\mathcal{L}_2$ -Lipschitz continuous, it follows by Algorithm 3.4

$$\begin{aligned} \left\| N_i(u_i^{n-1}, v_i^{n-1}) - N_i(u_i^n, v_i^n) \right\|_i & \leq \beta_1 \left\| u_i^{n-1} - u_i^n \right\|_1 + \xi_1 \left\| v_i^{n-1} - v_i^n \right\|_2 \\ & \leq \beta_1 \left(1 + \frac{1}{n} \right) \mathcal{L}_1(R_1(x_1^{n-1}), R_1(x_1^n)) \\ & \quad + \xi_1 \left(1 + \frac{1}{n} \right) \mathcal{L}_2(R_1(x_2^{n-1}), R_1(x_2^n)) \\ & \leq \beta_1 k_1 \left(1 + \frac{1}{n} \right) \left\| x_1^{n-1} - x_1^n \right\|_1 + \xi_1 \mu_1 \left(1 + \frac{1}{n} \right) \left\| x_2^{n-1} - x_2^n \right\|_2. \end{aligned} \quad (3.22)$$

Since η_1 is τ_1 -Lipschitz continuous, it follows from relation (3.21) considered for $i = 1$ that

$$\begin{aligned} \left[\sigma_1 + \frac{\delta_1}{\rho_1^{n+1}} \right] \|x_1^n - x_1^{n+1}\|_1 &\leq \left[\tau_1 \beta_1 k_1 \left(1 + \frac{1}{n} \right) + \frac{\|T_1\|}{\rho_1^n} \right] \|x_1^{n-1} - x_1^n\|_1 \\ &\quad + \tau_1 \xi_1 \mu_1 \left(1 + \frac{1}{n} \right) \|x_2^{n-1} - x_2^n\|_2. \end{aligned} \quad (3.23)$$

Hence,

$$\|x_1^n - x_1^{n+1}\|_1 \leq \theta_{1n} \|x_1^{n-1} - x_1^n\|_1 + \vartheta_{1n} \|x_2^{n-1} - x_2^n\|_2, \quad (3.24)$$

with

$$\theta_{1n} = \frac{[\rho_1^{n+1} \tau_1 \beta_1 k_1 (1 + 1/n) + \rho_1^{n+1} \|T_1\| / \rho_1^n]}{[\rho_1^{n+1} \sigma_1 + \delta_1]}, \quad \vartheta_{1n} = \frac{\rho_1^{n+1} \tau_1 \xi_1 \mu_1 (1 + 1/n)}{[\rho_1^{n+1} \sigma_1 + \delta_1]}. \quad (3.25)$$

Similarly, by the assumptions on F_2 , N_2 , η_2 , R_2 , A_2 , and T_2 , it follows from relation (3.21) considered for $i = 2$ that

$$\|x_2^n - x_2^{n+1}\|_2 \leq \theta_{2n} \|x_1^{n-1} - x_1^n\|_1 + \vartheta_{2n} \|x_2^{n-1} - x_2^n\|_2, \quad (3.26)$$

with

$$\theta_{2n} = \frac{[\rho_2^{n+1} \tau_2 \beta_2 k_2 (1 + 1/n)]}{[\rho_2^{n+1} \sigma_2 + \delta_2]}, \quad \vartheta_{2n} = \frac{\rho_2^{n+1} \tau_2 \xi_2 \mu_2 (1 + 1/n) + \rho_2^{n+1} \|T_2\| / \rho_2^n}{[\rho_2^{n+1} \sigma_2 + \delta_2]}. \quad (3.27)$$

Adding the inequalities (3.24) and (3.26), we obtain

$$\begin{aligned} \|x_1^n - x_1^{n+1}\|_1 + \|x_2^n - x_2^{n+1}\|_2 &\leq (\theta_{1n} + \theta_{2n}) \|x_1^{n-1} - x_1^n\|_1 + (\vartheta_{1n} + \vartheta_{2n}) \|x_2^{n-1} - x_2^n\|_2 \\ &\leq \Lambda_n \left(\|x_1^{n-1} - x_1^n\|_1 + \|x_2^{n-1} - x_2^n\|_2 \right), \end{aligned} \quad (3.28)$$

where $\Lambda_n = \max\{\theta_{1n} + \theta_{2n}, \vartheta_{1n} + \vartheta_{2n}\}$.

On $X_1 \times X_2$, let us consider the norm $\|\cdot\|_*$ defined by $\|(x, y)\|_* = \|x\|_1 + \|y\|_2$ for all $(x, y) \in X_1 \times X_2$, that is, $(X_1 \times X_2, \|\cdot\|_*)$ is a Banach space. It follows from relation (3.28) that

$$\|(x_1^n, x_2^n) - (x_1^{n+1}, x_2^{n+1})\|_* \leq \Lambda_n \|(x_1^{n-1}, x_2^{n-1}) - (x_1^n, x_2^n)\|_*. \quad (3.29)$$

Taking account of the assumptions, it is easy to see that $\Lambda_n \rightarrow \Lambda$ as $n \rightarrow \infty$. From condition (C_1) (relation (3.17)), we know that $0 < \Lambda < 1$. Hence, there exist $\Lambda_0 \in (0, 1)$ and $n_0 > 0$ such that $\Lambda_n \leq \Lambda_0 < 1$ for all $n \geq n_0$. Therefore, it follows from (3.29) that

$$\|(x_1^n, x_2^n) - (x_1^{n+1}, x_2^{n+1})\|_* \leq \Lambda_0 \|(x_1^{n-1}, x_2^{n-1}) - (x_1^n, x_2^n)\|_*. \quad (3.30)$$

This implies that $\{(x_1^n, x_2^n)\}$ is a Cauchy sequence in $X_1 \times X_2$. Thus, (x_1^n, x_2^n) converges strongly to some $(\bar{x}_1, \bar{x}_2) \in K_1 \times K_2$.

By Algorithm 3.4 and the Lipschitz continuity assumption on R_1, R_2, A_1 , and A_2 , we have

$$\begin{aligned} \|u_i^{n+1} - u_i^n\|_1 &\leq \left(1 + \frac{1}{n+1}\right) \mathcal{L}_1(R_i(x_1^{n+1}), R_i(x_1^n)) \leq k_i \left(1 + \frac{1}{n+1}\right) \|x_1^{n+1} - x_1^n\|_1, \\ \|v_i^{n+1} - v_i^n\|_2 &\leq \left(1 + \frac{1}{n+1}\right) \mathcal{L}_2(A_i(x_2^{n+1}), A_i(x_2^n)) \leq \mu_i \left(1 + \frac{1}{n+1}\right) \|x_2^{n+1} - x_2^n\|_2. \end{aligned} \quad (3.31)$$

It follows, for each $i \in I$, that $\{u_i^n\}$ is a Cauchy sequence in X_1^* and $\{v_i^n\}$ is a Cauchy sequence in X_2^* . Thus, there exists $(\bar{u}_i, \bar{v}_i) \in X_1^* \times X_2^*$ such that (u_i^n, v_i^n) converges strongly to $(\bar{u}_i, \bar{v}_i)$.

Noting that $u_1^n \in R_1(x_1^n)$, it follows that

$$\begin{aligned} d(\bar{u}_1, R_1(x_1^n)) &\leq \|\bar{u}_1 - u_1^n\|_1 + d(u_1^n, R_1(x_1^n)) + \mathcal{L}_1(R_1(x_1^n), R_1(\bar{x}_1)) \\ &\leq \|\bar{u}_1 - u_1^n\|_1 + k_1 \|\bar{x}_1 - x_1^n\|_1 \longrightarrow 0 \quad \text{as } n \longrightarrow \infty. \end{aligned} \quad (3.32)$$

Hence, we must have $\bar{u}_1 \in R_1(\bar{x}_1)$. Similarly, one can show that $\bar{u}_2 \in R_2(\bar{x}_1)$, $\bar{v}_1 \in A_1(\bar{x}_2)$, and $\bar{v}_2 \in A_2(\bar{x}_2)$. By Algorithm 3.4, we have that, for all $i \in I$,

$$\begin{aligned} \rho_i^{n+1} &\left[F_i(x_i^{n+1}, y_i) + \left\langle N_i(u_i^n, v_i^n) - w_i, \eta_i(y_i, x_i^{n+1}) \right\rangle_i + \varphi(x_i^{n+1}, y_i) - \varphi(x_i^{n+1}, x_i^{n+1}) \right] \\ &+ \left\langle T_i(y_i) - T_i(x_i^{n+1}), x_i^{n+1} - x_i^n \right\rangle \geq 0 \quad \forall y_i \in K_i, \quad n = 0, 1, 2, \dots \end{aligned} \quad (3.33)$$

Since N_i is (β_i, ξ_i) -mixed Lipschitz continuous and η_i is continuous in the second argument, one has, for each $y_i \in K_i$,

$$\begin{aligned} &\left| \left\langle N_i(u_i^n, v_i^n) - w_i, \eta_i(y_i, x_i^{n+1}) \right\rangle_i - \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i \right| \\ &\leq \left| \left\langle N_i(u_i^n, v_i^n) - N_i(\bar{u}_i, \bar{v}_i), \eta_i(y_i, x_i^{n+1}) \right\rangle_i \right| + \left| \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, x_i^{n+1}) - \eta_i(y_i, \bar{x}_i) \right\rangle_i \right| \\ &\leq \|N_i(u_i^n, v_i^n) - N_i(\bar{u}_i, \bar{v}_i)\| \|\eta_i(y_i, x_i^{n+1})\| + \left| \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, x_i^{n+1}) - \eta_i(y_i, \bar{x}_i) \right\rangle_i \right|. \end{aligned} \quad (3.34)$$

Hence,

$$\left| \left\langle N_i(u_i^n, v_i^n) - w_i, \eta_i(y_i, x_i^{n+1}) \right\rangle_i - \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i \right| \longrightarrow 0 \quad \text{as } n \longrightarrow \infty. \quad (3.35)$$

From the monotonicity of F_i and the linearity of T_i , one has

$$\begin{aligned} & \left\langle N_i(u_i^n, v_i^n) - w_i, \eta_i(y_i, x_i^{n+1}) \right\rangle_i - \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i \\ & + \psi_i(x_i^{n+1}, y_i) - \psi_i(x_i^{n+1}, x_i^{n+1}) + \frac{\|T_i\|}{\rho_i^{n+1}} \|y_i - x_i^{n+1}\| \|x_i^{n+1} - x_i^n\| \\ & \geq F_i(y_i, x_i^{n+1}) - \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i \quad \forall y_i \in K_i, \quad \forall n = 0, 1, 2, \dots \end{aligned} \quad (3.36)$$

Taking into account result (3.35) and since, for each $i \in I$, $F_i(y, \cdot)$ is a lower semicontinuous function, ψ_i is continuous, and the sequence $\{x_i^n\}$ converges strongly to $\bar{x}_i$, it follows from relation (3.36) by passing to the limit when n goes to infinity that

$$\psi_i(y_i, \bar{x}_i) - \psi_i(\bar{x}_i, \bar{x}_i) \geq F_i(y_i, \bar{x}_i) - \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i. \quad (3.37)$$

Now, for each $i \in I$ and for $t \in (0, 1]$ and $y_i \in K_i$, set $y_i(t) = ty_i + (1-t)\bar{x}_i$. Since K_i is convex, then $y_i(t) \in K_i$ for $t \in (0, 1]$; it follows that

$$\left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i(t), \bar{x}_i) \right\rangle_i + \psi_i(y_i(t), \bar{x}_i) - \psi_i(\bar{x}_i, \bar{x}_i) \geq F_i(y_i(t), \bar{x}_i). \quad (3.38)$$

Taking into account the convexity of $\psi_i(\cdot, \bar{x}_i)$, the fact that N_i is affine with respect to the first argument and $\eta_i(\bar{x}_i, \bar{x}_i) = 0$, one has

$$F_i(y_i(t), \bar{x}_i) \leq t(\psi_i(y_i, \bar{x}_i) - \psi_i(\bar{x}_i, \bar{x}_i)) + t \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i. \quad (3.39)$$

On the other hand, since $F_i(x_i, x_i) = 0$ for each $x_i \in K_i$ and $F_i(y_i, \cdot)$ is convex, it follows that

$$0 \leq F_i(y_i(t), y_i(t)) \leq tF_i(y_i(t), y_i) + (1-t)F(y_i(t), \bar{x}_i). \quad (3.40)$$

Hence, from relation (3.39), one has

$$0 \leq t[F_i(y_i(t), y_i) + (1-t)(\psi_i(y_i, \bar{x}_i) - \psi_i(\bar{x}_i, \bar{x}_i) + \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i)]. \quad (3.41)$$

Therefore, for all $t \in]0, 1]$, one has

$$0 \leq F_i(y_i(t), y_i) + (1-t)(\psi_i(y_i, \bar{x}_i) - \psi_i(\bar{x}_i, \bar{x}_i) + \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i). \quad (3.42)$$

Since for each $y_i \in K_i$, $x \mapsto F_i(x, y)$ is upper hemicontinuous, it follows by passing to limit when $t \rightarrow 0$ in the previous inequality that

$$F_i(\bar{x}_i, y_i) + \left\langle N_i(\bar{u}_i, \bar{v}_i) - w_i, \eta_i(y_i, \bar{x}_i) \right\rangle_i + \psi_i(\bar{x}_i, y_i) - \psi_i(\bar{x}_i, \bar{x}_i) \geq 0 \quad \forall y_i \in K_i, \quad (3.43)$$

which completes the proof. $\square$

4. Commentaries

In conclusion, the approach used in this paper lets us improve and extend some new results in literature related to the problem studied. To be more precise,

- (1) Theorems 3.2 and 3.5 improve recent results given by Ding [18, Theorem 3.1, Theorem 4.1], since the bifunction F_i is not needed to be δ_i -Lipschitz continuous nor weakly upper semicontinuous with respect to the first argument in all steps of the procedure, firstly when dealing with the existence of solutions for the auxiliary problem and secondly when studying the convergence of the algorithm;
- (2) we mention also that all the results obtained in [18] are under the assumption $\text{int}\{y_i \in K_i, \psi_i(y_i, y_i) < \infty\} = \text{int} K_i \neq \emptyset$; in our approach, this assumption is not needed;
- (3) in the special case where $F_i \equiv 0$, the results obtained improve those obtained by Ding and Wang [17], since the assumptions on the bifunctions ψ_i have been relaxed.

References

- [1] C. Baiocchi and A. Capelo, *Variational and Quasivariational Inequalities: Applications to Free Boundary Problems*, John Wiley & Sons, New York, NY, USA, 1984.
- [2] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [3] I. Konnov, "Generalized monotone equilibrium problems and variational inequalities," in *Handbook of Generalized Convexity and Generalized Monotonicity, Nonconvex Optimization and Applications*, N. Hadjisavvas, S. Komlosi, and S. Schaible, Eds., pp. 559–618, Springer, New York, NY, USA, 2005.
- [4] M. Patriksson, *Nonlinear Programming and Variational Inequality Problems: a unified approach*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1999.
- [5] A. Bnouhachem and M. A. Noor, "A new iterative method for variational inequalities," *Applied Mathematics and Computation*, vol. 182, no. 2, pp. 1673–1682, 2006.
- [6] P. L. Combettes and S. A. Hirstoaga, "Equilibrium programming using proximal-like algorithms," *Mathematical Programming*, vol. 78, no. 1, pp. 29–41, 1997.
- [7] X.-F. He, J. Chen, and Z. He, "Generalized projection method for a variational inequality system with different mapping in Banach spaces," *Computers & Mathematics with Applications*, vol. 58, no. 7, pp. 1391–1396, 2009.
- [8] B.-S. He, Z.-H. Yang, and X.-M. Yuan, "An approximate proximal-extragradient type method for monotone variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 300, no. 2, pp. 362–374, 2004.
- [9] A. Moudafi and M. Théra, "Proximal and dynamical approaches to equilibrium problems," in *Lecture Notes in Economics and Mathematical Systems*, vol. 477, pp. 187–201, Springer, Berlin, Germany, 1999.
- [10] M. A. Noor, "Projection-proximal methods for general variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 318, no. 1, pp. 53–62, 2006.
- [11] R. U. Verma, "General convergence analysis for two-step projection methods and applications to variational problems," *Applied Mathematics Letters*, vol. 18, no. 11, pp. 1286–1292, 2005.
- [12] X. P. Ding, "Existence of solutions and an algorithm for mixed variational-like inequalities in Banach spaces," *Journal of Optimization Theory and Applications*, vol. 127, no. 2, pp. 285–302, 2005.
- [13] X. P. Ding, J.-C. Yao, and L.-C. Zeng, "Existence and algorithm of solutions for generalized strongly nonlinear mixed variational-like inequalities in Banach spaces," *Computers & Mathematics with Applications*, vol. 55, no. 4, pp. 669–679, 2008.
- [14] G. Mastroeni, "On auxiliary principle for equilibrium problems," in *Equilibrium Problems and Variational Models*, P. Daniele, F. Giannessi, and A. Maugeri, Eds., pp. 289–298, Kluwer, Dordrecht, The Netherlands, 2003.
- [15] L. C. Zeng, S. Schaible, and J. C. Yao, "Iterative algorithm for generalized set-valued strongly nonlinear mixed variational-like inequalities," *Journal of Optimization Theory and Applications*, vol. 124, no. 3, pp. 725–738, 2005.

- [16] K. R. Kazmi and F. A. Khan, "Auxiliary problems and algorithm for a system of generalized variational-like inequality problems," *Applied Mathematics and Computation*, vol. 187, no. 2, pp. 789–796, 2007.
- [17] X. P. Ding and Z. B. Wang, "The auxiliary principle and an algorithm for a system of generalized set-valued mixed variational-like inequality problems in Banach spaces," *Journal of Computational and Applied Mathematics*, vol. 233, no. 11, pp. 2876–2883, 2010.
- [18] X.-P. Ding, "Auxiliary principle and approximation solvability for system of new generalized mixed equilibrium problems in reflexive Banach spaces," *Journal of Applied Mathematics and Mechanics*, vol. 32, no. 2, pp. 231–240, 2011.
- [19] B. S. Mordukhovich, B. Paniciucci, M. Pappalardo, and M. Passacantando, "Hybrid proximal methods for equilibrium problems," *Optimization Letters*. In press.
- [20] A. S. Antipin, "Computation of fixed points of extremal mappings by means of gradient-type methods," *Zhurnal Vychislitel cprime not Matematiki i Matematicheskoy Fiziki*, vol. 37, no. 1, pp. 42–53, 1997.
- [21] O. Chadli, Z. Chbani, and H. Riahi, "Equilibrium problems with generalized monotone bifunctions and applications to variational inequalities," *Journal of Optimization Theory and Applications*, vol. 105, no. 2, pp. 299–323, 2000.
- [22] S. B. Nadler, "Multi-valued contraction mappings," *Pacific Journal of Mathematics*, vol. 30, pp. 475–488, 1969.