

## Characterization of the Family I inorganic pyrophosphatase from *Pyrococcus horikoshii* OT3

SUNG-JONG JEON,<sup>1,2</sup> and KAZUHIKO ISHIKAWA<sup>3,4</sup>

<sup>1</sup> Department of Biotechnology and Bioengineering, Dong-Eui University, Busan, 614-714, Korea

<sup>2</sup> Department of Biomaterial Control, Dong-Eui University, Busan, 614-714, Korea

<sup>3</sup> Research Institute for Cell Engineering, National Institute of Advanced Industrial Science and Technology (AIST Kansai), 1-8-31 Midorigaoka, Ikeda, Osaka 563-8577, Japan

<sup>4</sup> Corresponding author (kazu-ishikawa@aist.go.jp)

Received January 8, 2005; accepted February 2, 2005; published online May 31, 2005

**Summary** A gene encoding for a putative Family inorganic pyrophosphatase (PPase, EC 3.6.1.1) from the hyperthermophilic archaeon *Pyrococcus horikoshii* OT3 was cloned and the biochemical characteristics of the resulting recombinant protein were examined. The gene (Accession No. 1907) from *P. horikoshii* showed some identity with other Family I inorganic pyrophosphatases from archaea. The recombinant PPase from *P. horikoshii* (*Ph*PPase) has a molecular mass of 24.5 kDa, determined by SDS-PAGE. This enzyme specifically catalyzed the hydrolysis of pyrophosphate and was sensitive to NaF. The optimum temperature and pH for PPase activity were 70 °C and 7.5, respectively. The half-life of heat inactivation was about 50 min at 105 °C. The heat stability of *Ph*PPase was enhanced in the presence of Mg<sup>2+</sup>. A divalent cation was absolutely required for enzyme activity, Mg<sup>2+</sup> being most effective; Zn<sup>2+</sup>, Co<sup>2+</sup> and Mn<sup>2+</sup> efficiently supported hydrolytic activity in a narrow range of concentrations (0.05–0.5 mM). The K<sub>m</sub> for pyrophosphate and Mg<sup>2+</sup> were 113 and 303 μM, respectively; and maximum velocity, V<sub>max</sub>, was estimated at 930 U mg<sup>-1</sup>.

**Keywords:** archaea, hyperthermophile.

### Introduction

Inorganic pyrophosphatases (PPases, EC 3.6.1.1) are widely distributed in all living cells and catalyze the hydrolysis of pyrophosphate (PP<sub>i</sub>) into two orthophosphates (P<sub>i</sub>) (Chen et al. 1990, Lundin et al. 1991). In many organisms, the bulk of PPase activity is soluble, localized in the cytosol and serves mainly to remove the PP<sub>i</sub> formed in biosynthetic reactions driven by nucleotide triphosphates (Kornberg 1962). Pyrophosphatases from the Bacteria, Archaea and Eukarya have been characterized. Bacterial and archaeal PPases exist as either homotetramers or homohexamers, with a subunit molecular mass of about 20 kDa. In contrast, eukaryotic PPases form mostly homodimers, with a subunit molecular mass of about 28–35 kDa. To date, the soluble PPases have been classified

into Family I and Family II PPases (Shintani et al. 1998, Young et al. 1998). Family I PPases have a highly conserved active site structure formed by 14–16 amino acid residues and three to four Mg<sup>2+</sup> ions and have similar catalytic properties (Harutyunyan et al. 1996, Heikinheimo et al. 1996, Baykov et al. 1999). Family II PPases are activated by Mn<sup>2+</sup> and display 10–20 times greater activity than Family I PPases (Shintani et al. 1998, Parfenyev et al. 2001).

Enzymes derived from microorganisms growing at extreme temperatures are of biotechnological use as highly thermostable biocatalysts. Thermostable PPases are commonly used to suppress pyrophosphorolysis in DNA sequencing reactions with DNA polymerases (Tabor and Richardson 1990, Vander et al. 1997). Some extremely thermostable PPases have been isolated and characterized from archaea (*Sulfolobus* (Wakagi et al. 1992), *Methanobacterium* (van Alebeek et al. 1994), *Methanococcus* (Kuhn et al. 2000), *Thermus* (Satoh et al. 1998) and *Thermoplasma* (Richter and Schafer 1992)). However, no information is available on the biochemical properties of thermostable PPase from hyperthermophilic archaeon *Pyrococcus* spp.

In the genome database of the hyperthermophilic archaeon *Pyrococcus horikoshii* OT3, we found the homologous PPase gene. The putative *Ph*PPase gene has been classified as a member of the Family I PPases (Young et al. 1998, Shintani et al. 1998), and the structure of *P. horikoshii* PPase has already been reported (Liu et al. 2004). In this study, we report on the cloning of *P. horikoshii* PPase and compare the properties of the recombinant *Ph*PPase with those of other microbial PPases.

### Materials and methods

#### Materials

The DNA primers and substrates were prepared by FASMA (Kanagawa, Japan). The plasmid pET-21a was purchased from Novagen (Madison, WI, USA). The KOD DNA polymerase

was purchased from Toyobo (Osaka, Japan). We obtained ATP, ADP and AMP from Sigma (St. Louis, MO). Sodium tripolyphosphate (PPP<sub>i</sub>), sodium pyrophosphate (PP<sub>i</sub>), sodium phosphate (P<sub>i</sub>), phosphoenolpyruvate (PEP) and *p*-nitrophenyl phosphate (pNP) were obtained from Wako Pure Chemical Industries (Tokyo, Japan).

#### Enzyme construction and expression

Chromosomal DNA of *P. horikoshii* OT3 was prepared as reported previously (Gonzalez et al. 1998). The gene (PH1907) was amplified by PCR using the chromosomal DNA as the template and two primers, PP1 (forward): 5'-AGGGGGTA CATATGAACCCGTTCCACGACCT-3'; and PP2 (reverse): 5'-AATAAAGTCCGACTCACTCCTTCTTGCCGA-3', which was designed based on an open reading frame (ORF) sequence for a protein of 178 amino acids. Amplification by PCR was carried out with KOD DNA polymerase at 94 °C for 30 s, 55 °C for 2 s, 74 °C for 30 s for 30 cycles. An amplified 0.53-kbp DNA fragment was digested with restriction enzymes *Nde*I and *Sal*I (the *Nde*I and *Sal*I sites are underlined in primers PP1 and PP2, respectively) and inserted into the pET-21a vector. The nucleotide sequences of the inserted genes were confirmed with an ABI PRISM 310 Genetic Analyzer (Perkin-Elmer, Wellesley, MA). *Escherichia coli* BL21(DE3) cells containing the expression plasmid were grown in 2YT medium (1% yeast extract, 1.6% tryptone, 0.5% NaCl) containing ampicillin (100 µg ml<sup>-1</sup>) to an OD<sub>600</sub> of approximately 0.6. Gene expression was then induced with 0.5 mM isopropyl-1-thio-β-D-galactopyranoside (IPTG) for 4 h at 37 °C.

#### Purification of the recombinant proteins

*Escherichia coli* cells containing the expressed recombinant enzyme were centrifuged and frozen at -70 °C. The thawed cells were then disrupted by sonication in buffer A (50 mM Tris-HCl, pH 8.0, 15 mM MgCl<sub>2</sub>, 0.1 mM EDTA). The suspension of disrupted cells was centrifuged at 27,000 *g* for 30 min and the supernatant fraction was heat-treated at 85 °C for 30 min, followed by recentrifugation. The supernatant was loaded on a HiTrap Q column (Amersham Biosciences, Piscataway, NJ) equilibrated in buffer A, and the bound protein was eluted with a linear gradient of NaCl (0 to 1.0 M in the same buffer). The protein solution was concentrated using a centricon 10 filter from Amicon (Millipore, Bedford, MA) and dialyzed against buffer B (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 15 mM MgCl<sub>2</sub>). The dialyzed solution was loaded on a HiPrep Sephacryl S-200 HR 26/60 column (Amersham Biosciences) and eluted with buffer B. The purity of the recombinant protein was assessed by SDS-PAGE. Protein concentration was determined using a Bio-Rad protein assay system (Bio-Rad, Hercules, CA) with bovine serum albumin as the standard.

#### Enzyme assay

In the standard assay, PPase activity was determined by the formation of inorganic phosphate at 60 °C in the reaction mix-

ture containing 50 mM Tris-HCl (pH 7.5 at 60 °C), 5 mM MgCl<sub>2</sub> and 1 mM sodium phosphate. The reaction was started by the addition of enzyme, and 100-µl samples were taken at 30-s time intervals and put on ice. The phosphate present was determined by adding 1 ml of phosphate reagent (1% ammonium heptamolybdate (w/v), 0.83 M sulfuric acid, 8% iron (II) sulfate (w/v)) and the absorbance of the phosphomolybdate complex was measured at 660 nm after 30 min (van Alebeek et al. 1994). Activities were developed with 0.1 µmol of standard sodium phosphate. The activity towards other substrates such as sodium tripolyphosphate (PPP<sub>i</sub>), phosphoenolpyruvate (PEP), *p*-nitrophenyl phosphate (pNP), ATP, ADP and AMP was determined by the same method as above except that PP<sub>i</sub> was replaced by each substrate. One unit of activity is defined as the amount of enzyme required to hydrolyze 1 µmol PP<sub>i</sub> per min at 60 °C.

## Results and discussion

#### Identification of inorganic pyrophosphatase gene

In the *Pyrococcus horikoshii* OT3 genome database ([http://www.bio.nite.go.jp/dogan/MicroTop?GENOME\\_ID=ot3\\_G1](http://www.bio.nite.go.jp/dogan/MicroTop?GENOME_ID=ot3_G1)) (Kawarabayasi et al. 1998), we identified an ORF (Accession No. PH1907) encoding a putative inorganic pyrophosphatase (*Ph*PPase). The putative *Ph*PPase gene encodes a protein of 178 amino acids, with a predicted molecular mass of 20,833 Da and an isoelectric point of 4.81. From the data of sequence homology (Shintani et al. 1998, Young et al. 1998), *Ph*PPase is classified as a member of Family I PPase. The deduced amino acid sequence showed a high identity with other archaeal PPases from *Pyrococcus furiosus* (94%) (Robb et al. 2001), *Methanopyrus kandleri* (69%) (Slesarev et al. 2002), *Methanobacterium thermoautotrophicum* (61%) (Smith et al. 1997) and *Thermoplasma acidophilum* (54%) (Richter and Schafer 1992). Structural and multiple sequence alignments reveal a group of 14 conserved, mostly polar, amino acid residues (Kankare 1994). The 14 amino acids have been found to be identical in all other *Ph*PPase amino acid sequences.

The ORF (PH1907) was amplified by PCR from a *P. horikoshii* genomic DNA, cloned and sequenced to confirm the sequences in the database. The gene was expressed in *E. coli* cells, and the recombinant protein was purified as described in the Materials and methods. Expression and subsequent purification yielded 3.2 mg of *Ph*PPase from a 1-l culture.

#### Molecular properties

The molecular mass of the protein from PH1907 was estimated to be about 24.5 kDa by SDS-PAGE (Figure 1). This value is inconsistent with the size (20,833 Da) calculated from the amino acid sequence. This discrepancy may be explained by the many positively charged amino acid residues of *Ph*PPase. Recently, a three-dimensional crystal structure of *Ph*PPase showed that the enzyme is an oligomeric hexamer (Liu et al. 2004).




Figure 1. Purification of recombinant *Pyrococcus horikoshii* pyrophosphatase (*PhPPase*). Lane 1, molecular mass markers; Lane 2, crude extract of induced cells; Lane 3, supernatant of crude extract after heat treatment at 85 °C for 30 min; Lane 4, HiTrap Q column peak fractions; and Lane 5, Sephacryl S-200 HR 26/60 column peak fractions. The gel was stained with Coomassie brilliant blue.

#### Catalytic properties

We measured the activity of the recombinant *PhPPase* in the presence of 50 mM Tris-HCl (pH 7.5), 5 mM  $MgCl_2$  and 1 mM sodium pyrophosphate. The recombinant *PhPPase* showed maximal activity in the temperature range from 60 to 75 °C, with a temperature optimum of 70 °C (Figure 2a). The optimal pH for the *PhPPase* was 7.5 at 60 °C. Heat inactivation of the enzyme was estimated by measuring the residual PPase activity after heat treatment at two different temperatures in the presence or absence of 5 mM  $Mg^{2+}$  (Figure 2b). The recombinant *PhPPase* showed high thermostability, and the half-life of heat inactivation was about 50 min at 105 °C. The heat stabil-

ity of *PhPPase* was much higher than that reported previously for other thermostable PPases from *Aquifex aeolicus* (90 min at 95 °C) (Hoe et al. 2001) and *Sulfolobus acidocaldarius* (150 min at 95 °C) (Hansen et al. 1999). Furthermore, the heat stability of *PhPPase* was enhanced by the addition of  $Mg^{2+}$  to the incubation mixture, as reported for the other PPases (Hachimori et al. 1979, Ichiba et al. 1988). The difference between the optimum temperature of *PhPPase* and the optimum growth temperature of *P. horikoshii* suggests that *PhPPase* requires co-factors for its thermophilicity in cells.

#### Divalent cation requirement

Family I PPase has shown strong metal cation-dependency, with  $Mg^{2+}$  conferring the highest efficiency (Cooperman 1982) and sensitivity to inhibition by  $Ca^{2+}$  (Yang and Wensel 1992). In contrast, Family II PPase prefers  $Mn^{2+}$  over  $Mg^{2+}$  (Parfenyev et al. 2001). *Pyrococcus horikoshii* pyrophosphatase is dependent on the presence of divalent cations for catalytic activity, with the highest activity in the presence of  $Mg^{2+}$ . Other cations ( $Zn^{2+}$ ,  $Co^{2+}$ ,  $Mn^{2+}$ ) could efficiently replace  $Mg^{2+}$ , but the effectiveness of the latter cations is limited to a narrow range of concentrations (0.05–0.5 mM) (Figure 3). Neither  $Ni^{2+}$  nor  $Ca^{2+}$  could activate the enzyme for catalysis. These results support the conclusion that *PhPPase* belongs to the Family I PPases. The enzyme responded sensitively to changes in the cation concentration of the reaction mixture. In particular, a drastic decrease in enzyme activity was observed at  $Zn^{2+}$ ,  $Co^{2+}$  and  $Mn^{2+}$  concentrations above 0.5 mM (Figure 3). The inhibition at high cation concentrations is probably due to the presence of free cations in the assay (Celis and Romero 1987).

#### Substrate specificity

The substrate specificity of *PhPPase* was investigated (Table 1) and found to be high with respect to  $PP_i$ . The release activities of inorganic phosphate ( $P_i$ ) from  $PPP_i$ , ATP and ADP were determined to be 2.7, 5.9 and 2.9% of that from  $PP_i$ , respectively. Hydrolysis of AMP, PEP and pNP by *PhPPase* was not observed.

The reaction rate of *PhPPase* at different  $PP_i$  and  $Mg^{2+}$  concentrations did not follow normal Michaelis-Menten kinetics.




Figure 2. Thermophilicity and thermostability of *Pyrococcus horikoshii* pyrophosphatase (*PhPPase*). (a) The PPase activity was determined at the indicated temperatures, as described in the Materials and methods. Negative control reactions in the absence of the enzyme were performed in parallel. (b) Enzyme (1  $\mu M$  *PhPPase* in 50 mM Tris-HCl, pH 7.5) was incubated at 100 °C in the presence (○) or absence (●) of  $Mg^{2+}$  and at 105 °C in the presence (◻) or absence (■) of  $Mg^{2+}$ , and the residual PPase activity of samples was measured at 60 °C.




Figure 3. Effect of divalent cation concentration on pyrophosphate (PP<sub>i</sub>) hydrolysis. Assay conditions were as follows: 50mM Tris-HCl, pH 7.5, 1 mM sodium pyrophosphate (NaPP<sub>i</sub>), and concentrations of divalent cations were varied as indicated. Symbols: ● = Mg<sup>2+</sup>; ■ = Zn<sup>2+</sup>; ▲ = Co<sup>2+</sup>; and ◆ = Mn<sup>2+</sup>.

Table 1. Substrate specificity of the *Pyrococcus horikoshii* pyrophosphatase (*PhPPase*). Activities were determined as described in Materials and methods. Each substrate was tested at a concentration of 1 mM in the presence of 5 mM Mg<sup>2+</sup>.

Substrate	Relative activity (%)
PP <sub>i</sub>	100
PPP <sub>i</sub>	2.7
ATP	5.9
ADP	2.9
AMP	0
PEP	0
pNP	0




Figure 4. Inhibition of *PhPPase* activity by NaF. The specific activity was determined at the indicated NaF concentrations, as described in Materials and methods.

Therefore, kinetic data were analyzed according to the procedure of Hill by the EZ-Fit program (Perrella 1988). At pH 7.5 and 60 °C, the values of  $K_m$  and maximum velocity ( $V_{max}$ ) for PP<sub>i</sub> were determined to be  $113 \pm 12 \mu\text{M}$  and  $930 \pm 40 \text{U mg}^{-1}$ , respectively, which corresponds to a  $k_{cat}$  of  $744 \text{s}^{-1}$ . An apparent  $K_m$  for Mg<sup>2+</sup> was determined to be  $303 \pm 9 \mu\text{M}$ .

#### Effect of fluoride

The influence of fluoride, a potent inhibitor of PPases, was examined. A plot of activity for NaF concentration (Figure 4) indicated that 0.5 mM NaF inhibits PPase activity by 75%. The  $I_{50}$  (concentration for 50% inhibition) was 0.1 mM. The fluoride sensitivity of *PhPPase* is higher than that of bacterial and archaeal PPases: *E. coli* (91%, 1 mM) (Josse 1966), *Thiobacillus thiooxidans* (81%, 1.25 mM) (Tominaga and Mori 1977), *Rhodospseudomonas palustris* (50%, 5 mM) (Schwarm et al. 1986) and *Methanobacterium thermoautotrophicum* (50%, 0.9 mM) (van Alebeek et al. 1994).

In conclusion, we have identified a gene that codes for PPase protein from the *P. horikoshii* genome and also compared the properties of recombinant *PhPPase* with those of other PPases. The *PhPPase* is more similar to bacterial and archaeal enzymes than to eukaryotic enzymes, based on the molecular mass of a single subunit. Cytoplasmic PPase has no function in energy conservation. Its function seems to keep the cellular PP<sub>i</sub> concentration low, through the hydrolysis of PP<sub>i</sub>. Whether *PhPPase* is located in the cytoplasm has not been reported. This work increased our understanding of the metabolism of a hyperthermophilic archaeon. The hyperthermostability of *PhPPase* suggests that it can be used to suppress pyrophosphorolysis in sequencing reactions using thermostable DNA polymerases (Tabor and Richardson 1990, Vander Horn et al. 1997).

#### Acknowledgment

This work was performed as part of the fellowship program supported by the New Energy Industrial Technology Development Organization (NEDO).

#### Reference

- Baykov, A.A., B.S. Cooperman, A. Goldman and R. Lahti. 1999. Cytoplasmic inorganic pyrophosphatase. In *Inorganic Polyphosphates*. Vol. 23. Ed. H.C. Schroder. Springer-Verlag, Berlin, pp 127–150.
- Celis, H. and I. Romero. 1987. The phosphate-pyrophosphate exchange and hydrolytic reactions of the membrane-bound pyrophosphatase of *Rhodospirillum rubrum*: effects of pH and divalent cations. *J. Bioenerg. Biomembr.* 19:255–272.
- Chen, J., A. Brevet, M. Fromant, F. Lévêque, J.M. Schmitter, S. Blanquet and P. Plateau. 1990. Pyrophosphatase is essential for growth of *Escherichia coli*. *J. Bacteriol.* 172:5686–5689.
- Cooperman, B.S. 1982. The mechanism of action of yeast inorganic pyrophosphatase. *Methods Enzymol.* 87:526–548.
- Gonzalez, J.M., Y. Masuchi, F.T. Robb, J.W. Ammerman, D.J. Maeder, M. Yanagibayashi, J. Tamaoka and C. Kato. 1998. *Pyrococcus horikoshii* sp. nov., a hyperthermophilic archaeon isolated from a hydrothermal vent at the Okinawa Trough. *Extremophiles* 2:123–130.

- Hachimori, A., Y. Shiroya, A. Hirato, T. Miyahara and T. Samejima. 1979. Effects of divalent cations on thermophilic inorganic pyrophosphatase. *J. Biochem.* 86:121–130.
- Hansen, T., C. Urbanke, V.M. Leppanen, A. Goldman, K. Brandenburg and G. Schafer. 1999. The extreme thermostable pyrophosphatase from *Sulfolobus acidocaldarius*: enzymatic and comparative biophysical characterization. *Arch. Biochem. Biophys.* 363:135–147.
- Harutyunyan, E.H., I.P. Kuranova, B.K. Vainshtein, W.E. Hohne, V.S. Lamzin, Z. Dauter, A.V. Teplyakov and K.S. Wilson. 1996. X-ray structure of yeast inorganic pyrophosphatase complexed with manganese and phosphate. *Eur. J. Biochem.* 239:220–228.
- Heikinheimo, P., J. Lehtonen, A. Baykov, R. Lahti, B.S. Cooperman and A. Goldman. 1996. The structural basis for pyrophosphatase catalysis. *Structure* 4:1491–1508.
- Hoe, H.S., H.K. Kim and S.T. Kwon. 2001. Expression in *Escherichia coli* of the thermostable inorganic pyrophosphatase from the *Aquifex aeolicus* and purification and characterization of the recombinant enzyme. *Protein Expr. Purif.* 23:242–248.
- Ichiba, T., T. Shibasaki, E. Izuka, A. Hachimori and T. Samejima. 1988. Cation-induced thermostability of yeast and *Escherichia coli* pyrophosphatases. *Biochem. Cell Biol.* 66:25–31.
- Josse, J. 1966. Constitutive inorganic pyrophosphatase of *Escherichia coli*. I. Purification and catalytic properties. *J. Biol. Chem.* 241:1938–1947.
- Kankare, J., G.S. Neal, T. Salminen, T. Glumoff, B.S. Cooperman, R. Lahti and A. Goldman. 1994. The structure of *E. coli* soluble inorganic pyrophosphatase at 2.7 Å resolution. *Protein Eng.* 7:823–830.
- Kawarabayasi, Y., M. Sawada, H. Horikawa et al. 1998. Complete sequence and gene organization of the genome of a hyper-thermophilic archaeobacterium, *Pyrococcus horikoshii* OT3. *DNA Res.* 5:55–76.
- Kornberg, A. 1962. On the metabolic significance of phosphorolytic and pyrophosphorolytic reactions. In *Horizons in Biochemistry*. Eds. M. Kasha and B. Pullman. Academic Press, New York, pp 251–264.
- Kuhn, N.J., A. Wadson, S. Ward and T.W. Young. 2000. *Methanococcus jannaschii* ORF mj0608 codes for a class C inorganic pyrophosphatase protected by Co<sup>2+</sup> or Mn<sup>2+</sup> ions against fluoride inhibition. *Arch. Biochem. Biophys.* 379:292–298.
- Liu, B., M. Bartlam, R. Gao, W. Zhou, H. Pang, Y. Liu, Y. Feng and Z. Rao. 2004. Crystal structure of the hyperthermophilic inorganic pyrophosphatase from the archaeon *Pyrococcus horikoshii*. *Biophys. J.* 86:420–427.
- Lundin, M., H. Baltscheffsky and H. Ronne. 1991. Yeast PPA2 gene encodes a mitochondrial inorganic pyrophosphatase that is essential for mitochondrial function. *J. Biol. Chem.* 266:12,168–12,172.
- Parfenyev, A.N., A. Salminen, P. Halonen, A. Hachimori, A.A. Baykov and R. Lahti. 2001. Quaternary structure and metal ion requirement of family II pyrophosphatases from *Bacillus subtilis*, *Streptococcus gordonii*, and *Streptococcus mutans*. *J. Biol. Chem.* 276:24,511–24,518.
- Perrella, F.W. 1988. EZ-FIT: a practical curve-fitting microcomputer program for the analysis of enzyme kinetic data on IBM-PC compatible computers. *Anal. Biochem.* 174:437–447.
- Richter, O.M. and G. Schafer. 1992. Cloning and sequencing of the gene for the cytoplasmic inorganic pyrophosphatase from the thermoacidophilic archaeobacterium *Thermoplasma acidophilum*. *Eur. J. Biochem.* 209:351–355.
- Robb, F.T., D.L. Maeder, J.R. Brown, J. DiRuggiero, M.D. Stump, R.K. Yeh, R.B. Weiss and D.M. Dunn. 2001. Genomic sequence of hyperthermophile, *Pyrococcus furiosus*: implications for physiology and enzymology. *Methods Enzymol.* 330:134–157.
- Satoh, T., T. Samejima, M. Watanabe, S. Nogi, Y. Takahashi, H. Kaji, A. Teplyakov, G. Obmolova, I. Kuranova and K. Ishii. 1998. Molecular cloning, expression, and site-directed mutagenesis of inorganic pyrophosphatase from *Thermus thermophilus* HB8. *J. Biochem.* 124:79–88.
- Schwarm, H.M., H. Vigerschow and K. Knobloch. 1986. Purification and properties of a soluble inorganic pyrophosphatase from *Rhodospirillum rubrum*. *Biol. Chem. Hoppe-Seyler* 367:119–126.
- Shintani, T., T. Uchiumi, T. Yonezawa, A. Salminen, A.A. Baykov, R. Lahti and A. Hachimori. 1998. Cloning and expression of a unique inorganic pyrophosphatase from *Bacillus subtilis*: evidence for a new family of enzymes. *FEBS Lett.* 439:263–266.
- Slesarev, A.I., K.V. Mezhevaya, K.S. Makarova et al. 2002. The complete genome of hyperthermophile *Methanopyrus kandleri* AV19 and monophyly of archaeal methanogens. *Proc. Natl. Acad. Sci. USA* 99:4644–4649.
- Smith, D.R., L.A. Doucette-Stamm, C. Deloughery et al. 1997. Complete genome sequence of *Methanobacterium thermoautotrophicum* deltaH: functional analysis and comparative genomics. *J. Bacteriol.* 179:7135–7155.
- Tabor, S. and C.C. Richardson. 1990. DNA sequence analysis with a modified bacteriophage T7 DNA polymerase: effect of pyrophosphorylation and metal ions. *J. Biol. Chem.* 265:8322–8328.
- Tominaga, N. and T. Mori. 1977. Purification and characterization of inorganic pyrophosphatase from *Thiobacillus thiooxidans*. *J. Biochem.* 81:477–483.
- van Alebeek, G.J., J.T. Keltjens and C. van der Drift. 1994. Purification and characterization of inorganic pyrophosphatase from *Methanobacterium thermoautotrophicum* (strain delta H). *Biochim. Biophys. Acta* 1206:231–239.
- Vander Horn, P.B., M.C. Davis, J.J. Cunniff et al. 1997. Thermo-Sequenase DNA polymerase and *T. acidophilum* pyrophosphatase: new thermostable enzyme for DNA sequencing. *BioTechniques* 22:758–765.
- Wakagi, T., C.H. Lee and T. Oshima. 1992. An extremely stable inorganic pyrophosphatase purified from the cytosol of a thermoacidophilic archaeobacterium, *Sulfolobus acidocaldarius* strain 7. *Biochim. Biophys. Acta* 1120:289–296.
- Yang, Z. and T.G. Wensel. 1992. Inorganic pyrophosphatase from bovine retinal rod outer segments. *J. Biol. Chem.* 267:24,634–24,640.
- Young, T.W., N.J. Kuhn, A. Wadson, S. Ward, D. Burges and G.D. Cooke. 1998. *Bacillus subtilis* ORF yybQ encodes a manganese-dependent inorganic pyrophosphatase with distinctive properties: the first of a new class of soluble pyrophosphatase? *Microbiology* 144:2563–2571.



**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

