Review Article

Archaea in Symbioses

Christoph Wrede,^{1,2} Anne Dreier,^{1,3} Sebastian Kokoschka,¹ and Michael Hoppert^{1,3}

¹ *Institute of Microbiology and Genetics, Georg-August-Universität Göttingen, Grisebachstraße 8, 37077 Göttingen, Germany*

² *Hannover Medical School, Institute of Functional and Applied Anatomy, Carl-Neuberg-Straße 1, 30625 Hannover, Germany*

³ *Courant Centre Geobiology, Georg-August-Universität Göttingen, Goldschmidtstraße 3, 37077 Göttingen, Germany*

Correspondence should be addressed to Anne Dreier, anne.dreier@gmx.de

Received 5 September 2012; Accepted 19 November 2012

Academic Editor: Martin Krüger

Copyright © 2012 Christoph Wrede et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

During the last few years, the analysis of microbial diversity in various habitats greatly increased our knowledge on the kingdom Archaea. At the same time, we became aware of the multiple ways in which Archaea may interact with each other and with organisms of other kingdoms. The large group of euryarchaeal methanogens and their methane oxidizing relatives, in particular, take part in essential steps of the global methane cycle. Both of these processes, which are in reverse to each other, are partially conducted in a symbiotic interaction with different partners, either ciliates and xylophagous animals or sulfate reducing bacteria. Other symbiotic interactions are mostly of unknown ecological significance but depend on highly specific mechanisms. This paper will give an overview on interactions between Archaea and other organisms and will point out the ecological relevance of these symbiotic processes, as long as these have been already recognized.

1. Introduction

Symbiotic interactions between various groups of prokaryotes as well as between prokaryotes and eukaryotic organisms were one essential driving force of evolution, including the development of differentiated tissues in multicellular organisms [1]. Apart from the essential key events of endosymbiosis, leading to mitochondria and chloroplasts, a multitude of symbiotic interactions at various levels is an ongoing process [2]. Interestingly, most of these interactions are contributed by Bacteria, frequently by Proteobacteria [3]. Regarding parasitic or pathogenic interactions in particular, the outer envelope of the bacterial cell mediates highly specific contact to its host. Surface structures like pili, lipopolysaccharides, and outer membrane proteins may rapidly adapt to modified host tissue structures, mainly with respect to deleterious host-pathogen interactions [4]. How about Archaea? Up to now, no clearly identifiable pathogenic interactions between an Archaeon and its host have been detected, though some archaeal commensals may be indirectly involved in bacterial infections [5]. On the other hand, mutualistic symbioses have been well described, some of them with high relevance to global environmental cycles

[6]. Here we will present a short overview on interaction mechanisms known so far and relevant symbioses between Archaea and other organisms. We use the term symbiosis here in a broader sense (see Table 1); in most if not all cases the benefit of both interaction partners could not be proven, though none of these interactions appeared to be detrimental for one of the partners [7].

2. Mechanisms for Interaction with Host Cells

Apart from vertically transmitted endosymbionts, any interaction between host and symbiont depends on the surface-surface recognition. A variety of proteinaceous and polysaccharide-based surface structures are known to be involved. Surface layers are common in all Archaeal groups known so far. Though their function in adhesion is up to now poorly understood, it may be expected that in particular the glycosylated surface layers are involved in adhesion. In fact, like in Bacteria, extracellular polysaccharides are known as adhesive matrix for biofilm formation [8]. As it has been described for Bacteria, filamentous protein appendages are important for adhesion of Archaeal cells. Some of them appear to be unique for particular phylotypes, like the hami,

TABLE 1: Some examples for symbioses between Archaea and other organisms (red Archaea, green Bacteria, blue-unicellular Eukarya, and metazoans).

Symbiotic partners	Examples	Interaction	Key features
Archaeon-Archaeon	<i>Ignicoccus hospitalis</i> / <i>Nanaoarchaeum equitans</i>	Cell-cell contact	Transfer of essential biological macromolecules from host to symbiont
Archaea-Bacteria	SM1/ <i>Thiothrix</i> ANME-Archaea/sulfate reducing Bacteria	Cell appendages Cell-cell contact/ no direct contact	Sulfur redox cycling? Anaerobic methane oxidation coupled to sulfate reduction
Archaea-unicellular Eukarya	Methanogenic Archaea/Ciliates, Archamoebae	Endosymbiotic	Methanogen/hydrogenosome association, transfer of hydrogen, and/or C ₁ /C ₂ compounds
Archaea-metazoans	Porifera-associated Thaumarchaeota Diverse associations between insect and vertebrate guts	Tissue associated Gut microbial communities	Ammonia oxidation Methanogens as a terminal part of the anaerobic food chain

highly complex proteinaceous appendages appearing like hooks. These structures are supposedly involved in the formation of a tight biofilm consisting of the euryarchaeon SM1 and a filamentous *Thiothrix*-related sulfur-oxidizing proteobacterium [9, 10] and seem to be unrelated to any other known surface appendage. Remarkably, the filamentous protein assembly ends up in a terminal hook. In addition, short prickles branch from the main filament. Though it is not known in which way interaction at the molecular level may work, a tight binding of the cells to each other and to various surfaces was shown. The filaments and an exopolysaccharide supposedly excreted by the Archaeon provide the matrix for the formation of a tight consortium between the Archaeon and the sulfur oxidizer. The shape of the consortia may vary but exhibit sort of a “string of pearl” appearance. Each pearl is in the millimeter order of magnitude and is colonized by cells of the anaerobic SM1 Archaeon in the core and a shell of the aerobic sulfur oxidizer. SM1-related sequences have been detected in low saline sulfidic water worldwide, thus a certain ecological significance is likely. The tight association is beneficial for both symbiotic partners when the Archaeon uses the sulfate generated by the sulfur oxidizer for dissimilatory sulfate reduction. The *Thiothrix* “shell” will provide anoxic conditions inside the consortium [11].

In this context another apparently unique surface structure should be mentioned. Though not involved in symbiotic interaction, hollow tubes (cannulae) composed of glycoproteins interconnect the cells of the hyperthermophilic *Pyrodictum occultum* [12]. Another uncommon structure has been described recently by cryoelectron microscopy of a microbial biofilm [13]. Cells of a member of the archaeal

group Thermoplasmatales form protuberances penetrating cells of the ultrasmall archaeal Richmond mine acidophilic organism (ARMAN).

In addition to these unique structures, interaction is frequently brought about by appendages that are also common to Bacteria. Several pilus types involved in recognition of and attachment to surfaces have been detected in Bacteria so far [4]. Yet, by far most of the archaeal pili have similarities to just the bacterial type IV pilin. Intriguingly, the archaeal flagellin is homologous to the bacterial pilin protein. The archaeal rotating flagellum is homologous to the bacterial type IV pilus secretion apparatus. Consequently, no homologies between the bacterial flagellar genes and archaeal sequences could be detected up to now [14, 15]. Also the genes of the machinery for pilus assembly have been detected in the archaeal genomes. The involvement of the appendages in attachment may differ in various groups. The flagella of *Pyrococcus furiosus* are probably motility organelles, but are also important for biofilm formation and surface attachment [16]. Flagella and pili are also necessary for the aggregate formation and surface adherence of *Sulfolobus solfataricus* [17]. As in bacteria, the pili are responsible for primary adhesion on surfaces and initiate biofilm formation. The environmental conditions for biofilm formation have been extensively studied for several *Sulfolobus* strains. Basically, temperature, pH, and iron concentrations, which are also relevant in the natural (hot spring) habitat, strongly influence biofilm development. In particular, pH and iron concentration may synergistically act on biofilm development, but in different ways in various *Sulfolobus* strains [18]. In an artificial archaeal biofilm formed by *Pyrococcus furiosus* and *Methanopyrus kandleri*,

the latter adheres to the surface (mica, glass, and others), whereas *Pyrococcus* adheres to *Methanopyrus* via flagella and/or direct contact between cells [19]. *Haloferax volcanii* uses nonmotile pili for surface attachment [20].

Pili are also involved in interactions between the Archaea and eukaryote hosts. *Methanobrevibacter*'s polar pilus-like fibers are responsible for the attachment of cells to the hindgut epithelium cuticle of the termite *Reticulitermes flavipes* [21].

Generally, a dual nature of cellular appendages (motility and attachment) is not uncommon and has been also repeatedly described for Bacteria [22, 23]. This feature is also true among Archaea, as the mentioned examples may illustrate. However, some types of cellular appendages have not been detected in Archaea. The type III secretion system (TTSS), in particular, is the essential export mechanism for bacterial flagellins and is also an important pathogenicity factor. Specific bacterial proteins are delivered to a eukaryote host after recognition by the TTSS via a hollow channel. This very specific interaction to eukaryotes may have been developed at a time in evolution, when specific signaling between pathogens and multicellular eukaryotes was evolutionary useful [24].

3. Archaeon-Archaeon Interaction

The symbiosis between the host *Ignicoccus hospitalis* and *Nanoarchaeum equitans* is well described at the structural level. *Ignicoccus*, (Desulfurococcales, Crenarchaeota) is an anaerobic, hyperthermophilic obligate chemolithoautotrophic hydrogen oxidizing Archaeon. Interestingly, cells belonging to the genus *Ignicoccus* are surrounded by a dual membrane, which appears to be a similarity to most Bacteria. However, the archaeal "outer membrane" is distinct from the composition of the known bacterial outer membranes. Most interestingly, the outer membrane of *Ignicoccus* hosts the H₂: sulfur oxidoreductase and ATPase protein complexes, that is, membrane energization takes place at this membrane and not at the inner (normally referred to as cytoplasmic) membrane as it is common in all Bacteria with a double membrane cell envelope [25]. Typical porins, homologous to those in bacterial outer membranes, are missing, which also implies that the *Ignicoccus* outer membrane is not homologous to the outer membrane of Bacteria. Instead, in *Ignicoccus hospitalis*, a unique pore-forming complex (Ihomp1) consists of nine monomers of a small unique alpha-helical protein [26]; other membrane proteins appear to be involved in the *Ignicoccus*/*Nanoarchaeum* symbiosis as well [27]. The symbiont *Nanoarchaeum equitans* depends obligately on the *Ignicoccus* host. The *Nanoarchaeum* cells are directly attached to the outer membrane of *Ignicoccus*. The extremely reduced genome (490 Kbp) lacks genes for essential biosynthetic pathways, such as lipid, amino acid, and nucleotide biosynthesis. Thus biological macromolecules must be provided by the *Ignicoccus* host; even transfer of ATP from host to symbiont has been discussed [28].

The relationship between *Ignicoccus* and *Nanoarchaeum* does not appear to be a true mutualistic symbiosis: though the growth parameters of either infected or uninfected

Igni-coccus cultures (containing infected cells of different degrees and uninfected cells) are the same, attached *Nanoarchaeum* cells significantly reduce the ability of *Ignicoccus* to reproduce [29].

Up to now, direct interactions between two archaeal partners appear to be extremely rare. Other species of the genus *Ignicoccus* are free living and could not be infected with the *Nanoarchaeum equitans* symbiont [29]. Though it is unlikely that interactions within the kingdom of Archaea are an exception, it has to be taken into account that interactions between largely unculturable organisms are difficult to detect. In an artificial binary biofilm between *Pyrococcus furiosus* and *Methanopyrus kandleri* hydrogen produced by *Pyrococcus* is utilized by *Methanopyrus*, which implies that mutualistic benefits may lead to stable aggregations between Archaea [18]. Upcoming in situ techniques may uncover interactions between Archaea in the near future [13].

4. Archaea-Bacteria Interactions

Under anaerobic conditions, organic compounds are degraded by the anaerobic food chain whereby the product of one group serves as a substrate for the next group within this chain. Methanogenic Archaea terminate the chain by degrading C₁ and C₂ substrates to methane and carbon dioxide.

The conversion of higher organic acids to acetate and hydrogen is endergonic, unless the hydrogen partial pressure is kept low. This may be achieved by the activity of hydrogenotrophic methanogens. This necessary coupling of hydrogen formation and uptake by syntrophic microbial consortia is termed "interspecies hydrogen transfer." A well-known consortium, "*Methanobacillus omelianskii*," was isolated several times from anaerobic sediments and sewage sludge and was regarded as a pure culture of an anaerobe converting ethanol to acetate and methane [30]. In fact, the culture consisted of a methanogenic archaeon and a Gram-negative Bacterium [31, 32]. Since then a multitude of syntrophic associations have been described, for example, with the fermentative *Acetobacterium* or *Syntrophobacter* [33, 34], with *Desulfovibrio* under low sulfate concentrations [35], but also under thermophilic conditions with *Thermoanaerobacter*, *Desulfotomaculum*, and *Pelotomaculum* [36–38] and with hydrogenotrophic methanogens as syntrophic partners. These examples show the diversity of interactions with respect to organisms and metabolic properties. Though stable aggregates and specific interactions between the syntrophic partners have been observed [39], syntrophy in interspecies hydrogen transfer is generally highly variable and may depend on the availability of substrates [40].

An important process of methane oxidation in anoxic sediments is conducted by consortia of Euryarchaeota and sulfate reducing Bacteria (SRB). The anaerobic oxidation of methane (AOM) has been first postulated by Reeburgh [41]. Up to 90% of methane produced in marine sediments is anaerobically oxidized [42], which makes AOM to an essential process in global methane turnover. However, quantitative modeling based on existing data of the few sampling sites at the ocean floor is still difficult and

the contribution of the process to global methane cycling is still a matter of debate [43]. In ocean systems, methane is either generated by methanogenesis in sediments, or abiotically by serpentinization, and may derive from methane hydrates and fossil reservoirs. In cases of high methane fluxes from large reservoirs, the AOM is usually associated with the precipitation of carbonates and sulfides. This has been particularly observed at sites of intense methane seepage, such as marine mud volcanoes and cold methane seeps; also fossil seeps were identified [44–46]. These precipitates are mostly found in sediments as carbonate-cemented plates or large tabular constructions, but also as grapestone-like concretions or even as giant columnar structures, up to several tens of meters in height, buried in the sediment [47–49]. Under anaerobic conditions, below the chemocline of anoxic ocean basins, these precipitates may form tower-like constructions in the water column, reaching several meters in height [50–52]. In most cases, tube-like or columnar towers exhibit cavities that are perfused by methane and seawater. The inner faces of these concretions are covered by remarkably complex biofilms [53–55], dominated by various representatives of the ANME Archaea (ANME: anaerobic methanotroph). The three known ANME groups are not monophyletic. ANME-1 are distantly related to Methanomicrobiales [56], while ANME-2 and ANME-3 are distantly related to Methanosarcinales [57, 58]. A fourth group has been described as ANME-2d or GoM Arc I; this group is not monophyletic with the other ANME-2 subgroups [59–61]. AOM metabolism for this novel group has not yet been proven [42]. ANME-1 and ANME-2 are the most diverse groups detected in a multitude of habitats and appear to be most relevant for AOM in anoxic environments. ANME-1 cells exhibit a cylinder-shaped morphology with an external sheath and were found only in loose association with SRB of the *Desulfococcus/Desulfosarcina* (DSS) group [53]. ANME-2 cells are coccoid and are frequently detected in consortia with SRB [55, 62]. In ANME-2a/SRB-aggregates, both cell types appear to be randomly intermixed, while ANME-2c/SRB aggregates reveal a shell-like structure with SRB at the outer shell of the aggregate. ANME-2 are usually associated with SRB of the DSS group [63, 64], but also associations with alpha-Proteobacteria, beta-Proteobacteria, or *Desulfobulbus*-related SRB and ANME-2 cells without contact to other bacteria were reported [65–71].

There is up to now no indication that the metabolism of the SRB in AOM is distinct from free-living sulfate reducing bacteria. The metabolic pathway of the ANME archaea is clearly related to methanogenesis. Intriguingly, ANME Archaea use this pathway in the reverse direction, while reducing equivalents are transferred to SRB [42, 72]. Until now, it seems that AOM with ANME Archaea is feasible just in syntrophy with sulfate reduction. A recently discovered thermophilic ANME group closely affiliated to ANME1 (ANME 1c), though may conduct AOM in contact to hydrothermal vent systems without SRB and with Fe^{3+} as putative electron acceptor. However, conclusive evidence is still missing in this case [73].

The methyl-coenzyme M reductase (MCR) catalyses in methanogenic archaea the terminal step of methane

formation. In reversal, MCR is needed in a reversed methanogenic pathway for the initial step of AOM; also most of the other enzyme steps of methanogenesis operate in the reverse direction [74–76]. However, direct evidence for the reverse operation of this pathway is still lacking.

Intermediates for a necessary transfer of reducing equivalents between the syntrophic partners are still unknown. *In vitro* feeding studies excluded hydrogen, formate, acetate, methanol, and even more uncommon compounds like methylsulfides or humic acids [77–81]. The energy yield of AOM is still extremely low, compared with other anaerobic processes [82].

Recent findings indicate that the ANME Archaea are capable of both methane oxidation and sulfate reduction with elemental sulfur as an intermediate [83]. The reduced product HS_2^- may be the disproportionated by the symbiotic sulfate reducers to sulfate and HS^- . Thus, the symbioses may be less obligate than originally thought.

ANME-2 Archaea in consortia also conduct nitrogen fixation [84]. Nanometer secondary ion mass spectrometry (nano-SIMS) analysis implied the flow of nitrogen compounds from the Archaea to the sulfate reducers. Remarkably, the energy consuming nitrogen fixation is possible even under the conditions of the extremely low energy yield of AOM, though growth rates of the organisms were reduced by a factor of 20. Since AOM is a mayor sink of methane in marine sediments, nitrogen fixation by AOM may be as well a relevant process in the global nitrogen cycle. By this way, carbon, nitrogen, and sulfur cycles are linked by AOM.

Other recently described AOM processes may also be independent of archaeal groups. AOM with iron (Fe^{3+}) and manganese (Mn^{3+} , Mn^{4+}) has been described for enrichment cultures from marine sediment samples, but a direct involvement of either archaeal or bacterial phylotypes is speculative [85]. A nitrite/nitrate dependent AOM is conducted by Bacteria (NC10 phylum, candidate *Methylomirabilis oxyfera*; [86]). This process is clearly distinct from ANME/SRB AOM and appears to be homologous to the aerobic methane oxidation of methanotrophic Bacteria.

Another cell-cell interaction between the giant filamentous thaumarchaeote candidate *Giganthauma karukerense* and a sulfur oxidizing gamma-Proteobacterium has been described recently [87]. A closed cell monolayer of the proteobacteria covers the surface of the large thaumarchaeote filament. It is not known in how far the cells may interact physiologically. It might be possible that the sulfur oxidizer reduces the sulfide concentration in the immediate vicinity of the host cell.

5. Interaction between Archaea and Eukarya

With respect to hitherto known mutualistic symbioses with eukaryotes, most but not all Archaea are members of the methanogenic Euryarchaeota. Methanogens are essential in the degradation of organic substrates under anaerobic conditions to methane and carbon dioxide, as terminal part of the anaerobic food chain. It is reasonable to assume that organisms with guts as anaerobic niches of nutrient decomposition harbor also methanogens as commensals.

Remarkably, a single methanogen phylotype, *Methanobrevibacter smithii*, is known to be the predominant Archaeon in the human gut microflora [88]. Symbioses between Archaea and eukaryotes, however, are not restricted to the gut anaerobic food chain. Many of the anaerobic protozoa, either free living or gut symbionts themselves, contain methanogenic Archaea as endosymbionts. These free-living protozoa are widespread in sapropels. Instead of mitochondria, they contain hydrogenosome organelles lacking a tricarboxylic acid cycle [89]. Hydrogenosomes are descendants of mitochondria. In these organelles, ATP is generated in a fermentative pathway by conversion of acetyl-CoA to acetate; the reducing power is released as molecular hydrogen [90]. Hydrogenosomes are a prerequisite for the occurrence of endosymbiotic methanogens, and hydrogenotrophic methanogens use hydrogen and carbon dioxide or formate as substrates for methanogenesis [91]. Also acetoclastic methanogens may take benefit from acetate generated by the hydrogenosome [92]. Regularly, these symbionts are transmitted vertically in the protists. Consequently, the phylotypes of the methanogens differ with respect to the protist groups and their habitats. Methanogens from free-living ciliates in freshwater habitats are related to Methanomicrobiales, whereas endosymbionts in ciliates from millipedes, cockroaches, and even frogs are Methanobacteriales [93]. However, during culturing, ciliates may also tend to lose their symbionts and also uptake of *Methanobacterium formicicum* by the ciliate *Trimyema compressum* has been reported [94]. Mechanisms of interactions between the methanogens and protists are largely unknown. Several morphological peculiarities have been described. In the free-living ciliate *Metopus contortus* polymorphic endosymbionts were observed; some cells appear to lose their cell walls and become directly attached to hydrogenosomes [95]. Also variations in size (putatively due to enlargement of cells) and stellate shape of the endosymbiont with close contact to hydrogenosomes have been observed repeatedly [96].

These results show that a peculiar crosstalk between the symbiotic partners is necessary. Like in well-studied symbiosis between eukaryotes and bacterial endosymbionts (including intracellular pathogens), protection against lysosomal digestion or cytoskeletal rearrangements of the vacuole containing the endosymbiont requires elaborate signaling pathways between host and symbiont partners [97]. It is reasonable to assume that the Archaeal symbionts possess respective signaling mechanisms. However, the endosymbiotic associations between several groups of protists (Ciliata, and some representatives within groups of Archamoebae) are the only known endosymbioses so far. Thus, this way of interaction with eukaryotes does not appear to be a mainstream in archaeal life styles [98, 99].

Sponges, organisms at the evolutionary basis of the Metazoa, may be described as a diverse prokaryotic community in a eukaryote host, most of the prokaryotes with largely unknown function [100]. Though the majority of the organisms are free living in the sponge mesohyl, endosymbionts are common. Among cyanobacteria and heterotrophic bacteria, also fission yeasts have been described that are maternally transmitted via sponge eggs [101].

Archaea are ubiquitous in marine sponges, sometimes even dominant [102], though their ecological role is poorly understood. Since the composition of the archaeal community is distinct from seawater, a certain specificity of the sponge/Archaeon association must be assumed [103]. Also rather specific associations between certain archaeal phylotypes and sponges have been described. The association between a sponge and the Thaumarchaeota (formerly Crenarchaeota; [104, 105]) *Cenarchaeum symbiosum* has been first described for *Axinella mexicana* [106]. Three species of the Mediterranean *Axinella* harbor filamentous marine "group 1" Archaea colonizing the collagen surrounding the sponge spicules [107]. Marine Euryarchaeota are associated with the demosponge *Tentorium semisuberites* mesohyl [102]. The role of these symbioses is largely unknown, also with respect to the unknown ecological role of the marine Thaumarchaeota. Recent findings imply significance in the sponge nitrogen metabolism [103, 108, 109]. A vertical transmission of the ammonia-oxidizing Archaea also indicates the specificity of the symbiotic relationship [110]. Ammonia oxidizers may utilize ammonia excreted by the sponge as a metabolic end product and may thereby contribute to detoxification of the sponge tissue. This may be in particular of relevance in highly polluted areas, where high concentration of organic compounds and high ammonia concentrations affect marine biocoenoses [103].

The important role of Thaumarchaeota in nitrogen cycling, also with respect to symbioses, has been also identified in some marine mollusks: strains phylogenetically related to *Nitrosopumilus maritimus* were detected inside the tissue of the colonial ascidian *Cystodytes dellechiaiei*. Here, nitrification of the Archaeon could be determined in situ [111]. Recent studies on the diversity of ammonia oxygenase genes also show that ammonia oxidizing archaeal communities differ in various coral species and are also distinct from communities in the sediment or in the water column [112, 113]. It must be expected that symbioses between other groups of marine invertebrates and Thaumarchaeota are also of relevance, in particular with respect to ammonia oxidation [114].

Among arthropods, as the largest animal phylum, only in the groups of millipedes, cockroaches, termites, and scarabs relevant methane producing species are present [93]. Methanogens represent the terminal part of the anaerobic food chain in the guts of these insects (especially termites). In this symbiosis, these Archaea utilize the main degradation products hydrogen, carbon dioxide, and acetate released by the previous steps of anaerobic lignocellulose degradation [115]. All methanogens, including the methanogen endosymbiont-bearing ciliates, are in the hindguts of these arthropods [116]. Free-living methanogens adhere to the hindgut wall. Among these big groups, a correlation between a specific diet (e.g., plant litter) and methane production could not be found, and not all members of the mentioned groups contain methanogens. However, in the group of higher termites, soil feeding termites produce more methane and contain more methanogens (according to 16S rRNA analysis) than wood feeders [117]. In the soil feeding species *Cubitermes fungifaber*, the composition of the communities

vary across the species, which does not account for a pure vertical transmission of the gut community, but a strong influence of the community in food soil [118]. Remarkably, also a *Natronococcus*-related sequence could be retrieved from the gut. Related strains are obligate haloalkaliphilic organisms of the family Halobacteriaceae, isolated from soda lakes, and are aerobic heterotrophic Archaea [119]. The *Natronococcus*-related strain may be well adapted to the first section of the hindgut (P1 part). This section provides a highly alkaline environment, reaching a pH around 12. For *Cubitermes orthognatus*, an in depth analysis of archaeal communities in four sections of the hindgut revealed remarkable distinctions in particular between the alkaline P1 part and the following P3–P5. Whereas Methanosarcinaceae-related sequences dominated in P1, they were replaced by Methanobacteriaceae-related clones in all other posterior parts of the gut. Interestingly, also Thermoplasmatales and Crenarchaeota contributed up to 40% to the archaeal community in these parts. The ecological role of these archaeal groups have to be elucidated yet.

Methanogenesis in termites is a globally relevant source of methane, with 20–29 Tg methane per year [120]. Methanogens from all ruminants produce 91–107 Tg methane per year, which is the second largest methane source after wetlands. In ruminants, methanogens are in a similar way the terminal part of the anaerobic lignocellulolytic food chains as in termite hindguts and methanogen/ruminant symbioses have been extensively studied. Some methanogens like *Methanobacterium bryantii* or *Methanobrevibacter ruminantium* were isolated from rumen fluids and were extensively studied with respect to biochemistry and energetics in methanogens, including genome analysis of *Methanobrevibacter* [121, 122]. Abundant adhesin-like sequences in the *Methanobrevibacter* genome imply intensive interactions between the methanogen and other rumen microbes. In coculture experiments with *Butyrivibrio proteoclasticus*, several *Methanobrevibacter* adhesins were upregulated and co-aggregates of both cell types were observed [122]. Interestingly, formate utilisation genes were also upregulated. Butyrate, acetate (or lactate), formate, carbon dioxide, and hydrogen gas are major fermentation products of *Butyrivibrio* during growth on xylan [123, 124]. Hydrogen and formate may be utilized by *Methanobrevibacter* in this syntrophic interaction. In addition to free-living methanogens in the rumen fluid, methanogens which are extra- and intracellularly associated with ciliate protozoa are relevant contributors to methane production. In ruminants, more than one-third of the methane may be produced by these consortia [125].

One might consider that the presence or absence of methanogens in vertebrates generally depends on the diet or the presence of specific anatomical differentiations of the gut and all herbivorous animals harbor the entire anaerobic food chain. Systematic analysis of the methane production in guts of 253 vertebrate species revealed that methane production and hence the presence of relevant amounts of methanogens does depend on the phylogenetic lineage of the animal rather than on the diet or the anatomy of the digestive system [126]. In some phylogenetic lineages like ostriches,

intestinal methanogens got lost irrespective of the diet. Methanogens are also missing within the large lineages of Carnivora/Chiroptera/Eulipotyphla (formerly Insectivora), even in herbivorous pandas (Ursidae/Carnivora). Though in all other large lineages methane producers dominate, nonproducers occur also in several “branches” of these lineages. Generally, the results imply that once the methanogens got lost in the course of evolution, they did not reappear in the descending lineages [126]. One special case with respect to the bird digestive system has drawn attention recently. The hoatzin (*Opisthocomus hoazin*) is the only known example for foregut fermentation in birds similar to the ruminants [127, 128]. The rumen methanogens found in hoatzins are more closely related to ruminant strains than to methanogens found in feces of other birds, though the composition of the methanogen community and the phylotypes themselves were still distinct from those found in ruminants [129].

In the intestines of primates, including humans, Archaea are present. *Methanobrevibacter smithii* as the dominant species draws particular attention: a syntrophic interaction between *Methanobrevibacter* and *Bacteroides thetaiotaomicron*, as studied in gnotobiotic mice, may affect the energy balance of the host [130]. *Methanobrevibacter* utilizes the *Bacteroides* fermentation product formate. This syntrophy obviously determines the expression of *Bacteroides* enzymes: the pathway directed towards formate and acetate production is upregulated, whereas alternative pathways towards propionate and butyrate are downregulated. The ongoing human microbiome project will soon update our knowledge on archaeal diversity and putative function in humans.

6. Concluding Remarks

Interactions between Archaea and other organisms are definitely as specific as interactions with symbiotic Bacteria prokaryotes. Up to now, the mechanisms of surface recognition are still poorly understood. The prominent “model” pathogens *Escherichia coli*, *Salmonella typhimurium*, *Pseudomonas aeruginosa*, and *Vibrio* spp. greatly extended our knowledge on specific interactions of Proteobacteria with animal host tissue. However, model organisms of this kind are still missing in the archaeal world, due to the lack of easy manageable molecular tools for functional studies, in particular with respect to the generation of mutant strains. In addition, we are still far away from even a rough estimate of the true sizes of the large archaeal clades. Hence, we are still unable to explore the diverse ways how Archaea may interact with each other. The description of the few very diverse cases that we know—considering the fundamental differences, for example, between *Nanoarchaeum* and *Ignicoccus* or the SM1 and sulfur reducer interaction—gives us an impression on the diverse ways how Archaea may interact and how diverse the mechanisms may have to be expected (see Table 1).

The symbiotic interaction between prokaryotes also leads to the question if the first eukaryote may be an offspring of a symbiotic interaction between an Archaeum and a Bacterium ([131] and references therein). Though the different roles of ancient Archaea and Bacteria are still

speculative, it becomes more and more obvious that tight symbiosis between both prokaryotic cell types also direct us to the roots of eukaryote evolution.

Acknowledgments

The Authors are indebted to a DFG grant to M. Hoppert (Ho 1830/2-1) and a fellowship of the Studienstiftung des Deutschen Volkes to A. Dreier. This is Courant Research Centre Geobiology publication no. 112.

References

- [1] G. B. West, W. H. Woodruff, and J. H. Brown, "Allometric scaling of metabolic rate from molecules and mitochondria to cells and mammals," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, supplement 1, pp. 2473–2478, 2002.
- [2] N. Klitgord and D. Segrè, "Environments that induce synthetic microbial ecosystems," *PLoS Computational Biology*, vol. 6, no. 11, Article ID e1001002, 2010.
- [3] N. Dubilier, C. Bergin, and C. Lott, "Symbiotic diversity in marine animals: the art of harnessing chemosynthesis," *Nature Reviews Microbiology*, vol. 6, no. 10, pp. 725–740, 2008.
- [4] D. G. Thanassi, J. B. Bliska, and P. J. Christie, "Surface organelles assembled by secretion systems of Gram-negative bacteria: diversity in structure and function," *FEMS Microbiology Reviews*, vol. 36, no. 6, pp. 1046–1082, 2012.
- [5] H. P. Horz and G. Conrads, "The discussion goes on: what is the role of Euryarchaeota in humans?" *Archaea*, vol. 2010, Article ID 967271, 8 pages, 2010.
- [6] C. Moissl-Eichinger and H. Huber, "Archaeal symbionts and parasites," *Current Opinion in Microbiology*, vol. 14, no. 3, pp. 364–370, 2011.
- [7] J. N. Reeve and C. Schleper, "Archaea: very diverse, often different but never bad?" *Current Opinion in Microbiology*, vol. 14, no. 3, pp. 271–273, 2011.
- [8] A. Koerdt, S. Jachlewski, A. Ghosh, J. Wingender, B. Siebers, and S. V. Albers, "Complementation of *Sulfolobus solfataricus* PBL2025 with an α -mannosidase: effects on surface attachment and biofilm formation," *Extremophiles*, vol. 16, no. 1, pp. 115–125, 2011.
- [9] C. Moissl, C. Rudolph, and R. Huber, "Natural communities of novel archaea and bacteria with a string-of-pearls-like morphology: molecular analysis of the bacterial partners," *Applied and Environmental Microbiology*, vol. 68, no. 2, pp. 933–937, 2002.
- [10] C. Moissl, R. Rachel, A. Briegel, H. Engelhardt, and R. Huber, "The unique structure of archaeal 'hami', highly complex cell appendages with nano-grappling hooks," *Molecular Microbiology*, vol. 56, no. 2, pp. 361–370, 2005.
- [11] C. Rudolph, C. Moissl, R. Henneberger, and R. Huber, "Ecology and microbial structures of archaeal/bacterial strings-of-pearls communities and archaeal relatives thriving in cold sulfidic springs," *FEMS Microbiology Ecology*, vol. 50, no. 1, pp. 1–11, 2004.
- [12] S. Nickell, R. Hegerl, W. Baumeister, and R. Rachel, "Pyrodictum cannulae enter the periplasmic space but do not enter the cytoplasm, as revealed by cryo-electron tomography," *Journal of Structural Biology*, vol. 141, no. 1, pp. 34–42, 2003.
- [13] B. J. Baker, L. R. Comolli, G. J. Dick et al., "Enigmatic, ultrasmall, uncultivated Archaea," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 19, pp. 8806–8811, 2010.
- [14] S. V. Albers, Z. Szabó, and A. J. M. Driessen, "Protein secretion in the Archaea: multiple paths towards a unique cell surface," *Nature Reviews Microbiology*, vol. 4, no. 7, pp. 537–547, 2006.
- [15] M. Pohlschroder, A. Ghosh, M. Tripepi, and S. V. Albers, "Archaeal type IV pilus-like structures-evolutionarily conserved prokaryotic surface organelles," *Current Opinion in Microbiology*, vol. 14, no. 3, pp. 357–363, 2011.
- [16] D. J. Näther, R. Rachel, G. Wanner, and R. Wirth, "Flagella of *Pyrococcus furiosus*: multifunctional organelles, made for swimming, adhesion to various surfaces, and cell-cell contacts," *Journal of Bacteriology*, vol. 188, no. 19, pp. 6915–6923, 2006.
- [17] S. Fröls, M. Ajon, M. Wagner et al., "UV-inducible cellular aggregation of the hyperthermophilic archaeon *Sulfolobus solfataricus* is mediated by pili formation," *Molecular Microbiology*, vol. 70, no. 4, pp. 938–952, 2008.
- [18] A. Koerdt, J. Gödeke, J. Berger, K. M. Thormann, and S. V. Albers, "Crenarchaeal biofilm formation under extreme conditions," *PLoS ONE*, vol. 5, no. 11, Article ID e14104, 2010.
- [19] S. Schopf, G. Wanner, R. Rachel, and R. Wirth, "An archaeal bi-species biofilm formed by *Pyrococcus furiosus* and *Methanopyrus kandleri*," *Archives of Microbiology*, vol. 190, no. 3, pp. 371–377, 2008.
- [20] M. Tripepi, S. Imam, and M. Pohlschröder, "*Haloferax volcanii* flagella are required for motility but are not involved in PibD-dependent surface adhesion," *Journal of Bacteriology*, vol. 192, no. 12, pp. 3093–3102, 2010.
- [21] J. R. Leadbetter and J. A. Breznak, "Physiological ecology of *Methanobrevibacter cuticularis* sp. nov. and *Methanobrevibacter curvatus* sp. nov., isolated from the hindgut of the termite *Reticulitermes flavipes*," *Applied and Environmental Microbiology*, vol. 62, no. 10, pp. 3620–3631, 1996.
- [22] V. B. Tran, S. M. J. Fleiszig, D. J. Evans, and C. J. Radke, "Dynamics of flagellum-and pilus-mediated association of *Pseudomonas aeruginosa* with contact lens surfaces," *Applied and Environmental Microbiology*, vol. 77, no. 11, pp. 3644–3652, 2011.
- [23] A. K. Zgair and S. Chhibber, "Adhesion of *Stenotrophomonas maltophilia* to mouse tracheal mucus is mediated through flagella," *Journal of Medical Microbiology*, vol. 60, no. 7, pp. 1032–1037, 2011.
- [24] U. Gophna, E. Z. Ron, and D. Graur, "Bacterial type III secretion systems are ancient and evolved by multiple horizontal-transfer events," *Gene*, vol. 312, no. 1-2, pp. 151–163, 2003.
- [25] U. Küper, C. Meyer, V. Müller, R. Rachel, and H. Huber, "Energized outer membrane and spatial separation of metabolic processes in the hyperthermophilic Archaeon *Ignicoccus hospitalis*," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 7, pp. 3152–3156, 2010.
- [26] T. Burghardt, D. J. Näther, B. Junglas, H. Huber, and R. Rachel, "The dominating outer membrane protein of the hyperthermophilic Archaeum *Ignicoccus hospitalis*: a novel pore-forming complex," *Molecular Microbiology*, vol. 63, no. 1, pp. 166–176, 2007.
- [27] T. Burghardt, B. Junglas, F. Siedler, R. Wirth, H. Huber, and R. Rachel, "The interaction of *Nanoarchaeum equitans* with *Ignicoccus hospitalis*: proteins in the contact site between two

- cells," *Biochemical Society Transactions*, vol. 37, no. 1, pp. 127–132, 2009.
- [28] E. Waters, M. J. Hohn, I. Ahel et al., "The genome of *Nanoarchaeum equitans*: insights into early archaeal evolution and derived parasitism," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 22, pp. 12984–12988, 2003.
 - [29] U. Jahn, M. Gallenberger, W. Paper et al., "*Nanoarchaeum equitans* and *Ignicoccus hospitalis*: new insights into a unique, intimate association of two archaea," *Journal of Bacteriology*, vol. 190, no. 5, pp. 1743–1750, 2008.
 - [30] H. A. Barker, "Studies on the methane fermentation. V. Biochemical activities of *Methanobacterium omelianskii*," *Journal of Biological Chemistry*, vol. 137, no. 1, pp. 153–167, 1941.
 - [31] M. P. Bryant, E. A. Wolin, M. J. Wolin, and R. S. Wolfe, "*Methanobacillus omelianskii*, a symbiotic association of two species of bacteria," *Archiv für Mikrobiologie*, vol. 59, no. 1–3, pp. 20–31, 1967.
 - [32] C. A. Reddy, M. P. Bryant, and M. J. Wolin, "Characteristics of S organism isolated from *Methanobacillus omelianskii*," *Journal of Bacteriology*, vol. 109, no. 2, pp. 539–545, 1972.
 - [33] B. Eichler and B. Schink, "Oxidation of primary aliphatic alcohols by *Acetobacterium carbinolicum* sp. nov., a homoacetogenic anaerobe," *Archives of Microbiology*, vol. 140, no. 2–3, pp. 147–152, 1984.
 - [34] C. Wallrabenstein, E. Hauschild, and B. Schink, "*Syntrophobacter pfennigii* sp. nov., new syntrophically propionate-oxidizing anaerobe growing in pure culture with propionate and sulfate," *Archives of Microbiology*, vol. 164, no. 5, pp. 346–352, 1995.
 - [35] M. P. Bryant, L. L. Campbell, C. A. Reddy, and M. R. Crabill, "Growth of desulfovibrio in lactate or ethanol media low in sulfate in association with H₂ utilizing methanogenic bacteria," *Applied and Environmental Microbiology*, vol. 33, no. 5, pp. 1162–1169, 1977.
 - [36] A. Ben-Bassat, R. Lamed, and J. G. Zeikus, "Ethanol production by thermophilic bacteria: metabolic control of end product formation in *Thermoanaerobium brockii*," *Journal of Bacteriology*, vol. 146, no. 1, pp. 192–199, 1981.
 - [37] R. Klemps, H. Cypionka, F. Widdel, and N. Pfennig, "Growth with hydrogen, and further physiological characteristics of *Desulfotomaculum species*," *Archives of Microbiology*, vol. 143, no. 2, pp. 203–208, 1985.
 - [38] H. Imachi, Y. Sekiguchi, Y. Kamagata, S. Hanada, A. Ohashi, and H. Harada, "*Pelotomaculum thermopropionicum* gen. nov., sp. nov., an anaerobic, thermophilic, syntrophic propionate-oxidizing bacterium," *International Journal of Systematic and Evolutionary Microbiology*, vol. 52, no. 5, pp. 1729–1735, 2002.
 - [39] T. Shimoyama, S. Kato, S. Ishii, and K. Watanabe, "Flagellum mediates symbiosis," *Science*, vol. 323, no. 5921, p. 1574, 2009.
 - [40] A. J. M. Stams and C. M. Plugge, "Electron transfer in syntrophic communities of anaerobic bacteria and archaea," *Nature Reviews Microbiology*, vol. 7, no. 8, pp. 568–577, 2009.
 - [41] W. S. Reeburgh, "Methane consumption in Cariaco Trench waters and sediments," *Earth and Planetary Science Letters*, vol. 28, no. 3, pp. 337–344, 1976.
 - [42] K. Knittel and A. Boetius, "Anaerobic oxidation of methane: progress with an unknown process," *Annual Review of Microbiology*, vol. 63, pp. 311–334, 2009.
 - [43] P. Regnier, A. W. Dale, S. Arndt, D. E. LaRowe, J. Mogollón, and P. Van Cappellen, "Quantitative analysis of anaerobic oxidation of methane (AOM) in marine sediments: a modeling perspective," *Earth-Science Reviews*, vol. 106, no. 1–2, pp. 105–130, 2011.
 - [44] G. Aloisi, C. Pierre, J. M. Rouchy, J. P. Foucher, and J. Woodside, "Methane-related authigenic carbonates of Eastern Mediterranean Sea mud volcanoes and their possible relation to gas hydrate destabilisation," *Earth and Planetary Science Letters*, vol. 184, no. 1, pp. 321–338, 2000.
 - [45] J. Peckmann, E. Gischler, W. Oschmann, and J. Reitner, "An Early Carboniferous seep community and hydrocarbon-derived carbonates from the Harz Mountains, Germany," *Geology*, vol. 29, no. 3, pp. 271–274, 2001.
 - [46] G. Aloisi, I. Bouloubassi, S. K. Heijs et al., "CH₄-consuming microorganisms and the formation of carbonate crusts at cold seeps," *Earth and Planetary Science Letters*, vol. 203, no. 1, pp. 195–203, 2002.
 - [47] E. De Boever, D. Birgel, V. Thiel et al., "The formation of giant tubular concretions triggered by anaerobic oxidation of methane as revealed by archaeal molecular fossils (Lower Eocene, Varna, Bulgaria)," *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 280, no. 1–2, pp. 23–36, 2009.
 - [48] E. De Boever, D. Birgel, P. Muchez, J. Peckmann, L. Dimitrov, and R. Swennen, "Fabric and formation of grapestone concretions within an unusual ancient methane seep system (Eocene, Bulgaria)," *Terra Nova*, vol. 23, no. 1, pp. 56–61, 2011.
 - [49] S. B. Ziegenbalg, D. Birgel, L. Hoffmann-Sell, C. Pierre, J. M. Rouchy, and J. Peckmann, "Anaerobic oxidation of methane in hypersaline Messinian environments revealed by ¹³C-depleted molecular fossils," *Chemical Geology*, vol. 292–293, no. 1, pp. 140–148, 2012.
 - [50] J. Peckmann, A. Reimer, U. Luth et al., "Methane-derived carbonates and authigenic pyrite from the northwestern Black Sea," *Marine Geology*, vol. 177, no. 1–2, pp. 129–150, 2001.
 - [51] A. Y. Lein, M. V. Ivanov, N. V. Pimenov, and M. B. Gulin, "Geochemical peculiarities of the carbonate constructions formed during microbial oxidation of methane under anaerobic conditions," *Microbiology*, vol. 71, no. 1, pp. 78–90, 2002.
 - [52] W. Michaelis, R. Seifert, K. Nauhaus et al., "Microbial reefs in the black sea fueled by anaerobic oxidation of methane," *Science*, vol. 297, no. 5583, pp. 1013–1015, 2002.
 - [53] J. Reitner, A. Reimer, G. Schumann, V. Thiel, and J. Peckmann, "Methane-derived carbonate build-ups and associated microbial communities at cold seeps on the lower Crimean shelf (Black Sea)," *Facies*, vol. 51, no. 1–4, pp. 66–79, 2005.
 - [54] C. Wrede, C. Heller, J. Reitner, and M. Hoppert, "Correlative light/electron microscopy for the investigation of microbial mats from Black Sea Cold Seeps," *Journal of Microbiological Methods*, vol. 73, no. 2, pp. 85–91, 2008.
 - [55] C. Wrede, V. Krukenberg, A. Dreier, J. Reitner, C. Heller, and M. Hoppert, "Detection of metabolic key enzymes of methane turnover processes in cold seep microbial biofilms," *Geomicrobiology Journal*, vol. 30, no. 3, pp. 214–227, 2013.
 - [56] K. U. Hinrichs, J. M. Hayes, S. P. Sylva, P. G. Brewert, and E. F. DeLong, "Methane-consuming archaeobacteria in marine sediments," *Nature*, vol. 398, no. 6730, pp. 802–805, 1999.
 - [57] V. J. Orphan, K. U. Hinrichs, W. Ussler III et al., "Comparative analysis of methane-oxidizing archaea and sulfate-reducing bacteria in anoxic marine sediments," *Applied and Environmental Microbiology*, vol. 67, no. 4, pp. 1922–1934, 2001.
 - [58] H. Niemann, T. Lösekann, D. De Beer et al., "Novel microbial communities of the Haakon Mosby mud volcano and their

- role as a methane sink," *Nature*, vol. 443, no. 7113, pp. 854–858, 2006.
- [59] H. J. Mills, R. J. Martinez, S. Story, and P. A. Sobecky, "Characterization of microbial community structure in Gulf of Mexico gas hydrates: comparative analysis of DNA- and RNA-derived clone libraries," *Applied and Environmental Microbiology*, vol. 71, no. 6, pp. 3235–3247, 2005.
 - [60] K. G. Lloyd, L. Lapham, and A. Teske, "An anaerobic methane-oxidizing community of ANME-1b archaea in hypersaline gulf of Mexico sediments," *Applied and Environmental Microbiology*, vol. 72, no. 11, pp. 7218–7230, 2006.
 - [61] R. J. Martinez, H. J. Mills, S. Story, and P. A. Sobecky, "Prokaryotic diversity and metabolically active microbial populations in sediments from an active mud volcano in the Gulf of Mexico," *Environmental Microbiology*, vol. 8, no. 10, pp. 1783–1796, 2006.
 - [62] C. Heller, M. Hoppert, and J. Reitner, "Immunological localization of coenzyme M reductase in anaerobic methane-oxidizing archaea of ANME 1 and ANME 2 type," *Geomicrobiology Journal*, vol. 25, no. 3–4, pp. 149–156, 2008.
 - [63] A. Boetius, K. Ravensschlag, C. J. Schubert et al., "A marine microbial consortium apparently mediating anaerobic oxidation methane," *Nature*, vol. 407, no. 6804, pp. 623–626, 2000.
 - [64] L. Schreiber, T. Holler, K. Knittel, A. Meyerdierks, and R. Amann, "Identification of the dominant sulfate-reducing bacterial partner of anaerobic methanotrophs of the ANME-2 clade," *Environmental Microbiology*, vol. 12, no. 8, pp. 2327–2340, 2010.
 - [65] V. J. Orphan, C. H. House, K. U. Hinrichs, K. D. McKee-gan, and E. F. DeLong, "Multiple archaeal groups mediate methane oxidation in anoxic cold seep sediments," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 11, pp. 7663–7668, 2002.
 - [66] E. Durisch-Kaiser, L. Klausner, B. Wehrli, and C. Schubert, "Evidence of intense archaeal and bacterial methanotrophic activity in the Black Sea water column," *Applied and Environmental Microbiology*, vol. 71, no. 12, pp. 8099–8106, 2005.
 - [67] G. Eller, L. Kanel, and M. Krüger, "Cooccurrence of aerobic and anaerobic methane oxidation in the water column of Lake Plußsee," *Applied and Environmental Microbiology*, vol. 71, no. 12, pp. 8925–8928, 2005.
 - [68] C. J. Schubert, M. J. L. Coolen, L. N. Neretin et al., "Aerobic and anaerobic methanotrophs in the Black Sea water column," *Environmental Microbiology*, vol. 8, no. 10, pp. 1844–1856, 2006.
 - [69] T. Treude, V. Orphan, K. Knittel, A. Gieseke, C. H. House, and A. Boetius, "Consumption of methane and CO₂ by methanotrophic microbial mats from gas seeps of the anoxic Black Sea," *Applied and Environmental Microbiology*, vol. 73, no. 7, pp. 2271–2283, 2007.
 - [70] A. Pernthaler, A. E. Dekas, C. T. Brown, S. K. Goffredi, T. Embaye, and V. J. Orphan, "Diverse syntrophic partnerships from deep-sea methane vents revealed by direct cell capture and metagenomics," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 19, pp. 7052–7057, 2008.
 - [71] K. Knittel, T. Lösekann, A. Boetius, R. Kort, and R. Amann, "Diversity and distribution of methanotrophic archaea at cold seeps," *Applied and Environmental Microbiology*, vol. 71, no. 1, pp. 467–479, 2005.
 - [72] T. M. Hoehler, M. J. Alperin, D. B. Albert, and C. S. Martens, "Field and laboratory studies of methane oxidation in an anoxic marine sediment: evidence for a methanogen-sulfate reducer consortium," *Global Biogeochemical Cycles*, vol. 8, no. 4, pp. 451–463, 1994.
 - [73] S. D. Wankel, M. M. Adams, D. T. Johnston, C. M. Hansel, S. B. Joye, and P. R. Girguis, "Anaerobic methane oxidation in metalliferous hydrothermal sediments: influence on carbon flux and decoupling from sulfate reduction," *Environmental Microbiology*, vol. 14, no. 10, pp. 2726–2740, 2012.
 - [74] M. Krüger, A. Meyerdierks, F. O. Glöckner et al., "A conspicuous nickel protein in microbial mats that oxidize methane anaerobically," *Nature*, vol. 426, no. 6968, pp. 878–881, 2003.
 - [75] S. J. Hallam, N. Putnam, C. M. Preston et al., "Reverse methanogenesis: testing the hypothesis with environmental genomics," *Science*, vol. 305, no. 5689, pp. 1457–1462, 2004.
 - [76] S. Scheller, M. Goenrich, R. Boecher, R. K. Thauer, and B. Jaun, "The key nickel enzyme of methanogenesis catalyses the anaerobic oxidation of methane," *Nature*, vol. 465, no. 7298, pp. 606–608, 2010.
 - [77] K. Nauhaus, A. Boetius, M. Krüger, and F. Widdel, "In vitro demonstration of anaerobic oxidation of methane coupled to sulphate reduction in sediment from a marine gas hydrate area," *Environmental Microbiology*, vol. 4, no. 5, pp. 296–305, 2002.
 - [78] K. Nauhaus, T. Treude, A. Boetius, and M. Krüger, "Environmental regulation of the anaerobic oxidation of methane: a comparison of ANME-I and ANME-II communities," *Environmental Microbiology*, vol. 7, no. 1, pp. 98–106, 2005.
 - [79] F. Widdel, A. Boetius, and R. Rabus, "Anaerobic biodegradation of hydrocarbons including methane," in *The Prokaryotes*, M. Dworkin, S. Falkow, E. Rosenberg, K. H. Schleifer, and E. Stackebrandt, Eds., pp. 1028–1049, Springer, New York, NY, USA, 2006.
 - [80] J. J. Moran, E. J. Beal, J. M. Vrentas, V. J. Orphan, K. H. Freeman, and C. H. House, "Methyl sulfides as intermediates in the anaerobic oxidation of methane," *Environmental Microbiology*, vol. 10, no. 1, pp. 162–173, 2008.
 - [81] G. Wegener, H. Niemann, M. Elvert, K. U. Hinrichs, and A. Boetius, "Assimilation of methane and inorganic carbon by microbial communities mediating the anaerobic oxidation of methane," *Environmental Microbiology*, vol. 10, no. 9, pp. 2287–2298, 2008.
 - [82] R. K. Thauer and S. Shima, "Methane as fuel for anaerobic microorganisms," *Annals of the New York Academy of Sciences*, vol. 1125, pp. 158–170, 2008.
 - [83] J. Milucka, T. G. Ferdelman, L. Polerecky et al., "Zero-valent sulphur is a key intermediate in marine methane oxidation," *Nature*, vol. 491, no. 7425, pp. 541–546, 2012.
 - [84] A. E. Dekas, R. S. Poretsky, and V. J. Orphan, "Deep-Sea archaea fix and share nitrogen in methane-consuming microbial consortia," *Science*, vol. 326, no. 5951, pp. 422–426, 2009.
 - [85] E. J. Beal, C. H. House, and V. J. Orphan, "Manganese- and iron-dependent marine methane oxidation," *Science*, vol. 325, no. 5937, pp. 184–187, 2009.
 - [86] K. F. Ettwig, M. K. Butler, D. Le Paslier et al., "Nitrite-driven anaerobic methane oxidation by oxygenic bacteria," *Nature*, vol. 464, no. 7288, pp. 543–548, 2010.
 - [87] F. Muller, T. Brissac, N. Le Bris, H. Felbeck, and O. Gros, "First description of giant Archaea (Thaumarchaeota) associated with putative bacterial ectosymbionts in a sulfidic marine habitat," *Environmental Microbiology*, vol. 12, no. 8, pp. 2371–2383, 2010.

- [88] P. B. Eckburg, E. M. Bik, C. N. Bernstein et al., "Microbiology: diversity of the human intestinal microbial flora," *Science*, vol. 308, no. 5728, pp. 1635–1638, 2005.
- [89] J. H. P. Hackstein, A. Akhmanova, B. Boxma, H. R. Harhangi, and F. G. J. Voncken, "Hydrogenosomes: eukaryotic adaptations to anaerobic environments," *Trends in Microbiology*, vol. 7, no. 11, pp. 441–447, 1999.
- [90] M. Müller, M. Mentel, J. J. van Hellemond et al., "Biochemistry and evolution of anaerobic energy metabolism in eukaryotes," *Microbiology and Molecular Biology Reviews*, vol. 76, no. 2, pp. 444–495, 2012.
- [91] J. J. A. Van Bruggen, K. B. Zwart, and J. G. F. Hermans, "Isolation and characterization of *Methanoplanus endosymbiosus* sp. nov., an endosymbiont of the marine sapropelic ciliate *Metopus contortus* Quennerstedt," *Archives of Microbiology*, vol. 144, no. 4, pp. 367–374, 1986.
- [92] N. Narayanan, B. Krishnakumar, V. N. Anupama, and V. B. Manilal, "*Methanoseta* sp., the major archaeal endosymbiont of *Metopus* es," *Research in Microbiology*, vol. 160, no. 8, pp. 600–607, 2009.
- [93] J. H. P. Hackstein and T. A. van Alen, "Methanogens in the gastro-intestinal tract of animals," in *Microbiology Monographs: (Endo)Symbiotic Methanogenic Archaea*, J. H. P. Hackstein, Ed., pp. 115–142, Springer, Heidelberg, Germany, 2010.
- [94] S. Wagener, C. F. Bardele, and N. Pfennig, "Functional integration of *Methanobacterium formicicum* into the anaerobic ciliate *Trimyema compressum*," *Archives of Microbiology*, vol. 153, no. 5, pp. 496–501, 1990.
- [95] B. J. Finlay and T. Fenchel, "Polymorphic bacterial symbionts in the anaerobic ciliated protozoon *Metopus*," *FEMS Microbiology Letters*, vol. 79, no. 2-3, pp. 187–190, 1991.
- [96] B. J. Finlay, T. M. Embley, and T. Fenchel, "A new polymorphic methanogen, closely related to *Methanocorpusculum parvum*, living in stable symbiosis within the anaerobic ciliate *Trimyema* sp," *Journal of General Microbiology*, vol. 139, no. 2, pp. 371–378, 1993.
- [97] R. Gross, F. Vavre, A. Heddi, G. D. D. Hurst, E. Zchori-Fein, and K. Bourtzis, "Immunity and symbiosis," *Molecular Microbiology*, vol. 73, no. 5, pp. 751–759, 2009.
- [98] A. Moya, J. Peretó, R. Gil, and A. Latorre, "Learning how to live together: genomic insights into prokaryote-animal symbioses," *Nature Reviews Genetics*, vol. 9, no. 3, pp. 218–229, 2008.
- [99] F. Husník, T. Chrudimský, and V. Hypška, "Multiple origins of endosymbiosis within the Enterobacteriaceae (gamma-Proteobacteria): convergence of complex phylogenetic approaches," *BMC Biology*, vol. 9, no. 87, pp. 1–17, 2011.
- [100] N. S. Webster and L. L. Blackall, "What do we really know about sponge-microbial symbioses," *The ISME Journal*, vol. 3, no. 1, pp. 1–3, 2009.
- [101] M. Maldonado, N. Cortadellas, M. I. Trillas, and K. Rützler, "Endosymbiotic yeast maternally transmitted in a marine sponge," *The Biological Bulletin*, vol. 209, no. 2, pp. 94–106, 2005.
- [102] T. Pape, F. Hoffmann, N. V. Quéric, K. Von Juterzenka, J. Reitner, and W. Michaelis, "Dense populations of Archaea associated with the demosponge *Tentorium semisuberites* Schmidt, 1870 from Arctic deep-waters," *Polar Biology*, vol. 29, no. 8, pp. 662–667, 2006.
- [103] A. S. Turque, D. Batista, C. B. Silveira et al., "Environmental shaping of sponge associated archaeal communities," *PLoS ONE*, vol. 5, no. 12, Article ID e15774, 2010.
- [104] C. Brochier-Armanet, B. Boussau, S. Gribaldo, and P. Forterre, "Mesophilic crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota," *Nature Reviews Microbiology*, vol. 6, no. 3, pp. 245–252, 2008.
- [105] A. Spang, R. Hatzepichler, C. Brochier-Armanet et al., "Distinct gene set in two different lineages of ammonia-oxidizing archaea supports the phylum Thaumarchaeota," *Trends in Microbiology*, vol. 18, no. 8, pp. 331–340, 2010.
- [106] C. M. Preston, K. Y. Wu, T. F. Molinski, and E. F. Delong, "A psychrophilic crenarchaeon inhabits a marine sponge: *Cenarchaeum symbiosum* gen. nov., sp. nov.," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 93, no. 13, pp. 6241–6246, 1996.
- [107] H. Margot, C. Acebal, E. Toril, R. Amils, and J. L. Fernandez Puentes, "Consistent association of crenarchaeal Archaea with sponges of the genus *Axinella*," *Marine Biology*, vol. 140, no. 4, pp. 739–745, 2002.
- [108] M. Han, F. Liu Zhang, Z. Li, and H. Lin, "Bacterial and Archaeal symbionts in the south China Sea sponge *Phakellia fusca*: community structure, relative abundance, and ammonia-oxidizing populations," *Marine Biotechnology*, vol. 14, no. 6, pp. 701–713, 2012.
- [109] R. Radax, T. Rattei, A. Lanzen et al., "Metatranscriptomics of the marine sponge *Geodia barretti*: tackling phylogeny and function of its microbial community," *Environmental Microbiology*, vol. 14, no. 5, pp. 1308–1324, 2012.
- [110] D. Steger, P. Ettinger-Epstein, S. Whalan et al., "Diversity and mode of transmission of ammonia-oxidizing archaea in marine sponges," *Environmental Microbiology*, vol. 10, no. 4, pp. 1087–1094, 2008.
- [111] M. Martínez-García, P. Stief, M. Díaz-Valdés et al., "Ammonia-oxidizing Crenarchaeota and nitrification inside the tissue of a colonial ascidian," *Environmental Microbiology*, vol. 10, no. 11, pp. 2991–3001, 2008.
- [112] J. M. Beman, K. J. Roberts, L. Wegley, F. Rohwer, and C. A. Francis, "Distribution and diversity of archaeal ammonia monooxygenase genes associated with corals," *Applied and Environmental Microbiology*, vol. 73, no. 17, pp. 5642–5647, 2007.
- [113] N. Siboni, E. Ben-Dov, A. Sivan, and A. Kushmaro, "Geographic specific coral-associated ammonia-oxidizing archaea in the northern Gulf of Eilat (Red Sea)," *Microbiology Ecology*, vol. 64, no. 1, pp. 18–24, 2012.
- [114] M. Pester, C. Schleper, and M. Wagner, "The Thaumarchaeota: an emerging view of their phylogeny and ecophysiology," *Current Opinion in Microbiology*, vol. 14, no. 3, pp. 300–306, 2011.
- [115] M. Pester and A. Brune, "Hydrogen is the central free intermediate during lignocellulose degradation by termite gut symbionts," *The ISME Journal*, vol. 1, no. 6, pp. 551–565, 2007.
- [116] J. H. P. Hackstein and C. K. Stumm, "Methane production in terrestrial arthropods," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 91, no. 12, pp. 5441–5445, 1994.
- [117] A. Brauman, J. Doré, P. Eggleton, D. Bignell, J. A. Breznak, and M. D. Kane, "Molecular phylogenetic profiling of prokaryotic communities in guts of termites with different feeding habits," *FEMS Microbiology Ecology*, vol. 35, no. 1, pp. 27–36, 2001.
- [118] S. E. Donovan, K. J. Purdy, M. D. Kane, and P. Eggleton, "Comparison of Euryarchaea strains in the guts and food-soil of the soil-feeding termite *Cubitermes fungifaber* across

- different soil types,” *Applied and Environmental Microbiology*, vol. 70, no. 7, pp. 3884–3892, 2004.
- [119] B. J. Tindall, H. N. M. Ross, and W. D. Grant, “*Natronobacterium* gen. nov. and *Natronococcus* gen. nov., two new genera of haloalkaliphilic archaeobacteria,” *Systematic and Applied Microbiology*, vol. 5, no. 1, pp. 41–57, 1984.
- [120] K. L. Denman, G. Brasseur, A. Chidthaisong et al., “Couplings between changes in the climate system and biogeochemistry,” in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, S. Solomon, D. Qin, M. Manning et al., Eds., pp. 500–587, Cambridge University Press, New York, NY, USA.
- [121] W. B. Whitman and R. S. Wolfe, “Activation of the methylreductase system from *Methanobacterium bryantii* by ATP,” *Journal of Bacteriology*, vol. 154, no. 2, pp. 640–649, 1983.
- [122] S. C. Leahy, W. J. Kelly, E. Altermann et al., “The genome sequence of the rumen methanogen *Methanobrevibacter ruminantium* reveals new possibilities for controlling ruminant methane emissions,” *PloS one*, vol. 5, no. 1, p. e8926, 2010.
- [123] R. B. Hespell, R. Wolf, and R. J. Bothast, “Fermentation of xylans by *Butyrivibrio fibrisolvens* and other ruminal bacteria,” *Applied and Environmental Microbiology*, vol. 53, no. 12, pp. 2849–2853, 1987.
- [124] T. L. Miller and S. E. Jenesel, “Enzymology of butyrate formation by *Butyrivibrio fibrisolvens*,” *Journal of Bacteriology*, vol. 138, no. 1, pp. 99–104, 1979.
- [125] B. J. Finlay, G. Esteban, K. J. Clarke, A. G. Williams, T. M. Embley, and R. P. Hirt, “Some rumen ciliates have endosymbiotic methanogens,” *FEMS Microbiology Letters*, vol. 117, no. 2, pp. 157–161, 1994.
- [126] J. H. P. Hackstein, “Eukaryotic molecular biodiversity: systematic approaches for the assessment of symbiotic associations,” *Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology*, vol. 72, no. 1, pp. 63–76, 1997.
- [127] A. Grajal, S. D. Strahl, R. Parra, M. G. Dominguez, and A. Neher, “Foregut fermentation in the hoatzin, a neotropical leaf-eating bird,” *Science*, vol. 245, no. 4923, pp. 1236–1238, 1989.
- [128] F. Godoy-Vitorino, R. E. Ley, Z. Gao et al., “Bacterial community in the crop of the hoatzin, a neotropical folivorous flying bird,” *Applied and Environmental Microbiology*, vol. 74, no. 19, pp. 5905–5912, 2008.
- [129] A. D. G. Wright, K. S. Northwood, and N. E. Obispo, “Rumen-like methanogens identified from the crop of the folivorous South American bird, the hoatzin (*Opisthocomus hoazin*),” *The ISME Journal*, vol. 3, no. 10, pp. 1120–1126, 2009.
- [130] B. S. Samuel and J. I. Gordon, “A humanized gnotobiotic mouse model of host-archaeal-bacterial mutualism,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, no. 26, pp. 10011–10016, 2006.
- [131] T. Thiergart, G. Landan, M. Schenk, T. Dagan, and W. F. Martin, “An evolutionary network of genes present in the eukaryote common ancestor polls genomes on eukaryotic and mitochondrial origin,” *Genome Biology and Evolution*, vol. 4, no. 4, pp. 466–485, 2012.