

Clinical Study

Dexmedetomidine versus Magnesium Sulfate as Adjunct during Anesthesia for Laparoscopic Colectomy

Pierre Zarif,¹ Ahmed Abdelaal Ahmed Mahmoud,² Mohamed Mohamed Abdelhaq,¹ Hany M. S. Mikhail,³ and Ahmed Farag³

¹Department of Anesthesiology, Faculty of Medicine, Cairo University, Giza, Egypt

²Department of Anesthesiology, Faculty of Medicine, Beni-Suef University, Beni Suef, Egypt

³Department of General Surgery, Faculty of Medicine, Cairo University, Giza, Egypt

Correspondence should be addressed to Ahmed Abdelaal Ahmed Mahmoud; carnitin7@yahoo.com

Received 12 December 2015; Revised 11 February 2016; Accepted 15 February 2016

Academic Editor: Yukio Hayashi

Copyright © 2016 Pierre Zarif et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objectives. To compare dexmedetomidine versus magnesium during laparoscopic colectomy. **Patients and Methods.** 51 patients were randomly allocated into 3 groups: group C (control) received saline infusion, group D dexmedetomidine 1 g/kg and then 0.4 g/kg/hr, and group M MgSO₄ 2 g and then 15 g/kg/min. Intraoperative hemodynamics were measured before and 1 min after intubation (T1 and T2), before and 5 min after peritoneal insufflation (T3 and T4), before and 5 min after 30° Trendelenburg position (T5 and T6), 5 min after resuming flat position (T7), 5 min after peritoneal deflations (T8), after extubation (T9), and at time of admission to PACU (T10). Recovery time and degree of sedation were assessed. **Results.** HR and MAP were significantly higher in T2, T4, and T6 compared to T1, T3, and T5, respectively, in all groups with lower measurements in groups D and M compared to group C. Mean of collective measurements was significantly higher in group C. Recovery time and sedation score were significantly higher in groups D and M. Time to Aldrete score of ≥9 was significantly longer in groups D and M. **Conclusion.** Both drugs ameliorate the pressor responses during LC with a nonsignificant difference. This study is registered with PACTR201602001481308.

1. Introduction

Nowadays laparoscopic surgery is going to be the first choice for surgical management of various indications, especially with the well-trained laparoscopic surgeon. The benefits of minimal access techniques include less pain, early mobilization, shorter hospital stay, and better cosmetic results, which have further increased its applications [1, 2].

During general anesthesia laryngoscopy, tracheal intubation and extubation are the critical events provoking transient but marked sympathoadrenal response manifesting by hypertension and tachycardia. In addition, in laparoscopic surgery, CO₂ is routinely used to create pneumoperitoneum, which causes increased plasma level of catecholamine and vasopressin. Elevation of intra-abdominal pressure with raised diaphragm causes various adverse effects on the cardiovascular system such as decreased cardiac output, elevated arterial

pressure, and increased systemic and pulmonary vascular resistance leading to hypertension and tachycardia [3, 4].

Severe increases in arterial pressure can be a risk factor in patients with preexisting hypertension, ischaemic heart disease, or increased intracranial pressure. Opioids, alpha-2-adrenergic agonists, beta-blocking agents, or vasodilators are often used to avoid circulatory response to pneumoperitoneum [5].

Magnesium is well known to block the release of catecholamines from both adrenergic nerve terminals and the adrenal gland, and intravenous magnesium sulfate inhibits catecholamine release associated with laryngoscopy. Moreover, magnesium produces vasodilator effect by acting directly on blood vessels, and high-dose magnesium attenuates vasopressin-stimulated vasoconstriction [6, 7].

On the other hand, dexmedetomidine is an alpha-2-adrenergic agonist; it has properties of analgesia,

sympatholytic effect, and sedation without respiratory depression. It decreases opioid requirements and stress response to surgery ensuring a stable hemodynamic state. Its distribution half-life is around six min, so it can be used to attenuate the stress response to laryngoscopy [8–10].

Our study aimed at assessment of hemodynamic changes during laparoscopic colectomy under pneumoperitoneum in 30° Trendelenburg position and the impact of intraoperative dexmedetomidine and magnesium sulfate infusions on these changes.

2. Patients and Methods

The study was carried out at Cairo University Hospital since Jan 2010 till Aug 2013. After approval of the study protocol by the Local Ethical Committee and obtaining patients' written fully informed consents, all patients assigned for laparoscopic colorectal resection were enrolled in the study. Patients who required preoperative neoadjuvant therapy, are generally unfit for resection surgery, or had inoperable lesion requiring relieving or dysfunctioning colostomy were excluded from the study. Patients with cardiopulmonary diseases, renal or liver impairment, or allergy to any of the used drugs were also excluded from the study. Morbidly obese with body mass index (BMI) >35 kg/m² was not enrolled in the study.

Patients were randomly, using sealed envelopes, allocated to 3 groups: group C assigned to receive saline infusion as placebo, group D assigned to receive dexmedetomidine infusion, and group M assigned to receive intravenous infusion of magnesium sulfate.

Fifteen minutes before surgery, all patients were premedicated with midazolam 1–2 mg and given drugs as follows: group C: saline infusion 20 mL over 15 min; group D: dexmedetomidine loading 1 µg/kg in 20 mL normal saline over 15 min; and Group M: magnesium sulfate loading 2 g, 20 mL over 15 min. Baseline data (*T*₀) including systolic blood pressure (SBP), diastolic blood pressure (DBP), mean arterial blood pressure (MAP), and heart rate (HR) were noninvasively determined and monitored and then anesthesia was conducted with fentanyl 1 µg/kg and propofol 2 mg/kg, followed by cisatracurium (0.15–0.2 mg/kg) to facilitate tracheal intubation, and then maintenance of loaded drugs was continued as follow: Group C: saline infusion as placebo; Group D: dexmedetomidine infusion at rate of 0.4 µg/kg/hr; and Group M: magnesium sulfate infusion at rate of 15 µg/kg/min.

Controlled ventilation was adjusted to an end tidal CO₂ concentration of 30–35 mmHg and to ensure SpO₂ of 97%. Anesthesia was maintained in all groups by sevoflurane 2 MAC with 50% oxygen and 50% air and maintenance doses of cisatracurium. All groups received ketorolac 30 mg IV slowly and paracetamol 1 g IV at the beginning of surgery and Ringer acetate solution was given at a rate of 10 mL/kg/hr during anesthesia. Infusions and sevoflurane inhalation were stopped just at the end of surgery, then atropine sulfate 0.02 mg/kg and neostigmine 0.04 mg/kg were administered IV for reversal of muscle relaxation, and the patient was extubated. Following extubation, patients were maintained on supplemental O₂ until being awake in the recovery room.

Any pain in the recovery room was managed by pethidine 1 mg/kg IV if needed. The recovery time was estimated in all groups as the time elapsed since stoppage of sevoflurane and infusions, till the time the patient achieves a modified Aldrete scoring of ≥9. The degree of sedation was assessed at the end of end of surgery and 30 and 120 minutes thereafter using the Brussels Sedation Scale (BSS), [11] where 1 = sedated and unarousable, 2 = sedated but responding to painful not auditory stimuli, 3 = sedated but responding to auditory stimuli, 4 = awake and calm, and 5 = agitated.

Intraoperative hemodynamic measures were determined before intubation (*T*₁), 1 min after intubation (*T*₂), before and 5 min after peritoneal insufflation (*T*₃ and *T*₄), before and 5 min after tilting the table to 30° Trendelenburg position (*T*₅ and *T*₆), 5 min after resuming flat position (*T*₇), 5 min after peritoneal deflation (*T*₈), after extubation (*T*₉), and at time of admission to PACU (*T*₁₀). Mean of collective intraoperative changes of HR, SBP, DBP, and MAP was calculated collectively for measurements *T*₁–*T*₉.

3. Statistical Analysis and Sample Size Calculation

Sample power was calculated according to Kraemer and Theimann [12] using their proposed figure. Considering that laparoscopic colectomy is the unusual procedure for resection of colorectal cancer and that the hemodynamic changes during laparoscopic surgery under pneumoperitoneum are a frequent event and also considering a standard nomogram [13], we determined a sample size of 17 patients per group to obtain a study power of 80% to detect a difference at the significance level of 5%.

Collected data was presented as mean (±SD), numbers, and percentages, as appropriate. Categorical variables were analyzed using chi-square (χ²) test. Continuous variables were analyzed using unpaired Student's *t*-test or univariate two-group repeated measures "mixed-design" analysis of variance (ANOVA) with post hoc Dunnett's test as appropriate. Nominal and nonnormally distributed variables were analyzed using Mann-Whitney *U* test. Statistical analysis was done by using the computer program SPSS (Statistical Package for Social Sciences), Version 20, 2011. *P* value < 0.05 was considered statistically significant.

4. Results

The study included 51 patients, 36 males and 15 females, with mean age of 61.2 ± 7.3, range: 45–68 years. There were eight patients of normal weight, 25 patients were overweight, and 18 patients were obese with a total mean BMI of 29.3 ± 3.1, range: 22.6–34.1 kg/m². Details of patients' demographic data are shown in Table 1. There was a nonsignificant (*P* > 0.05) difference between studied groups with regard to demographic data.

Throughout the duration of surgery, HR showed variability during times of stress with significantly (*P* < 0.05) higher HR records at *T*₂, *T*₄, and *T*₆ compared to that recorded at *T*₁, *T*₃, and *T*₅, respectively, in all studied patients.

TABLE 1: Patients' demographic data.

		Group C	Group D	Group M	P value
Age (years)	Strata				
	<50	1 (5.9%)	3 (17.6%)	2 (11.8%)	$P_1 = 0.067$
	50–60	3 (17.6%)	3 (17.6%)	4 (23.5%)	$P_2 = 0.086$
	>60	13 (76.5%)	11 (64.7%)	11 (64.7%)	$P_3 = 0.421$
	Total	62.5 ± 4.9	60.9 ± 7.1	60.3 ± 7.3	$P_1 = 0.156$ $P_2 = 0.193$ $P_3 = 0.791$
Gender	Males	11 (64.7%)	13 (76.5%)	12 (70.6%)	$P_1 = 0.086$
	Females	6 (35.3%)	4 (23.5%)	5 (29.4%)	$P_2 = 0.106$ $P_3 = 0.519$
BMI data	Weight	83.4 ± 6.4	84.5 ± 7.5	84.6 ± 8	$P_1 = 0.482$ $P_2 = 0.307$ $P_3 = 0.619$
	Height	169.2 ± 4.2	169.7 ± 4.3	169.6 ± 4	$P_1 = 0.308$ $P_2 = 0.295$ $P_3 = 0.456$
	Strata				
	Normal	3 (17.6%)	2 (11.8%)	3 (17.6%)	$P_1 = 0.071$
	Overweight	7 (41.2%)	10 (58.8%)	8 (47.1%)	$P_2 = 0.532$
Obese	7 (41.2%)	5 (29.4%)	6 (35.3%)	$P_3 = 0.102$	
Total	29.2 ± 2.8	29.4 ± 3.3	29.5 ± 3.4	$P_1 = 0.285$ $P_2 = 0.231$ $P_3 = 0.495$	
ASA	Grade I	11 (64.7%)	12 (70.6%)	13 (76.5%)	$P_1 = 0.077$
	Grade II	4 (23.5%)	2 (11.8%)	3 (17.6%)	$P_2 = 0.059$
	Grade III	2 (11.8%)	3 (17.6%)	1 (5.9%)	$P_3 = 0.119$

Data are presented as mean \pm SD and numbers; percentages are in parenthesis. BMI: body mass index; P_1 : significance of difference between groups C and D; P_2 : significance of difference between groups C and M; P_3 : significance of difference between groups D and M.

TABLE 2: Mean HR records determined throughout the operative time.

	T0	T1	T2	T3	T4	T5	T6	T7	T8	T9	T10
Group C	82.9 ± 4	66 ± 2.7	69.5 ± 3.8	70.8 ± 4	74.2 ± 3.6	71.8 ± 3	75 ± 5	71.1 ± 3.5	73.2 ± 4	74.7 ± 4	76.9 ± 5.2
		$P = 0.001$		$P = 0.001$		$P = 0.009$					
Group D	83.4 ± 4.5	61.2 ± 2.9	65 ± 5.3	62.7 ± 3.8	66.8 ± 5.8	63.4 ± 4.8	67.1 ± 3.8	64.2 ± 2	64.8 ± 2.9	66.7 ± 3.8	71.8 ± 3.4
		$P = 0.041$		$P = 0.011$		$P = 0.026$					
Group M	83.7 ± 3.1	62.1 ± 2.1	66.8 ± 3.9	63.1 ± 3.9	67.1 ± 6.1	64.2 ± 4.9	69.2 ± 7.4	65.7 ± 5.5	65.8 ± 3.9	67.4 ± 5.5	72.4 ± 3
		$P = 0.005$		$P = 0.029$		$P = 0.036$					
Intergroup difference	P_1	=0.001	=0.029	=0.0008	=0.001	0.0008	=0.001	=0.0008	=0.0006	=0.001	=0.031
	P_2	=0.022	=0.112	=0.002	=0.001	=0.0008	=0.044	=0.004	=0.0006	=0.007	=0.003
	P_3	=0.387	=0.255	=0.925	=0.859	=0.604	=0.379	=0.649	=0.475	=0.777	=0.534

P : significance between times identified; P_1 : significance of difference between groups C and D; P_2 : significance of difference between groups C and M; P_3 : significance of difference between groups D and M.

The ameliorative effect of studied drugs was evident and manifested as higher significance of differences between these measures in group C than in groups D and M. Concerning the HR records throughout operative time till PACU transfer, HR records were significantly lower in groups D and M compared to group C at all times of recording with nonsignificantly

($P > 0.05$) lower HR records in group D compared to group M (Table 2).

Throughout the duration of surgery, MAP showed variability during times of stress with significantly ($P < 0.05$) higher MAP measurements at T2, T4, and T6 compared to that measured at T1, T3, and T5, respectively, in all studied

TABLE 3: Mean (\pm SD) MAP records determined throughout the operative time.

	T0	T1	T2	T3	T4	T5	T6	T7	T8	T9	T10
Group C	94.6 \pm 3.7	73.9 \pm 2.9	80.5 \pm 2.8	75.1 \pm 3	78.4 \pm 2.6	74.9 \pm 2.1	78.3 \pm 3.5	75.9 \pm 3.2	76.6 \pm 3	79 \pm 3.3	87.7 \pm 4.1
		$P = 0.0007$		$P = 0.003$		$P = 0.0009$					
Group D	95.7 \pm 3.8	66.8 \pm 2.8	72.3 \pm 3.7	68.2 \pm 3.3	72.4 \pm 4.6	67.6 \pm 4.1	73.6 \pm 2.7	72.7 \pm 2.1	73.6 \pm 2.5	75.4 \pm 3.1	84.4 \pm 2.7
		$P = 0.001$		$P = 0.019$		$P = 0.001$					
Group M	94.1 \pm 3.8	68 \pm 3.7	74.4 \pm 2.6	69.6 \pm 2.5	73.4 \pm 4.8	69 \pm 3.2	73.5 \pm 5.1	72.7 \pm 3.3	73.4 \pm 2.7	76.4 \pm 3.2	85.1 \pm 3.5
		$P = 0.0007$		$P = 0.002$		$P = 0.002$					
Intergroup difference	P_1	=0.001	=0.0009	=0.0009	=0.002	=0.0009	=0.002	=0.028	=0.036	=0.035	=0.029
	P_2	=0.001	=0.001	=0.0007	=0.007	=0.0009	=0.017	=0.041	=0.033	=0.042	=0.047
	P_3	=0.136	=0.112	=0.177	=0.445	=0.224	=0.756	=0.932	=0.813	=0.868	=0.435

P : significance between times identified; P_1 : significance of difference between groups C and D; P_2 : significance of difference between groups C and M; P_3 : significance of difference between groups D and M.

TABLE 4: Mean (\pm SD) of collective hemodynamic variable estimated from T1 to T9 period.

	Group C	Group D	Group M
HR (beats/min)	71.8 \pm 2	64.7 \pm 1.3	65.7 \pm 1.8
		$P_1 = 0.0006$	$P_2 = 0.0008$ $P_3 = 0.046$
SBP (mmHg)	84.5 \pm 1.7	80.2 \pm 1.3	79.8 \pm 1.8
		$P_1 = 0.0009$	$P_2 = 0.0009$ $P_3 = 0.234$
DBP (mmHg)	72.9 \pm 2.2	67.1 \pm 1.1	68.5 \pm 2.3
		$P_1 = 0.0008$	$P_2 = 0.0009$ $P_3 = 0.059$
MAP (mmHg)	76.8 \pm 1.6	71.5 \pm 0.8	72.3 \pm 1.4
		$P_1 = 0.0007$	$P_2 = 0.0009$ $P_3 = 0.076$

P_1 : significance of difference between groups C and D; P_2 : significance of difference between groups C and M; P_3 : significance of difference between groups D and M.

patients. However, both studied drugs provided better blood pressure stabilization as manifested by the lower significance of differences between these measures in groups D and M than in group C. MAP measurements determined throughout operative time till PACU transfer were significantly lower in groups D and M compared to group C at all times of recording with nonsignificant ($P > 0.05$) difference in favor of group D (Table 3).

Mean of collective HR records from T1 to T9 was significantly ($P < 0.05$) higher in group C compared to groups D and M with significantly ($P < 0.05$) lower HR records in group D compared to group M. Mean of collective blood pressure measurements since T1 to T9 was significantly ($P < 0.05$) higher in group C compared to groups D and M with nonsignificantly ($P > 0.05$) lower measurements in group D compared to group M (Table 4).

Studied patients showed nonsignificant ($P > 0.05$) difference concerning operative and surgical postoperative data (Table 5). Both studied drugs significantly prolonged postanesthetic recovery with significantly ($P < 0.05$) higher

frequency of sedated patients at the end of surgery and 15 and 30 minutes thereafter compared to group C. However, at 15 and 30 minutes after surgery, the frequency of awaked patients showed nonsignificant ($P > 0.05$) difference between the three groups, but the frequency of agitated patients was significantly ($P < 0.05$) higher in group C compared to other groups. Mean time till reaching modified Aldrete score of ≥ 9 was significantly ($P < 0.05$) longer in groups D and M compared to group C with a nonsignificant ($P > 0.05$) difference between both groups (Table 6).

5. Discussion

The key element in laparoscopic surgery is the creation of pneumoperitoneum and the physiologic effect CO₂. Moreover, pelvic and lower abdominal laparoscopic surgeries have a special concern wherein the body must be positioned in Trendelenburg position with leg elevation up to 30° to give room for the surgeon to deal with pelvic organs. This positioning imposed more hemodynamic burden secondary to the more diaphragmatic splinting action and increased venous return [14, 15].

In this study, the patients were assigned for laparoscopic colorectal resection so the body must be positioned in Trendelenburg position. The results showed that HR and MAP measurements were significantly higher in T2, T4, and T6 compared to T1, T3, and T5, respectively, in all groups with lower measurements in groups D and M compared to group C. Also, the mean of collective HR records was significantly higher in group C compared to groups D and M ($P < 0.05$), with significantly ($P < 0.05$) lower HR records in group D compared to group M. Mean of collective blood pressure measurements was significantly ($P < 0.05$) higher in group C compared to groups D and M with nonsignificantly ($P > 0.05$) lower measurements in group D compared to group M. Studied patients also showed that both studied drugs significantly prolonged postanesthetic recovery compared to group C ($P < 0.05$). However, at 15 and 30 minutes after surgery, the frequency of awaked patients showed nonsignificant difference between the three groups. But the frequency of agitated patients was significantly higher

TABLE 5: Operative and immediate postoperative surgical data.

	Group C	Group D	Group M	P value
Operative time (min)	123.6 ± 15.7	129.4 ± 12.4	128.8 ± 23.8	$P_1 = 0.186$ $P_2 = 0.569$ $P_3 = 0.850$
Amount of blood loss (mL)	309.1 ± 94.1	281.5 ± 50.3	295.9 ± 97.2	$P_1 = 0.480$ $P_2 = 0.501$ $P_3 = 0.687$
Need for blood transfusion				
Yes	5 (29.4%)	3 (17.6%)	4 (23.5%)	$P_1 = 0.079$
No	12 (70.6%)	14 (82.4%)	13 (76.5%)	$P_2 = 0.236$ $P_3 = 0.318$
Time till 1st flatus passage (days)	1.8 ± 0.7	2 ± 0.7	1.7 ± 0.8	$P_1 = 0.448$ $P_2 = 0.713$ $P_3 = 0.332$
Time till 1st oral fluid intake (days)	3 ± 0.8	2.6 ± 0.7	3.1 ± 0.9	$P_1 = 0.198$ $P_2 = 0.480$ $P_3 = 0.170$
Time till 1st oral solid food (days)	3.1 ± 1	3.5 ± 0.6	3.3 ± 0.9	$P_1 = 0.083$ $P_2 = 0.408$ $P_3 = 0.396$
Surgical complications				
Yes	3 (17.6%)	4 (23.5%)	2 (11.8%)	$P_1 = 0.467$
No	14 (82.4%)	13 (76.5%)	15 (88.2%)	$P_2 = 0.691$ $P_3 = 0.106$
PO hospital stay (days)	6.9 ± 1.2	7.1 ± 1.5	7 ± 1.6	$P_1 = 0.703$ $P_2 = 0.908$ $P_3 = 0.581$

Data are presented as mean ± SD and numbers; percentages are in parenthesis. P_1 : significance of difference between groups C and D; P_2 : significance of difference between groups C and M; P_3 : significance of difference between groups D and M.

in group C compared to other groups. Mean time till reaching modified Aldrete score of ≥ 9 was significantly ($P < 0.05$) longer in groups D and M compared to group C with a nonsignificant ($P > 0.05$) difference between both groups.

The results of this study about effect of magnesium sulfate on hemodynamics intraoperative correlate well with Lee and Kwon [16], who found that preoperative intravenous magnesium sulfate attenuated arterial pressure increases during the pre-delivery period and was recommended as an adjuvant during general anesthesia for caesarean section to avoid perioperative pressor response resulting from inadequate anesthesia, analgesia, or both of them [16]. Jee et al. [5] also found that intravenous magnesium sulfate before pneumoperitoneum attenuates arterial pressure increases during laparoscopic cholecystectomy and this attenuation is apparently related to reductions in the release of catecholamine, vasopressin, or both of them [5]. Kalra et al. [17] assessed which of magnesium or clonidine attenuates hemodynamic stress response to pneumoperitoneum better and found that systolic blood pressure was significantly higher in control group as compared to study groups during pneumoperitoneum with no significant difference between magnesium and clonidine given in dose of 1 $\mu\text{g}/\text{kg}$. However, the difference was better in patients receiving clonidine in dose of 1.5 $\mu\text{g}/\text{kg}$ [17].

The effect of magnesium on hemodynamics due to interact in the activation of membrane Ca-ATPase and Na-K-ATPase is involved in transmembrane ion exchanges during depolarization and repolarization phases, thus acting as a cell membrane stabilizer and also as an intracytoplasmic organelles stabilizer [7]. This calcium inhibitory effect of Mg causes central arteriolar vasodilatation and acts against vasospasm. Another mechanism could involve the reduction of catecholamine release with sympathetic stimulation, thereby decreasing the stress response to surgery. The analgesic effect of magnesium due to it blocks N-methyl-D-aspartate (NMDA) receptor which plays a significant role in the mechanisms underlying central sensitization in the spinal cord and is crucial for the establishment of several pain states [6, 7].

On the other hand, the results of this study about effect of dexmedetomidine on hemodynamics intraoperative correlate well with Kato et al. [18] who found that dexmedetomidine weakens arterial pressure preservation and HR responses after thigh cuff deflation, suggesting attenuated cardiovascular reflexes [18]. Klinger et al. [19] assessed the hemodynamic impact of dexmedetomidine administration in a large cohort of patients undergoing routine noncardiac surgery and found that a significantly higher percentage of patients in dexmedetomidine group met the composite

TABLE 6: Recovery data of studied patients.

	Group C	Group D	Group M	<i>P</i> value
<i>Brussels Sedation Scale (BSS)</i>				
At end of surgery				
I	5 (29.4%)	3 (17.6%)	2 (11.8%)	$P_1 = 0.069$
II	12 (70.6%)	14 (82.4%)	15 (88.2%)	$P_2 = 0.005$
15 min after end of surgery				$P_3 = 0.575$
II	2 (11.8%)	5 (29.4%)	6 (35.3%)	$P_1 = 0.005$
III	8 (47.1%)	9 (52.9%)	8 (47.1%)	$P_2 = 0.005$
IV	7 (41.1%)	3 (17.7%)	3 (17.7%)	$P_3 = 0.239$
30 min after end of surgery				
II	0	1 (5.9%)	2 (11.8%)	$P_1 = 0.006$
III	2 (11.8%)	3 (17.7%)	4 (23.5%)	$P_2 = 0.003$
IV	9 (52.9%)	10 (58.8%)	9 (52.9%)	$P_3 = 0.093$
V	6 (35.3%)	3 (17.7%)	2 (11.8%)	
Time till reaching Aldrete score ≥ 9				
5 min	2 (11.8%)	0	0	
10 min	3 (17.6%)	1 (5.9%)	2 (11.8%)	
15 min	6 (35.3%)	2 (11.8%)	1 (5.9%)	$P_1 = 0.0009$
20 min	3 (17.6%)	4 (23.5%)	3 (17.6%)	$P_2 = 0.0008$
25 min	1 (5.9%)	6 (35.3%)	5 (29.4%)	$P_3 = 0.075$
30 min	1 (5.9%)	3 (17.6%)	4 (23.5%)	
>30 min	1 (5.9%)	1 (5.9%)	2 (11.8%)	
Total time	16.5 \pm 8.1	23.2 \pm 6.4	24.1 \pm 7.5	$P_1 = 0.0008$ $P_2 = 0.0007$ $P_3 = 0.180$

Data are presented as mean \pm SD and numbers; percentages are in parenthesis. P_1 : significance of difference between groups C and D; P_2 : significance of difference between groups C and M; P_3 : significance of difference between groups D and M.

endpoint criteria with no significant difference in the overall incidence of intraoperative hypotension and concluded that dexmedetomidine administration was associated with hemodynamic stability with lowering of hemodynamic stress response without more hypotension or bradycardia [19].

Concerning laparoscopic surgery, the obtained results supported what was previously reported by Smania et al. [20] who found that dexmedetomidine efficiently blocks the hemodynamic responses to nociceptive stimuli when combined with inhaled isoflurane for anesthesia of children submitted to laparoscopic video appendectomy [20].

The effect of dexmedetomidine on hemodynamics is due to decrease of sympathetic outflow from the locus coeruleus. Its sympatholytic effect leads to decrease of mean arterial blood pressure (MAP) and heart rate (HR) by reducing norepinephrine release. Its analgesic actions are mediated by releasing of substance P from the dorsal horn of the spinal cord [8].

As a further support for efficacy of both drugs, Bryskin and Weldon [21] used a combination of dexmedetomidine and magnesium sulfate for hemodynamic control during laparoscopic resection of pheochromocytoma and reported that cardiovascular stability was achieved [21].

Unfortunately, there was no comparative study between the hemodynamic effects of dexmedetomidine and magnesium sulfate during laparoscopic surgery; however, other comparative studies provided evidence for their superiority to other drugs used for the same target, where Salman et al. [22] compared dexmedetomidine with remifentanyl in desflurane-based ambulatory gynecologic laparoscopic surgery and demonstrated that dexmedetomidine infusion causes a relatively slow recovery with reduced postoperative nausea, vomiting, and analgesic requirements, and similar hemodynamics compared to remifentanyl, and may be an alternative to remifentanyl in ambulatory anesthesia [22].

6. Conclusion

Intraoperative infusion of either dexmedetomidine or magnesium sulfate could ameliorate the pressor responses to anesthetic and surgical manipulations during laparoscopic colectomy under pneumoperitoneum in 30° Trendelenburg position.

Disclosure

Research is affiliated to Department of Anesthesia, Faculty of Medicine, Cairo University.

Competing Interests

The authors declare that they have no competing interests.

References

- [1] S. Pither, L. S. Bayonne Manou, J. M. Mandji Lawson, T. D. Tchanchou, R. Tchoua, and J. P. Ponties, "Surgical approaches to hysterectomy," *Sante*, vol. 21, no. 2, pp. 79–81, 2011.
- [2] T. Hatakeyama, M. Nakanishi, Y. Murayama et al., "Laparoscopic resection for colorectal cancer improves short-term outcomes in very elderly colorectal cancer patients," *Surgical Laparoscopy, Endoscopy and Percutaneous Techniques*, vol. 23, no. 6, pp. 532–535, 2013.
- [3] Y. Wang, Y.-T. Wang, Y.-Y. Jiang, Y.-H. Lin, and Z.-D. Zou, "Effect of different position and CO₂ pneumoperitoneum on deep vein hemodynamics of lower limb," *Zhonghua Wei Chang Wai Ke Za Zhi*, vol. 12, no. 4, pp. 361–363, 2009.
- [4] S. M. Meftahuzzaman, M. M. Islam, K. K. Chowdhury et al., "Haemodynamic and end tidal CO₂ changes during laparoscopic cholecystectomy under general anaesthesia," *Mymensingh Medical Journal*, vol. 22, no. 3, pp. 473–477, 2013.
- [5] D. Jee, D. Lee, S. Yun, and C. Lee, "Magnesium sulphate attenuates arterial pressure increase during laparoscopic cholecystectomy," *British Journal of Anaesthesia*, vol. 103, no. 4, pp. 484–489, 2009.
- [6] S. Herroeder, M. E. Schönherr, S. G. De Hert, and M. W. Hollmann, "Magnesium—essentials for anesthesiologists," *Anesthesiology*, vol. 114, no. 4, pp. 971–993, 2011.
- [7] S.-H. Do, "Magnesium: a versatile drug for anesthesiologists," *Korean Journal of Anesthesiology*, vol. 65, no. 1, pp. 4–8, 2013.
- [8] J. Afonso and F. Reis, "Dexmedetomidine: current role in anesthesia and intensive care," *Revista Brasileira de Anestesiologia*, vol. 62, no. 1, pp. 118–133, 2012.
- [9] E. Farag, M. Argalious, A. Abd-Elsayed, Z. Ebrahim, and D. J. Doyle, "The use of dexmedetomidine in anesthesia and intensive care: a review," *Current Pharmaceutical Design*, vol. 18, no. 38, pp. 6257–6265, 2012.
- [10] H. Ihmsen and T. I. Saari, "Dexmedetomidine. Pharmacokinetics and pharmacodynamics," *Anaesthetist*, vol. 61, no. 12, pp. 1059–1066, 2012.
- [11] O. Detriche, J. Berré, J. Massaut, and J.-L. Vincent, "The Brussels sedation scale: use of a simple clinical sedation scale can avoid excessive sedation in patients undergoing mechanical ventilation in the intensive care unit," *British Journal of Anaesthesia*, vol. 83, no. 5, pp. 698–701, 1999.
- [12] H. C. Kraemer and S. Theimann, *How Many Subjects?: Statistical Power Analysis in Research*, SAGE Publications, Newbury Park, Calif, USA, 1987.
- [13] K. R. Murphy and B. Myers, *Statistical Power Analysis: A Simple and General Model for Traditional and Modern Hypothesis*, Lawrence Erlbaum Associates, New York, NY, USA, 2nd edition, 2003.
- [14] B. K. Parikh, V. R. Shah, P. R. Modi, B. P. Butala, and G. P. Parikh, "Anaesthesia for laparoscopic kidney transplantation: influence of trendelenburg position and CO₂ pneumoperitoneum on cardiovascular, respiratory and renal function," *Indian Journal of Anaesthesia*, vol. 57, no. 3, pp. 253–258, 2013.
- [15] Y. Y. Jo, J. Y. Lee, M. G. Lee, and H. J. Kwak, "Effects of high positive end-expiratory pressure on haemodynamics and cerebral oxygenation during pneumoperitoneum in the Trendelenburg position," *Anaesthesia*, vol. 68, no. 9, pp. 938–943, 2013.
- [16] D. H. Lee and I. C. Kwon, "Magnesium sulphate has beneficial effects as an adjuvant during general anaesthesia for Caesarean section," *British Journal of Anaesthesia*, vol. 103, no. 6, pp. 861–866, 2009.
- [17] N. K. Kalra, A. Verma, A. Agarwal, and H. D. Pandey, "Comparative study of intravenously administered clonidine and magnesium sulfate on hemodynamic responses during laparoscopic cholecystectomy," *Journal of Anaesthesiology Clinical Pharmacology*, vol. 27, no. 3, pp. 344–348, 2011.
- [18] J. Kato, Y. Ogawa, W. Kojima, K. Aoki, S. Ogawa, and K. Iwasaki, "Cardiovascular reflex responses to temporal reduction in arterial pressure during dexmedetomidine infusion: a double-blind, randomized, and placebo-controlled study," *British Journal of Anaesthesia*, vol. 103, no. 4, pp. 561–565, 2009.
- [19] R. Y. Klinger, W. D. White, B. Hale, A. S. Habib, and E. Bennett-Guerrero, "Hemodynamic impact of dexmedetomidine administration in 15,656 noncardiac surgical cases," *Journal of Clinical Anesthesia*, vol. 24, no. 3, pp. 212–220, 2012.
- [20] M. C. Smania, J. P. Piva, and P. C. R. Garcia, "Dexmedetomidine in anesthesia of children submitted to videolaparoscopic appendectomy: a double-blind, randomized and placebo-controlled study," *Revista da Associacao Medica Brasileira*, vol. 54, no. 4, pp. 308–313, 2008.
- [21] R. Bryskin and B. C. Weldon, "Dexmedetomidine and magnesium sulfate in the perioperative management of a child undergoing laparoscopic resection of bilateral pheochromocytomas," *Journal of Clinical Anesthesia*, vol. 22, no. 2, pp. 126–129, 2010.
- [22] N. Salman, S. Uzun, F. Coskun, M. A. Salman, A. E. Salman, and U. Aypar, "Dexmedetomidine as a substitute for remifentanyl in ambulatory gynecologic laparoscopic surgery," *Saudi Medical Journal*, vol. 30, no. 1, pp. 77–81, 2009.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

