Research Article

Complexes of Cu(II) Ions and Noncovalent Interactions in Systems with L-Aspartic Acid and Cytidine-5'-Monophosphate

Romualda Bregier-Jarzebowska, Anna Gasowska, and Lechosław Lomozik

Faculty of Chemistry, Adam Mickiewicz University, Grunwaldzka 6, 60-780 Poznań, Poland

Correspondence should be addressed to Lechosław Lomozik, lomozik@amu.edu.pl

Received 19 February 2008; Revised 15 May 2008; Accepted 19 June 2008

Recommended by Henryk Kozłowski

Interactions between aspartic acid (Asp) and cytidine-5-monophosphate (CMP) in metal-free systems as well as the coordination of Cu(II) ions with the above ligands were studied. The composition and overall stability constants of the species formed in those systems were determined by the potentiometric method, and the interaction centres in the ligands were identified by the spectral methods UV-Vis, EPR, NMR, and IR. In metal-free systems, the formation of adducts, in which each ligand has both positive and negative reaction centres, was established. The main reaction centres in Asp are the oxygen atoms of carboxyl groups and the nitrogen atom of the amine group, while the main reaction centre in CMP at low pH is the N(3) atom. With increasing pH, the efficiency of the phosphate group of the nucleotide in the interactions significantly increases, and the efficiency of carboxyl groups in Asp decreases. The noncovalent reaction centres in the ligands are simultaneously the potential sites of metal-ion coordination. The mode of coordination in the complexes formed in the ternary systems was established. The sites of coordination depend clearly on the solution pH. In the molecular complexes $ML \cdots L$, metallation involves the oxygen atoms of the carboxyl groups of the amino acid, while the protonated nucleotide is in the outer coordination sphere and interacts noncovalently with the anchoring $CuH_x(Asp)$ species. The influence of the metal ions on the weak interactions between the biomolecules was established.

Copyright © 2008 Romualda Bregier-Jarzebowska et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

Interactions between metal ions and nucleic acids or their fragments affect the character of many biological processes including that of genetic information transfer [1–6]. The effective centres of coordination with metal ions are the donor nitrogen atoms N(3) from pyrimidine bases and the oxygen atoms from the phosphate groups of the nucleotide. These centres are also the sites of noncovalent interactions with the other bioligands present in living organisms, such as small organic polycations, polyamines, or amino acids [7, 8].

Aspartic acid (Asp) is a naturally occurring amino acid. Along with glutamic acid, it acts as a neurotransmitter in the central nervous system [9–14]. Aspartic acid takes part in thermogenic processes induced by prostaglandin E_1 (PGE_1) [15] and is a component of the active centres of some enzymes. It influences the solubility and ionic character of proteins, protects the liver against the toxic effect

of drugs, participates in the generation of ribonucleotides thereby enhancing the effectiveness of the immunological system of the organism, and prevents the destruction of neurons and the brain. The presence of metal ions in living organisms modifies the character of bioprocesses. The reactions between the amino acid and the metal ions are considered as models of the processes which take place at the molecular level in the metal/protein system.

Although studies of the systems of metals with dicarboxylic amino acids have been carried out since the 1970s, [16–19], no definite conclusions as to the mode of coordination have been obtained, in particular in the ternary systems, which is related to the fact that aspartic acid has three functional groups (one amine group and two carboxyl ones). To our best knowledge, no information has been reported on interactions in the metal-free systems of aspartic acid/nucleotide or on the character of interactions in ternary systems including metal ions.

This paper presents the results of a study on the coordination of Cu(II) ions with aspartic acid and cytidine-5'-monophosphate (CMP) and the interactions of these bioligands in metal-free systems.

2. EXPERIMENTAL

Cytidine 5'-monophosphate, $C_9H_{14}N_3O_8P$, and L-aspartic acid, $C_4H_7NO_4$, were purchased from Sigma-Aldrich and were used without further purification. $Cu(NO_3)_2$ bought in POCH Gliwice (Poland) was twice recrystallized from H_2O before use. The method of determination of Cu(II) concentration in a parent solution of a concentration of about 2.4×10^{-2} M was described earlier [20, 21]. Potentiometric studies were performed on a Methrom 702 SM Titrino with a glass electrode Methrom 6.0233.100 calibrated in terms of hydrogen ion concentration [22] with a preliminary use of borax (pH = 9.225) and phthalate (pH = 4.002) standard buffers. The concentrations of the CMP and Asp were $5 \cdot 10^{-3}$ in the metal-free systems and from $1 \cdot 10^{-3}$ to $2.5 \cdot 10^{-3}$ M in the systems with Cu(II). The ratio of ligand1:ligand2 in the metal-free systems was 1:1, metal:ligand1 was 1:2.5, and metal:ligand1:ligand2 were 1:1:1 or 1:2.5:2.5 in the ternary systems (ligand1-Asp, ligand2-CMP). Potentiometric titrations were performed at the ionic strength $\mu = 0.1$ M (KNO_3), at $20 \pm 1^\circ C$ under helium, using as a titrant CO_2 -free NaOH solution (about 0.2 M). For each system a series of 10 titrations was made; the initial volume of the sample was 30 cm^3 . No precipitate formation was observed in the entire pH range studied. Calculations were performed using 100–350 points for each job. The selection of the models and the determination of the stability constants of the complexes were made using the SUPERQUAD program [23], whereas the distribution of particular forms was determined by the HALTAFALL program [24]. The computer procedures used for the purpose, choice of the models, and the criteria of verification of results are described in [25–29]. The samples for ^{13}C NMR and ^{31}P NMR investigation were prepared by dissolving appropriate amounts of ligands and $Cu(NO_3)_2$ in D_2O and adjusting pH by the addition of NaOD (or $(C_2H_5)_4NOH$) and DNO_3 , correcting pH-readings (a pH-meter C5-501 made by Elmetron) according to the formula: $pD = pH_{\text{readings}} + 0.40$ [30]. The concentration of the ligands in the samples was 0.05 M, and the concentration ratio of Cu(II) to CMP and Asp was 1:200:200. ^{13}C NMR spectra were recorded on an NMR Gemini 300 VT Varian spectrometer using dioxane as an internal standard. The positions of ^{13}C NMR signals were converted to the TMS scale. ^{31}P NMR spectra were taken on an NMR Varian Unity 300 spectrometer with H_3PO_4 as a standard. UV-Vis spectra were taken on a UV 160 Shimadzu spectrometer for the ligand and metal concentrations of the same value as in the samples for potentiometric titrations. Electron spin resonance (EPR) spectra were taken at 77 K in a water-glycol solution (3:1, v/v) on a Radiopan SE/X 2547 spectrometer ($C_{Cu^{2+}} = 0.002$ M) at the ratio of metal:amino acid ratio 1:4 and metal:nucleotide:amino acid ratio 1:2.5:2.5. IR

SCHEME 1: Chemical formulae of the bioligands studied.

measurements were carried out using a Bruker ISS 66vS spectrophotometer.

The hydrolysis constants for the metal ion Cu(II) were taken from [31] and were fully employed in the calculations.

The ligands studied are presented in Scheme 1.

3. RESULTS AND DISCUSSION

In the systems of polyamine/nucleotide studied earlier at our laboratory, noncovalent interactions were observed in the pH ranges in which one ligand was deprotonated (nucleotide) and the other was protonated (polyamine), and the molecular complex formation was a result of an ion-ion or ion-dipole reaction [26, 32–35]. In the systems amino acid/nucleotide, the pH ranges of protonation of both ligands overlap and each ligand has both positive and negative reaction centres. The formation of a molecular complex can be described by the equation: $H_xAsp + H_y(CMP) \rightleftharpoons (Asp)_{H(x+y-n)}(CMP) + nH^+$. The release of a proton in this reaction permits the use of the potentiometric method for determination of the composition and stability constants of the adducts. Analogous procedures were used for the investigation of the coordination compounds. The modes of interactions were determined on the basis of the spectroscopic measurements in the pH ranges in which particular complexes dominate, as established on the basis of the equilibrium study.

3.1. Asp/nucleotide metal-free systems

Table 1 presents the composition, overall stability constants ($\log \beta$), and the equilibrium constants of the formation ($\log K_e$) of molecular complexes appearing in Asp/CMP systems, determined from the computer analysis of the potentiometric titration data.

As it was described earlier [28, 29, 36], the occurrence of noncovalent interactions between the ligands and the formation of molecular complexes in the system studied is indicated by the coincidence of the titration curves obtained experimentally and those obtained by computer simulation (with the use of the determined β values) as given in Figure 1.

Deprotonation of the phosphate group of the nucleotide begins at low pH ($\log K_3 \sim 0.4$, [37, 38]), beyond the range of the study. Subsequent stages of deprotonation correspond

FIGURE 1: Experimental and simulated titration curves for the Asp/CMP system: dotted line: experimental curve; solid line: simulated curve (an adduct formation was taken into account); dashed line: simulated curve (an adduct formation was not taken into account).

FIGURE 2: Distribution diagram for the Asp/CMP system; percentage of the species refers to total ligands. (1) H_3Asp ; (2) H_2Asp ; (3) $HAsp$; (4) Asp ; (5) H_2CMP ; (6) $HCMP$; (7) CMP ; (8) $(Asp)H_4(CMP)$; (9) $(Asp)H_3(CMP)$; (10) $(Asp)H_2(CMP)$; (11) $(Asp)H(CMP)$; $C_{Asp} = 6 \times 10^{-3} M$; $C_{CMP} = 6 \times 10^{-3} M$.

TABLE 1: Overall stability constants ($\log \beta$) and equilibrium constants ($\log K_e$) of adducts formation in Asp/CMP system.

Species	$\log \beta$	$\log K_e$
H_3Asp	15.42 (3)	2.06
H_2Asp	13.36 (1)	3.73
$HAsp$	9.63 (1)	9.63
H_2CMP	10.90 (1) [20]	4.47
$HCMP$	6.42 (2) [20]	6.42
$(Asp)H_4(CMP)$	27.13 (4)	2.87
$(Asp)H_3(CMP)$	23.37 (3)	2.84
$(Asp)H_2(CMP)$	18.79 (3)	2.74
$(Asp)H(CMP)$	12.01 (3)	2.38

to the abstraction of a proton from the endocyclic nitrogen atom N(3) from CMP and of another proton from the phosphate group. The deprotonation of the aspartic acid molecule begins with the proton abstraction from the carboxyl group $C_{(1)}$, followed by dissociation of the $-C_{(4)}OOH$ and the $-NH_3^+$ group [39, 40].

Because of the different stoichiometric compositions of particular species, the overall stability constants $\log \beta$ cannot be directly applied in analysis of the character of interactions. Therefore, the efficiency of bonding was estimated on the basis of the equilibrium constants calculated, for example, for the species $(Asp)H_2(CMP)$: $\log K_e = \log \beta_{(Asp)H_2(CMP)} - \log \beta_{(HAsp)} - \log \beta_{H(CMP)}$. On the basis of the protonation constants of the ligands (Table 1) and the pH ranges of occurrence of individual species (Figure 2), the substrates in the adduct formation reactions were identified.

The complex $(Asp)H_4(CMP)$ appears at pH below 4.0 (Figure 2) in the range in which one of the carboxyl groups of Asp and partly the $-PO_4^{2-}$ group from CMP are deprotonated [41–44]. In the ^{13}C NMR spectra, the chemical

shifts of the signals assigned to C(2) and C(4) from the vicinity of N(3) of the nucleotide changed by 0.982 and 0.760 ppm (pH 3.0), respectively, (Table 2) indicate that the protonated N(3)H from CMP is a positive centre of weak interactions.

The lack of significant changes in the ^{31}P NMR spectrum suggests that the partly protonated (however negative) phosphate group is not active, which is a result of the repulsion from the negative carboxyl group present in the neighbourhood of $-NH_3^+$ group of Asp. The change in the chemical shift of the signal assigned to $C_{(1)}$ by 0.049 ppm suggests that the negative centre of interaction is the deprotonated carboxyl group from Asp. (As established earlier, the energy of the noncovalent interactions does not correspond directly with the chemical shift value [27, 29, 34]). The protonated carboxyl group is not involved in the interactions as evidenced by the lack of changes in the chemical shifts of the NMR signals assigned to the carbon $C_{(4)}$ atom. In the $(Asp)H_3(CMP)$ adduct, dominant at a pH close to 4, the N(3)H group of CMP remains a positive centre of interactions as indicated by the changes in the NMR chemical shifts, while the group $-C_{(4)}OO^-$ becomes a negative centre of interaction (Table 2), which corresponds to the deprotonation of the second carboxyl group. The phosphate group remains inactive. The involvement of only one centre from each ligand in the interactions is confirmed by similar values of the equilibrium constants of the tri- and tetraprotonated complexes of $\log K_e = 2.84$ and 2.87, respectively.

With increasing pH, the deprotonation of the N(3)H group from the pyrimidine ring of CMP takes place. The $(Asp)H_2(CMP)$ adduct starts forming from a pH close to 4 and reaches its maximum concentration at a pH of about 5.5. The changes in the chemical shifts of the carbon atoms C(2) and C(4), neighbouring the N(3) atom of the nucleotide at pH 5.5, in the region of domination of the $(Asp)H_2(CMP)$ adduct, are 0.440 and 0.593 ppm. This observation indicates

TABLE 2: Differences between ^{13}C NMR and ^{31}P NMR chemical shifts for the ligands in the Asp/CMP system in relation to the free ligands [ppm].

pH	Asp					CMP				P
	$\text{C}_{(1)}$	$\text{C}_{(2)}$	$\text{C}_{(3)}$	$\text{C}_{(4)}$	$\text{C}_{(2)}$	$\text{C}_{(4)}$	$\text{C}_{(5)}$	$\text{C}_{(6)}$	$\text{C}_{(5')}$	
3.0	0.049	0.025	0.022	0.002	0.982	0.760	0.057	0.048	0.044	0.018
4.2	0.000	0.013	0.070	0.054	0.861	0.661	0.040	0.047	0.060	0.025
5.5	0.053	0.047	0.027	0.040	0.440	0.593	0.068	0.054	0.099	0.040
8.0	0.454	1.048	0.227	1.322	0.191	0.198	0.068	0.064	0.111	0.105

that N(3) is a centre of interactions, however, since it is already deprotonated, an inversion of interactions takes place and the N(3) atom becomes a negative centre of interaction. The ^{31}P NMR spectrum of $(\text{Asp})\text{H}_2(\text{CMP})$ at pH 5.5 does not show any significant changes in the chemical shift of the phosphorus atom (Table 2), which indicates a low efficiency of the phosphate group.

Above the physiological pH, the formation of $(\text{Asp})\text{H}(\text{CMP})$ begins and it dominates at a pH of about 8. In these complexes, the phosphate group from the nucleotide is engaged in noncovalent interactions with amino acid as proved by the changes in the position of the signal of ^{31}P NMR, by 0.105 ppm at pH = 8. Changes in the chemical shifts of the C(2) and C(4) atoms from CMP by 0.191 and 0.198 ppm, respectively, indicate that the deprotonated N(3) atom is another negative centre of interaction. On the other hand, as follows from the changes in the chemical shift of the signal assigned to $\text{C}_{(2)}$ (1.048 ppm), the protonated amine group from Asp (positive, $\log K_1 = 9.6$) takes part in the interaction. In the molecule of the nucleotide, there are no positive reaction centres, thus the intermolecular interactions of $-\text{COO}^-$ are impossible, (similar to the intramolecular interactions). Because of the lack of another active centre in the Asp molecule, no increase in the $\log K_e$ value of the monoprotonated complex, relative to that of diprotonated one, is observed. The unexpected changes in the chemical shift of the signals assigned to $\text{C}_{(1)}$ and $\text{C}_{(4)}$ from Asp in the ^{13}C NMR spectrum (Table 2) are a consequence of the interaction of the carboxyl groups (hard base) with Na^+ ions (hard acids) present in the systems studied. In the additional ^{13}C NMR spectrum taken for the $(\text{Asp})\text{H}(\text{CMP})$ adduct in the system containing $(\text{C}_2\text{H}_5)_4\text{NOH}$ instead of NaOH, the changes were insignificant (at pH 8: 0.007 and 0.020 ppm for $\text{C}_{(1)}$ and $\text{C}_{(4)}$, resp.). The positions of the signals assigned to $\text{C}_{(1)}$ and $\text{C}_{(4)}$ from Asp in the spectra of the Na^+ free system Asp/CMP were compared to those in the spectra of Asp, without sodium ions. (This results also explains the unexpected shifts in the signals assigned to carbon atoms from carboxylate groups in some other studied systems.) The above conclusions are confirmed by analysis of IR spectra recorded in the same conditions as NMR ones. A comparison of the position of the IR band assigned to $-\text{COO}^-$ of the amino acid (1615 and 1584 cm^{-1}) and the $(\text{Asp})\text{H}(\text{CMP})$ adduct (1617 and 1586 cm^{-1}) shows that these groups are not involved in the ligand-ligand interactions (Figure 3), because no changes in the band positions were observed.

In all adducts studied above, noncovalent interaction occurs with the inversion of interaction sites at a pH close to 3.5 and close to 7, as illustrated in Scheme 2. The pH values of the inversion correspond to the values of the protonation constants, and changes in the mode of interaction are a result of the deprotonation of the second carboxyl group of Asp, endocyclic N(3)H, and the phosphate group of CMP. No significant changes are noted in the acid-base equilibria of the ligands as is usually the case when metal-ligand bonds are formed, which confirms that the interactions are weak, noncovalent.

3.2. Cu/Asp binary systems

The stability constants of Cu(II) complexes with Asp were determined in the same conditions in which the heteroligand complexes in the ternary systems are formed (Table 3). The results are in agreement with the earlier reported data [39, 40, 45, 46].

In the pH range from 2.5 to 6, a protonated species $\text{CuH}(\text{Asp})$ is formed, while at pH 5.5, the dominant complex is $\text{Cu}(\text{Asp})$, binding about 80% of copper ions. The species $\text{Cu}(\text{Asp})_2$ dominates at a pH from 7 to 10, while the hydroxocomplex $\text{Cu}(\text{Asp})(\text{OH})$ begins forming from a pH close to 6 reaching a maximum concentration above pH 10.5.

The UV-Vis and EPR spectral parameters (Table 4) at pH 3, at which the $\text{CuH}(\text{Asp})$ complex dominates: $\lambda_{\text{max}} = 756\text{ nm}$, $g_{\parallel} = 2.413$ and $A_{\parallel} = 134$, indicate that the oxygen atoms from the Asp carboxyl groups are exclusively involved in the coordination of the copper ions [26, 32, 47, 48], while the protonated amine group $-\text{NH}_3^+$ is blocked to coordination and does not take part in the interactions ($\{\text{O}_x\}$ chromophore). Conclusions concerning the mode of coordination were drawn on the basis of an analysis of the relation between the spectral parameters and the number of donor atoms in the inner sphere of Cu(II) coordination. For tetragonal and square pyramidal species, the ground state is normally $d_{x^2-y^2}$ or rarely d_{xy} . As earlier established for Cu-N_x ($x = 1-6$) and $\text{Cu-N}_x\text{O}_y$ ($x = 0-4$, $y = 0-6$) in planar bonding, the value of $g_{\parallel}$ decreases and that of $A_{\parallel}$ increases [47, 49]. In general, the value of $g_{\parallel}$ is changed in the order $\text{CuO}_4 > \text{CuO}_2\text{N}_2 > \text{CuON}_3 > \text{CuN}_4$. Weak bonding of the donor atoms at the axial position is a result of the interactions between the 4s and 4p metal orbitals and the ligand orbitals. The hyperfine coupling constant was experimentally observed to decrease with increasing electron density of the 4s orbital [47].

SCHEME 2: Noncovalent interaction in the Asp/CMP system.

TABLE 3: Overall stability constants ($\log \beta$) and equilibrium constants ($\log K_e$) for the complexes of Cu(II) ions with Asp or CMP and Cu(II) ions with Asp and CMP.

Equilibrium	$\log \beta$	$\log K_e$
$\text{Cu} + \text{CMP} \rightleftharpoons \text{Cu}(\text{CMP})$	2.71 (7) [20]	2.71
$\text{Cu} + \text{CMP} + \text{H}_2\text{O} \rightleftharpoons \text{Cu}(\text{CMP})(\text{OH}) + \text{H}^+$	-4.26 (6) [20]	
$\text{Cu} + \text{H}^+ + \text{Asp} \rightleftharpoons \text{CuH}(\text{Asp})$	12.78 (5)	3.15
$\text{Cu} + \text{Asp} \rightleftharpoons \text{Cu}(\text{Asp})$	8.76 (3)	8.76
$\text{Cu} + 2\text{Asp} \rightleftharpoons \text{Cu}(\text{Asp})_2$	15.35 (4)	6.59
$\text{Cu} + \text{Asp} + \text{H}_2\text{O} \rightleftharpoons \text{Cu}(\text{Asp})(\text{OH}) + \text{H}^+$	0.79 (9)	
$\text{Cu} + 4\text{H}^+ + \text{Asp} + \text{CMP} \rightleftharpoons \text{Cu}(\text{Asp})\text{H}_4(\text{CMP})$	32.86 (5)	
$\text{Cu} + 3\text{H}^+ + \text{Asp} + \text{CMP} \rightleftharpoons \text{Cu}(\text{Asp})\text{H}_3(\text{CMP})$	29.14 (5)	5.46
$\text{Cu} + 2\text{H}^+ + \text{Asp} + \text{CMP} \rightleftharpoons \text{Cu}(\text{Asp})\text{H}_2(\text{CMP})$	25.28 (4)	6.08
$\text{Cu} + \text{H}^+ + \text{Asp} + \text{CMP} \rightleftharpoons \text{Cu}(\text{Asp})\text{H}(\text{CMP})$	20.12 (4)	4.94
$\text{Cu} + \text{H}^+ + 2\text{Asp} + \text{CMP} \rightleftharpoons \text{Cu}(\text{Asp})_2\text{H}(\text{CMP})$	29.87 (4)	
$\text{Cu} + \text{Asp} + \text{CMP} + \text{H}_2\text{O} \rightleftharpoons \text{Cu}(\text{Asp})(\text{CMP})(\text{OH}) + \text{H}^+$	4.69 (4)	

The carboxyl group involvement is confirmed by the changes in the ^{13}C NMR chemical shifts of the carbon atoms from the two carboxyl groups: $\text{C}_{(1)}$ and $\text{C}_{(4)}$ by 1.536 and 1.163 ppm, respectively (Table 5).

The increase in the equilibrium constant of the formation of $\text{Cu}(\text{Asp})$ by approximately 5.5 $\log K_e$ unit relative to the value for the protonated complex (Table 3) points to the involvement of the deprotonated amine group $-\text{NH}_2$ (for pH above 5) in the coordination of $\text{Cu}(\text{Asp})$ species, which is consistent with the model proposed by Chaberek and Martell, later confirmed in [39, 50]. Moreover, in the electronic absorption spectrum, the λ_{max} is shifted toward higher energy—from 756 to 700 nm, which points to the engagement of the nitrogen atom from Asp in metal coordination, (besides the oxygen atoms). The {N, Ox} type coordination was also confirmed by the EPR results as at

pH = 5, $g_{\parallel} = 2.301$ and $A_{\parallel} = 171$. The attachment of a subsequent molecule of amino acid to the anchoring $\text{Cu}(\text{Asp})$, according to the equation $\text{Cu}(\text{Asp}) + \text{Asp} \rightleftharpoons \text{Cu}(\text{Asp})_2$, leads to a decrease in the equilibrium constant by about 2 $\log K_e$ units, so by a value suggesting the same mode of coordination in the 1:1 and 1:2 complexes, taking into consideration the statistical relations. The spectral parameters obtained from the UV-Vis and EPR spectra for the $\text{Cu}(\text{Asp})_2$ complex are $\lambda_{\text{max}} = 630$ nm, $g_{\parallel} = 2.257$ nm, and $A_{\parallel} = 187$ (Table 4) indicating (in agreement with Gamp et al. [48]) the formation of the {N2, Ox} chromophore with the participation of the deprotonated nitrogen and oxygen atoms from the carboxyl groups of the amino acid, which also follows from the changes in the ^{13}C NMR spectra. At pH 8, at which the $\text{Cu}(\text{Asp})_2$ species reaches a maximum concentration, the changes in chemical shifts of

TABLE 4: Vis and EPR spectral data for Cu(II)/Asp and Cu(II)/Asp/CMP systems.

Species	pH	$\lambda_{\max}$ [nm]	ϵ [$\text{M}^{-1}\text{cm}^{-1}$]	EPR	
				$g_{\parallel} = (\text{dm}^3/\text{mol}\cdot\text{cm}^3)$	$A_{\parallel} (10^{-4}\text{cm}^{-1})$
CuH(Asp)	3	756	27	2.413	134
Cu(Asp)	5	700	40	2.301	171
Cu(Asp) ₂	8	630	76	2.257	187
Cu(Asp)H ₄ (CMP)	3	780	25	2.409	138
Cu(Asp)H ₂ (CMP)	4.5	710	29	2.275	179
Cu(Asp)H(CMP)	6	700	30	2.269	182
Cu(Asp) ₂ H(CMP)	8	660	49	2.261	182
Cu(Asp)(CMP)(OH)	10.5	630	96	2.258	183

FIGURE 3: IR spectra of Asp and (Asp)H(CMP) adduct pH = 8; $C_{\text{Asp}} = 0.2 \text{ M}$, $C_{\text{CMP}} = 0.2 \text{ M}$.

the carbon atoms $C_{(1)}$, $C_{(2)}$, and $C_{(4)}$ are 0.595, 0.602, and 0.755 nm, respectively.

The Cu(Asp)(OH) complex occurs in the same pH range as that of the dominant Cu(Asp)₂ species and, therefore, no spectra could be taken.

3.3. Cu(II)/Asp/CMP system

Analysis of the equilibria in the ternary systems was performed using the protonation constants and overall stability constants ($\log\beta$) of the complexes forming in the binary systems of Cu(II)/CMP [20]. In the system

Cu(II)/Asp/CMP, the following complexes were found: the protonated ones Cu(Asp)H₄(CMP), Cu(Asp)H₃(CMP), Cu(Asp)H₂(CMP), Cu(Asp)₂H(CMP), and the hydroxo-complex Cu(Asp)(CMP)(OH). Table 3 presents the results of a computer analysis of titration curves in ternary systems with Cu(II) ions, Asp, and CMP. The Cu(Asp)H₄(CMP) species was already observed to form at a pH below 2, binding over 80% of Cu(II) ions at pH 3 (Figure 4). Taking into regard the number of hydrogen atoms in the species studied, the position of the d-d bands, and the EPR parameters (at pH 3, $\lambda_{\max} = 780 \text{ nm}$, $g_{\parallel} = 2.409$, and $A_{\parallel} = 138$, Table 4), one can conclude that the Cu(II) ion coordination occurs only by oxygen atoms from Asp.

The composition (in particular the number of hydrogen atoms blocking the coordination) of the Cu(Asp)H₄(CMP) complex and the pH range of its occurrence suggest that it is a molecular complex. Conclusions concerning the mode of interactions could be achieved on the basis of spectroscopic studies. The UV-Vis and EPR data ($\lambda_{\max} = 780 \text{ nm}$, $g_{\parallel} = 2.409$, and $A_{\parallel} = 138$, Table 4) indicate bonding of copper(II) ion only via one oxygen atom (coordination Cu-O in an inner sphere) with the involvement in metallation of a deprotonated carboxyl group $C_{(1)}$ from the amino acid (the second carboxyl group is protonated and blocked for coordination). On the basis of this essential finding, the next stages of the analysis were performed. In the ¹³C NMR spectrum of this species (pH = 3), the change in the chemical shift of $C_{(1)}$ from Asp is 1.371 ppm. Moreover, the changes in the signals assigned to the Asp carbon atoms $C_{(2)}$ (0.460 ppm) and $C_{(4)}$ (1.094 ppm) and originated from the nucleotide carbon atoms $C(2)$ and $C(4)$ by 0.463 and 0.528 ppm, respectively, and the change in the ³¹P NMR of CMP by 0.166 ppm confirm clearly the hypothesis relating to the intermolecular interactions between the protonated complex of Cu(II) with Asp and the protonated CMP molecule located in the outer coordination sphere. The noncovalent interaction centre becomes the protonated Asp amine group, the CMP phosphate group, and N(3)H.

With increasing pH and deprotonation of the carboxyl group $C_{(4)}$ in the Asp molecule, the Cu(Asp)H₃(CMP) species is formed. The protonated nucleotide with the donor atoms blocked is probably involved in noncovalent interactions with the anchoring CuH(Asp) species. The pH

range of formation of this species overlaps the ranges of formation of other ones (Figure 4), which makes it impossible to perform spectral studies and to explain their mode of coordination.

The $\text{Cu(Asp)H}_2\text{(CMP)}$ complex begins to form from a pH close to 3 in parallel with the deprotonation of the nitrogen atom N(3) of the nucleotide, and at pH 4.5, it binds about 70% of Cu(II) ions. The position of the absorption band in the UV-Vis spectrum in the pH range of the complex domination $\lambda_{\text{max}} = 710 \text{ nm}$ and the values of the EPR parameters $g_{\parallel} = 2.275$ and $A_{\parallel} = 179$ (Table 4) [48, 51] indicate formation of the {N, Ox} type chromophore and it provides the basis of information concerning the character of interaction. In the NMR spectrum of $\text{Cu(Asp)H}_2\text{(CMP)}$, the changes in the chemical shifts of the carbon atoms $\text{C}_{(1)}$ and $\text{C}_{(4)}$ from Asp (0.069 and 0.080 ppm, resp.) and the carbon atoms $\text{C}_{(2)}$ and $\text{C}_{(4)}$ from the nucleotide (1.042 and 0.932 ppm, resp.), together with the changes in the ^{31}P NMR signal (0.080 ppm), point to the involvement in metallation of the deprotonated N(3) atom, the phosphate group of CMP, and the oxygen atoms from the carboxyl group of Asp. As follows from the value of $\log K_e = 6.08$ for $\text{Cu(Asp)H}_2\text{(CMP)}$ (so for the reaction of HCMP attachment to the anchoring CuHAsp), much higher than that for Cu(CMP) formation $\log K_e = 2.71$ [20], there is a noncovalent intramolecular ligand-ligand interaction in the $\text{Cu(Asp)H}_2\text{(CMP)}$ complex that additionally stabilises it (the presence of weak interactions is confirmed by changes in the chemical shift of $\text{C}_{(2)}$ from Asp (0.643 ppm) located in the proximity of the NH_3^+ group). The equilibrium constant of the $\text{Cu(Asp)H}_2\text{(CMP)}$ complex formation is by about 3 orders of magnitude higher than that of the adduct $\text{(Asp)H}_2\text{(CMP)}$, which is a result of significant differences in the character of the bond (in the metal-free system the bond is weak, noncovalent, but in the copper complex both ligands bind the metal ions with additional intramolecular interaction).

The Cu(Asp)H(CMP) complex begins to form from a pH close to 4, and at pH = 6, it binds 80% of the Cu(II) ions. In the pH range of its domination $\lambda_{\text{max}} = 700 \text{ nm}$, the EPR parameters are $g_{\parallel} = 2.269$ and $A_{\parallel} = 182$, which implies the involvement of one donor nitrogen atom and oxygen atoms from the ligand molecules in the coordination ({N, Ox} chromophore). The NMR spectra reveal changes in the positions of signals coming from the carbon atoms of the Asp carboxyl groups ($\text{C}_{(1)}$ 0.769 ppm and $\text{C}_{(4)}$ 0.809 ppm), the phosphorus atom of CMP (^{31}P NMR 0.190 ppm), and the carbon atom located close to the Asp amine group ($\text{C}_{(2)}$ 0.782 ppm). Moreover, changes in positions of the signal originated from the CMP carbon atoms located in the proximity of N(3) ($\text{C}_{(2)}$, 0.486 ppm and $\text{C}_{(4)}$, 0.772 ppm) are observed. The question of which nitrogen atoms (either from the Asp amine group or the endocyclic N(3) atom) from CMP are involved in the metallation has been solved by comparison of the thermodynamic stability of the two possible species. The first species corresponds to the formation of a stable system coupled with five- and six-membered rings (Scheme 3), while the second corresponds to the formation of an unstable seven-membered ring and a

FIGURE 4: Distribution diagram for the Cu(II)/Asp/CMP system; percentage of the species refers to total metal. (1) H_3Asp ; (2) H_2Asp ; (3) HAsp ; (4) Asp ; (5) $\text{H}_2\text{(CMP)}$; (6) H(CMP) ; (7) CMP ; (8) Cu(Asp)_2 ; (9) Cu(Asp)(OH) ; (10) $\text{Cu(Asp)H}_4\text{(CMP)}$; (11) $\text{Cu(Asp)H}_3\text{(CMP)}$; (12) $\text{Cu(Asp)H}_2\text{(CMP)}$; (13) Cu(Asp)H(CMP) ; (14) $\text{Cu(Asp)}_2\text{H(CMP)}$; (15) Cu(Asp)(CMP)(OH) ; $\text{C}_{\text{Cu}^{2+}} = 1 \times 10^{-3} \text{ M}$; $\text{C}_{\text{Asp}} = 2.5 \times 10^{-3} \text{ M}$; $\text{C}_{\text{CMP}} = 2.5 \times 10^{-3} \text{ M}$.

SCHEME 3: Tentative mode of interaction in the Cu(Asp)H(CMP) complex.

macrocholate structure with a coordination via the $-\text{PO}_4^{2-}$ group and N(3) atom from CMP.

In the pH range 7–8, the dominant species is $\text{Cu(Asp)}_2\text{H(CMP)}$. The positions of the d-d bands and the EPR parameters (pH = 8, Table 4) imply the formation of an {N2, Ox} type chromophore. The Cu(Asp)H(CMP) molecule accepts another amino acid molecule. As follows from the changes in NMR signals assigned to the carbon atoms of the carboxyl group ($\text{C}_{(1)}$ 0.809 ppm and $\text{C}_{(4)}$ 0.689 ppm) and from those in the proximity of the Asp amine group ($\text{C}_{(2)}$ 0.749 ppm) as well as the changes in the chemical shifts assigned to the $\text{C}_{(2)}$ and $\text{C}_{(4)}$ atoms from the neighbourhood of N(3) in CMP by 0.860 and 0.858 ppm, respectively, and changes in the chemical shifts in the ^{31}P NMR spectrum of the nucleotide (0.129 ppm), the coordination involves the oxygen atoms of the three carboxyl groups and the amine group from the Asp molecule and the endocyclic N(3) atom together with the phosphate group from CMP (monofunctional character of the second Asp ligand). The hydroxocomplex Cu(Asp)(CMP)(OH) begins to form from a pH close to 8, and at pH = 10.5, this

TABLE 5: Differences between ^{13}C NMR and ^{31}P NMR chemical shifts for the ligands in the Cu(II)/Asp and Cu(II)/Asp/CMP systems in relation to metal-free systems [ppm].

Systems	pH	Asp				CMP					P
		C ₍₁₎	C ₍₂₎	C ₍₃₎	C ₍₄₎	C(2)	C(4)	C(5)	C(6)	C(5')	
Cu(II)/Asp	3.0	1.536	0.129	0.122	1.163	—	—	—	—	—	—
	5.5	0.050	0.070	0.020	0.050	—	—	—	—	—	—
	8.0	0.595	0.602	0.169	0.755	—	—	—	—	—	—
Cu(II)/Asp/CMP	3.0	1.371	0.460	0.147	1.094	0.463	0.528	0.075	0.085	0.057	0.166
	4.5	0.069	0.643	0.113	0.080	1.042	0.932	0.052	0.047	0.051	0.080
	6.0	0.769	0.782	0.142	0.809	0.486	0.772	0.062	0.046	0.076	0.190
	8.0	0.809	0.749	0.152	0.689	0.860	0.858	0.071	0.064	0.062	0.129

species dominates (the equilibrium $\text{Cu}(\text{Asp})_2\text{H}(\text{CMP}) + \text{OH}^- \rightleftharpoons \text{Cu}(\text{Asp})(\text{CMP})(\text{OH}) + \text{H}(\text{Asp})$). The maximum of the band in the UV-Vis spectrum is at 630 nm, and the EPR parameters are $g_{\parallel} = 2.258$ and $A_{\parallel} = 183$, which corresponds to the $\{\text{N}2, \text{Ox}\}$ chromophore with coordination via the nitrogen atom N(3), the phosphate group from CMP, the oxygen atom from the Asp amine group, oxygen atoms from the carboxyl groups of the amino acid, and oxygen atoms from the OH group.

4. CONCLUSIONS

The main reaction centres in complexes formed as a result of noncovalent interactions in metal-free systems at low pH are the nitrogen N(3)H group from CMP and carboxyl groups from Asp. With increasing pH, the efficiency of the interactions of the Asp carboxyl group decreases, while that of the amine group increases. For a pH up to about 5, the protonated N(3)H from the nucleotide is the positive centre of weak interactions and at a higher pH, an inversion occurs and it becomes a negative centre. Above a pH of about 7, the involvement of the phosphate group occurs in the interaction. The change in the mode of interactions corresponds to the pH ranges of protonation of particular ligands. No significant acid-base shifts are observed in the systems studied, which confirms the hypothesis that the interactions are weak. Introduction of the metal ion into the binary system changes the character of the reaction in the ternary system relative to that in the metal-free systems. Particularly important is a significant increase in the efficiency of the phosphate group from CMP in the noncovalent interactions in the $\text{Cu}(\text{Asp})\text{H}_4(\text{CMP})$ molecular complex. In the $(\text{Asp})\text{H}_4(\text{CMP})$ adduct which forms in the same pH range, the $-\text{PO}_4\text{H}^-$ group is ineffective.

In the ternary systems, Cu(II)/Asp/CMP coordination is realised through the carboxyl groups from Asp, and starting from a pH close to 4, also the phosphate groups from CMP and the endocyclic N(3) atom are involved in interactions. At low pH, the carboxyl groups from the amino acid are the main metal binding site, but with increasing pH, their efficiency decreases to the advantage of the phosphate groups, as follows from the analysis of the $\log K_e$ values. The presence of metal ions changes the character of noncovalent

interactions, whereas the introduction of an additional ligand CMP into the binary system Cu(II)/Asp changes the mode of coordination. For instance, in the Cu(Asp) complex, the coordination is realised via the oxygen atoms from the carboxyl groups and the amine group (five-membered ring), while in the ternary complex Cu(Asp)H(CMP) occurring in the same pH range, the involvement of the phosphate group from CMP in metal binding leads to a change in the mode of coordination and the formation of a structure with a system of coupled five- and six-membered rings.

ACKNOWLEDGMENT

This work was supported by the Polish Ministry of Science and Higher Education.

REFERENCES

- [1] A. Sigel and H. Sigel, Eds., *Metal Ions in Biological Systems, Interaction of Metal Ions with Nucleotides, Nucleic Acids, and Their Constituents*, vol. 32, Marcel Dekker, New York, NY, USA, 1996.
- [2] S. J. Lippart and J. M. Berg, *Principles of Bioinorganic Chemistry*, University Science Books, Mill Valley, Calif, USA, 1994.
- [3] A. S. Mildvan, "Role of magnesium and other divalent cations in ATP-utilizing enzymes," *Magnesium*, vol. 6, no. 1, pp. 28–33, 1987.
- [4] R. B. Martin, "Metal ions binding to nucleoside and nucleotides," in *Frontiers in Bioinorganic Chemistry*, A. V. Xavier, Ed., p. 71, VCH, Weinheim, Germany, 1986.
- [5] L. G. Marzilli, "Metal complex of nucleic acid derivatives and nucleotides: binding sites and structures," in *Metal Ions in Genetic Information Transfer*, G. L. Eichhorn and L. G. Marzilli, Eds., pp. 47–85, Elsevier, New York, NY, USA, 1981.
- [6] H. Sigel, "Complexes of metal ions with various nucleic acids components," in *Handbook of Metal-Ligand Interactions in Biological Fluids*, G. Berthon, Ed., pp. 451–465, Marcel Dekker, New York, NY, USA, 1995.
- [7] R. B. Martin, "Nucleoside sites for transition metal ion binding," *Accounts of Chemical Research*, vol. 18, no. 2, pp. 32–38, 1985.
- [8] B. de Castro, J. Pereira, P. Gameiro, and F. C. Castro, "Multinuclear NMR and potentiometric studies on the interaction of zinc and cadmium with cytidine and glycylglycine. The effect of the anion," *Journal of Inorganic Biochemistry*, vol. 45, no. 1, pp. 53–64, 1992.

- [9] K. C. Tewari, J. Lee, and N. C. Li, "Zinc and mercury (II) interaction with cytidine and glycylglycine," *Transactions of the Faraday Society*, vol. 66, pp. 2069–2075, 1970.
- [10] F. M. Vaccarino, M. L. Schwartz, D. Hartigan, and J. F. Leckman, "Basic fibroblast growth factor increases the number of excitatory neurons containing glutamate in the cerebral cortex," *Cerebral Cortex*, vol. 5, no. 1, pp. 64–78, 1995.
- [11] D. R. Curtis and J. M. Crawford, "Central synaptic transmission-microelectrophoretic studies," *Annual Review of Pharmacology*, vol. 9, pp. 209–240, 1969.
- [12] M. Aprison, R. P. Shank, R. A. Davidoff, and R. R. Werman, "The distribution of glycine, a neurotransmitter suspect in the central nervous system of several vertebrate species," *Life Sciences*, vol. 7, no. 11, pp. 583–590, 1968.
- [13] R. Werman, R. A. Davidoff, and M. H. Aprison, "Inhibitory of glycine on spinal neurons in the cat," *Journal of Neurophysiology*, vol. 31, no. 1, pp. 81–95, 1968.
- [14] P. E. Chen, M. T. Geballe, P. J. Stansfeld, et al., "Structural features of the glutamate binding site in recombinant NR1/NR2A N-methyl-D-aspartate receptors determined by site-directed mutagenesis and molecular modeling," *Molecular Pharmacology*, vol. 67, no. 5, pp. 1470–1484, 2005.
- [15] M. Monda, A. Viggiano, A. Sullo, and V. De Luca, "Aspartic and glutamic acids increase in the frontal cortex during prostaglandin E_1 hyperthermia," *Neuroscience*, vol. 83, no. 4, pp. 1239–1243, 1998.
- [16] A. Gergely and I. Sovago, "Log β , ΔH and ΔS values of mixed complexes of Cu(II) with histamine and some aliphatic aminoacids," *Journal of Inorganic Nuclear Chemistry*, vol. 35, no. 12, pp. 4355–4365, 1973.
- [17] I. Nagypál and A. Gergely, "Studies on transition-metal-peptide complexes. Part 2. Equilibrium study of the mixed complexes of copper(II) with aliphatic dipeptides and aminoacids," *Journal of the Chemical Society, Dalton Transactions*, no. 11, pp. 1109–1111, 1977.
- [18] M. M. Petit-Ramel and M. R. Páris, "Polarimetric study of metallic complexes of amino acids. II. Mixed complexes of copper with two amino acids," *Bulletin de la Société Chimique de France*, vol. 7, pp. 2791–2796, 1968 (French).
- [19] H. C. Freeman and R.-P. Martin, "Potentiometric study of equilibria in aqueous solution between copper (II) ions, L (or D)-histidine and L-threonine and their mixtures," *The Journal of Biological Chemistry*, vol. 244, no. 18, pp. 4823–4830, 1969.
- [20] A. Gasowska and L. Lomozik, "Cobalt(II), nickel(II) and copper(II) complexes with adenosine 5'-monophosphate and cytidine 5'-monophosphate in aqueous solutions and in solid," *Polish Journal of Chemistry*, vol. 73, no. 3, pp. 465–474, 1999.
- [21] L. Lomozik, "Complex compounds of Cu(II) and Zn(II) with N,N-dimethylglycine and N,N-diethylglycine in water and in water-methanol system," *Monatshefte für Chemie*, vol. 115, no. 3, pp. 261–270, 1984.
- [22] H. M. Irving, M. G. Miles, and L. D. Pettit, "A study of some problems in determining the stoichiometric proton dissociation constants of complexes by potentiometric titrations using a glass electrode," *Analytica Chimica Acta*, vol. 38, pp. 475–485, 1967.
- [23] P. Gans, A. Sabatini, and A. Vacca, "SUPERQUAD: an improved general program for computation of formation constants from potentiometric data," *Journal of the Chemical Society, Dalton Transactions*, no. 6, pp. 1195–1200, 1985.
- [24] N. Ingri, W. Kakolowicz, L. G. Sillen, and B. Warnqvist, "High-speed computers as a supplement to graphical methods—V Haltafall, a general program for calculating the composition of equilibrium mixtures," *Talanta*, vol. 14, no. 11, pp. 1261–1286, 1967.
- [25] L. Lomozik, M. Jaskolski, and A. Wojciechowska, "A multistage verification procedure for the selection of models in the studies of complex formation equilibria," *Polish Journal of Chemistry*, vol. 65, pp. 1797–1807, 1991.
- [26] A. Gasowska, "Interaction centres of pyrimidine nucleotides: cytidine-5'-diphosphate (CDP) and cytidine-5'-triphosphate (CTP) in their reactions with tetramines and Cu(II) ions," *Journal of Inorganic Biochemistry*, vol. 99, no. 8, pp. 1698–1707, 2005.
- [27] A. Gasowska, L. Lomozik, and R. Jastrzab, "Mixed-ligand complexes of copper(II) ions with AMP and CMP in the systems with polyamines and non-covalent interaction between bioligands," *Journal of Inorganic Biochemistry*, vol. 78, no. 2, pp. 139–147, 2000.
- [28] L. Lomozik and A. Gasowska, "Investigations of binding sites and stability of complexes formed in ternary Cu(II)/adenosine or cytidine/putrescine systems," *Journal of Inorganic Biochemistry*, vol. 62, no. 2, pp. 103–115, 1996.
- [29] L. Lomozik, A. Gasowska, and L. Bolewski, "Noncovalent interactions in polyamine/nucleoside (or diaminocarboxylate) systems studied by potentiometric and NMR techniques," *Journal of the Chemical Society. Perkin Transactions*, vol. 2, no. 6, pp. 1161–1166, 1997.
- [30] P. K. Glasoe and F. A. Long, "Use of glass electrodes to measure acidities in deuterium oxide," *Journal of Physical Chemistry*, vol. 64, no. 1, pp. 188–190, 1960.
- [31] R. N. Sylva and M. R. Davidson, "The hydrolysis of metal ions. Part 1. Copper(II)," *Journal of the Chemical Society, Dalton Transactions*, no. 2, pp. 232–235, 1979.
- [32] A. Gasowska, "Interaction centres of purine nucleotides: adenosine-5'-diphosphate and adenosine-5'-triphosphate in their reactions with tetramines and Cu(II) ions," *Journal of Inorganic Biochemistry*, vol. 96, no. 2–3, pp. 346–356, 2003.
- [33] L. Lomozik and R. Jastrzab, "Copper(II) complexes with uridine, uridine 5'-monophosphate, spermidine, or spermine in aqueous solution," *Journal of Inorganic Biochemistry*, vol. 93, no. 3–4, pp. 132–140, 2003.
- [34] L. Lomozik and A. Gasowska, "Complexes of copper(II) with spermine and non-covalent interactions in the systems including nucleosides or nucleotides," *Journal of Inorganic Biochemistry*, vol. 72, no. 1–2, pp. 37–47, 1998.
- [35] L. Lomozik, A. Gasowska, R. Bregier-Jarzebowska, and R. Jastrzab, "Coordination chemistry of polyamines and their interactions in ternary systems including metal ions, nucleosides and nucleotides," *Coordination Chemistry Reviews*, vol. 249, no. 21–22, pp. 2335–2350, 2005.
- [36] L. Lomozik, A. Gasowska, and G. Krzysko, "Interactions of 1,12-diamino-4,9-dioxadodecane (OSpm) and Cu(II) ions with pyrimidine and purine nucleotides: adenosine-5'-monophosphate (AMP) and cytidine-5'-monophosphate (CMP)," *Journal of Inorganic Biochemistry*, vol. 100, no. 11, pp. 1781–1789, 2006.
- [37] S. S. Massoud and H. Sigel, "Metal ion coordinating properties of pyrimidine-nucleoside 5'-monophosphates (CMP, UMP, TMP) and of simple phosphate monoesters, including D-ribose 5'-monophosphate. Establishment of relations between complex stability and phosphate basicity," *Inorganic Chemistry*, vol. 27, no. 8, pp. 1447–1453, 1988.
- [38] H. Sigel, S. S. Massoud, and R. Tribolet, "Comparison of the metal ion coordinating properties of tubercidin 5'-monophosphate (7-deaza-AMP) with those of adenosine 5'-monophosphate (AMP) and 1,N⁶-ethenoadenosine

- 5'-monophosphate (ϵ -AMP). Definite evidence for metal ion-base backbinding to N-7 and extent of macrochelate formation in M(AMP) and M(ϵ -AMP)," *Journal of the American Chemical Society*, vol. 110, no. 20, pp. 6857–6865, 1988.
- [39] S. Chaberek Jr. and A. E. Martell, "Stability of metal chelates. III. Iminopropionic-acetic and aspartic acids," *Journal of the American Chemical Society*, vol. 74, no. 23, pp. 6021–6025, 1952.
- [40] I. Nagypal, A. Gergely, and E. Farkas, "Thermodynamic study of the parent and mixed complexes of aspartic acid, glutamic acid and glycine with copper(II)," *Journal of Inorganic Nuclear Chemistry*, vol. 36, no. 3, pp. 699–706, 1974.
- [41] C. M. Frey and J. E. Stuehr, "Interactions of divalent metal ions with inorganic and nucleoside phosphates. I. Thermodynamics," *Journal of the American Chemical Society*, vol. 94, no. 25, pp. 8898–8904, 1972.
- [42] M. Matthies and G. Zundel, "Hydration and self-association of adenosine triphosphate, adenosine diphosphate, and their 1:1 complexes with magnesium(II) at various pH values: Infrared investigations," *Journal of the Chemical Society, Perkin Transactions*, vol. 2, no. 14, pp. 1824–1830, 1977.
- [43] K. H. Scheller and H. Sigel, "A proton nuclear magnetic resonance study of purine and pyrimidine nucleoside 5'-diphosphates. Extent of macrochelate formation in monomeric metal ion complexes and promotion of self-stacking by metal ions," *Journal of the American Chemical Society*, vol. 105, no. 18, pp. 5891–5900, 1983.
- [44] D. A. Nation, Q. Lu, and A. E. Martell, "Molecular recognition of nucleotides by the protonated macrocyclic ligand 3,6,9,17,20,23-hexaazatricyclo[23.3.1.1^{11,15}] triacontal(29), 11(30),12,14,25,27-hexaene and by its Cu(II) complexes; catalysis of the conversion of ATP to ADP," *Inorganica Chimica Acta*, vol. 263, no. 1-2, pp. 209–217, 1997.
- [45] A. Wojciechowska, L. Lomozik, and S. Zieliński, "Potentiometric and spectral studies of complex formation of La(III), Pr(III) and Lu(III) with aspartic acid and asparagine," *Monatshefte für Chemie*, vol. 118, no. 12, pp. 1317–1324, 1987.
- [46] L. Lomozik, A. Wojciechowska, and M. Jaskólski, "Complex formation studies on copper(II) and zinc(II) with asparagine and aspartic acid in aqueous solution," *Monatshefte für Chemie*, vol. 114, no. 11, pp. 1185–1188, 1983.
- [47] R. Barbucci and M. J. M. Campbell, "An investigation of the structures of copper(II) polyamine complexes in aqueous solution by a combined evaluation of the EPR and thermodynamic parameters," *Inorganica Chimica Acta*, vol. 16, pp. 113–120, 1976.
- [48] H. Gampp, H. Sigel, and A. D. Zuberbühler, "Apical interactions in copper(II) complexes. Stability and structure of the binary and ternary copper(II) complexes formed with L-alaninamide and diethylenetriamine in aqueous solution," *Inorganic Chemistry*, vol. 21, no. 3, pp. 1190–1195, 1982.
- [49] L. Lomozik, L. Bolewski, and R. Dworczak, "Complex formation in copper(II) ternary systems involving polyamines and diaminocarboxylates studied by potentiometric and spectroscopic techniques," *Journal of Coordination Chemistry*, vol. 41, no. 4, pp. 261–274, 1997.
- [50] L. Lomozik and A. Wojciechowska, "Comparative studies of the coordination mode in complexes of copper with asparagine and copper with aspartic acid in solution and solid state," *Polyhedron*, vol. 8, no. 1, pp. 1–6, 1989.
- [51] L. Lomozik, L. Bolewski, and R. Bregier-Jarzebowska, "Stability and structure studies of copper(II) complexes with polyamines and related ligands in aqueous solution," *Polish Journal Chemistry*, vol. 69, pp. 197–205, 1995.