

Research Article

Structural Basis for pH-Dependent Oligomerization of Dihydropyrimidinase from *Pseudomonas aeruginosa* PAO1

Jen-Hao Cheng,¹ Chien-Chih Huang,¹ Yen-Hua Huang,¹ and Cheng-Yang Huang ^{1,2}

¹School of Biomedical Sciences, Chung Shan Medical University, No. 110, Sec. 1, Chien-Kuo N. Rd., Taichung, Taiwan

²Department of Medical Research, Chung Shan Medical University Hospital, No. 110, Sec. 1, Chien-Kuo N. Rd., Taichung, Taiwan

Correspondence should be addressed to Cheng-Yang Huang; cyhuang@csmu.edu.tw

Received 17 October 2017; Revised 20 November 2017; Accepted 3 December 2017; Published 30 January 2018

Academic Editor: Luigi Casella

Copyright © 2018 Jen-Hao Cheng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dihydropyrimidinase, a dimetalloenzyme containing a carboxylated lysine within the active site, is a member of the cyclic amidohydrolase family, which also includes allantoinase, dihydroorotase, hydantoinase, and imidase. Unlike all known dihydropyrimidinases, which are tetrameric, pseudomonas dihydropyrimidinase forms a dimer at neutral pH. In this paper, we report the crystal structure of *P. aeruginosa* dihydropyrimidinase at pH 5.9 (PDB entry 5YKD). The crystals of *P. aeruginosa* dihydropyrimidinase belonged to space group C222₁ with cell dimensions of $a = 108.9$, $b = 155.7$, and $c = 235.6$ Å. The structure of *P. aeruginosa* dihydropyrimidinase was solved at 2.17 Å resolution. An asymmetric unit of the crystal contained four crystallographically independent *P. aeruginosa* dihydropyrimidinase monomers. Gel filtration chromatographic analysis of purified *P. aeruginosa* dihydropyrimidinase revealed a mixture of dimers and tetramers at pH 5.9. Thus, *P. aeruginosa* dihydropyrimidinase can form a stable tetramer both in the crystalline state and in the solution. Based on sequence analysis and structural comparison of the dimer-dimer interface between *P. aeruginosa* dihydropyrimidinase and *Thermus* sp. dihydropyrimidinase, different oligomerization mechanisms are proposed.

1. Introduction

Dihydropyrimidinase is a key enzyme for pyrimidine catabolism [1, 2]. Dihydropyrimidinase catalyzes the reversible cyclization of dihydrouracil to *N*-carbamoyl- β -alanine in the second step of the pyrimidine degradation pathway (Figure 1). Dihydropyrimidinase can also detoxify xenobiotics with an imide functional group, ranging from linear imides to heterocyclic imides [3–9]. Homologous enzymes from microorganisms are known as hydantoinase, used as biocatalyst for hydrolysis of 5-monosubstituted hydantoins in the synthesis of D- and L-amino acids [10, 11]. Optically pure amino acids have been widely used as intermediates for semisynthesis of antibiotics, active peptides, hormones, antifungal agents, pesticides, and sweeteners. Dihydropyrimidinase and hydantoinase generally possess a similar active site, but their overall sequence identity and substrate specificity may differ [3, 12]. For example, hydantoinase purified from *Agrobacterium* species has no 5,6-dihydropyrimidine amidohydrolase activity [13]. Dihydropyrimidinases from the yeast *Saccharomyces kluyveri*

and the slime mold *Dictyostelium discoideum* do not hydrolyze hydantoin [14]. Thus, several bacterial hydantoinases are still named and identified as dihydropyrimidinase because of their catalytic activity toward natural substrates, namely, dihydrouracil and dihydrothymine. These bacterial enzymes include *Pseudomonas aeruginosa* and *Thermus* sp. dihydropyrimidinases [15, 16].

Dihydropyrimidinase, hydantoinase, imidase, allantoinase, and dihydroorotase belong to the cyclic amidohydrolase family because of their functional and structural similarities [17]. Members of this enzyme family catalyze the ring-opening hydrolysis of the cyclic amide bond of each substrate in either five- or six-membered rings. Even if these enzymes have similar functions, they have relatively low amino acid sequence identity. In addition, the substrate selectivity and specificity of these enzymes highly differ [18, 19]. Most of the active sites of dihydropyrimidinases, hydantoinases, allantoinases, and dihydroorotases contain four histidines, one aspartate, and one carboxylated lysine residue, which are required for metal binding and catalytic




FIGURE 1: The physiological reaction of dihydropyrimidinase. Dihydropyrimidinase catalyzes the reversible cyclization of dihydrouracil to *N*-carbamoyl- β -alanine in the second step of the pyrimidine degradation pathway.

activity [8, 15, 18, 20, 21]. The presence of a carboxylated lysine in hydantoinase is also required for the self-assembly of the binuclear metal center [12, 20, 22] and increases the nucleophilicity of the hydroxide for catalysis [23]. The global architecture of the dihydropyrimidinase monomer consists of two domains, namely, a large domain with a classic (β/α)₈-barrel structure core embedding the catalytic dimetal center and a small β -sandwich domain [16, 22, 24, 25].

All known dihydropyrimidinases are tetramers except pseudomonas enzymes. Hydantoinase from *P. putida* YZ-26 functions as a dimer [26, 27]. Recently, we identified that dihydropyrimidinase from *P. aeruginosa* PAO1 also forms a dimer [28]. In addition, the crystal structure of *P. aeruginosa* PAO1 dihydropyrimidinase indicated that several residues crucial for tetramerization are not found in *P. aeruginosa* dihydropyrimidinase [28]. In this study, we found that the oligomerization of *P. aeruginosa* PAO1 dihydropyrimidinase is a pH-dependent process. At pH 5.9, *P. aeruginosa* PAO1 dihydropyrimidinase mainly formed a tetramer. To confirm this result and determine how this enzyme can also form a tetramer, we also determined the crystal structure of *P. aeruginosa* PAO1 dihydropyrimidinase at 2.17 Å resolution at acidic environment. Structural comparison indicated that although *P. aeruginosa* PAO1 dihydropyrimidinase can also form a tetramer, the residues being crucial for tetramerization are different from those in *Thermus* sp. dihydropyrimidinases.

2. Materials and Methods

2.1. Cloning, Protein Expression, and Purification. Construction of the *P. aeruginosa* dihydropyrimidinase expression plasmid has been reported [15]. Recombinant *P. aeruginosa* dihydropyrimidinase was expressed and purified using the protocol described previously [15]. The protein purified from the soluble supernatant by Ni²⁺-affinity chromatography (HiTrap HP; GE Healthcare Bio-Sciences, Piscataway, NJ, USA) was eluted with Buffer A (20 mM Tris-HCl, 250 mM imidazole, and 0.5 M NaCl, pH 7.9) and dialyzed against a dialysis buffer (20 mM HEPES and 100 mM NaCl, pH 7.0; Buffer B). Protein purity remained > 97% as determined by SDS-PAGE (Mini-PROTEAN Tetra System; Bio-Rad, CA, USA).

2.2. Gel Filtration Chromatography. Gel filtration chromatography was carried out by the AKTA-FPLC system (GE Healthcare Bio-Sciences, Piscataway, NJ, USA). In brief, purified protein (5 mg/mL) in Buffer C (20 mM MES and

TABLE 1: Data collection and refinement statistics.

Data collection	
Crystal	<i>P. aeruginosa</i> dihydropyrimidinase
Wavelength (Å)	0.975
Resolution (Å)	30–2.17
Space group	C222 ₁
Cell dimension (Å)	$a = 108.9$, $\alpha = 90$ $b = 155.7$, $\beta = 90$ $c = 235.6$, $\gamma = 120$
Completeness (%)	99.8 (100)*
$\langle I/\sigma I \rangle$	15.13 (3.7)
R_{sym} or R_{merge} (%)	0.122 (0.599)
Redundancy	7.1 (7.3)
Refinement	
Resolution (Å)	30–2.17
Number of reflections	100197
$R_{\text{work}}/R_{\text{free}}$	0.1759/0.2312
Number of atoms	
Protein	1912
Water	312
RMS deviation	
Bond lengths (Å)	0.0151
Bond angles (°)	1.6495
Ramachandran plot	
In preferred regions	1345 (94.19%)
In allowed regions	68 (4.76%)
Outliers	15 (1.05%)
PDB entry	5YKD

*Values in parentheses are for the highest resolution shell.

100 mM NaCl, pH 5.9) was applied to a Superdex 200 prep grade column (GE Healthcare Bio-Sciences, Piscataway, NJ, USA) equilibrated with the same buffer [29]. The column was operated at a flow rate of 0.5 mL/min, and the proteins were detected at 280 nm. The column was calibrated with proteins of known molecular weight: thyroglobulin (670 kDa), γ -globulin (158 kDa), ovalbumin (44 kDa), myoglobin (17 kDa), and vitamin B₁₂ (1.35 kDa).

2.3. Crystallography. Before crystallization, *P. aeruginosa* dihydropyrimidinase was concentrated to 20 mg/mL in Buffer C. Crystals were grown at room temperature by hanging drop vapor diffusion in 10% PEG 8000, 100 mM HEPES, 200 mM calcium acetate, pH 5.9. Data collection and refinement statistics for the crystal of *P. aeruginosa* dihydropyrimidinase are shown in Table 1. Data were collected using an ADSC Quantum-315r CCD area detector at SPXF beamline BL13C1 at NSRRRC (Taiwan, ROC). All data integration and scaling were carried out using HKL-2000 [30]. There were four *P. aeruginosa* dihydropyrimidinase monomers per asymmetric unit. The crystal structure of *P. aeruginosa* dihydropyrimidinase was solved at 2.17 Å resolution with the molecular replacement software AMoRe [31] using the dihydropyrimidinase (PDB entry 5E5C) [28] as




FIGURE 2: Crystal structure of *P. aeruginosa* dihydropyrimidinase. (a) Ribbon diagram of a *P. aeruginosa* dihydropyrimidinase tetramer. Each *P. aeruginosa* dihydropyrimidinase monomer is color-coded. Two zinc ions in the active site are presented as black spheres. (b) Ribbon diagram of a *P. aeruginosa* dihydropyrimidinase monomer with the secondary structures labeled.




FIGURE 3: Structural comparison. (a) Superposition of the active site of dihydropyrimidinases. Their active sites contain four histidines, one aspartate, and one carboxylated lysine residue, which are required for metal binding and catalytic activity. Dihydropyrimidinases from *P. aeruginosa* (PDB entry 5E5C; green), *Thermus* sp. (PDB entry 1GKQ; salmon), *Tetraodon nigroviridis* (PDB entry 4H01; pale yellow), and the structure (PDB entry 5YKD; purple blue) in this study are shown. The architecture of these active sites is similar. (b) Superposition of the active site of members of the amidohydrolase family. Their active sites contain four histidines, one aspartate, and one carboxylated lysine residue, which are required for metal binding and catalytic activity. *P. aeruginosa* dihydropyrimidinase (PDB entry 5YKD; purple blue), *Escherichia coli* allantoinase (PDB entry 3E74; bright orange), *Burkholderia pickettii* hydantoinase (PDB entry 1NFG; aquamarine), and *E. coli* dihydroorotase (PDB entry 1J79; brown) are shown. The architecture of these active sites is similar.

model. After molecular replacement, model building was carried out using XtalView [32]. CNS was used for molecular dynamics refinement [33]. The final structure was refined to an R -factor of 0.1759 and an R_{free} of 0.2312. Atomic coordinates and related structural factors have been deposited in the PDB with accession code 5YKD.

3. Results and Discussion

3.1. Structure of the *P. aeruginosa* Dihydropyrimidinase Monomer. Crystals of *P. aeruginosa* dihydropyrimidinase

were grown at room temperature by hanging drop vapor diffusion in 10% PEG 8000, 100 mM HEPES, 200 mM calcium acetate, pH 5.9. The crystals of *P. aeruginosa* dihydropyrimidinase grown under this condition belonged to space group $C222_1$ with cell dimensions of $a = 108.9$, $b = 155.7$, and $c = 235.6$ Å. The crystal structure of *P. aeruginosa* dihydropyrimidinase was solved at 2.17 Å resolution (Table 1). The unit cell contained eight molecules. An asymmetric unit of the crystal contained four crystallographically independent *P. aeruginosa* dihydropyrimidinase monomers, in which two zinc ions were found in the active site per




FIGURE 4: Gel filtration chromatographic analysis. Gel filtration chromatography was carried out by the AKTA-FPLC system in Buffer C (20 mM MES and 100 mM NaCl, pH 5.9). The corresponding peaks show the eluting *P. aeruginosa* dihydropyrimidinase. The column was calibrated with proteins of known molecular weight: thyroglobulin (670 kDa), γ -globulin (158 kDa), ovalbumin (44 kDa), myoglobin (17 kDa), and vitamin B₁₂ (1.35 kDa).




FIGURE 5: The structure of *P. aeruginosa* dihydropyrimidinase tetramer. An asymmetric unit contains four crystallographically independent *P. aeruginosa* dihydropyrimidinase monomers B-A-C-D. Crystallographically related tetramer B-A-C'-D' was formed and further stabilized via many hydrogen bonds and salt bridges. This tetramerization structure was similar to that of *Thermus* sp. dihydropyrimidinase (PDB entry 1GKQ).

monomer (Figure 2(a)). The majority of the electron density for *P. aeruginosa* dihydropyrimidinase exhibited good quality, and no discontinuity was observed. Briefly, the overall structure of each *P. aeruginosa* dihydropyrimidinase unit consists of 17 α -helices, 19 β -sheets, and two zinc ions (Figure 2(b)). At pH 5.9, the architecture of the *P. aeruginosa* dihydropyrimidinase monomer consists of two domains, namely, a large domain with a classic $(\beta/\alpha)_8$ -barrel structure core embedding the catalytic dimetal center and a small β -sandwich domain.

3.2. Structural Comparison. The overall structure and architecture of the active site of *P. aeruginosa* dihydropyrimidinase are similar to those of other dihydropyrimidinases (Figure 3(a)) and other members of the amidohydrolase family of enzymes, such as hydantoinases, dihydroorotases,

and allantoinases (Figure 3(b)). The active sites of these enzymes contain four histidines, one aspartate, and one carboxylated lysine residue, which are required for metal binding and catalytic activity [12, 14, 15, 19, 20, 34, 35].

3.3. pH-Dependent Oligomerization of *P. aeruginosa* Dihydropyrimidinase. It was noted that the crystals of the dimeric *P. aeruginosa* dihydropyrimidinase belonged to space group $P3_121$ grown at the condition of 28% PEG 6000, 100 mM HEPES, 200 mM lithium acetate, pH 7.5 [28]. Due to the different crystallization condition, we attempted to test whether the oligomerization of *P. aeruginosa* dihydropyrimidinase is pH-dependent. All known dihydropyrimidinases are tetramers. However, pseudomonas dihydropyrimidinase/hydantoinase forms a dimer at neutral pH [26–28]. Given that the structure implies that

TABLE 2: The formation of hydrogen bonds at the dimer-dimer interface of *P. aeruginosa* dihydropyrimidinase.

Subunit 1	Distance [Å]	Subunit 2
A: K374 [NZ]	3.00	B: E14 [OE1]
A: H13 [NE2]	2.88	B: E14 [OE1]
A: R386 [NH2]	3.86	B: E14 [OE2]
A: R386 [NH1]	2.81	B: E15 [OE2]
A: R386 [NH2]	2.83	B: E15 [OE2]
A: R468 [NH2]	3.61	B: Q306 [OE1]
A: R253 [NH1]	3.27	B: S307 [O]
A: R253 [NH2]	3.13	B: S307 [O]
A: R467 [NH1]	2.92	B: V354 [O]
A: R468 [NE]	2.95	B: G357 [O]
A: R468 [NH2]	3.09	B: G357 [O]
A: R468 [NH2]	3.40	B: R358 [O]
A: R467 [NH1]	3.24	B: L359 [O]
A: E14 [OE1]	3.09	B: K374 [NZ]
A: E14 [OE1]	2.47	B: H13 [NE2]
A: E15 [OE2]	2.70	B: R386 [NH1]
A: S307 [O]	3.30	B: R253 [NH1]
A: S307 [O]	3.55	B: R253 [NH2]
A: V354 [O]	2.91	B: R467 [NH1]
A: G357 [O]	2.94	B: R468 [NH2]
A: G357 [O]	2.94	B: R468 [NE]
A: R358 [O]	3.56	B: R468 [NH2]
A: L359 [O]	3.16	B: R467 [NH1]
C': H13 [NE2]	2.79	D': E14 [OE1]
C': K374 [NZ]	3.25	D': E14 [OE1]
C': R386 [NH1]	2.85	D': E15 [OE1]
C': R386 [NH2]	2.59	D': E15 [OE2]
C': R468 [NH2]	3.26	D': Q306 [OE1]
C': R253 [NH1]	3.13	D': S307 [O]
C': R253 [NH2]	3.16	D': S307 [O]
C': R468 [NE]	2.71	D': G357 [O]
C': R468 [NH2]	3.11	D': R358 [O]
C': E14 [OE1]	2.88	D': H13 [NE2]
C': E14 [OE1]	2.89	D': K374 [NZ]
C': E15 [OE2]	2.88	D': R386 [NH1]
C': E15 [OE2]	2.73	D': R386 [NH2]
C': Q306 [OE1]	3.53	D': R468 [NH2]
C': S307 [O]	3.21	D': R253 [NH1]
C': S307 [O]	3.59	D': R253 [NH2]
C': G357 [O]	2.65	D': R468 [NE]
C': R358 [O]	3.33	D': R468 [NH2]

P. aeruginosa dihydropyrimidinase may also form a tetramer in the crystalline state at pH 5.9 (Figure 2(a)), we performed biochemical verification to confirm the oligomerization state. To confirm whether or not the oligomerization of *P. aeruginosa* dihydropyrimidinase is pH-dependent, we conducted gel filtration chromatography at pH 5.9. As shown in Figure 4, the results revealed that two species with

TABLE 3: The formation of salt bridges at the dimer-dimer interface of *P. aeruginosa* dihydropyrimidinase.

Subunit 1	Distance [Å]	Subunit 2
A: K374 [NZ]	3.00	B: E14 [OE1]
A: H13 [NE2]	2.88	B: E14 [OE1]
A: R386 [NH2]	3.86	B: E14 [OE2]
A: H13 [NE2]	3.75	B: E14 [OE2]
A: R386 [NH1]	3.55	B: E15 [OE1]
A: R386 [NH1]	2.81	B: E15 [OE2]
A: R386 [NH2]	2.83	B: E15 [OE2]
A: E14 [OE1]	3.09	B: K374 [NZ]
A: E14 [OE1]	2.47	B: H13 [NE2]
A: E14 [OE2]	3.93	B: H13 [NE2]
A: E15 [OE1]	3.69	B: R386 [NH1]
A: E15 [OE2]	3.00	B: R386 [NH2]
A: E15 [OE2]	2.70	B: R386 [NH1]
C': H13 [NE2]	2.79	D': E14 [OE1]
C': K374 [NZ]	3.25	D': E14 [OE1]
C': H13 [NE2]	3.86	D': E14 [OE2]
C': R386 [NH1]	2.85	D': E15 [OE1]
C': R386 [NH2]	3.84	D': E15 [OE1]
C': R386 [NH1]	2.96	D': E15 [OE2]
C': R386 [NH2]	2.59	D': E15 [OE2]
C': E14 [OE1]	2.88	D': H13 [NE2]
C': E14 [OE1]	2.89	D': K374 [NZ]
C': E14 [OE2]	3.78	D': H13 [NE2]
C': E15 [OE1]	3.34	D': R386 [NH1]
C': E15 [OE2]	2.88	D': R386 [NH1]
C': E15 [OE2]	2.73	D': R386 [NH2]

elution volume of 63.25 and 69.26 mL did coexist. The molecular mass of a *P. aeruginosa* dihydropyrimidinase monomer, as calculated from the amino acid sequence, is 53 kDa. Assuming that these two forms of *P. aeruginosa* dihydropyrimidinase have a shape and partial specific volume similar to the standard proteins, the native molecular masses of *P. aeruginosa* dihydropyrimidinase were estimated to be 105 and 180 kDa, approximately 1.9 and 3.5 times the molecular mass of a *P. aeruginosa* dihydropyrimidinase monomer, respectively. In comparison at pH 7.5, gel filtration chromatographic analysis of *P. aeruginosa* dihydropyrimidinase revealed a single peak; the native molecular mass was estimated to be 117 kDa [28]. The two forms of this enzyme obtained from the gel filtration chromatography at pH 5.9 had similar specific activity (data not shown). Thus, *P. aeruginosa* dihydropyrimidinase did exist as a mixture of dimers and tetramers at pH 5.9.

3.4. Structural Insights into Dimer of Dimer (Tetramer) Formation of Dihydropyrimidinase. In this study, we have identified that *P. aeruginosa* dihydropyrimidinase did exist as a mixture of dimers and tetramers at pH 5.9. To assess how *P. aeruginosa* dihydropyrimidinase can form a stable tetramer, the dimer-dimer interface was analyzed. In the




FIGURE 6: Comparison of the tetrameric structures of *Thermus* sp. dihydropyrimidinase and *P. aeruginosa* dihydropyrimidinase. (a) Structural analysis of the dimer-dimer interface of *P. aeruginosa* dihydropyrimidinase. The distance (Å) of the residues is shown. (b) Many residues crucial for forming hydrogen bonds at the dimer-dimer interface of *P. aeruginosa* dihydropyrimidinase were not found in the dimer-dimer interface of *Thermus* sp. dihydropyrimidinase.

M S L L I R G A T V V T H E E S Y R A D V L C A N G L I Q A I G E N L E T P S G C D V L D G G G Q Y L M P G G I D P H T	60
M P L L I K N G E I I T A D S R Y K A D I Y A E G E T I T R I G Q N L E A P P G T E V I D A T G K Y V F P G F I D P H V	60
H M Q L P F M G T V A S E D F F S G T A A G L A G G T T S I I D F V I P N P R Q S L L E A F H T W R G W A Q - K S A A D	119
H I Y L P F M A T F A K D T H E T G S K A A L M G G T T Y I E M C C P S R N D D A L E G Y Q L W K S K A E G N S Y C D	120
Y G F H V A I T W W S D E V A R E M G E L V A Q H G V N S F K H F M A Y K N A I M A A D D T L V A S F E R C L E L G A V	179
Y T F H M A V S K F D E K T E G Q L R E I V A D - G I S S F K I F L S Y K N F F G V D D G E M Y Q T L R L A K E L G V I	179
P T V H A E N G E L V F H L Q Q K L L A Q G L T G P E A H P L S R P P Q V E G E A A S R A I R I A E T L G T P L Y L V H	239
V T A H C E N A E L V G R L Q Q K L L S E G K T G P E W H E P S R P E A V E A E G T A R F A T F L E T T G A T G Y V V H	239
I S S R E A L D E I A Y A R A K G Q P V Y G E V L A G H L L L D D S V Y R H P D W A T A A G Y V M S P P F R P V E H Q E	299
L S C K P A L D A A M A A K A R G V P I Y I E S V I P H F L L D K T - Y A E R G G V E A M K Y I M S P P L R D K R N Q K	298
A L W R G L Q S G N L H T T A T D H C C F C A E Q K A M G R D D F S K I P N G T A G I E D R M A L L W D A G V N S G R L	359
V L W D A L A Q G F I D T V G T D H C P F D T E Q K L L G K E A F T A I P N G I P A I E D R V N L L Y T Y G V S R G R L	358
S M H E F V A L T S T N T A K I F N L F P R K G A I R V G A D A D L V L W D P Q G S R T L S A A T H H Q R V D F N I F E	419
D I H R F V D A A S T K A A K L F G L F P R K G T I A V G S D A D L V V Y D P Q Y R G T I S V K T Q H V N N D Y N G F E	418
G R T V R G I P S H T I S Q G K L L W A A G D L R A E P G A G R Y V E R - P A Y P S V Y E V L G R R A E R Q R P V A V E	478
G F E I D G R P S V V T V R G K V A V R D G Q F V G E K C W G K L L R E P M Y F -	458
R <i>Pseudomonas aeruginosa</i> PAO1	479
- <i>Thermus</i> sp.	

FIGURE 7: Sequence alignment of dihydropyrimidinases from *P. aeruginosa* and *Thermus* sp. The amino acids that are involved in dimer-dimer interface of *P. aeruginosa* and *Thermus* sp. dihydropyrimidinase are boxed, respectively.

crystal of *P. aeruginosa* dihydropyrimidinase, the four molecules formed two pairs of dimers, B-A and C-D, respectively (Figure 5). Since the two dimers of *P. aeruginosa* dihydropyrimidinase associate via few contacts to create the

tetramer, it was thought that the tetrameric state may be possibly due to crystal packing forces. We noted that in the crystal, another crystallographically related tetramer B-A-C^I-D^I (Figure 5) was formed and further stabilized via many

hydrogen bonds and salt bridges (Tables 2 and 3). This tetramerization structure was similar to that of *Thermus* sp. dihydropyrimidinase (PDB entry 1GKQ).

We also compared the residues important for tetramerization located at the B-A-C'-D' dimer-dimer interface with those of *Thermus* sp. dihydropyrimidinase (Figure 6). Although their overall structures are similar, the important residues for tetramer (dimer B-C' with dimer A-D') formation are quite different. For the tetramer formation of *P. aeruginosa* dihydropyrimidinase, many hydrogen bonds with close distance were found: these bonds (<3 Å) include K374(A)-E14(B), H13(A)-E14(B), R386(A)-E14(B), R386(A)-E15(B), R467(A)-V354(B), R468(A)-G357(B), E14(A)-H13(B), E15(A)-R386(B), V354(A)-R467(B), G357(A)-R468(B), H13(C')-E14(D'), R386(C')-E15(D'), R468(C')-G357(D'), E14(C')-H13(D'), E14(C')-K374(D'), E15(C')-R386(D'), and G357(C')-R468(D'); however, these residues were not found for the tetramer formation of *Thermus* sp. dihydropyrimidinase (Figure 6). Only A13-D14 hydrogen bond was found in *Thermus* sp. dihydropyrimidinase (i.e., H13-E14 in *P. aeruginosa* dihydropyrimidinase). Thus, the dimer-dimer interface between *P. aeruginosa* dihydropyrimidinase and *Thermus* sp. dihydropyrimidinase was significantly different (Figure 7). Comparison by superimposition indicated that many Arg residues (R253, R358, R386, R467, and R468) found in *P. aeruginosa* dihydropyrimidinase, but not in *Thermus* sp. dihydropyrimidinase, may play a crucial role for the pH-dependent oligomerization. If consider the pK_a , a much better candidate is His13, which is involved in intermolecular interactions and, dependent on the environment of its side chain, which may easily change protonation state between pH 5.9 and pH 7.5. However, this speculation needs to be confirmed by further biochemical experiments.

3.5. Different Mechanisms for Tetramer Formation of Dihydropyrimidinases. In this study, we identified *P. aeruginosa* dihydropyrimidinase can be a tetramer both in the crystalline state and in solution (Figure 4). The structure of the tetrameric *Thermus* sp. dihydropyrimidinase and *P. aeruginosa* dihydropyrimidinase was compared (Figure 6). Many important residues for *Thermus* sp. dihydropyrimidinase tetramer formation are different from those for *P. aeruginosa* dihydropyrimidinase (Figure 7). On the basis of these results, we concluded that *P. aeruginosa* dihydropyrimidinase could form a tetramer, but its oligomerization mechanism differed from those of other dihydropyrimidinases such as *Thermus* sp. dihydropyrimidinase.

Conflicts of Interest

The authors declare that they have no conflicts of interest regarding the publication of this paper.

Acknowledgments

The authors thank the technical services provided by the Synchrotron Radiation Protein Crystallography Facility of the National Core Facility Program for Biotechnology,

Ministry of Science and Technology, and the National Synchrotron Radiation Research Center, a national user facility supported by the Ministry of Science and Technology, Taiwan, ROC. This research was supported by a grant from the Ministry of Science and Technology, Taiwan (MOST 106-2320-B-040-004) to Cheng-Yang Huang.

References

- [1] K. D. Schnackerz and D. Dobritzsch, "Amidohydrolases of the reductive pyrimidine catabolic pathway purification, characterization, structure, reaction mechanisms and enzyme deficiency," *Biochimica et Biophysica Acta*, vol. 1784, no. 3, pp. 431-444, 2008.
- [2] D. P. Wallach and S. Grisolia, "The purification and properties of hydroxypyrimidine hydrase," *Journal of Biological Chemistry*, vol. 226, no. 1, pp. 277-288, 1957.
- [3] C. C. Hsu, L. Y. Lu, and Y. S. Yang, "From sequence and structure of sulfotransferases and dihydropyrimidinases to an understanding of their mechanisms of action and function," *Expert Opinion on Drug Metabolism & Toxicology*, vol. 6, no. 5, pp. 591-601, 2010.
- [4] C. Y. Huang and Y. S. Yang, "Discovery of a novel N-iminylamidase activity: substrate specificity, chemoselectivity and catalytic mechanism," *Protein Expression and Purification*, vol. 40, no. 1, pp. 203-211, 2005.
- [5] C. Y. Huang and Y. S. Yang, "A novel cold-adapted imidase from fish *Oreochromis niloticus* that catalyzes hydrolysis of maleimide," *Biochemical and Biophysical Research Communications*, vol. 312, no. 12, pp. 467-472, 2003.
- [6] C. Y. Huang, S. K. Chiang, Y. S. Yang, and Y. J. Sun, "Crystallization and preliminary X-ray diffraction analysis of thermophilic imidase from pig liver," *Acta Crystallographica Section D Biological Crystallography*, vol. 59, no. 5, pp. 943-945, 2003.
- [7] C. Y. Huang, Y. P. Chao, and Y. S. Yang, "Purification of industrial hydantoinase in one chromatographic step without affinity tag," *Protein Expression and Purification*, vol. 30, no. 1, pp. 134-139, 2003.
- [8] C. Y. Huang and Y. S. Yang, "The role of metal on imide hydrolysis: metal content and pH profiles of metal ion-replaced mammalian imidase," *Biochemical and Biophysical Research Communications*, vol. 297, no. 4, pp. 1027-1032, 2002.
- [9] Y. S. Yang, S. Ramaswamy, and W. B. Jakoby, "Rat liver imidase," *Journal of Biological Chemistry*, vol. 268, no. 15, pp. 10870-10875, 1993.
- [10] H. E. Schoemaker, D. Mink, and M. G. Wubbolts, "Dispelling the myths-biocatalysis in industrial synthesis," *Science*, vol. 299, no. 5613, pp. 1694-1697, 2003.
- [11] J. Altenbuchner, M. Siemann-Herzberg, and C. Syldatk, "Hydantoinases and related enzymes as biocatalysts for the synthesis of unnatural chiral amino acids," *Current Opinion in Biotechnology*, vol. 12, no. 6, pp. 559-563, 2001.
- [12] C. Y. Huang, C. C. Hsu, M. C. Chen, and Y. S. Yang, "Effect of metal binding and posttranslational lysine carboxylation on the activity of recombinant hydantoinase," *Journal of Biological Inorganic Chemistry*, vol. 14, no. 1, pp. 111-121, 2009.
- [13] S. M. Runser and P. C. Meyer, "Purification and biochemical characterization of the hydantoin hydrolyzing enzyme from *Agrobacterium* species. A hydantoinase with no 5,6-dihydropyrimidine amidohydrolase activity," *European Journal of Biochemistry*, vol. 213, no. 3, pp. 1315-1324, 1993.

- [14] Z. Gojkovic, L. Rislund, B. Andersen et al., "Dihydropyrimidine amidohydrolases and dihydroorotases share the same origin and several enzymatic properties," *Nucleic Acids Research*, vol. 31, no. 6, pp. 1683–1692, 2003.
- [15] C. Y. Huang, "Inhibition of a putative dihydropyrimidinase from *Pseudomonas aeruginosa* PAO1 by flavonoids and substrates of cyclic amidohydrolases," *PLoS One*, vol. 10, no. 5, Article ID e0127634, 2015.
- [16] J. Abendroth, K. Niefind, and D. Schomburg, "X-ray structure of a dihydropyrimidinase from *Thermus* sp. at 1.3 Å resolution," *Journal of Molecular Biology*, vol. 320, no. 1, pp. 143–156, 2002.
- [17] L. Holm and C. Sander, "An evolutionary treasure: unification of a broad set of amidohydrolases related to urease," *Proteins: Structure, Function, and Genetics*, vol. 28, no. 1, pp. 72–82, 1997.
- [18] W. F. Peng and C. Y. Huang, "Allantoinase and dihydroorotase binding and inhibition by flavonols and the substrates of cyclic amidohydrolases," *Biochimie*, vol. 101, pp. 113–122, 2014.
- [19] C. C. Wang, H. W. Tsau, W. T. Chen, and C. Y. Huang, "Identification and characterization of a putative dihydroorotase KPN01074, from *Klebsiella pneumoniae*," *Protein Journal*, vol. 29, no. 6, pp. 445–452, 2010.
- [20] Y. Y. Ho, Y. H. Huang, and C. Y. Huang, "Chemical rescue of the post-translationally carboxylated lysine mutant of allantoinase and dihydroorotase by metal ions and short-chain carboxylic acids," *Amino Acids*, vol. 44, no. 4, pp. 1181–1191, 2013.
- [21] D. R. Evans and H. I. Guy, "Mammalian pyrimidine biosynthesis: fresh insights into an ancient pathway," *Journal of Biological Chemistry*, vol. 279, no. 32, pp. 33035–33038, 2004.
- [22] Y. C. Hsieh, M. C. Chen, C. C. Hsu, S. I. Chan, Y. S. Yang, and C. J. Chen, "Crystal structures of vertebrate dihydropyrimidinase and complexes from *Tetraodon nigroviridis* with lysine carbamylation: metal and structural requirements for post-translational modification and function," *Journal of Biological Chemistry*, vol. 288, no. 42, pp. 30645–30658, 2013.
- [23] V. Kumar, N. Saxena, M. Sarma, and K. V. Radha Kishan, "Carboxylated lysine is required for higher activities in hydantoinases," *Protein & Peptide Letters*, vol. 18, no. 7, pp. 663–669, 2011.
- [24] S. Martinez-Rodriguez, A. I. Martinez-Gomez, J. M. Clemente-Jimenez et al., "Structure of dihydropyrimidinase from *Sinorhizobium meliloti* CECT4114: new features in an amidohydrolase family member," *Journal of Structural Biology*, vol. 169, pp. 200–208, 2010.
- [25] B. Lohkamp, B. Andersen, J. Piskur, and D. Dobritzsch, "The crystal structures of dihydropyrimidinases reaffirm the close relationship between cyclic amidohydrolases and explain their substrate specificity," *Journal of Biological Chemistry*, vol. 281, no. 19, pp. 13762–13776, 2006.
- [26] X. Y. Zhang, L. X. Niu, Y. W. Shi, and J. M. Yuan, "The flexibility of the non-conservative region at the C terminus of D-hydantoinase from *Pseudomonas putida* YZ-26 is extremely limited," *Applied Biochemistry and Biotechnology*, vol. 144, no. 3, pp. 237–247, 2008.
- [27] L. Niu, X. Zhang, Y. Shi, and J. Yuan, "Subunit dissociation and stability alteration of D hydantoinase deleted at the terminal amino acid residue," *Biotechnology Letters*, vol. 29, no. 2, pp. 303–308, 2007.
- [28] C. T. Tzeng, Y. H. Huang, and C. Y. Huang, "Crystal structure of dihydropyrimidinase from *Pseudomonas aeruginosa* PAO1: insights into the molecular basis of formation of a dimer," *Biochemical and Biophysical Research Communications*, vol. 478, no. 3, pp. 1449–1455, 2016.
- [29] Y. H. Huang, Y. Lien, C. C. Huang, and C. Y. Huang, "Characterization of *Staphylococcus aureus* primosomal DnaD protein: highly conserved C-terminal region is crucial for ssDNA and PriA helicase binding but not for DnaA protein-binding and self-tetramerization," *PLoS One*, vol. 11, no. 6, Article ID e0157593, 2016.
- [30] Z. Otwinowski and W. Minor, "Processing of X-ray diffraction data collected in oscillation mode," *Methods in Enzymology*, vol. 276, pp. 307–326, 1997.
- [31] J. Navaza, "AMoRe: an automated package for molecular replacement," *Acta Crystallographica Section A Foundations of Crystallography*, vol. 50, no. 2, pp. 157–163, 1994.
- [32] D. E. McRee, "XtalView/Xfit—A versatile program for manipulating atomic coordinates and electron density," *Journal of Structural Biology*, vol. 125, no. 2–3, pp. 156–165, 1999.
- [33] A. T. Brunger, P. D. Adams, G. M. Clore et al., "Crystallography & NMR system: a new software suite for macromolecular structure determination," *Acta Crystallographica Section D Biological Crystallography*, vol. 54, no. 5, pp. 905–921, 1998.
- [34] Y. H. Huang and C. Y. Huang, "Creation of a putative third metal binding site in type II dihydroorotases significantly enhances enzyme activity," *Protein & Peptide Letters*, vol. 22, no. 12, pp. 1117–1122, 2015.
- [35] Y. Y. Ho, H. C. Hsieh, and C. Y. Huang, "Biochemical characterization of allantoinase from *Escherichia coli* BL21," *Protein Journal*, vol. 30, no. 6, pp. 384–394, 2011.



Hindawi

Submit your manuscripts at
www.hindawi.com

