Research Article

Serine Protease Variants Encoded by *Echis ocellatus* Venom Gland cDNA: Cloning and Sequencing Analysis

S. S. Hasson,¹ R. A. Mothana,² T. A. Sallam,³ M. S. Al-balushi,¹ M. T. Rahman,¹
and A. A. Al-Jabri¹

¹ Division of Immunology, Department of Microbiology and Immunology, College of Medicine and Health Sciences, Sultan Qaboos University, P.O. Box 35, Muscat, 123, Oman

² Department of Pharmacognosy, College of Pharmacy, King Saud University, P.O. Box 2457, Riyadh 11451, Saudi Arabia

³ Department of Community Health, Faculty of Medical Sciences, Al-Baha University, Al-Baha, P.O. Box 2457, Al-Baha 11451, Saudi Arabia

Correspondence should be addressed to S. S. Hasson, shyahasson@yahoo.co.uk

Received 28 May 2010; Accepted 20 July 2010

Academic Editor: Fuli Yu

Copyright © 2010 S. S. Hasson et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Envenoming by *Echis* saw-scaled viper is the leading cause of death and morbidity in Africa due to snake bite. Despite its medical importance, there have been few investigations into the toxin composition of the venom of this viper. Here, we report the cloning of cDNA sequences encoding four groups or isoforms of the haemostasis-disruptive Serine protease proteins (SPs) from the venom glands of *Echis ocellatus*. All these SP sequences encoded the cysteine residues scaffold that form the 6-disulphide bonds responsible for the characteristic tertiary structure of venom serine proteases. All the *Echis ocellatus* EoSP groups showed varying degrees of sequence similarity to published viper venom SPs. However, these groups also showed marked intercluster sequence conservation across them which were significantly different from that of previously published viper SPs. Because viper venom SPs exhibit a high degree of sequence similarity and yet exert profoundly different effects on the mammalian haemostatic system, no attempt was made to assign functionality to the new *Echis ocellatus* EoSPs on the basis of sequence alone. The extraordinary level of interspecific and intergeneric sequence conservation exhibited by the *Echis ocellatus* EoSPs and analogous serine proteases from other viper species leads us to speculate that antibodies to representative molecules should neutralise (that we will exploit, by epidermal DNA immunization) the biological function of this important group of venom toxins in vipers that are distributed throughout Africa, the Middle East, and the Indian subcontinent.

1. Introduction

Envenoming resulting from snake bites is an important public health hazard in many regions, particularly in tropical and subtropical countries [1, 2]. The saw-scaled viper *Echis ocellatus* is the most abundant [3] and medically important viper species in West Africa [4]. Envenoming by saw-scaled viper (*Viperidae*: *Echis*) species is thought to be responsible for more snakebite deaths worldwide than any other snake genus [5]. In northern Nigeria, *E. ocellatus* is responsible for 95% of all envenoming by snakes [6], causing several hundred deaths annually [7]. The precise incidence of snakebite is difficult to determine and is often grossly underestimated, but in some areas of the Nigerian savannas, victims of *E. ocellatus* envenoming may occupy more than

10% of hospital beds [8]. In the Benue valley of Nigeria, for example, the estimated incidence is 497 per 100 000 population per year with 10%–20% untreated mortality [9]. Local effects of *Echis* viper envenoming include pain, swelling, blistering, and haemorrhage which, in severe cases, can lead to necrosis, permanent disfigurement, and even amputation of the affected limb [10]. Systemic effects include potentially lethal consumption coagulopathy, haemorrhage and hypovolaemic shock [10].

Snake venoms contain a great variety of toxic proteases [11, 12]. Many of these components are proteases, for example, metalloproteases [13], serine proteases [14], phospholipases A₂ [15] and C-type lectins [16] and mediate their toxicity by either stimulating or inhibiting the haemostatic system of human victims or experimental animals, resulting

in clinical complications of blood clotting or uncontrolled haemorrhage [12, 17–19]. Several of these proteinases cleave plasma proteins of the victims in a specific manner with varying degrees of substrate specificity. Thus, while some serine proteases have both fibrinogenolytic and fibrinolytic activities, others have only fibrinogenolytic activity and are called “thrombin-like” proteases [19–25]. Approximately 100 snake venom toxins have been identified as “thrombin-like” enzymes activating the blood coagulation factor [26]. These “thrombin-like” proteases hydrolyze fibrinogen specifically and release either fibrinopeptide A or B or both [27] resulting in the disruption of the blood coagulation system by producing abnormal fibrin clots composed of short polymers that are rapidly dispersed and no longer cross-linked by activated factor XIII [28].

Another group of serine proteases of *Batroxobin*, *Crotalase*, and Ancrod venoms affect other substrates, for example, plasminogen [27] by cleaving fibrinogen in manner distinct from that of thrombin. Other venom serine proteases function like mammalian kallikrein (or kininogenase) releasing bradykinin from kininogen [29–31] and are called “kallikrein-like” proteases [29], an example of this is halystase [32], a kallikrein-like serine protease isolated from *A. halys blomhoffii* venom, which cleaves the β chain at Arg⁴² and slowly degrades the α chain of fibrinogen to generate a product that is no longer converted to normal fibrin clots by thrombin; this results in both reduction of blood pressure as well as inhibiting fibrinogen clotting in the victims. Another kallikrein-like serine protease with potent biological activity but with different physicochemical properties from those of halystase has been isolated from the venoms of *A. caliginosus*, *C. atrox*, *C. viridis*, and *Trimeresurus mucrosquamatus* [29, 30, 32–34]. The latter showed both a strong β -fibrinogenolytic and kallikrein-like activities, cleaving β -chain of fibrinogen molecules specifically and releasing bradykinin from kininogen, respectively. Moreover, the purified enzymes indicated that they have specificities different from thrombin and thrombin-like proteases of snake venom reported previously by decreasing fibrinogen levels in plasma and prolonging bleeding without formation of fibrin clots. They also exhibit amidase activity against *N*-benzoyl-Pro-Phe-Arg-*p*-nitroanilide, which is a specific synthetic substrate for kallikrein-like proteases.

In addition, there have been a few reports on venom serine proteases with a unique activity, such as ACC-C, a protein C activator isolated from the *A. contortrix* venom [35] (which inhibits blood coagulation by inactivating the activated forms of factor V and VIII), a plasminogen activator such as TSV-PA isolated from the *T. stejnegeri* venom [36, 37], PA-BJ, a platelet aggregating enzyme isolated from the *B. jararaca* and *Trimeresurus mucrosquamatus* venoms [38], and RVV-V, a factor V-activating enzyme isolated from the *V. russelli* venom [39].

These data indicate that snake venom serine proteases comprise an enzyme superfamily with multifunctional activities that may have diverged or have undergone gene duplication resulting in alteration of their biological properties during the process of evolution thus acquiring special functions [40, 41]. Although a considerable amount of data

is now available, no standardised grouping of these venom serine proteases has yet been documented. However, in 2001 Wang et al. [27] compared sequences of 40 serine proteinases isolated from different snake venoms, using a constructed phylogram in which such sequences were clustered into three groups designated as coagulating enzymes, kininogenases, and plasminogen activators.

No Serine proteinases have yet been purified from venom of the West African saw-scaled viper *Echis ocellatus*, in particular or for members of the *Echis* genus in general. However, the fact that the serine protease superfamily was important in the venom of the Viperidae suggested that such enzymes should be present in the venom of *E. ocellatus* and that serine protease-specific antibodies are likely to be an important factor in *E. ocellatus* envenoming. We therefore screened the *E. ocellatus* cDNA library in order to isolate and characterise different isoforms or variants of this enzyme superfamily.

2. Materials and Methods

2.1. Animals. Adult *E. ocellatus* (Nigeria) carpet viper used in this study was maintained in the herpetarium, Liverpool School of Tropical Medicine, Liverpool, UK.

2.2. Extraction of Total Venom Gland RNA and Construction of cDNA Libraries. Venom glands were dissected from three *Echis ocellatus* snakes. The vipers were sacrificed 3 days after venom extraction when toxin gene transcription rates are at a peak. Glands were homogenized under liquid Nitrogen and total RNA extracted using guanidinium thiocyanate-phenol-chloroform as described previously [15]. Lambda phage cDNA libraries for *E. ocellatus* were constructed by RT-PCR using the SMART cDNA library construction kit (Clontech, California, USA). The lambda phage of the *E. ocellatus* was packaged using Gigapack III Gold Packaging Extract (Stratagene) and boiled for 5 min prior to being used as targets of polymerase chain reaction (PCR) amplification.

2.3. Isolation and Analysis of cDNA Sequences. A PCR strategy [42] was used to isolate sequences encoding serine proteinases from the cDNA libraries. A sense primer (5V-**GGA-TCC-ATG-GTG-CTG-ATC-AGA-GTG-CTA-ATC-GCA-3V**) and an antisense primer (5V-**CTC-GAG-TGG-GGG-GCA-AGT-CGC-AGT-TGT-ATT-TCC-3V**) complementary to highly conserved amino-terminal signal peptide (M-V-L-I-R-V) and to the less conserved carboxy-terminal (T-T-A-T-C-P-P) domains of published serine proteinases DNA sequences of related viper species were synthesized commercially (Sigma-Genosys, UK). A TAG stop codon was inserted in the 3' primer and BamH1 and Xho1 restriction endonuclease sites (bold) were included in the 5' and 3' primers, respectively, to facilitate future subcloning into mammalian expression plasmids. PCR was performed using an initial denaturation (95°C—6 minutes) and annealing (55°C—1 minute) step, followed by 35 cycles (1 minute each) of extension (74°C), denaturation (94°C), and annealing (55°C), and a terminal extension step (7 minutes)

at 72°C in a thermal cycler (Gene Cycler, BioRad, Hercules, CA, USA). The inclusion of water-only controls with each PCR reaction allowed us to monitor and prevent cross-over contamination. The amplicons were subcloned into the TA cloning vector, pCR 2.1-TOPO, (Invitrogen, Groningen, The Netherlands) and used to transform chemically competent *E. coli* cells (TOP10F', Invitrogen) under ampicillin selection. Plasmid DNA was extracted (Mini-spin prep kit, Qiagen, Hilden, Germany) and digested with BamH1 and Xho1 at 37°C to select plasmids containing inserts of the predicted size for DNA sequencing. DNA sequencing was carried out by the dideoxy-nucleotide chain-termination method in a Beckman Coulter CEQk 2000 XL DNA Analysis System. To confirm that the cDNA sequences encoded CTLs, the predicted amino acid sequences were subjected to BLAST searches of the GenBank, PDB, SwissProt, PIR, and PRF databases. All the cDNAs exhibited significant sequence homology to Serine proteinases of related vipers. The CLUSTALW program [43] with PAM 250 residue weight matrix was used to align deduced amino acid sequences representing each *E. ocellatus* Serineproteinases isoforms with analogues in venoms from related Viperidae species as illustrated in Table 1. Serine proteinase (CAB62591) from *V. lebetina* [44], Serine protease 1 (AAR24534) from *B. gabonica* [45], Thrombin-like enzyme pre. (AAK12273) from *D. acutus* [46], Venom serine protease 5 (AAN52350) from *T. stejnegeri* [47], Serine proteinase 3 pre. (O13063) from *gramineus* [48], Serine proteinase A Precursor (Q9PTU8) from *B. jararaca* [46, 49], Serine proteinase 2A pre. (O13060) from *T. gramineus* [45, 48], Serine protease (AAP42416) from *B. jararacussu* [50], KN-BJ2 (BAA20283) from *B. jararaca* [51], Serine proteinase 1 pre. (AAG10788) from *T. jerdonii* [52], Thrombin-like serine protease (AAL68708) from *G. ussuriensis* [53], and, finally, Serine protease catroxtase I pre. (AAL77226) from *C. atrox* [54]. The phylogenetic trees constructed from the above alignments were generated by a neighbour-joining [55] algorithm in Lasergene software (DNASTAR, USA). The predicted antigenic profile [56] of the published and new *Echis ocellatus* serine protease (*EoSer*) isoforms analysed here was determined using Protean Software (DNASTAR).

3. Results

3.1. Isolation of cDNAs Encoding *E. Ocellatus* Serine Protease. PCR screening of the *Echis ocellatus* venom gland cDNA libraries resulted in a total of 14 *E. ocellatus* (*Eo*) cDNAs whose sequences matched (BLAST searches) those of published Serine proteases. The cDNAs consisted of 822 nucleotides (Figures 1 and 2(a)) and were predicted to encode an open reading frame proteins of 264 amino acids (28.5 kDa) (Figure 2(b)). Alignment of the predicted amino acid sequences of the 14 specific cDNAs encoding the *EoSP* proteins (Figure 2(b)) revealed sequence variations. The sequence similarity between the *EoSP* variants proteins was less than 60% for the mature protein-coding region but over 90% for regions coding both the signal peptide and the carboxyl-terminal end. Where two or more identical sequences were obtained from any one of these libraries, a

FIGURE 1: PCR product of the *E. ocellatus* serine proteases. Analysis of PCR amplification products by 0.7% agarose gel electrophoresis. Bands were visualised using the ultraviolet transillumination. Lane 2: represents the amplified PCR product (circled) of about 800 bp from *E. ocellatus* venom glands cDNA compared with Lane 1: 1 kb ladder DNA-marker bands, of known molecular weight. Lanes 3 and 4 represent a H₂O negative control and a SOD positive control, respectively.

single representative cDNA was used for subsequent analysis. Structural properties analysis (Emin algorithm-DNASTAR, USA) (Figure 3) was used to categorise the 14 Serine protease sequences into four distinct groups, based solely on sequence alignment.

3.2. BLAST Search of the Predicted Amino Acid Sequence. Accession numbers assigned to the new *Echis ocellatus* Serine protease sequences are as follows: “group 1” *EoSer*-1 (GU562413), “group 2” *EoSer*-3 (GU592440), “group 3” *EoSer*-17 (GU592441), and “group 4” *EoSer*-7 (GU592439). The predicted amino acid sequences of the *EoSP*-01, *EoSP*-03, *EoSP*-07, and *EoSP*-17 were submitted to BLAST searches of the genetic data bases and their similarity to published viper serine protease (Table 1) confirmed that the *EoSP* cDNAs encoded serine proteases.

3.3. Comparison of *E. Ocellatus* cDNAs with Analogous Serine Proteases from Other Viper Species. All the *EoSer*-variants contained the serine protease-consensus 24 amino acid signal peptide sequence (Figure 4, arrows), including the six-amino acids-activated motif. The signal peptide residues were followed by a protease domain of 236 residues. The deduced primary structures of all *EoSP* cDNA clones include the requisite, highly conserved, 12 cysteine residues that form the 6-disulphide bonds responsible for the characteristic tertiary structure of venom serine proteases. The complete amino acid sequences of the *EoSP* variants were aligned with those of other venom serine proteases (Figure 4). Viper

(a)

FIGURE 2: Continued.

FIGURE 2: (a) The nucleotide sequence of the fourteen *E. ocellatus* venom gland cDNAs resulting from PCR amplification. (b) Deduced amino acid sequences of *E. ocellatus* venom gland serine protease cDNAs.

venom SP sequences in the genetic databases were compared with the *E. o* groups (Table 2 and Figure 4) by BLAST. Groups 1–4 represent novel, highly similar, SP isomers with less than 65% sequence similarity to analogues in related viper species. Group 4 showed the greatest sequence similarity (80% and 82%) to the Serine protease of the African *V. lebetina* and *B. gabonica* vipers, respectively. Of all the *EoSP* clusters seemed to represent a SP sequence which showed the highest sequence similarity range between 62% to 70% to the SP of the vipers. None of the clusters showed more than 72% sequence similarity to the partial peptide sequences for the Thrombin-like serine protease isolated from the venom of

the *G. ussuriensis* viper [52]. Similarly, the Serine protease catroaxase I pre. of *C. atrox* venom showed no greater than 65% sequence similarity to any of the *EoSP* sequences.

3.4. Predicted Antigenic Profile Analysis of *E. ocellatus* Serine Proteases with Analogous Molecules. Since the main focus of our research is to develop toxin neutralising antibodies by immunisation with DNA encoding specific toxins in venoms of the most medically important African vipers [15, 59, 60], we next compared the algorithm-predicted immunogenicity of the *E. ocellatus* serine protease cluster cDNA sequences with those of all the published SPs from vipers of African

FIGURE 3: Differentiation of the fourteen cDNA-encoding *E. ocellatus* venom gland serine proteases. The predicted surface probabilities (Emin algorithm, DNASTAR, USA) of the 14 *E. ocellatus* serine protease cDNAs were aligned. The boxed areas indicate group specific structural motifs.

TABLE 1: Percent sequence similarity between *E. ocellatus* serine proteases and analogous molecules from related viper species.

Species	Accession no.	References	Serine protease	<i>EoSP</i> -1	<i>EoSP</i> -3	<i>EoSP</i> -7	<i>EoSP</i> -17
<i>V. lebetina</i>	CAB62591	Siigur et al. [44]	Serine proteinase	65	66	80	64
<i>B. gabonica</i>	AAR24534	Francischetti et al. [45]	Serine protease 1	62	63	82	65
<i>D. acutus</i>	AAK12273	Liang et al. [46]	Thrombin-like enzyme pre.	67	67	71	70
<i>T. stejnegeri</i>	AAN52350	Lee and Zhang [47]	Venom serine protease 5	66	66	65	69
<i>T. gramineus</i>	O13063	Deshimaru et al. [48]	Serine proteinase 3 pre.	71	71	61	76
<i>B. jararaca</i>	Q9PTU8	Murayama, [49]	Serine proteinase A pre.	66	68	66	74
<i>T. gramineus</i>	O13060	Deshimaru et al. [48]	Serine proteinase 2A pre.	65	65	73	70
<i>B. jararacussu</i>	AAP42416	Kashima et al. [50]	Serine protease	63	63	72	68
<i>B. jararaca</i>	BAA20283	Serrano et al. [51]	KN-BJ2	69	68	62	69
<i>T. jerdonii</i>	AAG10788	Lu et al. [52]	Serine proteinase 1 pre.	65	65	71	70
<i>G. ussuriensis</i>	AAL68708	Zhao et al. [53]	Thrombin-like serine protease	67	67	72	71
<i>C. atrox</i>	AAL77226	Tsai et al. [54]	Serine protease catroxase I pre.	66	65	63	65

FIGURE 4: Amino acid sequence similarity between *EoSP* Variants and serine proteases from related vipers. The residues shaded in black correspond to residues that are identical to *EoSP-01*. The asteriks [*] represented the twelve conserved cysteine residues. The catalytic traid **His/Arg** (67), **Asp** (110) and **Ser** (208) are represented in red circles. Activated peptide where the mature protein cleaved is represented by green rectangle.

origin (Figure 5). The predicted antigenic profiles of the published and new *E. ocellatus* serine proteases were analysed as shown in Figure 5 using Protean Software (DNASTAR, USA) [53]. The deduced signal peptide domains of the *EoSP* variants are separated by a vertical dotted line, as these would normally be cleaved from the native proteins during posttranslational. The thin vertical boxes depict the residues comprising the catalytic traid, H/R/N, D/G/N, and S/P/N/T (67, 110, and 208), that show the greatest immunogenic

domains conservation common to all the new and published African viper venom SPs sequences as demonstrated in Figure 5.

4. Discussion

Serine proteases are a major component of viper venoms and are thought to disrupt several distinct elements of the blood coagulation system of envenomed victims.

FIGURE 5: Comparison of antigenic profile of the *EoSer* variants with analogous serine proteases used in Figure 4. The top horizontal scale represents the number of amino acid residues. The conserved signal peptide is separated from the mature protein by a vertical dotted line. The three vertical boxes were drawn to indicate the conserved catalytic traid regions described in the text.

TABLE 2: Comparison of amino acid motifs which are responsible for the potent effects and characterisation of some published venom serine proteases with the four *EoSP* cDNAs.

Amino acid	TSV-PA	Batroxobin	Ancrod	<i>EoSP</i> -1	<i>EoSP</i> -17	<i>EoSP</i> -3	<i>EoSP</i> -7	References
H/R	H ⁵⁷	H ⁵⁷	H ⁵⁷	H ⁶⁷	H ⁶⁷	H ⁶⁷	R ⁶⁷	Braud et al. [57]
D	D ¹⁰²	D ¹⁰²	D ¹⁰²	D ¹¹²	D ¹¹²	D ¹¹²	D ¹¹²	
S	S ¹⁹⁵	S ¹⁹⁵	S ¹⁹⁵	S ²⁰⁸	S ²⁰⁸	S ²⁰⁸	T ²⁰⁸	
H	H ¹⁹²	G ¹⁹²	N ¹⁹²	K ²⁰⁵	L ²⁰⁵	K ²⁰⁵	K ²⁰⁵	
F	F ¹⁹³	G ¹⁹³	S ¹⁹³	G ²⁰⁶	G ²⁰⁶	G ²⁰⁶	A ²⁰⁶	Guinto et al. [58]
D	D ¹⁸⁹	D ¹⁸⁹	D ¹⁸⁹	G ²⁰²	D ²⁰²	G ²⁰²	D ²⁰²	
P	p ²²⁵	p ²²⁵	p ²²⁵	p ²³⁵	p ²³⁵	p ²³⁵	p ²³⁵	
P	p ²¹⁹	p ²¹⁹	p ²¹⁹	p ²²⁸	V ²²⁸	p ²²⁸	p ²²⁸	
D	D ⁹⁶	N ⁹⁶	R ⁹⁶	Y ¹⁰⁶	Y ¹⁰⁶	Y ¹⁰⁶	Y ¹⁰⁶	Lee and Zhang [47]
D	D ⁹⁷	V ⁹⁷	T ⁹⁷	T ¹⁰⁷	T ¹⁰⁷	T ¹⁰⁷	T ¹⁰⁷	
E	E ⁹⁸	I ⁹⁸	S ⁹⁸	L ¹⁰⁸	L ¹⁰⁸	K ¹⁰⁸	R ¹⁰⁸	

HDS: Catalytic Traid; H/F: substrate specificity; D & P: Architecture of water channel; P: Evolutionary region to kallikrein; DDE: substrate specificity to plasminogen.

A detailed understanding of the functions of these enzymes is important for both acquiring a full understanding of the pathology of envenoming and because these venom proteins have shown a vital role in treating blood coagulation disorders.

In general, serine proteinases including fibrinogenolytic enzymes are very abundant in *Viperidae* venoms in which they may account for 20% of their total protein content [61]. The unique specificity of snake venom proteinases makes them potentially useful in research of fibrinogen-depletion and limited proteolysis [62, 63]. This may be due to the existence of multiple forms of serine proteases in the venom of a single viper species which is likely to contribute to the diverse biological effects exerted by the whole venom. Therefore, screening the *E. ocellatus* cDNA library to isolate different isoforms or variants of serine proteases was the aim of this research work. The results obtained in this work provide the first molecular sequence data for *E. ocellatus* serine proteases they also reveal that the serine protease composition of *E. ocellatus* is as complex as that of the better characterised *Viperidae* species. The utilization of PCR amplification of *E. ocellatus* venom gland cDNA with the new viper serine protease-specific primers was successful and produced fourteen cDNAs sequences that were identified (BLAST) as belonging to the serine protease enzyme family. All *EoSP* cDNAs were of similar total length (approximately 0.80 kb, Figure 1) and encoded 260 amino acids (Figure 2(b)) with a predicted molecular weight of 28.5 kDa. To differentiate between the isolated *EoSP* clones a surface probability algorithm was used to assign the 14 *E. ocellatus* serine protease cDNAs into four main groups (Figure 3). A single representative clone from each group was chosen for further analyses as described earlier. The sequence similarity between the *EoSP* variants proteins was less than 60% for the mature protein-coding region but over 90% for regions coding both the signal peptide and the carboxyl-terminal end. Thus the latter two regions are highly conserved, which explains why the PCR experiment to amplify the cDNAs-encoding *EoSP* clones was successful.

The *EoSP* cDNA sequences were confirmed by BLAST searches as encoding serine proteases (Figure 4). The greatest sequence similarity was between *EoSer-7* and *B. gabonica* and *V. labetina* (80% and 85%) with the remaining *EoSP* cDNAs showing 60%–76% sequence similarity with other snake venom serine proteinases as illustrated in Table 1. From the proteins with known biological activity, sequence similarities of the *EoSP* variants (i.e., *EoSer-01*, *EoSer-03*, *EoSer-07* and *EoSer-17*) were 62%–69% with the kinin-releasing and fibrinogen-clotting serine protease (KN-BJ) from venom of *B. jararaca* [51] (Table 1). The putative 18 amino acid signal-peptide of the *EoSP* variants was as conserved (over 90% sequence similarity) as that in the serine proteases of other viper species (Figure 4, arrows). Following the signal peptide all the *EoSP* variants contained the predicted six-amino acid cleavage (activation) site **Q-K/T/M/E-S-S-E-L/P** (Figure 4 in green) as proposed for batroxobin [64]; thus cleavage generates a hydrophilic zymogen peptide, based on the processing site of pre-peptides of mammalian serine proteinases [65–67]. Comparison of the *EoSP* variants with

analogous members of the serine protease family revealed that all *EoSP* variants encoded the presumed catalytic triad, which is common to venom serine proteases H67, D110 and S208 as shown in Figure 4. Such residues were highly conserved in groups 1–3, except proteins of group 4 (Figures 2(b) and 4) which contain R instead of H at the same position (Figure 4). Furthermore, comparison of the *EoSP* amino acid sequence alignment with analogous venom serine proteases (Figure 4) revealed a conserved consensus active site of L-T/S-A-A-H/R/N-C corresponding to position 63–68, as previously determined [68]. Most SVSPs are likely to be glycoproteins showing a variable number of N- or O-glycosylation sites in sequence positions that differ from one SVSP to the other [69]. Using the primary structure of *EoSP* variants (Figure 4) the putative N-linked glycosylation sites, Asn-X-Thr/Ser [45], were found and are located at two different positions. *EoSer-01*, *EoSer-03*, and *EoSer-17* [N⁴⁴-X⁴⁵-S⁴⁶ and N²⁵⁷-X²⁵⁸-T²⁵⁹] and *EoSer-07* [N¹²⁴-R¹²⁵-T¹²⁶ and N²⁵⁷-T²⁵⁸-T²⁵⁹]. Although such motifs are thought to be needed for protein stabilization rather than for the catalytic function of the venom enzymes [30], confirmation of the roles of such motifs in venom proteases remain to be investigated. All serine proteases have a common pattern of 6-disulfide bridges [69, 70]. They contain twelve cysteine residues, ten of which form five disulfide bonds, based on the homology with trypsin [64]; the remaining two cysteines form a unique and conserved bridge among SVSPs, involving Cys245e (chymotrypsinogen numbering), found in the C-terminal extension [35].

From the results obtained this was found in all *EoSP* clones (Figure 2(b)) that encoded the common 12 cysteine residues in which are strongly conserved forming putative disulphide bridges which are located at Cys³¹, Cys³², C⁶⁸, C¹⁰⁰, C¹⁴⁵, C¹⁶⁵, C¹⁷⁶, C²⁰⁴, C²¹⁴, C²²⁹, and C²⁶⁰ (Figure 4). This suggests that the *EoSP* proteins possess a similar tertiary structure to that of other serine proteases which are well characterized.

Despite such sequence and structural conservation, viper venom serine proteases show very divergent effects on haemostasis as previously stated. In some cases certain amino acid sequences have been shown to be responsible for such effects as demonstrated in Table 2. Although such table gives a preliminary prediction of the functional characterization of the *EoSP* cDNAs in comparison with well-known characterized venom serine proteases, it cannot be considered as a functional confirmation or even a categorization strategy to differentiate between the four *EoSP* cDNAs. However, from Figure 4 and Table 2 it can be generally concluded that such comparison demonstrates that the enzymes encoded by the four *EoSP* cDNAs confer multiple haemostasis-disruptive activities to *E. ocellatus* venom. Furthermore, the sequence and predicted structural similarities of these four *EoSP* groups suggest that an antibody generated to one group may be capable of neutralizing the other group of *EoSP*s. To examine this permeability the sequences of *EoSP* groups were subjected to a more specific algorithm that predicted amino acid motifs of high immunogenicity. A protein structure-predicting algorithm [56] has been used (i) to identify domains of strong antigenic potential in the

toxin gene product and (ii) to determine whether these domains are conserved in analogous venom toxin gene products of related vipers. The signal peptide was separated from the mature protein by dotted line as would be cleaved posttranslationally. The peaks shown by the *EoSPs* profile indicate the numerous domains predicted to have a surface location and potential for antibody induction. Although the antigenic peaks of the catalytic traid of the *EoSPs* showed less similarity with that of the analogous venom SPs particularly those at residues 67 and 110, many antigenic residue similarities of *EoSPs* are shared with other SVSPs of related vipers. Therefore, it is likely that antibodies raised by *EoSP* DNA immunisation are likely to possess considerable cross-reactivity and might competitively inhibit the function of these domains in the similar venom toxins of related vipers. However, binding of antibodies specific to conserved antigenic domains without a known function are equally as likely to disrupt protein function by virtue of steric hindrance. The veracity of these speculations need to be confirmed experimentally and thus is a focus of our current research.

In conclusion, the predicted Jameson-Wolf antigenic profiles (DNASTAR, USA) of the *EoSP* variants aligned with very low identity to their (BLAST) analogous serine proteases. This observation strongly suggests that an antibody raised by immunisation with group one *EoSP* DNA is likely to be less effective against the gene products of groups 2, 3, or 4. Therefore additional antibodies generated against antigenic index that showed less conservation will be required.

Acknowledgments

Funding for this project was provided by the Wellcome Trust (RAH, Grant no. 061325), the University of Science and Technology, Yemen, and the Gunter Trust (S. Hasson). The authors would like to thank Dr. R. A. Harrison, Prof. R. D. G. Theakston, and Mr. Paul Rowley for their assistance during extraction of the venom glands from snakes and Dr. A. Nasidi, Federal Ministry of Health, Nigeria, for obtaining the snakes.

References

- [1] J. M. Gutiérrez, R. D. G. Theakston, and D. A. Warrell, "Confronting the neglected problem of snake bite envenoming: the need for a global partnership," *PLoS Medicine*, vol. 3, no. 6, article e150, 2006.
- [2] WHO, *Rabies and Envenomings: A Neglected Public Health Issue*, Report of a consultative meeting, WHO, Geneva, Switzerland, 2007.
- [3] J. F. Trape, G. Pison, E. Guyavarch, and Y. Mane, "High mortality from snakebite in south-eastern Senegal," *Transactions of the Royal Society of Tropical Medicine and Hygiene*, vol. 95, no. 4, pp. 420–423, 2001.
- [4] S. C. Wagstaff and R. A. Harrison, "Venom gland EST analysis of the saw-scaled viper, *Echis ocellatus*, reveals novel $\alpha\beta 1$ integrin-binding motifs in venom metalloproteinases and a new group of putative toxins, renin-like aspartic proteases," *Gene*, vol. 377, no. 1-2, pp. 21–32, 2006.
- [5] N. R. Casewell, R. A. Harrison, W. Wüster, and S. C. Wagstaff, "Comparative venom gland transcriptome surveys of the saw-scaled vipers (*Viperidae: Echis*) reveal substantial intra-family gene diversity and novel venom transcripts," *BMC Genomics*, vol. 10, article 564, 2009.
- [6] S. S. Hasson, A. A. Al-Jabri, T. A. Sallam, M. S. Al-Balushi, and R. A. A. Mothana, "Antisnake venom activity of *Hibiscus aethiopicus* L. against *Echis ocellatus* and *Naja n. nigricollis*," *Journal of Toxicology*, vol. 2010, Article ID 837864, 8 pages, 2010.
- [7] J. P. Chippaux, "The treatment of snake bites: analysis of requirements and assessment of therapeutic efficacy in tropical Africa," in *Perspectives in Molecular Toxicology*, A. Menez, Ed., pp. 457–472, Wiley, New York, NY, USA, 2002.
- [8] P. Revault, "Ecology of *Echis ocellatus* and peri-urban bites in Ouagadougou," *Toxicon*, vol. 34, no. 2, p. 144, 1996.
- [9] A. G. Habib, S. B. Abubakar, I. S. Abubakar et al., "Envenoming after carpet viper (*Echis ocellatus*) bite during pregnancy: timely use of effective antivenom improves maternal and foetal outcomes," *Tropical Medicine and International Health*, vol. 13, no. 9, pp. 1172–1175, 2008.
- [10] D. A. Warrell, N. McD. Davidson, and B. M. Greenwood, "Poisoning by bites of the saw scaled or carpet viper (*Echis carinatus*) in Nigeria," *Quarterly Journal of Medicine*, vol. 46, no. 181, pp. 33–62, 1977.
- [11] M. L. D. Weinberg, L. F. Felicori, C. A. Bello et al., "Biochemical properties of a bushmaster snake venom serine proteinase (LV-Ka), and its kinin releasing activity evaluated in rat mesenteric arterial rings," *Journal of Pharmacological Sciences*, vol. 96, no. 3, pp. 333–342, 2004.
- [12] A. B. Sallau and M. A. Ibrahim, "Characterization of phospholipase A2 (PLA2) from *Echis ocellatus* venom," *African Journal of Biochemistry Research*, vol. 2, no. 4, pp. 98–101, 2008.
- [13] J.-M. Howes, R. D. G. Theakston, and G. D. Laing, "Antigenic relationships and relative immunogenicities of isolated metalloproteinases from *Echis ocellatus* venom," *Toxicon*, vol. 45, no. 5, pp. 677–680, 2005.
- [14] S. Liu, M.-Z. Sun, C. Sun, B. Zhao, F. T. Greenaway, and Q. Zheng, "A novel serine protease from the snake venom of *Agkistrodon blomhoffii ussuriensis*," *Toxicon*, vol. 52, no. 7, pp. 760–768, 2008.
- [15] K. Bharati, S. S. Hasson, J. Oliver, G. D. Laing, R. D. G. Theakston, and R. A. Harrison, "Molecular cloning of phospholipases A2 from venom glands of *Echis* carpet vipers," *Toxicon*, vol. 41, no. 8, pp. 941–947, 2003.
- [16] Q. Lu, A. Navdaev, J. M. Clemetson, and K. J. Clemetson, "Snake venom C-type lectins interacting with platelet receptors. Structure-function relationships and effects on haemostasis," *Toxicon*, vol. 45, no. 8, pp. 1089–1098, 2005.
- [17] K. Stocker, "Snake venom proteins affecting hemostasis and brinolysis," in *Medical Use of Snake Venom Proteins*, K. F. Stocker, Ed., pp. 97–160, CRC Press, Boca Raton, Fla, USA, 1990.
- [18] T. Matsui, Y. Fujimura, and K. Titani, "Snake venom proteases affecting hemostasis and thrombosis," *Biochimica et Biophysica Acta*, vol. 1477, no. 1-2, pp. 146–156, 2000.
- [19] F. S. Markland Jr., "Preface: snake venoms and hemostasis," *Toxin Reviews*, vol. 25, no. 4, pp. 319–321, 2006.
- [20] F. S. Markland Jr., "Inventory of α - and β -fibrinogenases from snake venoms. For the Subcommittee on Nomenclature of Exogenous Hemostatic Factors of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis," *Thrombosis and Haemostasis*, vol. 65, no. 4, pp. 438–443, 1991.

- [21] H. Pirkle, "Thrombin-like enzymes from snake venoms: an updated inventory," *Thrombosis and Haemostasis*, vol. 79, no. 3, pp. 675–683, 1998.
- [22] H. Pirkle and K. Stocker, "Thrombin-like enzymes from snake venoms: an inventory. For the Subcommittee on Nomenclature of Exogenous Hemostatic Factors of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis," *Thrombosis and Haemostasis*, vol. 65, no. 4, pp. 444–450, 1991.
- [23] H. Pirkle and I. Theodor, "Thrombin-like venom enzymes: structure and function," *Advances in Experimental Medicine and Biology*, vol. 281, pp. 165–176, 1991.
- [24] A. V. Pérez, A. Rucavado, L. Sanz, J. J. Calvete, and J. M. Gutiérrez, "Isolation and characterization of a serine proteinase with thrombin-like activity from the venom of the snake *Bothrops asper*," *Brazilian Journal of Medical and Biological Research*, vol. 41, no. 1, pp. 12–17, 2008.
- [25] H. C. Castro, R. B. Zingali, M. G. Albuquerque, M. Pujol-Luz, and C. R. Rodrigues, "Snake venom thrombin-like enzymes: from reptilase to now," *Cellular and Molecular Life Sciences*, vol. 61, no. 7-8, pp. 843–856, 2004.
- [26] I. Panfoli, D. Calzia, S. Ravera, and A. Morelli, "Inhibition of hemorrhagic snake venom components: old and new approaches," *Toxins*, vol. 2, pp. 417–427, 2010.
- [27] Y.-M. Wang, S.-R. Wang, and I.-H. Tsai, "Serine protease isoforms of *Deinagkistrodon acutus* venom: cloning, sequencing and phylogenetic analysis," *Biochemical Journal*, vol. 354, no. 1, pp. 161–168, 2001.
- [28] R. A. Hutton and D. A. Warrell, "Action of snake venom components on the haemostatic system," *Blood Reviews*, vol. 7, no. 3, pp. 176–189, 1993.
- [29] J. B. Bjarnason, A. Barish, and G. S. Direnzo, "Kallikrein-like enzymes from *Crotalus atrox* venom," *Journal of Biological Chemistry*, vol. 258, no. 20, pp. 12566–12573, 1983.
- [30] Y. Komori, T. Nikai, and H. Sugihara, "Comparison of the lethal components in *Vipera aspis aspis* and *Vipera aspis zinnikeri* venom," *Journal of Natural Toxins*, vol. 7, no. 2, pp. 101–108, 1998.
- [31] L. F. Felicori, C. T. Souza, D. T. Velarde et al., "Kallikrein-like proteinase from *bushmaster* snake venom," *Protein Expression and Purification*, vol. 30, no. 1, pp. 32–42, 2003.
- [32] T. Matsui, Y. Sakurai, Y. Fujimura et al., "Purification and amino acid sequence of halystase from snake venom of *Agkistrodon halys blomhoffii*, a serine protease that cleaves specifically fibrinogen and kininogen," *European Journal of Biochemistry*, vol. 252, no. 3, pp. 569–575, 1998.
- [33] S. Iwanaga, G. Oshima, and T. Suzuki, "Proteinases from the venom of *Agkistrodon halys blomhoffii*," *Methods in Enzymology*, vol. 45, pp. 459–468, 1976.
- [34] C.-C. Hung and S.-H. Chiou, "Fibrinogenolytic proteases isolated from the snake venom of Taiwan Habu: serine proteases with kallikrein-like and angiotensin-degrading activities," *Biochemical and Biophysical Research Communications*, vol. 281, no. 4, pp. 1012–1018, 2001.
- [35] M. A. A. Parry, U. Jacob, R. Huber, A. Wisner, C. Bon, and W. Bode, "The crystal structure of the novel snake venom plasminogen activator TSV-PA: a prototype structure for snake venom serine proteinases," *Structure*, vol. 6, no. 9, pp. 1195–1206, 1998.
- [36] Y. Zhang, A. Wisner, Y. Xiong, and C. Bon, "A novel plasminogen activator from snake venom: purification, characterization, and molecular cloning," *Journal of Biological Chemistry*, vol. 270, no. 17, pp. 10246–10255, 1995.
- [37] Y. Zhang, A. Wisner, R. C. Maroun, V. Choumet, Y. Xiong, and C. Bon, "*Trimeresurus stejnegeri* snake venom plasminogen activator: site-directed mutagenesis and molecular modeling," *Journal of Biological Chemistry*, vol. 272, no. 33, pp. 20531–20537, 1997.
- [38] S. M. T. Serrano, R. Mentele, C. A. M. Sampaio, and E. Fink, "Purification, characterization, and amino acid sequence of a serine proteinase, PA-BJ, with platelet-aggregating activity from the venom of *Bothrops jararaca*," *Biochemistry*, vol. 34, no. 21, pp. 7186–7193, 1995.
- [39] F. Tokunaga, F. Nagasawa, S. Tamura, T. Miyata, S. Iwanaga, and W. Kisiel, "The factor V-activating enzyme (RVV-V) from Russell's viper venom. Identification of isoproteins RVV-V(α), -V β , and -V γ and their complete amino acid sequences," *Journal of Biological Chemistry*, vol. 263, no. 33, pp. 17471–17481, 1988.
- [40] J. J. Calvete, C. Marcinkiewicz, D. Monleón et al., "Snake venom disintegrins: evolution of structure and function," *Toxicon*, vol. 45, no. 8, pp. 1063–1074, 2005.
- [41] L. Sanz, A. Bazaa, N. Marrakchi et al., "Molecular cloning of disintegrins from *Cerastes vipera* and *Macrovipera lebetina transmediterranea* venom gland cDNA libraries: insight into the evolution of the snake venom integrin-inhibition system," *Biochemical Journal*, vol. 395, no. 2, pp. 385–392, 2006.
- [42] D. Israel, "A PCR-based method for high stringency screening of DNA libraries," *Nucleic Acids Research*, vol. 21, no. 11, pp. 2627–2631, 1993.
- [43] J. D. Thompson, D. G. Higgins, and T. J. Gibson, "CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice," *Nucleic Acids Research*, vol. 22, no. 22, pp. 4673–4680, 1994.
- [44] E. Siigur, A. Aaspõllu, and J. Siigur, "Sequence diversity of *Vipera lebetina* snake venom gland serine proteinase homologs—result of alternative-splicing or genome alteration," *Gene*, vol. 263, no. 1-2, pp. 199–203, 2001.
- [45] I. M. B. Francischetti, V. My-Pham, J. Harrison, M. K. Garfield, and J. M. C. Ribeiro, "*Bitis gabonica* (Gaboon viper) snake venom gland: toward a catalog for the full-length transcripts (cDNA) and proteins," *Gene*, vol. 337, supplement, pp. 55–69, 2004.
- [46] N. S. Liang, T. T. Liu, H. P. He, Y. A. Xie, Z. Q. Meng, and A. M. Liang, "Cloning and sequence analysis of the new cDNA of thrombin-like enzyme from *Deinagkistrodon acutus*," *Guangxi Yi Ke Da Xue Xue Bao*, vol. 19, no. 1, pp. 27–30, 2002.
- [47] W. H. Lee and Y. Zhang, "Molecular cloning and sequence comparison of serine proteases from the venom of *Trimeresurus stejnegeri*," Unpublished, Direct Submission, BLAST Search, 2002.
- [48] M. Deshimaru, T. Ogawa, K.-I. Nakashima et al., "Accelerated evolution of crotalinae snake venom gland serine proteases," *FEBS Letters*, vol. 397, no. 1, pp. 83–88, 1996.
- [49] N. Murayama, "Thrombin-like snake venom serine protease," Direct Submission, Unpublished, BLAST Search, 1999.
- [50] S. Kashima, P. G. Roberto, A. M. Soares et al., "Analysis of *Bothrops jararacussu* venomous gland transcriptome focusing on structural and functional aspects: I-gene expression profile of highly expressed phospholipases A₂," *Biochimie*, vol. 86, no. 3, pp. 211–219, 2004.
- [51] S. M. T. Serrano, Y. Hagiwara, N. Murayama et al., "Purification and characterization of a kinin-releasing and fibrinogen-clotting serine proteinase (KN-BJ) from the venom of *Bothrops jararaca*, and molecular cloning and sequence analysis of

- its cDNA," *European Journal of Biochemistry*, vol. 251, no. 3, pp. 845–853, 1998.
- [52] Q.-M. Lu, Y. Jin, D.-S. Li, W.-Y. Wang, and Y.-L. Xiong, "Characterization of a thrombin-like enzyme from the venom of *Trimeresurus jerdonii*," *Toxicon*, vol. 38, no. 9, pp. 1225–1236, 2000.
- [53] Y. Zhao, K. Fang, and K. Sun, "cDNA for thrombin-like serine protease from venom gland of *Agkistrodon ussuriensis*," Unpublished, Direct Submission, BLAST Search, 2001.
- [54] I. H. Tsai, J. C. Hsu, and Y. M. Wang, "Catroxtase I and II, the serine proteases of *Crotalus Atrox* venom: cloning, complete sequencing and functional characterization," Unpublished, Direct Submission, BLAST Search, 2002.
- [55] N. Saitou and M. Nei, "The neighbor-joining method: a new method for reconstructing phylogenetic trees," *Molecular Biology and Evolution*, vol. 4, no. 4, pp. 406–425, 1987.
- [56] B. A. Jameson and H. Wolf, "The antigenic index: a novel algorithm for predicting antigenic determinants," *Computer Applications in the Biosciences*, vol. 4, no. 1, pp. 181–186, 1988.
- [57] S. Braud, M. A. Parry, R. Maroun, C. Bon, and A. Wisner, "The contribution of residues 192 and 193 to the specificity of snake venom serine proteinases," *The Journal of Biological Chemistry*, vol. 275, pp. 1823–1828, 2000.
- [58] E. R. Cuinto, S. Caccia, T. Rose, K. Futterer, G. Waksman, and E. Di Cera, "Unexpected crucial rule of residue 225 in serine proteases," *Proceedings of the National Academy of Sciences*, vol. 96, pp. 1852–1857, 1999.
- [59] S. S. Hasson, R. D. G. Theakston, and R. A. Harrison, "Cloning of a prothrombin activator-like metalloproteinase from the West African saw-scaled viper, *Echis ocellatus*," *Toxicon*, vol. 42, no. 6, pp. 629–634, 2003.
- [60] R. A. Harrison, J. Oliver, S. S. Hasson, K. Bharati, and R. D. G. Theakston, "Novel sequences encoding venom C-type lectins are conserved in phylogenetically and geographically distinct *Echis* and *Bitis* viper species," *Gene*, vol. 315, no. 1-2, pp. 95–102, 2003.
- [61] A. Wisner, S. Braud, and C. Bon, "Snake venom proteinases as tools in hemostasis studies: structure-function relationship of a plasminogen activator purified from *Trimeresurus stejnegeri* venom," *Haemostasis*, vol. 31, no. 3–6, pp. 133–140, 2001.
- [62] D. C. I. Koh, A. Armugam, and K. Jeyaseelan, "Snake venom components and their applications in biomedicine," *Cellular and Molecular Life Sciences*, vol. 63, no. 24, pp. 3030–3041, 2006.
- [63] E. E. Gardiner and R. K. Andrews, "The cut of the clot(h): snake venom fibrinogenases as therapeutic agents," *Journal of Thrombosis and Haemostasis*, vol. 6, no. 8, pp. 1360–1362, 2008.
- [64] N. Itoh, N. Tanaka, S. Mihashi, and I. Yamashina, "Molecular cloning and sequence analysis of cDNA for batroxobin, a thrombin-like snake venom enzyme," *Journal of Biological Chemistry*, vol. 262, no. 7, pp. 3132–3135, 1987.
- [65] G. H. Swift, J. C. Dagorn, P. L. Ashley, S. W. Cummings, and R. J. MacDonald, "Rat pancreatic kallikrein mRNA: nucleotide sequence and amino acid sequence of the encoded preproenzyme," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 79, no. 23, pp. 7263–7267, 1982.
- [66] R. J. MacDonald, S. J. Stary, and G. H. Swift, "Two similar but nonallelic rat pancreatic trypsinogens. Nucleotide sequences of the cloned cDNAs," *Journal of Biological Chemistry*, vol. 257, no. 16, pp. 9724–9732, 1982.
- [67] R. J. MacDonald, G. H. Swift, C. Quinto et al., "Primary structure of two distinct rat pancreatic preproelastases determined by sequence analysis of the complete cloned messenger ribonucleic acid sequences," *Biochemistry*, vol. 21, no. 6, pp. 1453–1463, 1982.
- [68] S. Brenner, "The molecular evolution of genes and proteins: a tale of two serines," *Nature*, vol. 334, no. 6182, pp. 528–529, 1988.
- [69] S. M. T. Serrano and R. C. Maroun, "Snake venom serine proteinases: sequence homology vs. substrate specificity, a paradox to be solved," *Toxicon*, vol. 45, no. 8, pp. 1115–1132, 2005.
- [70] T. Nikai, A. Ohara, Y. Komori, J. W. Fox, and H. Sugihara, "Primary structure of a coagulant enzyme, bilineobin, from *Agkistrodon bilineatus* venom," *Archives of Biochemistry and Biophysics*, vol. 318, no. 1, pp. 89–96, 1995.