

Research Article

JAK3/STAT5/6 Pathway Alterations Are Associated with Immune Deviation in CD8⁺ T Cells in Renal Cell Carcinoma Patients

Elisabetta Cavalcanti,¹ Margherita Gigante,² Vito Mancini,¹ Michele Battaglia,¹ Pasquale Ditunno,¹ Carmela Capobianco,¹ Raffaele I. Cincione,² Francesco P. Selvaggi,¹ Wolfgang Herr,³ Walter J. Storkus,⁴ Loreto Gesualdo,² and Elena Ranieri²

¹ Department of Emergency and Organ Transplantation, University of Bari, Piazza G. Cesare 11, 70124 Bari, Italy

² Department of Biomedical Sciences and Bioagromed, University of Foggia, Via Napoli, 71100 Foggia, Italy

³ Hematology and Oncology, Johannes Gutenberg-University of Mainz, 55101 Mainz, Germany

⁴ Department of Dermatology, University of Pittsburgh School of Medicine, 200 Lothrop Street, Pittsburgh, PA 15213, USA

Correspondence should be addressed to Elena Ranieri, e.ranieri@unifg.it

Received 28 August 2009; Revised 16 December 2009; Accepted 12 January 2010

Academic Editor: Soldano Ferrone

Copyright © 2010 Elisabetta Cavalcanti et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

To investigate the molecular mechanisms underlying altered T cell response in renal cell carcinoma (RCC) patients, we compared autologous and allogeneic CD8⁺ T cell responses against RCC line from RCC patients and their HLA-matched donors, using mixed lymphocyte/tumor cell cultures (MLTCs). In addition, we analyzed the expression of molecules associated with cell cycle regulation. Autologous MLTC responder CD8⁺ T cells showed cytotoxic activity against RCC cell lines; however the analysis of the distribution of CD8⁺ T-cell subsets revealed that allogeneic counterparts mediate superior antitumor efficacy. In RCC patients, a decreased proliferative response to tumor, associated with defects in JAK3/STAT5/6 expression that led to increased p27KIP1 expression and alterations in the cell cycle, was observed. These data define a molecular pathway involved in cell cycle regulation that is associated with the dysfunction of tumor-specific CD8⁺ effector cells. If validated, this may define a therapeutic target in the setting of patients with RCC.

1. Introduction

Cancer is associated with immune deficiency but the biological mechanisms underlying immune failure remain poorly-defined. T cells are key elements in effective cancer immunity in RCC, a disease that has proven refractory to conventional treatment modalities, such as chemotherapy and radiotherapy [1]. On the contrary, RCC patients have been reported to be responsive to immunotherapeutic approaches establishing the concept that RCC is an immunogenic tumor [2, 3]. Given the perceived importance of CD8⁺ T cells in mediating antitumor immunity, and that Th1-skewed CD4⁺ Th cells are required to support durable CD8⁺ T-cell immunity [4, 5], it becomes particularly important to understand how dysfunctional or partially functional CD8⁺ T cells arise and what molecular mechanisms underlie their activation in patients with cancer. There is considerable interest in

gaining a better understanding of the composition of the antitumor CD8⁺ T cell population, with regard to its various functional subsets. Sallusto et al. [6] used CCR7 expression status to define T cell memory subsets. Central memory CD8⁺ T cells (T_{CM}) express CCR7, a chemokine receptor required for trafficking to secondary lymphoid organs. In contrast, this marker is significantly downregulated on effector memory CD8⁺ T cells (T_{EM}). T_{CM} have been shown to mediate superior anti-tumor activity compared with T_{EM} [7, 8]. There is a third T cell memory subset, terminally differentiated CD8⁺ T_{EM} cells (T_{EMRA}), that express CD45RA but lack CCR7 expression. Notably, T_{CM} have a higher proliferative potential and greater resistance to apoptosis, whereas T_{EM}/T_{EMRA} have a skewed TCR repertoire and are characterized by a “senescent” replication history [9, 10]. Furthermore, the results of gene expression analyses have revealed that T_{CM} are characterized by high basal and

cytokine-induced STAT5/6 phosphorylation, reflecting their capacity for enhanced self-renewal versus effector memory cells, in particular, T_{EMRA} [11].

The JAK3/STAT5/6 pathway is a crucial signal transduction component for many growth factor receptors and it has been shown to be necessary for the downregulation of several genes that inhibit the cell cycle. This pathway regulates the expression of cyclin-dependent kinase (Cdk) inhibitors p27, the hyperphosphorylation of the retinoblastoma protein (Rb) gene product, and the release of active E2F in T cell lines [12]. In vivo studies have provided evidence of the role of p27^{KIP1}, a cell cycle inhibitory protein, in T cell proliferation [13, 14]. High p27^{KIP1} levels were correlated with a decreased proliferative response to IL-4 and IL-12 in STAT6 deficient T cells [15, 16]. Therefore, the characterization of cell cycle regulatory proteins and the delineation of aberrations in CD8⁺ T cell function are important to understand in the context of tumor development and progression.

Inhibition of T-cell proliferation is an important aspect of immunosuppression as observed in cancer patients, and it represents a mechanism adopted by tumors to evade immune rejection [17]. Various studies have led to the identification of a pathway controlling the progression of cells from quiescence, through G1, and into S phase involving the activation of G1 cyclin-dependent kinases (cdk), inactivation of Rb and related proteins [18]. The transcription factors of the E2F family and Id (inhibitor of DNA binding) protein family play a central role in the regulation of cell growth [19]. In particular, E2F4 and Id2 are essential for Rb function [20] and are involved in independent functions that are orchestrated in order to allow for cell cycle progression. Id2 and E2F4 may compete for Rb binding; hence, Id2 may function via the release of restraining pocket proteins on E2F transcription. Deregulation of E2Fs transcriptional control contributes to oncogenic transformation in vitro and the development of metastasis in vivo [21].

In this paper, we have investigated RCC-specific CD8⁺ T cell response in RCC patients and HLA-matched normal donors that were generated in mixed lymphocyte/tumor cell cultures (MLTCs) in order to define molecular pathways associated with immune deviation in CD8⁺ T cell compartment in RCC patients. However, the generation of autologous RCC-reactive T cells requires the availability of established RCC cell lines, that can only be obtained from a minority of patients. Thus, we established tumor cell lines from the primary RCC tissue of patients undergoing nephrectomy. In such patients as well as HLA-matched normal donors, we studied the phenotypic and functional profile of autologous and allogeneic RCC-specific CD8⁺ T cells. In particular, we focused on the integrity of the JAK3/STAT5/6 signaling pathway and its relationship with cell cycle regulatory proteins, such as E2F4 and Id2, in order to better understand relevant mechanisms of dysfunction associated with CD8⁺ T cells isolated from RCC patients. T cells from RCC patients displayed decreased proliferative responses to tumor antigens, which was linked to defects in JAK3/STAT5/6 expression (and to increased p27^{KIP1} expression and altered cell cycle progression/proliferation). Interestingly, we observed a significant correlation between

RCC CD8⁺ T subset phenotypes and such signaling defects, providing a surrogate index for clinical immune deviation that might be employed in the monitoring of patient response to immunotherapy.

2. Material and Methods

2.1. Cell Lines. Cell lines were generated from primary kidney tissue explants of 13 patients with RCC. Freshly isolated tumor tissue was minced, then digested using an enzymatic cocktail as described previously [22]. Corresponding normal renal cell lines (proximal tubular epithelial cells, PTEC) were established from tumor-distal normal kidney tissue and cultured in complete medium. Epstein-Barr virus- (EBV-) transformed lymphoblastoid cell lines (LCL) were also generated from patient RCC1 PBMC (RCC1-LCL) and healthy donor-2 (donor-2-LCL) using the B95.8 (type 1) virus isolate. Allogeneic phytohemagglutinin- (PHA-) activated T cell blasts (PHA-B) from donor-2 and donor-3 (donor-2-PHA-B, donor-3-PHA-B) were prepared by stimulating PBMC in media containing 2 µg/mL PHA (Sigma) and IL-2 (250 IU/mL, Proleukin, Chiron, Emeryville, CA).

2.2. Isolation of PBMC and HLA Typing. Peripheral blood mononuclear cells (PBMCs) were obtained from RCC patients and HLA-matched healthy donors after written consent, under an IRB-approved protocol. All patients had histological confirmed RCC, clear cell subtype, and had not received preoperative therapy. To determine the HLA-class I genotype of donor PBMCs and RCC lines, individual alleles were amplified from genomic DNA with A/B/C allele-specific primers and were then completely sequenced (Table 1, Dr. B. Favoino, Laboratory of typing tissue).

PBMCs were isolated at the interface of Ficoll-Hypaque density gradients centrifugation (Sigma Chemical Co., St Louis, MO), per the manufacturer's instructions, washed twice in PBS 3 (Invitrogen-Life Technologies, Italy) and used in mixed lymphocyte/tumor cell cultures as described below.

2.3. Isolation of CD8⁺ T-Effector Cells. CD8⁺ T cells were isolated from the PBMC of autologous and allogeneic normal, healthy donors by positive selection using specific MACS magnetic beads (Miltenyi Biotec, Bergisch Gladbach, Germany) on MiniMACS columns, per the manufacturer's protocol and subsequently assessed for their functional and phenotypic profiles.

2.4. Mixed Lymphocyte/Tumor Cell Cultures (MLTCs). PBMCs from RCC patients or HLA-matched healthy donors were cocultured in 24-well plates (Costar, Corning, USA) at 10⁶ cells/well with irradiated RCC stimulator cells (10⁵ cells/well) in 2 mL of AIM-V medium (Life Technologies, Invitrogen, Italy) supplemented with 10% heat-inactivated pooled human serum (Sigma (medium M^b)). Recombinant human IL-2 was added on day 3 at a final concentration of 250 IU/mL (Proleukin, Chiron, and Emeryville, CA). Responder lymphocytes (10⁶ cells/well) were restimulated weekly with 10⁵ irradiated tumor cells in

TABLE 1: HLA-class I genotyping of RCC patients and healthy donors.

| | HLA-A | | HLA-B | | HLA-Cw | |
|----------|-------|------|-------|------|--------|------|
| RCC1 | 0301 | 2402 | 0702 | 0801 | 0702 | 0702 |
| Donor 1a | 0301 | 2402 | 0702 | 0801 | 0702 | 0702 |
| Donor 1b | 0301 | 2402 | 0702 | 0801 | 0702 | 0702 |
| RCC2 | 0301 | 2402 | 1806 | 3506 | 0401 | 0702 |
| Donor 2 | 0301 | 2402 | 1806 | 3506 | 0401 | 0702 |
| RCC3 | 2402 | 3302 | 3506 | 6506 | 0401 | 0702 |
| Donor 3 | 2402 | 3302 | 3506 | 6506 | 0401 | 0702 |
| RCC4 | 0301 | 2402 | 1806 | 3506 | 0401 | 0702 |
| Donor 4 | 0301 | 2402 | 1806 | 3506 | 0401 | 0702 |
| RCC5 | 0201 | 28 | 3506 | 5104 | 0401 | 0702 |
| Donor 5 | 0201 | 28 | 3506 | 5104 | 0401 | 0702 |

IL-2-containing medium M^b for 2 weeks before CD8⁺ T cells isolation. CD8⁺ T cell isolation was performed using immunomagnetic CD8⁺ beads (Miltenyi Biotec, Milan, Italy) and positively isolated T cells cultured for an additional 2 weeks. On day 35 of culture, CD8⁺ T cells responders were used as effector cells in functional and molecular analyses.

2.5. IFN- γ ELISPOT Assay. CD8⁺ T cell responders were assessed for their ability to secrete IFN- γ in response to specific stimulation using ELISPOT assays (Mabtech, Mariemont, OH), as previously described [23]. Determinations were performed in triplicate and spots were enumerated using an automatic plate reader (Zeiss-Kontron, Jena, Germany). In some cases, the HLA-restriction of T cell recognition was determined by addition of blocking antibodies (final concentration = 100 μ g/mL) to replicate ELISPOT wells. Antibodies included; W6/32, an anti-HLA class I, α -gap A3, an anti-HLA-A3, B1.23.2, an anti-HLA-B and HLA-CA24, an anti-HLA-A24, L243, an anti-HLA class II-DR. IgG2a isotype-matched control Abs were used as negative controls.

2.6. ⁵¹Chromium-Release Assay. Responder CD8⁺ T cells were evaluated for their ability to kill target cells including patient-derived RCC cell lines, PHA-blast or EBV-cells and K562 in standard 4 h ⁵¹Cr-release assays [23].

2.7. Phenotypic Analysis. T cells were washed and resuspended in FACS buffer (phosphate-buffered saline pH 7.2, 0.2% bovine serum albumin, and 0.02% sodium azide) containing 3% human serum and incubated with fluorochrome-conjugated mAbs for 15 minutes at 4°C, then washed with the same buffer before the performance of 6-color flow cytometry. For the surface staining of T cells, fluorescently labeled mAbs were used: PE-CD3, APC-CD8, FITC-CD45RA, PerCP-CD28, (all BD-9Pharmingen), Pe-Cy-7-CCR7 (R&D Systems), and APC-Cy-7-CD27. Flow data were acquired using a FACSort™ (Becton Dickinson) flow cytometer and analyzed using WinMDI version 2.8 software.

2.8. Cell Cycle Analysis. T cells were fixed in 70% ethanol for 24 hours, prior to being treated with 50 mg of RNase A/mL at 37°C for 30 minutes. After resuspension in 0.5 mL of propidium iodide solution (50 μ g/mL propidium iodide, 0.1% sodium citrate, and 0.1% Nonidet P-40) and a 30-minute incubation in the dark at 4°C, the cellular DNA content was analyzed by flow cytometry using WinMDI version 2.8 software.

2.8.1. Confocal Analysis. The expression of STAT5, STAT6, p-27^{KIP1}, E2F4, and Id2 in T cells was analyzed by confocal fluorescence microscopy using specific Abs (Santa Cruz Biotechnology, DBA, Italy). CD8⁺ T cells were allowed to adhere to poly-L-lysine-coated slides (Sigma-Aldrich, Italy) for 1 hour on ice, followed by fixation in 3.7% paraformaldehyde. After washing three times with PBS (pH 7.4), and being blocked with BSA 1% and PBS, cells were stained with anti-pSTAT5 Ab, anti-pSTAT6 Ab, anti-E2F4 Ab, anti-p27KIP1 Ab and anti-ID2 Ab. After washing, T cells were then incubated with fluorescently labeled secondary Abs: Alexa Fluor FITC- and TRITC-conjugated anti-Ig (Molecular Probes, Eugene, OR). Stained cells were then washed four times with PBS, air-dried, and mounted using Gel/Mount (Biomedica, Foster City, CA). Samples were analyzed using a laser scanning Leica TCS SP2 confocal microscope (Leica, Wetzlar, Germany) equipped with argon krypton (488 nm), green-neon (543 nm), and helium-neon (633 nm) lasers. The expression levels of pSTAT5/6, p-27^{KIP1}, E2F4, and Id2 were measured in at least 15 high-power fields (HPFs; 63 \times)/section by two independent observers blinded to the origin of the slides. The final count was the mean of the two measures \pm standard deviation (SD). In no case the interobserver variability exceeds 20%.

2.8.2. Statistical Analysis. Data are reported as mean \pm standard deviation (SD) and analyzed for differences using an unpaired Student's *t*-test. Statview software package (SAS Institute Inc. Cary, NC, USA; 5, 0 version) was used for all analyses. *P* values < .05 were considered statistically significant.

3. Results and Discussion

3.1. Selective Expansion of Autologous and Allogeneic Antitumor CD8⁺ T Cells by MLTC. In our initial set of experiments, we attempted to establish primary RCC lines from 13 patients who underwent radical or partial nephrectomy for unilateral renal cell carcinoma. However, given limited quantities of available tissue, we were only able to develop 5 RCC lines which exhibited with the appropriate morphology, histopathology, and immunohistochemical characteristics of (clear-cell) RCC. Autologous and HLA-matched allogeneic PBMCs isolated from normal, healthy donors (Table 1) were used as responder cells to these tumor cells in MLTC established in the presence of high dose rhIL-2. After 3 weeks of culture, responder cells were harvested and MACS-purified CD8⁺ T cells were restimulated in an identical fashion (to the primary induction) for additional 2 weeks. On day 35 of culture, CD8⁺ T cells were evaluated for specific anti-RCC reactivity using IFN- γ ELISPOT assays and ⁵¹Cr-release assays.

In a first example of the results obtained, we examined autologous and allogeneic CD8⁺ T cell responses against the RCC1 cell line. Interestingly for this model, we employed 2 allogeneic responders: donor 1a (a healthy sibling to the patient) and donor 1b (a healthy unrelated donor). As shown in Figure 1(a), on day 0 (T0) CD8⁺ T cells isolated from RCC1 patient's PBMC displayed an elevated (HLA class I-restricted) reactivity against the autologous RCC1 cell line versus unstimulated CD8⁺ T cells isolated from either of the healthy donor controls (Figures 1(b) and 1(c), resp.), which was statistically significant ($P < .03$). After MLTC, the generated patient's CD8⁺ T cells continued to recognize RCC1 tumor cells, but they failed to react against autologous nonmalignant kidney cells (PTECs) or autologous PHA blasts (Figure 2(a)). T cell reactivity against the RCC1 line was MHC class-I restricted, with mAbs directed against single HLA-A (i.e., HLA-A24) or common HLA-B/-C determinants inhibiting specific T cell recognition of tumor cells by 75.5% and 43.4%, respectively (Figure 2(b)). An analysis of T cells generated from the patient's sibling (i.e., MLTC2) revealed that the responder CD8⁺ T cells recognized the HLA-matched RCC cell line to a statistically greater degree than observed from the patient's own CD8⁺ T cells ($P < .05$), while responding against the K562 LAK/NK target cell line to a moderate level (Figure 2(c)). These CTL recognized the RCC1 cell line in a predominantly class I-restricted manner based on the ability of the anti-A24 and anti-HLA-B/C mAbs to inhibit responses by 52.3% and 70.7%, respectively. Finally, in the case of the HLA-matched, unrelated donor (i.e., MLTC3), we observed the greatest capacity (versus MLTC1 and MLTC2) of CD8⁺ T cells to respond to the RCC1 cell line ($P < .03$; Figure 2(d)). Anticlass I blocking studies identified HLA-A24 (28% inhibition) and HLA-A3 (57.8% inhibition) as the predominant HLA-restricting elements for MLTC3 T cells. In all the three anti-RCC1 T cell systems studied, ELISPOT assays showed that T lymphocytes specifically secreted IFN- γ and granzyme B, but not interleukin-5 (IL-5), supporting dominant Type1 T cell responsiveness (data not shown).




FIGURE 1: IFN- γ Elispot Assay CD8⁺ T cells isolated at day 0. RCC1 cell line as stimulator cells and CD8⁺ T cells isolated from PBMC of RCC1-patient (a) and their allogeneic donor-1a (b) and donor-1b (c) at day 0 were tested in 20 h IFN- γ -Elispot Assay. HLA restriction of anti-RCC reactivity was determined by adding mAbs specific for total HLA-class I and HLA-DR. Results represent the average (\pm SD) of triplicate wells and the presented data derive from 1 representative experiment of 5 performed. A paired *t*-test analysis was used to compare the frequencies of autologous versus allogeneic specific CD8⁺ T cells stimulated with RCC1 cell line, after subtraction of HLA class I reactivity; $P < .03$ was considered statistically significant (*).

Analysis of cytotoxic CD8⁺ T cell responses using ⁵¹Cr-release assays similarly indicated a hierarchy of MLTC3 (75% lysis at an E/T ratio of 20 : 1) > MLTC2 (55% lysis) > MLTC1 (40% lysis) T cell response against the RCC1 cell line (Figures 3(a)–3(c)). MLTC1 and MLTC2 T cells also mediated moderate cytolysis of K562 cells, but not RCC1-LCL (Figures 3(a) and 3(b)), while MLTC3 T cells recognized RCC1-LCLs (65% lysis at an E/T ratio of 20 : 1) but not K562 cells (Figure 3(c)).

3.2. Differential Turnover of the CD8⁺ Memory T Cell Subset in RCC Patients after MLTC Stimulation. Given the perceived importance of CD8⁺ T cells in mediating anti-tumor




FIGURE 2: IFN- γ ELISPOT analyses of RCC-specific CD8⁺ T cell responses after MLTC. RCC1 CD8⁺ T cells stimulated by MLTC were tested for their reactivity against HLA-matched target cells such as autologous RCC, PTEC, and PHA-activated PBMC, (a). In panels (b), (c), and (d), K562 cells served as targets. A paired *t*-test analysis was used to compare the frequencies of autologous responder-CD8⁺ T cells versus allogenic responder-CD8⁺ T cells generated from the patient's sibling ($*P < .05$) and unrelated donor ($^{\S}P < .03$), stimulated with RCC1 line. The HLA restriction of the anti-RCC T cell response was determined by addition of blocking mAbs specific for total HLA-class I, HLA-A24, HLA-A3, or HLA-B/-C alleles. Results represent the average (\pm SD) of triplicate wells and the presented data derive from 1 representative experiment of 5 performed.

immunity, we extended our characterization of autologous and allogenic RCC-specific CD8⁺ T cells generated from MLTC, using 6-color flow cytometry, focusing on markers of T cell activation, proliferation, and differentiation. In agreement with previous studies [6, 10], four distinct phenotypic subsets of responder CD8⁺ T cells could be identified using CD45RA and CCR7 as markers. Whereas naive T cells express both CD45RA and CCR7, memory T cells were characterized by the down regulated expression of CD45RA and could be additionally subdivided, based on differential expression of CCR7, into effector memory (T_{EM} : CD45RA⁻, CCR7⁻) and central memory subsets (T_{CM} : CD45RA⁻, CCR7⁺). Several recent reports have defined the CD45RA⁺/CCR7⁻ T cell subset (T_{EMRA}) as *terminally differentiated effector* T cells which lack CD27 and/or CD28 expression and are able to migrate to inflamed tissues, where they may exert vigorous *ex vivo* effector functions, including target cell lysis [24, 25].

CD8⁺ T cells were purified from PBMC isolated from 5 RCC patients as well as HLA-matched normal donors and analyzed for their phenotypes on day 0 (T0) and after 5 weeks of MLTC culture (T35; Table 2). Freshly isolated CD8⁺ T cells from RCC patients displayed a predominant memory phenotype comprised of T_{EM} and T_{EMRA} , while similarly isolated CD8⁺ T cells from healthy donors exhibited a predominantly naive (i.e., CD45⁺, CCR7⁺) phenotype, with lower percentages of T_{EMRA} cells and comparable levels of T_{CM} and T_{EM} cells. As expected, after 35 days of MLTC, the percentage of naive T cells (versus T0) declined in all subjects, and the percentage of T_{CM} increased, most notably in the normal donor MLTC ($59\% \pm 4.6$) versus the patient MLTC ($11 \pm 2.8\%$).

When taken together, these data are in line with previous reports in which tumor-reactive CD8⁺ T_{CM} mediate superior anti-tumor efficacy when compared with T_{EM} [7, 26]. In contrast, MLTC stimulation induced higher frequencies of





FIGURE 3: Cytotoxicity assay of RCC1-specific CTL at day 35 of the MLTC. Specific lysis was determined in a 4-hour ^{51}Cr -release assay using CD8^+ T cells collected from d 35 MLTC established from patient RCC1 (a) or HLA-matched normal donors (the patient's sibling (b) and an unrelated individual (c)). T cells were used at varying effector-to-target (E:T) cell ratios against target cells including RCC1, K562, and RCC1-LCL. Data are presented as percentages of specific lysis \pm SD.

TABLE 2: Flow cytometry analysis of CD45RA/CCR7 expression on CD8^+ T cells at day 0 and after 35 days of MLTC. Differential expression of the CD45RA, CCR7 cell surface molecules was analyzed on CD8^+ T cells at day 0 and after extended MLTC stimulation. CD8^+ T cells were separated into four subsets (T_N , T_{CM} , T_{EM} , T_{EMRA}) based on differential expression of the CD45RA and CCR7 markers. Results are the mean (\pm SD) of values obtained from 5 RCC patients and their corresponding HLA-matched normal donors.

| | CD8^+ Naive $\text{CD45RA}^+\text{CCR7}^+$ | | T_{CM} $\text{CD45RA}^-\text{CCR7}^+$ | | T_{EM} $\text{CD45RA}^-\text{CCR7}^-$ | | T_{EMRA} $\text{CD45RA}^+\text{CCR7}^-$ | |
|----------------|--|---------------|--|--------------|--|----------------|--|----------------|
| | Day 0 | Day 35 | Day 0 | Day 35 | Day 0 | Day 35 | Day 0 | Day 35 |
| RCC patients | 30.6 ± 4.0 | 4.4 ± 1.6 | 8.2 ± 2.3 | 11 ± 2.8 | 14 ± 3.8 | 52.6 ± 6.7 | 46 ± 5.3 | 30.6 ± 4.2 |
| Healthy donors | 49 ± 3.1 | 3.1 ± 1.6 | 10.8 ± 5.3 | 59 ± 4.6 | 20.2 ± 3.9 | 32.8 ± 7.2 | 17.4 ± 3.6 | 4.4 ± 1.8 |
| P value | .021 | .139 | .019 | .008 | .034 | .003 | .0017 | .0006 |



| | pE1 CD27 ⁺ CD28 ⁺ | | pE2 CD27 ⁺ CD28 ⁻ | | E CD27 ⁻ CD28 ⁻ | |
|--------------|--|--------|--|----------|--|----------|
| | Day 0 | Day 35 | Day 0 | Day 35 | Day 0 | Day 35 |
| RCC patients | 20 ± 2.3 | 13 ± 4 | 19 ± 3.8 | 62 ± 4.1 | 61 ± 5.3 | 25 ± 2.5 |
| P value | .018 | | .001 | | .005 | |

(d)

FIGURE 4: Characterization of the T_{EMRA} population in RCC patients. Based on CD28/CD27 expression, CD8⁺T cells were separated into three distinct populations: pE1 (RA⁺CCR7⁻27⁺28⁺), pE2 (RA⁺CCR7⁻27⁺28⁻), and E (RA⁺CCR7⁻27⁻28⁻). Results are depicted for patients RCC2 but are representative of all 5 RCC patients evaluated.

CD8⁺ T_{EM} (and predominantly T_{EMRA}) from RCC patients versus normal donors. Given this difference, we further characterized the T_{EMRA} subpopulation of CD8⁺ T cells generated from RCC patients versus normal donors with regard to comparative levels of expression of the CD27 and CD28 markers (Figure 4). In accordance with the model proposed by Rufer et al. [10], we identified three functionally distinct CD8⁺ T_{EMRA} (RA⁺CCR7⁻) subsets in RCC patients: pE1 (27⁺28⁺), and pE2 (27⁺28⁻), and E (27⁻28⁻) that have been defined “early”, “intermediate,” and “late” phenotypes, respectively, based on a proposed linear pathway of differentiation.

Our data suggest that activated CD8⁺ T cells isolated from RCC patients were more likely to be in a state of intermediate differentiation (pE2) when compared to normal donor CD8⁺ T cells. Indeed, we noted a profound increase in T cell expression of CD27 after MLTC stimulation (from (T0) 19% ± 3.8 to (T35) 62% ± 4.1). Given previous reports that the level of CD27 expressed by CD8⁺ T cells correlates inversely with the ability of these cells to mediate cytotoxic functions [27], our results intimate that the reduced cytotoxic activity observed for CD8⁺ T lymphocytes in RCC patients may be due to their lack of terminal (functional) differentiation.

3.3. Cell Cycle Dysfunction in Activated CD8⁺ T Cells from RCC Patients. To further investigate the relationship between cell cycling and T cell differentiation status, we evaluated the expression pattern of cell cycle proteins involved in the G1/S transition in CD8⁺ T cells isolated from RCC patients vs. normal donors on day 35 of MLTC. It has been well established that initiation of T lymphocytes-mediated immune responses involves two cellular processes: entry into the cell cycle (G0 → G1) for clonal proliferation and coordinated changes in surface and secreted molecules that mediate effector functions, associated with the differentiation of T lymphocytes into an effector or memory subpopulations. As shown in Figure 5, activated patient CD8⁺ T cells were mainly present in the G0/G1 phase (85.7 ± 2.08 vs. 63.3 ± 5.5) and less in the S phase (1.41 ± 0.18 vs. 12 ± 4.5) when compared with CD8⁺ T cells generated in normal donor MLTC. A statistically significant negative correlation was noted between phases G1 and S ($r = -0.95$, $P < .05$) in the patients’ T cells, suggesting that in RCC patients there was an arrest at this cell cycle transition point (Figure 6).

3.4. Association between Cell Division and Tumor Responsiveness in Activated RCC CD8⁺ T Cells. To better understand




FIGURE 5: Cell cycle analysis of responder-CD8⁺ T cells after MLTC. Cellular DNA content was analyzed in three RCC patients and their HLA-matched normal donors by flow cytometry, with data analyzed on WinMDI version 2.8 software.




FIGURE 6: Alterations in the cell cycle distribution of patient versus normal donor anti-tumor CD8⁺ T cells. A statistically significant, negative correlation was observed between G1 and S percentages in 3 RCC patients.

the association between T cell cycling and anti-tumor tumor responsiveness, we assessed day 0 and day 35 MLTC CD8⁺ T cells for their expression of p27^{KIP1} and STAT5/6, proteins known to be required for normal T cell proliferation [15, 28]. IL2-mediated downregulation of p27^{KIP1} is crucial for the normal progression of CD8⁺ T cells from the G1 phase to the S phase of the cell cycle. Activation of the JAK3/STAT5/6 pathway is required for p27^{KIP1} downregulation, hyperphosphorylation of RB, and release of active E2F in T cells. When we analyzed STAT5 expression (Figure 7(a)), we found that IL-2-induced STAT5 phosphorylation was readily detectable in normal T cells (64.5 ± 5.3 /HPF and 71 ± 6.1 /HPF resp.) but resulted markedly reduced in RCC patients ($22.1 \pm$

3.3 /HPF, $P < .001$). Interestingly, as displayed in Figure 7(b), we observed that STAT6 expression decreased in patients (40.2 ± 4.6 /HPF), but not normal donors (57.3 ± 6.8 /HPF and 60.8 ± 1.7 /HPF resp., $P < .0032$) MLTC CD8⁺ T cells.

This suggests that, despite the presence of high doses of rhIL-2, the inability of patient CD8⁺ T cells to proliferate in MLTC likely results from defects in the JAK3/STAT5/6 signaling pathway. To test this hypothesis, we evaluated p27^{KIP1} expression status in MLTC CD8⁺ T cells developed from RCC patients vs. normal donors. We observed that T cell expression of p27^{KIP1} was increased in T cells from patients (17.5 ± 3.2 /HPF) vs. normal, healthy donors (8 ± 1.55 /HPF and 10.2 ± 0.37 /HPF, $P = .0046$ resp.) (Figure 7(b)).

These data suggest that the integrity of the JAK3/STAT5/6 pathway is likely linked to T cell division and that activated T cells that fail to divide upon appropriate stimulation present with operational defects in this pathway.

In order to confirm that G1/S regulatory defects were present in MLTC CD8⁺ T cells developed from RCC patients, we next investigated these cells for expression of two key factors in the control of the G₁-S-phase transition, E2F4 and Id2, which might confer a proliferative advantage to tumor cells in vivo [21, 29]. As shown in Figure 8, E2F4 expression was reduced in MLTC CD8⁺ T cells from patients (11.2 ± 3.2 /HPF) vs. normal donors (14.71 ± 1.028 /HPF and 16.6 ± 2.8 /HPF, $P = .0018$ resp.). We also found that CD8⁺ T cells from patients expressed lower levels of Id2 vs. normal donor T cells (28.7 ± 4.3 /HPF vs. 56.9 ± 8.4 /HPF and 34.8 ± 6.5 /HPF, $P = .037$, resp.), confirming an arrest in the G₀ cell cycle phase among patient T cells (Figure 9).




FIGURE 7: Altered expression of pSTAT5, pSTAT6, and p27^{KIP1} in RCC patient CD8⁺ T cells. (a) Distribution of pSTAT5 evaluated by confocal microscopy on CD8⁺ T cells of RCC patients and their HLA-matched normal donors, using anti-STAT5 (green) antibodies. (b) The distribution of p-p27^{KIP1} and pSTAT6 were evaluated by confocal microscopy on CD8⁺ T cells of RCC patients and their HLA-matched normal donors, using anti-p-p27^{KIP1} (red) anti-STAT6 (green) antibodies. p27^{KIP1}/STAT6 interaction results in a yellow staining in the merge. Quantitative analysis of high-power fields (HPFs) in (a) and (b) was carried out by confocal microscopy software (Leica, TCS-SP2). The results were expressed as mean values \pm SD. A *P*-value < .05 was considered statistically significant. Data are from 1 of 5 experiments that all gave similar results.




FIGURE 8: Altered expression of E2F4 in RCC patient CD8⁺ T cells. CD8⁺ T cells were surface-labeled with anti-p-p27^{KIP1} (red) and anti-E2F4 (green) mAbs and the performance of confocal microscopy analysis. Quantitative analysis of HPF was carried out by confocal microscopy software (Leica, TCS-SP2; Student's *t*-test, $P < .05$). Data are from 1 of 5 experiments, that all gave comparable results.




FIGURE 9: Altered expression of Id2 in RCC patient CD8⁺ T cells. CD8⁺ T cells were surface-labeled with anti-Id2 (red) mAb and the performance of confocal microscopy analysis. Quantitative analysis of HPF was carried out by confocal microscopy software (Leica, TCS-SP2; Student's *t*-test, $P < .05$). Data are from 1 of 5 experiments that all gave similar results.

4. Conclusions

The experimental approaches used in our study provide valuable tools to assist in the elucidation of cellular mechanisms underlying the altered activation state of T cells in patients with cancer. We used RCC as a model cancer to explore this question, as it is well documented that patients with active, disseminated disease are typically characterized by “functionally inappropriate”, dysfunctional, or proapoptotic immune cells that are impotent in mediating clinically beneficial outcomes *in vivo* [30, 31].

We confirmed that it is possible to generate and expand tumor-reactive CD8⁺ T cells *in vitro* from the peripheral blood of RCC patients as well as normal donors [32]. Normal donor RCC-specific CD8⁺ T cells appeared to recognize a diverse array of antigens and displayed superior anti-tumor activity when compared with patient-derived CD8⁺ T cells [23]. Normal donor CD8⁺ T cells developed in MLTC cultures also demonstrated significant reactivity against K562 target cells (in contrast to patient MLTC-derived CD8⁺ T cells), supporting the conclusion that these allogenic MLTC-CD8⁺ T cells were capable of mediating HLA class I-restricted as well as HLA-unrestricted cytotoxicity.

A corollary analysis of the phenotypes of allogenic and autologous MLTC-CD8⁺ T cells predicated on a recently proposed pathway of T-cell lineage differentiation [6] revealed that the majority of freshly isolated CD8⁺ T cells from RCC patients display a memory phenotype characterized by very low frequencies of CD45RA⁻, CCR7⁺T_{CM} cells, high frequencies of CD45RA⁻, CCR7⁻T_{EM}, and terminally differentiated CD45RA⁺, CCR7⁻T_{EMRA}. Interestingly, we noted that after MLTC expansion, patient CD8⁺T cells are mainly composed of antigen-experienced T_{EM} and T_{EMRA} and are deficient in T_{CM}. In sharp contrast, MLTC-CD8⁺ T cells developed from normal donors contain high frequencies of T_{CM} population and far lower percentages of both T_{EM} and T_{EMRA} cells. These data are particularly important as tumor-reactive CD8⁺ T_{CM} have been previously suggested to provide superior anti-tumor protection vs. T_{EM} [7, 26]. Furthermore, it has been shown that T_{CM} exhibit greater expansion potential in concert with enhanced resistance to apoptosis, whereas T_{EM}/T_{EMRA} have a skewed TCR repertoire and proliferate poorly in response to cognate antigen [9]. Importantly, our study revealed that patients CD8⁺ T_{EMRA} expanded in MLTC are more likely to exhibit an intermediate state of differentiation (i.e., CD27⁺, CD28⁻), capable of mediating only partial effector functions and proliferative capacity. Normally, CD8⁺ T cell differentiation correlates with a progressive loss in expression of the CCR7, CD28, and CD27 markers and increased cytolytic capacity [33]. Hence, our results suggest that although CD8⁺ T cells in RCC patients are capable of reacting against tumor cells, their effector functions and longevity are suboptimal due to the absence of T_{CM} cells and their incomplete state of differentiation. These findings have important implications for cancer immunotherapy approaches [34], including DC-based vaccines, and in this regard recently we have demonstrated that dendritic cells (DCs) generated using IFN- α and loaded with tumor antigens are capable of “revitalizing”

or expanding a therapeutically preferred cohort of anti-tumor CD8⁺ T cells (T_{CM}) from the PBMC of RCC patients [35].

Other groups have investigated the profiles of human CD8⁺ T_{CM}, T_{EM}, T_{EMRA} using gene expression microarrays, showing that T_{CM} are characterized by a higher level of basal and cytokine-induced STAT5/6 phosphorylation, reflecting their superior capacity for self-renewal vs. effector memory cells, and in particular, T_{EMRA} [11]. Interestingly, we observed a down regulation of both STAT5/6 molecules in patient CD8⁺ T cells, in association with their inferior proliferative capacity, which is consistent with their known dysfunction *in vivo*. These data suggest a role for STAT pathway in the post activation survival of T cells and provide new insights into the nature of a late (cell cycle) proliferation block in the T cell compartment as a result of STAT6 deficiency. Indeed, in CD8⁺ T cells analyzed from RCC patients, we observed a down regulation in JAK3/STAT5/6 expression and corollary defects in cell cycle (G1/S) progression and cell proliferation. These results establish the crucial role of the JAK3/STAT5/6 signaling pathway in normal CD8⁺ T cell proliferation/differentiation, consistent with a recent report by Jin et al. [36].

T cell activation and proliferation require several appropriate signals to complete transitions through the cell cycle, from the G1 to the S phase. Interestingly, activated CD8⁺ T cells from RCC patients, which exhibit reduced proliferative responses against their own tumors, express increased levels of p27^{KIP1}, which is known to lead to G0 phase cell cycle arrest and a slowed/arrested progression to G1/S transition. These results are consistent with those reported by Zhang et al. [28] where p27^{KIP1} was found to inhibit antigen-driven T_{EM} expansion *in vivo*. Our report adds to a very limited literature in identifying a G1/S cell cycle regulatory defect in T cells isolated from cancer patients, and it provides a better understanding of mechanism(s) underlying tumor-induced immune deviation that occurs during cancer progression.

One of the key events during the G1 phase is activation of the transcription factor E2F4 [37]. E2F4 is a crucial switch in the control of homeostasis and tumorigenesis [21] and, together with Id2, serves as a critical regulator of cell cycle progression. Specifically, Id2 has been suggested to antagonize the regulation of the cell cycle by tumor suppressor proteins [38]. Our analyses revealed a dramatic reduction in E2F4 and Id2 expression in CD8⁺T cells from RCC patients vs. healthy donors after MLTC. These data provide evidence for the importance of Id2 in enforcing “normal” cell cycle progression in tumor-reactive T cells harvested from RCC patients.

In conclusion, we have shown for the first time that RCC patients exhibit a functionally suboptimal anti-tumor CD8⁺ T cell repertoire based on defects in (the JAK3/STAT5/6) intracellular signaling pathway that leads to T cell arrest in the G0/G1 phase of the cell cycle and prevention of their terminal differentiation. Furthermore, our data support the conclusion that CD8⁺ T cell overexpression of p27KIP1 and/or loss of Id2 expression are characteristics of dysfunctional T cells in RCC patients, making these potential therapeutic targets in the setting of patients with RCC.

Acknowledgments

The authors would like to thank Professor FP Schena for his support in the clinical aspects of this work and they are grateful to M. V. C. Pragnell, B. A., for language revision. E. Cavalcanti and M. Gigante equally contributed to this work.

References

- [1] B. I. Rini, S. C. Campbell, and W. K. Rathmell, "Renal cell carcinoma," *Current Opinion in Oncology*, vol. 18, no. 3, pp. 289–296, 2006.
- [2] C. Coppin, "Immunotherapy for renal cell cancer in the era of targeted therapy," *Expert Review of Anticancer Therapy*, vol. 8, no. 6, pp. 907–919, 2008.
- [3] S. Négrier, D. Perol, A. Ravaud, et al., "Randomized study of intravenous versus subcutaneous interleukin-2, and IFN α in patients with good prognosis metastatic renal cancer," *Clinical Cancer Research*, vol. 14, no. 18, pp. 5907–5912, 2008.
- [4] N. Imai, H. Ikeda, I. Tawara, and H. Shiku, "Tumor progression inhibits the induction of multifunctionality in adoptively transferred tumor-specific CD8⁺ T cells," *Journal of Immunology*, vol. 39, pp. 241–253, 2009.
- [5] E. M. Janssen, E. E. Lemmens, T. Wolfe, U. Christen, M. G. von Herrath, and S. P. Schoenberger, "CD4⁺ T cells are required for secondary expansion and memory in CD8⁺ T lymphocytes," *Nature*, vol. 421, no. 6925, pp. 852–856, 2003.
- [6] F. Sallusto, J. Geginat, and A. Lanzavecchia, "Central memory and effector memory T cell subsets: function, generation, and maintenance," *Annual Review of Immunology*, vol. 22, pp. 745–763, 2004.
- [7] C. A. Klebanoff, L. Gattinoni, P. Torabi-Parizi, et al., "Central memory self/tumor-reactive CD8⁺ T cells confer superior antitumor immunity compared with effector memory T cells," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, no. 27, pp. 9571–9576, 2005.
- [8] C. Berger, M. C. Jensen, P. M. Lansdorp, M. Gough, C. Elliott, and S. R. Riddell, "Adoptive transfer of effector CD8⁺ T cells derived from central memory cells establishes persistent T cell memory in primates," *Journal of Clinical Investigation*, vol. 118, no. 1, pp. 294–305, 2008.
- [9] J. Geginat, A. Lanzavecchia, and F. Sallusto, "Proliferation and differentiation potential of human CD8⁺ memory T-cell subsets in response to antigen or homeostatic cytokines," *Blood*, vol. 101, no. 11, pp. 4260–4266, 2003.
- [10] N. Rufer, A. Zippelius, P. Batard, et al., "Ex vivo characterization of human CD8⁺ T subsets with distinct replicative history and partial effector functions," *Blood*, vol. 102, no. 5, pp. 1779–1787, 2003.
- [11] T. Willinger, T. Freeman, H. Hasegawa, A. J. McMichael, and M. F. C. Callan, "Molecular signatures distinguish human central memory from effector memory CD8⁺ T cell subsets," *Journal of Immunology*, vol. 175, no. 9, pp. 5895–5903, 2005.
- [12] S. G. Rane and E. P. Reddy, "Janus kinases: components of multiple signaling pathways," *Oncogene*, vol. 19, pp. 5662–5579, 2000.
- [13] M. L. Fero, M. Rivkin, M. Tasch, et al., "A syndrome of multiorgan hyperplasia with features of gigantism, tumorigenesis, and female sterility in p27^{Kip1}-deficient Mice," *Cell*, vol. 85, no. 5, pp. 733–744, 1996.
- [14] K. Nakayama, N. Ishida, M. Shirane, et al., "Mice lacking p27^{Kip1} display increased body size, multiple organ hyperplasia, retinal dysplasia, and pituitary Tumors," *Cell*, vol. 85, no. 5, pp. 707–720, 1996.
- [15] M. H. Kaplan, C. Daniel, U. Schindler, and M. J. Grusby, "Stat proteins control lymphocyte proliferation by regulating p27^{Kip1} expression," *Molecular Biology of Eukaryotic Cells*, vol. 18, pp. 1996–2003, 1998.
- [16] R. Shen and M. H. Kaplan, "The homeostasis but not the differentiation of T cells is regulated by p27^{Kip1}," *Journal of Immunology*, vol. 169, no. 2, pp. 714–721, 2002.
- [17] L. Rivoltini, P. Canese, V. Huber, et al., "Escape strategies and reasons for failure in the interaction between tumour cells and the immune system: how can we tilt the balance towards immune-mediated cancer control?" *Expert Opinion on Biological Therapy*, vol. 5, pp. 463–476, 2005.
- [18] N. C. Lea, S. J. Orr, K. Stoeber, et al., "Commitment point during G₀ → G₁ that controls entry into the cell cycle," *Molecular and Cellular Biology*, vol. 23, no. 7, pp. 2351–2361, 2003.
- [19] S. Gaubatz, G. J. Lindeman, S. Ishida, et al., "E2F4 and E2F5 play an essential role in pocket protein-mediated G1 control," *Molecular Cell*, vol. 6, no. 3, pp. 729–735, 2000.
- [20] N. Dyson, "The regulation of E2F by pRB-family proteins," *Genes and Development*, vol. 12, no. 15, pp. 2245–2262, 1998.
- [21] Z. H. Fang and Z. C. Han, "The transcription factor E2F: a crucial switch in the control of homeostasis and tumorigenesis," *Histology and Histopathology*, vol. 21, no. 4–6, pp. 403–413, 2006.
- [22] M. Lahn, G. Köhler, C. Schmoor, et al., "Processing of tumor tissues for vaccination with autologous tumor cells," *European Surgical Research*, vol. 29, no. 4, pp. 292–302, 1997.
- [23] S. Kausche, T. Wehler, E. Schnürer, et al., "Superior antitumor in vitro responses of allogeneic matched sibling compared with autologous patient CD8⁺ T cells," *Cancer Research*, vol. 66, no. 23, pp. 11447–11454, 2006.
- [24] P. Romero, A. Zippelius, I. Kurth, et al., "Four functionally distinct populations of human effector-memory CD8⁺ T lymphocytes," *Journal of Immunology*, vol. 178, no. 7, pp. 4112–4119, 2007.
- [25] H. Takata and M. Takiguchi, "Three memory subsets of human CD8⁺ T cells differently expressing three cytolytic effector molecules," *Journal of Immunology*, vol. 177, no. 7, pp. 4330–4340, 2006.
- [26] L. Gattinoni, S. E. Finkelstein, C. A. Klebanoff, et al., "Removal of homeostatic cytokine sinks by lymphodepletion enhances the efficacy of adoptively transferred tumor-specific CD8⁺ T cells," *Journal of Experimental Medicine*, vol. 202, no. 7, pp. 907–912, 2005.
- [27] H. Tomiyama, H. Takata, T. Matsuda, and M. Takiguchi, "Phenotypic classification of human CD8⁺ T cells reflecting their function: inverse correlation between quantitative expression of CD27 and cytotoxic effector function," *European Journal of Immunology*, vol. 34, no. 4, pp. 999–1010, 2004.
- [28] S. Zhang, V. A. Lawless, and M. H. Kaplan, "Cytokine-stimulated T lymphocyte proliferation is regulated by p27^{Kip1}," *Journal of Immunology*, vol. 165, pp. 6270–6277, 2000.
- [29] M. A. Cannarile, N. A. Lind, R. Rivera, et al., "Transcriptional regulator Id2 mediates CD8⁺ T cell immunity," *Nature Immunology*, vol. 7, no. 12, pp. 1317–1325, 2006.
- [30] L. E. Van Den Hove, S. W. Van Gool, H. Van Poppel, et al., "Phenotype, cytokine production and cytolytic capacity of fresh (uncultured) tumour-infiltrating T lymphocytes in human renal cell carcinoma," *Clinical and Experimental Immunology*, vol. 109, no. 3, pp. 501–509, 1997.

- [31] M. Gigante, A. Blasi, A. Loverre, et al., "Dysfunctional DC subsets in RCC patients: ex vivo correction to yield an effective anti-cancer vaccine," *Molecular Immunology*, vol. 46, no. 5, pp. 893–901, 2009.
- [32] A. Dörrschuck, A. Schmidt, E. Schnürer, et al., "CD8⁺ cytotoxic T lymphocytes isolated from allogeneic healthy donors recognize HLA class Ia/Ib-associated renal carcinoma antigens with ubiquitous or restricted tissue expression," *Blood*, vol. 104, no. 8, pp. 2591–2599, 2004.
- [33] V. Appay, P. R. Dunbar, M. Callan, et al., "Memory CD8⁺ T cells vary in differentiation phenotype in different persistent virus infections," *Nature Medicine*, vol. 8, no. 4, pp. 379–385, 2002.
- [34] J. Copier, A. G. Dalglish, C. M. Britten, et al., "Improving the efficacy of cancer immunotherapy," *European Journal of Cancer*, vol. 45, no. 8, pp. 1424–1431, 2009.
- [35] M. Gigante, M. Mandic, A. K. Wesa, et al., "Interferon-alpha (IFN-alpha)-conditioned DC preferentially stimulate type-1 and limit Treg-type in vitro T-cell responses from RCC patients," *Journal of Immunotherapy*, vol. 31, pp. 254–262, 2008.
- [36] D. Jin, M. Takamoto, T. Hu, S. Taki, and K. Sugane, "STAT6 signalling is important in CD8⁺ T-cell activation and defence against toxoplasma gondii infection in the brain," *Immunology*, vol. 127, no. 2, pp. 187–195, 2009.
- [37] H. Müller, A. P. Bracken, R. Vernell, et al., "E2Fs regulate the expression of genes involved in differentiation, development, proliferation, and apoptosis," *Genes and Development*, vol. 15, no. 3, pp. 267–285, 2001.
- [38] A. Lasorella, A. Iavarone, and M. A. Israel, "Id2 specifically alters regulation of the cell cycle by tumor suppressor proteins," *Molecular and Cellular Biology*, vol. 16, no. 6, pp. 2570–2578, 1996.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

