

## Review Article

# Schistosome: Its Benefit and Harm in Patients Suffering from Concomitant Diseases

Yoshio Osada and Tamotsu Kanazawa

Department of Immunology and Parasitology, University of Occupational and Environmental Health, Kitakyushu 807-8555, Japan

Correspondence should be addressed to Yoshio Osada, yosada@med.uoeh-u.ac.jp

Received 1 July 2011; Accepted 6 September 2011

Academic Editor: Luis I. Terrazas

Copyright © 2011 Y. Osada and T. Kanazawa. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Schistosomiasis is an important tropical disease affecting approximately 200 million people worldwide. Because of its chronicity and robust immunomodulatory activity, the effects of schistosomes on other diseases, such as allergies, autoimmunity, and infectious diseases, have been studied extensively in both epidemiological and experimental settings. In this paper, we summarize the beneficial and harmful effects of schistosomes. The importance of controlling schistosomiasis is also discussed.

## 1. Introduction

*Schistosoma* spp., blood flukes, are parasitic helminths found mainly in developing countries with a tropical or subtropical climate and affect 200 million people worldwide [1]. *Schistosoma mansoni*, *japonicum*, and *mekongi* harbor in veins of the portal system and lay eggs in the blood vessels. The deposition of numerous eggs in the intestines and liver results in intestinal and hepatic granulomatous lesions, fibrosis, portal hypertension, and hepatosplenomegaly. In contrast, *Schistosoma haematobium* mainly harbors in the venous plexus of the bladder and/or rectal venous plexus. This worm usually causes bloody urine but is also considered to have an etiological relationship with bladder cancer [2]. Because of the extensive distribution of schistosomes and morbidity due to egg deposition, researchers have been interested in the influences of schistosome infections on concomitant diseases [3]. As mentioned, *S. haematobium* is an important carcinogenic factor of bladder cancer although the worm itself does not have mutagenic activity [4]. This parasite is also suggested to be a risk factor for the transmission of HIV [5]. These effects are attributable to pathological lesions caused by the parasites. On the other hand, most helminthic parasites including schistosomes are known to induce robust Th2-polarization. Especially in schistosome infections, egg deposition in the host tissues was reported to be the major stimulus of Th2 responses [6], and egg proteins (e.g.,

omega-1, IPSE, and peroxiredoxin) are involved in the Th2-biasing activity [7–11]. In addition, schistosome eggs have immunomodulatory potential inducing the alternative activation of macrophages [12] and regulatory T-cell expansion [13]. Because of its robust systemic Th2-inducing and immunomodulatory ability, this worm has been studied most extensively for its bystander effects on various immunological phenomena *in vivo*. In this paper, we summarize the effects of a concurrent infection of schistosomes on immunological disorders and parasitic/microbial infections.

## 2. Effects of Schistosomes on Immunological Disorders or Infectious Diseases in Mice and Humans

**2.1. Allergy.** Effects of helminths (including schistosomes) on allergic diseases have been studied extensively in both experimental and epidemiological settings. In experimental asthma or airway hypersensitivity models, schistosome infections and antigen administrations have been consistently shown to protect the animals from the diseases (Table 1). In most studies on the antiasthmatic effects of schistosomes, cellular infiltration (including eosinophils) into bronchoalveolar lavage fluid (BALF) was diminished and simultaneously IL-4, IL-5, and IgE levels were reduced. In contrast, an increase in Treg cells and augmentation of IL-10 production

were observed. Fallon and Mangan [14] designated this kind of Th2 response (Treg dominant and IL-10 dominant) a “modified Th2 response” as opposed to the conventional “allergic Th2 response.” They demonstrated that in infected mice, IL-10-producing CD1d<sup>high</sup> B cells induce Treg cells and consequently ameliorate allergic airway inflammation [15]. In the study, IL-10 was indispensable to the effects of schistosomes. The authors also showed the importance of B cells, and IL-10 in the suppression of systemic anaphylaxis by schistosomes [16]. Smits et al. [17] found that in adoptive transfer experiments, spleen cells (especially B cells and CD4<sup>+</sup> T cells) from chronically infected mice could confer protection against pulmonary infiltration by white blood cells, especially eosinophils. In addition, administration of anti-IL-10R antibody cancelled out the effects of the cell transfer. These studies seem to support the critical importance of B cells, Treg cells and IL-10 in schistosome-induced protection against airway allergic inflammation. In studies by another group, however, IL-10 signaling was not essential for the antiallergic immunomodulation by schistosomes [18, 19]. The reason for this discrepancy is unclear, but as the authors point out, other immunomodulatory factors may be able to compensate the absence of IL-10 [19].

Antiallergic effects of schistosomes have also been demonstrated in humans. In a study in Brazil, both asthmatic symptoms and skin reactivity to indoor allergens were reduced in *S. mansoni*-infected asthmatics compared to non-infected individuals [42]. In another study, higher expression levels of HLA-DR, IL-10R (in monocytes), CTLA-4 and CD40L (in CD4<sup>+</sup> T cells) were observed in *S. mansoni*-infected asthmatics [43]. According to that paper, the main sources of IL-10 were monocytes and Treg cells. A meta-analysis of current parasite infections and atopy [44] revealed that schistosomiasis was protective against allergic skin sensitization as well as some other helminths (*Ascaris lumbricoides*, *Trichuris trichiura*, hookworm). The same research group also reported a meta-analysis of interrelationships between helminths and asthma [45]. Hookworm infections were shown to be protective, but no beneficial effect was found for schistosome infections, probably due to the insufficient number of studies (only two). Collectively, anti-asthmatic effects of schistosomes have been confirmed in animal models and suppressive effects on allergic skin reactivity have been confirmed in humans. However, more epidemiological (especially intervention) studies are necessary to conclude whether schistosome infections have beneficial or detrimental effects on asthmatic patients.

**2.2. Autoimmunity.** Autoimmune disorders had been considered Th1-mediated diseases for a long time. As the Th2-biasing ability of parasitic helminths and consequent downregulation of Th1 responses have been well known, antiautoimmune effects of such parasites had been attributed to the downregulation of Th1 responses in infected animals. However, some major autoimmune diseases are now considered to be dependent on Th17, a newly found pathogenic T-cell subset that mainly produces IL-17 [46]. With this finding, the antiautoimmune properties of helminths have been revisited. In recent years, downmodulation of Th17 responses by para-

sitic helminths has been reported [24, 36, 47, 48]. If the suppressive activity on both Th1 and Th17 is common to parasitic helminths, helminths may become ideal sources of drug screening for treatment of autoimmune disorders. That is because both T-cell subsets are involved in the pathogenesis of some autoimmune diseases [49]. Moreover, the unstable nature of Th17 [49] reinforces this idea.

As summarized in Table 1, schistosomes suppress various autoimmunity models in rodents. An upregulation of IL-4 and downregulation of IFN- $\gamma$  responses are almost commonly observed. In addition, responses of IL-17 and/or TNF- $\alpha$ , both of which play pathological roles in autoimmune arthritis [24, 26] and hapten-induced colitis [36, 40], were also downregulated by schistosome infections. Moreover, our study in mice with collagen-induced arthritis (CIA) revealed that the disease-associated local augmentation of bone-destructive cytokines (i.e., IL-6 and RANKL) was abrogated in infected animals [24]. These results indicate that schistosomes have suppressive effects on both Th1/Th17 and inflammatory cytokines. In addition, schistosomes suppress other pathogenic mediators such as autoreactive antibodies. Schistosomes decreased levels of anti-insulin IgG [30], anti-TSHR IgG2a [41], and anticollagen IgG [24, 25]. This effect was also observed in humans, as Mutapi et al. recently reported that *S. haematobium* infection intensity was inversely related to autoreactive antinuclear antibody (ANA) levels [50]. The authors also found that antihelminthic treatment increased ANA levels. However, Rahima et al. reported the presence of antinuclear antibodies in *S. mansoni*-infected mice and that sera from patients with systemic lupus erythematosus (SLE) reacted with cercarial antigens [51], suggesting that schistosomes trigger some kinds of autoimmunity. In conclusion, large-scale cross-sectional studies may be necessary to reveal the interrelationships between schistosomiasis and autoimmunity.

It is reasonable to hypothesize that the downmodulation of proinflammatory cytokines and pathogenic antibodies is involved in the antiautoimmune activity of schistosomes, at least partially. As regulatory cytokines are known to downregulate proinflammatory cytokines and pathogenic antibodies, it is important to determine the “essential” regulatory cytokines in each disease models for elucidation of suppressive mechanisms. For this purpose, it is necessary to perform experiments of cytokine neutralization with specific antibodies or experiments using gene-targeted animals. In Th1/Th17-dependent autoimmunity, IL-4, the key cytokine of Th2 responses, may be responsible for the alleviation of the disease symptoms. For instance, by using gene-targeted mice, STAT6 (a key signaling molecule in the response to IL-4 and IL-13) was shown to be indispensable to the *S. mansoni* egg-induced suppression of experimental autoimmune encephalomyelitis (EAE) [28] and of TNBS-induced colitis [35]. In contrast, IL-4 and IL-13 were dispensable to the suppression of DSS-induced colitis (Th2 cytokine dominant and macrophage-mediated colitis) by male worms of *S. mansoni* [39]. In the same study, authors demonstrated that IL-10 and TGF- $\beta$  were also dispensable to the anticolic effects of schistosome, by using specific antibodies against IL-10R and TGF- $\beta$ . IL-10 was not a crucial regulatory cytokine

TABLE 1: Suppressive effects of schistosome on experimental allergy and autoimmunity in rodents.

Category of animal models	Diseases	Schistosome	Treatment	Proposed mechanisms	Refs	
Allergy	Asthma/Airway hypersensitivity or inflammation	Sm	Infection (male)	IL-5 ↓, IL-10↑	[20]	
			Infection, Egg i.p.	IL-4 ↓, IL-5 ↓, IgE ↓, Treg↑, Independent of IL-10	[18]	
			Infection (chronic)	B cells and CD4 <sup>+</sup> T cells, Dependent on IL-10	[17]	
			Infection and Adoptive transfer	IL-10-producing CD1d <sup>high</sup> B cells → Treg↑	[15]	
			Sm22.6, PIII	IL-4 ↓, IL-5 ↓, IgE ↓, Treg ↑, independent of IL-10	[19]	
	Systemic anaphylaxis	Sm	Sj	Infection (male, mixed)	IL-4 ↓, IL-5 ↓, IgE ↓, IL-10 ↑	[21]
				Infection and adoptive transfer	DC → IL-4 ↓, IL-5 ↓, IL-10 ↑	[22]
			Sj	SEA, Eggs (i.p., p.o.)	Treg↑	[23]
			Infection	IL-10-producing B cell	[16]	
Autoimmunity	Collagen-induced arthritis (CIA)	Sm	Infection	IL-17 ↓, TNF-α ↓, IL-6 ↓, RANKL ↓, anticollagen IgG ↓	[24]	
			Sj	Infection	IL-4 ↑, anticollagen IgG ↓	[25]
	Adjuvant-induced arthritis (AIA)	Sj	Sj16 i.p.	TNF-α ↓, IL-1β ↓, NO ↓, IL-10 ↑	[26]	
			Sm	Infection	IL-12p40 ↓, IFN-γ ↓, TNF-α ↓, IL-4 ↑	[27]
	Experimental autoimmune encephalomyelitis (EAE)	Sm	Egg i.p.	IFN-γ ↓, IL-4 ↑, TGF-β ↑, IL-10 ↑, dependent on STAT6	[28]	
			Sj	SEA i.p.	IFN-γ ↓, IL-4 ↑	[29]
	Type 1 diabetes in NOD mice	Sm	Infection, Egg i.p.	Inhibition of Ab class switch (anti-insulin IgG ↓)	[30]	
				SEA, SWA i.p.	NKT ↑	[31]
			SEA i.p.	Treg ↑	[32]	
	Streptozotocin-induced diabetes (multiple low dose)	Sm	Infection		[33]	
	TNBS-induced colitis	Sm	Infection	IL-2 ↑, IL-4 ↑	[34]	
				Egg i.p.	IFN-γ ↓, IL-4 ↑, dependent on STAT6	[35]
			SWA i.p.	IFN-γ ↓, IL-17 ↓, TGF-β ↑, IL-10 ↑	[36]	
				Sj	Egg i.p.	IFN-γ ↓, IL-4 ↑, IL-10 ↑, Treg ↑
			Egg i.p.	IFN-γ ↓, IL-4 ↑, TLR4 ↓	[38]	
	DSS-induced colitis	Sm	Infection (male)	Dependent on macrophages, independent of Treg, IL-10, IL-4, IL-13 and TGF-β	[39]	
				Infection	TNF-α ↓	[40]
Grave's hyperthyroidism	Sm	Infection	Anti-TSHR IgG2a ↓	[41]		

↓: down-regulation, ↑: up-regulation, Sm: *S. mansoni*, Sj: *S. japonicum*.

also in other helminthic infections, that is, piroxicam-induced colitis was suppressed by *Heligmosomoides polygyrus* [48], and EAE was suppressed by *Fasciola hepatica* [47], both in IL-10-deficient mice. In the latter study, however, TGF-β was shown to be responsible for anti-EAE effects of the worms [47]. Taken together, the involvements of IL-4, IL-10, and TGF-β in antiautoimmune effects of helminths depend on the disease models and helminth species. Further investigations using various autoimmunity models and gene-targeted

animals would be necessary for comprehensive elucidation of the suppressive mechanisms by regulatory cytokines in schistosome infections.

Regarding the participation of regulatory cell populations in schistosome-induced antiautoimmune effects, Treg cells, macrophages, and other types of cells (e.g., NKT cells) have been suggested. Although Treg cell population is known to expand by schistosome infection or SEA administration [13, 32], their involvement seems to depend on the disease

models. For instance, Cooke et al. have been studying anti-diabetic effects of *S. mansoni* using a spontaneous T1D model (NOD mouse) [30], and they demonstrated that Treg cells were essential in the suppression of T1D by cell transfer experiments [32]. The authors (Zacone et al.) showed that splenocytes from nontreated NOD mice successfully transmitted diabetes into NOD/SCID recipients, whereas splenocytes from SEA-treated NOD mice had a reduced capacity to transmit diabetes. They also showed that SEA had various immunomodulatory effects on dendritic cells (DCs), macrophages, and T cells of NOD mice [12, 32]: for example, increased expressions of TGF- $\beta$ , galectins, PD-L1, and so forth. In contrast to this T1D model, Smith et al. showed that depletion of Treg cells did not influence the suppressive effect of schistosomes on DSS-induced colitis [39]. They demonstrated that macrophages (but NOT alternatively activated macrophages) played an essential role in the amelioration of the colitis.

In some of the studies in Table 1, the injection of eggs or SEA was not effective [27, 39, 40]. Likewise, in our study of CIA, SEA injection was not effective [52]. Taken together with findings that mice infected with male worms become resistant to DSS-induced colitis [39], egg-derived substances are not sufficient to explain all of the immunomodulatory activities of schistosomes. Indeed, injection of soluble worm antigens (SWAs) could prevent T1D in NOD mice [31] and TNBS-induced colitis [36]. Therefore, differential effects of worms and eggs should be further elucidated for a precise understanding of the immunomodulatory mechanisms of schistosomes. In addition, further studies on differences between single-sex infections and mixed-sex infections may be necessary.

**2.3. Parasitic Infections.** In tropical developing countries, infections with multiple microbes/parasites are common. Consequently, the immune responses and/or pathological lesions caused by one pathogen may affect the outcome of other infections. Therefore, influences of parasitic infections on concomitant diseases have been studied. In this section, we focus on the effects of schistosomes on other parasitic infections. Malaria is the world's deadliest and most widely distributed parasitic disease. Consequently, there are more than a few experimental studies on the interrelationships between schistosomiasis and malaria (Table 2). The influence of schistosome infections depends on the species of the malarial parasites and mouse strains used in the experiments. For instance, in CBA/Ca mice, *S. mansoni* protected against *P. chabaudi* infection, worsened *P. yoelii* infection, and had no effect on *P. berghei* infection [53]. Even when the same parasite, *P. chabaudi*, was used, schistosomes exacerbated the parasitemia and mortality in C57BL/6 mice but ameliorated the outcome in A/J mice [54, 55]. Although these complicated outcomes should be taken into consideration, in general, schistosome infections seem to exacerbate rodent malaria, that is, increase of parasitemia, mortality, and hepatosplenomegaly [56]. Detrimental effects of schistosome infections on malarial outcome are also reported in humans [57, 58]. In Kenyan school children, even a light *S. mansoni* infection

was shown to exacerbate hepatosplenomegaly of malarial patients [57]. In a study in Senegal, heavy *S. mansoni* infections significantly increased the incidence of malarial attacks [58]. However, the modification of antibody responses by concomitant schistosomiasis is controversial [59, 60]. The mechanism responsible for the exacerbation (or amelioration) of malaria is clear in neither mice nor humans. Helmby et al. [54] suggested diminished production of TNF- $\alpha$  in schistosome-infected mice to have contributed to the increase in parasitemia of *P. chabaudi*. Yoshida et al. extensively analyzed possible mechanisms of protective effects of schistosomes against *P. chabaudi* in A/J mice [55]. They observed an enhanced Th1 response to *P. chabaudi* in schistosome-infected mice and that an anti-IFN- $\gamma$  antibody abrogated the schistosome-induced protective effect against *P. chabaudi*. As monoinfection with *S. mansoni* induced a Th2-dominant response, upregulation of IFN- $\gamma$  seemed to derive from the mixed infections of both parasites. They also observed the upregulation of iNOS gene expression in spleen from mice with mixed infections, suggesting an increase of splenic nitric oxide production to have contributed to the protection from *P. chabaudi*. Although the mechanism of the exacerbation of rodent malaria by schistosome is yet to be sufficiently analyzed, Treg cells have been shown to play an important role in the exacerbation of *P. yoelii* infection by preceding *H. polygyrus* infection [61]. Likewise, the induction/expansion of Treg cells by schistosome might be involved in the increased susceptibility to rodent malaria. The Th1 immune response protects against malarial parasites, but it also plays pathological roles in malaria, especially cerebral malaria [62]. Thus, schistosome as a representative Th2-biasing helminth is expected to suppress brain pathology. Indeed, in a model of cerebral malaria using mice infected with *P. berghei* ANKA, *S. mansoni* reduced the incidence of cerebral malaria and delayed death [63]. Moreover, the administration of IPSE/alpha-1, a Th2-inducing schistosomal egg protein [9, 10], also delayed death. Bucher et al. reported that brain pathology was reduced in schistosome-infected mice although they did not observe any beneficial effect on mortality [64].

Influence of schistosomes on other protozoan infections has also been investigated (Table 2). Regarding *Leishmania major*, reports vary, with Yoshida et al. finding no effects of schistosome on the outcome of *L. major* infection [67], and La Flamme et al. reporting an exacerbation of experimental leishmaniasis [68]. The reason for the discrepancy is not clear but might be intensity of the *S. mansoni* infection (20 cercariae in the former versus 70 cercariae in the latter). In the case of *L. donovani*, schistosome-preinfected mice failed to control growth of the protozoa in the liver [70]. In the coinfecting mice, hepatic egg granulomas were shown to provide a favorable microenvironment for the growth of the amastigotes. Likewise, *S. mansoni* exacerbated *Trypanosoma cruzi* infection [71] and *Toxoplasma gondii* infection [72]. In general, schistosome infections seem to be detrimental to animals infected with protozoan parasites.

In marked contrast to that for protozoan parasites, protective immunity against intestinal helminths is usually Th2 dependent. Therefore, the Th2-dominant environment

TABLE 2: Effects of schistosome on other parasitic infections in rodents.

Category	Parasites	Schistosome	Mouse strain	Effects	Refs.	
Malaria	<i>Plasmodium berghei</i>	Sm	CBA/Ca	No effect	[53]	
	<i>Plasmodium berghei</i> ANKA	Sm	Swiss albino	Parasitemia ↑, mortality ↑	[65]	
	<i>Plasmodium berghei</i> NK65	Sm	BALB/c	Parasitemia ↑, mortality ↑	[66]	
	<i>Plasmodium chabaudi</i>	Sm	CBA/Ca	Parasitemia ↓	[53]	
	<i>Plasmodium chabaudi</i> AS	Sm	A/J	Mortality ↓	[55]	
			C57BL/6	Parasitemia ↑, mortality ↑		
			Sm	C57BL/6	Parasitemia ↑	[54]
		<i>Plasmodium yoelii</i>	Sm	CBA/Ca	Parasitemia ↑	[53]
			Sm	BALB/c	Parasitemia ↑, mortality ↑	[56]
	Cerebral Malaria	<i>Plasmodium berghei</i> ANKA	Sm	ICR HSD	Cerebral malaria ↓	[63]
		Sm	C57BL/6	Brain pathology ↓, no effect on mortality	[64]	
Other protozoan infections	<i>Leishmania major</i>	Sm	BALB/c	No effect	[67]	
			C57BL/6	No effect		
			Sm	C57BL/6	Parasitemia ↑, lesion resolution delayed	[68]
	<i>Leishmania mexicana</i>	Sm	Outbred	Incubation period shortened	[69]	
	<i>Leishmania donovani</i>	Sm	C57BL/6	Parasitemia ↑	[70]	
	<i>Trypanosoma cruzi</i>	Sm	Albino	Parasitemia ↑, mortality ↑	[71]	
	<i>Toxoplasma gondii</i>	Sm	Albino	Mortality ↑	[72]	
Helminthic infections	<i>Hymenolepis diminuta</i>	Sm	NMRI	Expulsion ↑	[73]	
	<i>Strongyloides venezuelensis</i>	Sm	C57BL/6	Migration ↓	[67]	
		Sj	C57BL/6	Migration ↓, expulsion ↑	[69]	
	<i>Trichuris muris</i>	Sm	AKR	Expulsion ↑	[74]	

↓: downregulation, ↑: upregulation; Sm: *S. mansoni*, Sj: *S. japonicum*.

produced by schistosomes can be expected to protect against intestinal helminths. Indeed, as summarized in Table 2, schistosome infections protected mice from intestinal helminths. Likewise, in a human study in Brazil [75], *S. mansoni* egg counts were inversely correlated with *A. lumbricoides* and *T. trichiura*. (One exception was *Ancylostoma*, which was positively correlated with schistosome infections). Although protection against intestinal helminths is commonly observed in experimental settings, the mechanisms differ in each case. For instance, protection against a lung-migratory parasite, *Strongyloides venezuelensis*, mainly involved eosinophil-mediated killing of larvae in the lungs [76]. In addition, intestinal mucosal mastocytosis induced by *S. japonicum* infection rendered mice resistant to harboring of adult worms [76]. Antigen cross-reactivity between schistosomes and *S. venezuelensis* [67, 76] may have also contributed to the protection. In contrast, in the case of *Hymenolepis diminuta* and *Trichuris muris* (nonmigratory intestinal helminths), the protection seems to be dependent on accelerated expulsion from the intestines [73, 74]. Overall, schistosome infections appear to be beneficial to animals infected with intestinal helminths.

**2.4. Bacterial Infections and Vaccinations.** Prolonged bacteremia in schistosomiasis patients was first reported more

than 50 year ago, and the relationship between enterobacteria infections and schistosomiasis has been long studied [3]. The prolonged enterobacterial infection is referred to “prolonged septicemic enterobacteriosis.” The schistosome-induced exacerbation of enterobacterial infections has also been observed in experimental settings [77, 78]. As most studies were conducted before the “molecular immunology age,” the mechanisms of exacerbation of bacterial infections have not been sufficiently elucidated. However, the mechanisms are likely similar to those responsible for the schistosome-induced increase in susceptibility to protozoan parasites, that is, a reduction in Th1-dependent protective immunity as a consequence of augmented Th2 responses. Indeed, a Th1-inducing protozoan parasite, *L. donovani*, did not affect the growth of *Salmonella paratyphi* A in infected hamsters [78]. Moreover, impairment of the bactericidal function of macrophages from schistosome-infected mice was reported [79]. In addition to the immunomodulation by schistosomes, there are direct schistosome-bacteria interactions providing worm bodies as foci for bacterial multiplication [3, 80, 81]. Another considerable influence of schistosomes on bacterial infections is a reduction in vaccine efficacy. The *Mycobacterium bovis* BCG-induced protective response against *Mycobacterium tuberculosis* in mice was reduced by *S. mansoni* infection [82]. The authors also reported increased susceptibility to intravenous BCG

inoculations and lung pathology in *S. mansoni*-infected mice [83]. In these studies, decrease in IFN- $\gamma$  and nitric oxide in response to PPD were observed.

**2.5. Viral Infections and Vaccinations.** As hepatotropic viruses, that is, hepatitis B virus (HBV) and hepatitis C virus (HCV), cause liver cirrhosis during chronic infections, the synergistic exacerbation of hepatic pathology is expectable outcome of concurrent infections of HBV/HCV and schistosomes. Because of a lack of suitable animal models for infections of these hepatotropic viruses, major findings in schistosome-HBV/HCV coinfections have been obtained from epidemiological studies. Regarding HBV, schistosomiasis (especially the severe hepatosplenic form) was correlated with a higher frequency of HBV infection [3, 84, 85]. This observation could be explained by an increased susceptibility to HBV caused by schistosome infections. On the other hand, there are reports of no relationship between schistosomiasis and HBV [86, 87]. In addition, experiments with animals do not support increased susceptibility to HBV in schistosomiasis. Some evidence comes from an experiment using woodchucks infected with both schistosomes and woodchuck hepatitis virus (WHV) [88]. As HBV and WHV belong to the same family (family Hepadnaviridae), a concurrent infection by schistosomes and WHV in woodchucks is a good model of concurrent infections of schistosomes and HBV in humans. The authors reported no impact of the schistosome infection on WHV serum markers. Another paper on HBV transgenic mice [89] reported an inhibition of HBV replication during schistosome infection. In that study, the antiviral effects of schistosomes were attributed to IFN- $\gamma$  and nitric oxide. Overall, it seems premature to conclude the presence of certain positive or negative effects of schistosomes on HBV infection. Regarding HBV vaccines (both serum derived and recombinant), they were immunogenic in schistosomiasis patients although reduced responses to vaccination were observed in hepatosplenic schistosomiasis [90–92].

In contrast to the controversial effects on HBV infections, detrimental effects of schistosomes on HCV infections have been clearly demonstrated, that is, schistosomes weaken anti-HCV immune responses and worsen liver disease. According to studies in Egypt, patients with coinfections were characterized by a more advanced liver pathology, greater viral burden, higher levels of anti-HCV antibodies, and progression to chronic hepatitis [93–95]. Moreover, schistosomiasis was shown to be inversely correlated with HCV-specific CD4<sup>+</sup> T cells, CD8<sup>+</sup> T cells, and/or Th1 cytokine responses [95–100]. In addition to the modulatory effects of schistosomes on HCV-specific immune responses, SEA of *S. mansoni* [101] and *S. haematobium* [102] were shown to enhance *in vitro* viral replication in a hepatoblastoma cell line (HepG2) and peripheral blood mononuclear cells (PBMCs), respectively. Likewise, in other viral infections, schistosomes were shown to suppress specific CTL and cytokine responses and to prevent viral clearance [103–107]. It is interesting that the granulomas in *S. mansoni*-infected mice provide a microenvironment suitable for viral expansion [107], as in the case of a hepatic infection with the protozoan parasite *L. donovani* [70].

### 3. Concluding Remarks

Based on the experimental and epidemiological findings reviewed here, it can be concluded that schistosome infections are generally beneficial to patients with intestinal helminth infections and detrimental to patients with bacterial, viral, or protozoan infections. In most tropical or subtropical countries where schistosomiasis is endemic, more serious infectious diseases (e.g., HIV/AIDS, tuberculosis, and malaria) are also endemic. Therefore, control of schistosomiasis (especially infections of *S. mansoni* and *S. haematobium*) has been given low priority compared to control of such infectious diseases. However, if the detrimental bystander effects of schistosomes on concomitant bacterial, viral, or protozoan infections are properly considered, the importance of controlling schistosomiasis should be more emphasized.

In 2002, J. F. Bach summarized epidemiological trends of allergic and autoimmune diseases during recent several decades in developed countries [108]. According to the paper, the prevalence of the immunological disorders such as asthma, T1D, multiple sclerosis (MS), and Crohn's disease (CD) was increasing, whereas the prevalence of infectious diseases such as rheumatic fever, hepatitis A, tuberculosis, mumps, and measles was decreasing. These phenomena may be explained by the "hygiene hypothesis" in which microbial and helminthic infections prevent immunological disorders. Along with this hypothesis, schistosome infections are expected to prevent or alleviate symptoms of immunological disorders. According to the experimental studies until now, antiallergic and antiautoimmune effects of schistosomes are also plausible in humans. However, schistosomes could not be directly used for therapeutic treatment because of their pathogenicity. Instead, purified immunomodulatory products or recombinant proteins could be tested for clinical use. Indeed, considerable numbers of helminths' products have been shown to protect against experimental immunological disorders [109]. The immunomodulators exert their effects via Toll-like receptors (TLRs) and/or C-type lectin receptors (CLRs) [109]. Taken together with the finding that systemic administration of TLR agonists could prevent experimental autoimmunity and allergy [110], appropriately synthesized TLR agonists may be able to mimic or replace the prophylactic or therapeutic effects of schistosomes.

### Acknowledgments

Our studies introduced in this paper were supported by Grants-in-Aid for Scientific Research from the Japan Society for Promotion of Science (nos. 20590435 and 22590386), and Health and Labour Sciences Research Grants from the Ministry of Health, Labour and Welfare of Japan (H22-Kokui-Shitei-004). We acknowledge the technical assistance of Ms. Yukari Horie and the clerical assistance of Ms. Tomoko Fujitomi.

### References

- [1] P. J. Hotez, D. A. P. Bundy, K. Beegle et al., "Helminth infections: soil-transmitted helminth infections and schistosomiasis," in *Disease Control Priorities in Developing Countries*,

- D. T. Jamison, J. G. Breman, A. R. Measham et al., Eds., chapter 24, World Bank, Washington, DC, USA, 2nd edition, 2006.
- [2] B. J. Vennervald and K. Polman, "Helminths and malignancy," *Parasite Immunology*, vol. 31, no. 11, pp. 686–696, 2009.
- [3] P. P. Chieffi, "Interrelationship between schistosomiasis and concomitant diseases," *Memorias do Instituto Oswaldo Cruz*, vol. 87, pp. 291–296, 1992.
- [4] Y. Osada, T. Kumagai, K. Masuda, T. Suzuki, and T. Kanazawa, "Mutagenicity evaluation of *Schistosoma* spp. extracts by the *umu*-test and V79/HGPRT gene mutation assay," *Parasitology International*, vol. 54, no. 1, pp. 29–34, 2005.
- [5] E. F. Kjetland, P. D. Ndhlovu, E. Gomo et al., "Association between genital schistosomiasis and HIV in rural Zimbabwean women," *AIDS*, vol. 20, no. 4, pp. 593–600, 2006.
- [6] J. M. Grzych, E. Pearce, A. Cheever et al., "Egg deposition is the major stimulus for the production of Th2 cytokines in murine schistosomiasis *mansoni*," *Journal of Immunology*, vol. 146, no. 4, pp. 1322–1327, 1991.
- [7] B. Everts, G. Perona-Wright, H. H. Smits et al., "Omega-1, a glycoprotein secreted by *Schistosoma mansoni* eggs, drives Th2 responses," *Journal of Experimental Medicine*, vol. 206, no. 8, pp. 1673–1680, 2009.
- [8] S. Steinfeldt, J. F. Andersen, J. L. Cannons et al., "The major component in schistosome eggs responsible for conditioning dendritic cells for Th2 polarization is a T2 ribonuclease ( $\omega$ -1)," *Journal of Experimental Medicine*, vol. 206, no. 8, pp. 1681–1690, 2009.
- [9] G. Schramm, F. H. Falcone, A. Gronow et al., "Molecular characterization of an interleukin-4-inducing factor from *Schistosoma mansoni* eggs," *Journal of Biological Chemistry*, vol. 278, no. 20, pp. 18384–18392, 2003.
- [10] G. Schramm, K. Mohrs, M. Wodrich et al., "Cutting edge: IPSE/  $\alpha$ -1, a glycoprotein from *Schistosoma mansoni* eggs, induces IgE-dependent, antigen-independent IL-4 production by murine basophils *in vivo*," *Journal of Immunology*, vol. 178, no. 10, pp. 6023–6027, 2007.
- [11] S. Donnelly, C. M. Stack, S. M. O'Neill, A. A. Sayed, D. L. Williams, and J. P. Dalton, "Helminth 2-Cys peroxiredoxin drives Th2 responses through a mechanism involving alternatively activated macrophages," *FASEB Journal*, vol. 22, no. 11, pp. 4022–4032, 2008.
- [12] P. Zaccane, O. T. Burton, S. Gibbs et al., "Immune modulation by *Schistosoma mansoni* antigens in NOD mice: effects on both innate and adaptive immune systems," *Journal of Biomedicine and Biotechnology*, vol. 2010, Article ID 795210, 2010.
- [13] J. J. Taylor, M. Mohrs, and E. J. Pearce, "Regulatory T cell responses develop in parallel to Th responses and control the magnitude and phenotype of the Th effector population," *Journal of Immunology*, vol. 176, no. 10, pp. 5839–5847, 2006.
- [14] P. G. Fallon and N. E. Mangan, "Suppression of TH2-type allergic reactions by helminth infection," *Nature Reviews Immunology*, vol. 7, no. 3, pp. 220–230, 2007.
- [15] S. Amu, S. P. Saunders, M. Kronenberg, N. E. Mangan, A. Atzberger, and P. G. Fallon, "Regulatory B cells prevent and reverse allergic airway inflammation via Foxp3-positive T regulatory cells in a murine model," *Journal of Allergy and Clinical Immunology*, vol. 125, no. 5, pp. 1114.e8–1124.e8, 2010.
- [16] N. E. Mangan, R. E. Fallon, P. Smith, N. Van Rooijen, A. N. McKenzie, and P. G. Fallon, "Helminth infection protects mice from anaphylaxis via IL-10-producing B cells," *Journal of Immunology*, vol. 173, no. 10, pp. 6346–6356, 2004.
- [17] H. H. Smits, H. Hammad, M. van Nimwegen et al., "Protective effect of *Schistosoma mansoni* infection on allergic airway inflammation depends on the intensity and chronicity of infection," *Journal of Allergy and Clinical Immunology*, vol. 120, no. 4, pp. 932–940, 2007.
- [18] L. G. G. Pacífico, F. A. V. Marinho, C. T. Fonseca et al., "*Schistosoma mansoni* antigens modulate experimental allergic asthma in a murine model: a major role for CD4+ CD25+ Foxp3+ T cells independent of interleukin-10," *Infection and Immunity*, vol. 77, no. 1, pp. 98–107, 2009.
- [19] L. S. Cardoso, S. C. Oliveira, A. M. Góes et al., "*Schistosoma mansoni* antigens modulate the allergic response in a murine model of ovalbumin-induced airway inflammation," *Clinical and Experimental Immunology*, vol. 160, no. 2, pp. 266–274, 2010.
- [20] N. E. Mangan, N. Van Rooijen, A. N. J. McKenzie, and P. G. Fallon, "Helminth-modified pulmonary immune response protects mice from allergen-induced airway hyperresponsiveness," *Journal of Immunology*, vol. 176, no. 1, pp. 138–147, 2006.
- [21] H. M. Mo, J. H. Lei, Z. W. Jiang et al., "*Schistosoma japonicum* infection modulates the development of allergen-induced airway inflammation in mice," *Parasitology Research*, vol. 103, no. 5, pp. 1183–1189, 2008.
- [22] P. Liu, J. Li, X. Yang et al., "Helminth infection inhibits airway allergic reaction and dendritic cells are involved in the modulation process," *Parasite Immunology*, vol. 32, no. 1, pp. 57–66, 2010.
- [23] J. Yang, J. Zhao, Y. Yang et al., "*Schistosoma japonicum* egg antigens stimulate CD4+ CD25+ T cells and modulate airway inflammation in a murine model of asthma," *Immunology*, vol. 120, no. 1, pp. 8–18, 2007.
- [24] Y. Osada, S. Shimizu, T. Kumagai, S. Yamada, and T. Kanazawa, "*Schistosoma mansoni* infection reduces severity of collagen-induced arthritis via down-regulation of pro-inflammatory mediators," *International Journal for Parasitology*, vol. 39, no. 4, pp. 457–464, 2010.
- [25] Y. He, J. Li, W. Zhuang et al., "The inhibitory effect against collagen-induced arthritis by *Schistosoma japonicum* infection is infection stage-dependent," *BMC Immunology*, vol. 11, article 28, 2010.
- [26] X. Sun, Y. H. Liu, Z. Y. Lv et al., "rSj16, a recombinant protein of *Schistosoma japonicum*-derived molecule, reduces severity of the complete Freund's adjuvant-induced arthritis in rats' model," *Parasite Immunology*, vol. 32, no. 11–12, pp. 739–748, 2010.
- [27] A. C. La Flamme, K. Ruddenklau, and B. T. Bäckström, "Schistosomiasis decreases central nervous system inflammation and alters the progression of experimental autoimmune encephalomyelitis," *Infection and Immunity*, vol. 71, no. 9, pp. 4996–5004, 2003.
- [28] D. Sewell, Z. Qing, E. Reinke et al., "Immunomodulation of experimental autoimmune encephalomyelitis by helminth ova immunization," *International Immunology*, vol. 15, no. 1, pp. 59–69, 2003.
- [29] X. Zheng, X. Hu, G. Zhou et al., "Soluble egg antigen from *Schistosoma japonicum* modulates the progression of chronic progressive experimental autoimmune encephalomyelitis via Th2-shift response," *Journal of Neuroimmunology*, vol. 194, no. 1–2, pp. 107–114, 2008.
- [30] A. Cooke, P. Tonks, F. M. Jones et al., "Infection with *Schistosoma mansoni* prevents insulin dependent diabetes mellitus in non-obese diabetic mice," *Parasite Immunology*, vol. 21, no. 4, pp. 169–176, 1999.

- [31] P. Zacccone, Z. Feheérvári, F. M. Jones et al., “*Schistosoma mansoni* antigens modulate the activity of the innate immune response and prevent onset of type 1 diabetes,” *European Journal of Immunology*, vol. 33, no. 5, pp. 1439–1449, 2003.
- [32] P. Zacccone, O. Burton, N. Miller, F. M. Jones, D. W. Dunne, and A. Cooke, “*Schistosoma mansoni* egg antigens induce Treg that participate in diabetes prevention in NOD mice,” *European Journal of Immunology*, vol. 39, no. 4, pp. 1098–1107, 2009.
- [33] H. S. El-Wakil, T. S. Aboushousha, O. El Haddad, N. B. Gamil, T. Mansour, and H. El-Said, “Effect of *Schistosoma mansoni* egg deposition on multiple low doses streptozotocin induced insulin dependent diabetes,” *Journal of the Egyptian Society of Parasitology*, vol. 32, no. 3, pp. 987–1002, 2002.
- [34] T. G. Moreels, R. J. Nieuwendijk, J. G. De Man et al., “Concurrent infection with *Schistosoma mansoni* attenuates inflammation induced changes in colonic morphology, cytokine levels, and smooth muscle contractility of trinitrobenzene sulphonic acid induced colitis in rats,” *Gut*, vol. 53, no. 1, pp. 99–107, 2004.
- [35] D. E. Elliott, J. Li, A. Blum et al., “Exposure to schistosome eggs protects mice from TNBS-induced colitis,” *American Journal of Physiology*, vol. 284, no. 3, pp. G385–G391, 2003.
- [36] N. E. Ruysers, B. Y. De Winter, J. G. De Man et al., “Therapeutic potential of helminth soluble proteins in TNBS induced colitis in mice,” *Inflammatory Bowel Diseases*, vol. 15, no. 4, pp. 491–500, 2009.
- [37] H. M. Mo, W. Q. Liu, J. H. Lei, Y. L. Cheng, C. Z. Wang, and Y. L. Li, “*Schistosoma japonicum* eggs modulate the activity of CD4+ CD25+ Tregs and prevent development of colitis in mice,” *Experimental Parasitology*, vol. 116, no. 4, pp. 385–389, 2007.
- [38] Y. Zhao, S. Zhang, L. Jiang, J. Jiang, and H. Liu, “Preventive effects of *Schistosoma japonicum* ova on trinitrobenzenesulfonic acid-induced colitis and bacterial translocation in mice,” *Journal of Gastroenterology and Hepatology*, vol. 24, no. 11, pp. 1775–1780, 2009.
- [39] P. Smith, N. E. Mangan, C. M. Walsh et al., “Infection with a helminth parasite prevents experimental colitis via a macrophage-mediated mechanism,” *Journal of Immunology*, vol. 178, no. 7, pp. 4557–4566, 2007.
- [40] P. Bodammer, G. Waitz, M. Loebermann et al., “*Schistosoma mansoni* infection but not egg antigen promotes recovery from colitis in outbred NMRI mice,” *Digestive Diseases and Sciences*, vol. 56, no. 1, pp. 70–78, 2011.
- [41] Y. Nagayama, K. Watanabe, M. Niwa, S. M. McLachlan, and B. Rapoport, “*Schistosoma mansoni* and  $\alpha$ -galactosylceramide: prophylactic effect of Th1 immune suppression in a mouse model of Graves’ hyperthyroidism,” *Journal of Immunology*, vol. 173, no. 3, pp. 2167–2173, 2004.
- [42] M. Medeiros, J. P. Figueiredo, M. C. Almeida et al., “*Schistosoma mansoni* infection is associated with a reduced course of asthma,” *Journal of Allergy and Clinical Immunology*, vol. 111, no. 5, pp. 947–951, 2003.
- [43] R. R. Oliveira, K. J. Gollob, J. P. Figueiredo et al., “*Schistosoma mansoni* infection alters co-stimulatory molecule expression and cell activation in asthma,” *Microbes and Infection*, vol. 11, no. 2, pp. 223–229, 2009.
- [44] J. Feary, J. Britton, and J. Leonardi-Bee, “Atopy and current intestinal parasite infection: a systematic review and meta-analysis,” *Allergy*, vol. 66, no. 4, pp. 569–578, 2011.
- [45] J. Leonardi-Bee, D. Pritchard, and J. Britton, “Asthma and current intestinal parasite infection: systematic review and meta-analysis,” *American Journal of Respiratory and Critical Care Medicine*, vol. 174, no. 5, pp. 514–523, 2006.
- [46] H. Park, Z. Li, X. O. Yang et al., “A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17,” *Nature Immunology*, vol. 6, no. 11, pp. 1133–1141, 2005.
- [47] K. P. Walsh, M. T. Brady, C. M. Finlay, L. Boon, and K. H. G. Mills, “Infection with a helminth parasite attenuates autoimmunity through TGF- $\beta$ -mediated suppression of Th17 and Th1 responses,” *Journal of Immunology*, vol. 183, no. 3, pp. 1577–1586, 2009.
- [48] D. E. Elliott, A. Metwali, J. Leung et al., “Colonization with *Heligmosomoides polygyrus* suppresses mucosal IL-17 production,” *Journal of Immunology*, vol. 181, no. 4, pp. 2414–2419, 2008.
- [49] M. El-Behi, A. Rostami, and B. Ciric, “Current views on the roles of Th1 and Th17 cells in experimental autoimmune encephalomyelitis,” *Journal of Neuroimmune Pharmacology*, vol. 5, no. 2, pp. 189–197, 2010.
- [50] F. Mutapi, N. Imai, N. Nausch et al., “Schistosome infection intensity is inversely related to auto-reactive antibody levels,” *PLoS ONE*, vol. 6, no. 5, article e19149, 2011.
- [51] D. Rahima, R. Tarrab-Hazdai, M. Blank, R. Arnon, and Y. Shoenfeld, “Anti-nuclear antibodies associated with schistosomiasis and anti-schistosomal antibodies associated with SLE,” *Autoimmunity*, vol. 17, no. 2, pp. 127–141, 1994.
- [52] Y. Osada and T. Kanazawa, “Parasitic helminths: new weapons against immunological disorders,” *Journal of Biomedicine & Biotechnology*, vol. 2010, Article ID 743758, 9 pages, 2010.
- [53] M. Lwin, C. Last, G. A. T. Targett, and M. J. Doenhoff, “Infection of mice concurrently with *Schistosoma mansoni* and rodent malarias: contrasting effects of patent *S. mansoni* infections on *Plasmodium chabaudi*, *P. yoelii* and *P. berghei*,” *Annals of Tropical Medicine and Parasitology*, vol. 76, no. 3, pp. 265–273, 1982.
- [54] H. Helmby, M. Kullberg, and M. Troye-Blomberg, “Altered immune responses in mice with concomitant *Schistosoma mansoni* and *Plasmodium chabaudi* infections,” *Infection and Immunity*, vol. 66, no. 11, pp. 5167–5174, 1998.
- [55] A. Yoshida, H. Maruyama, T. Kumagai et al., “*Schistosoma mansoni* infection cancels the susceptibility to *Plasmodium chabaudi* through induction of type 1 immune responses in A/J mice,” *International Immunology*, vol. 12, no. 8, pp. 1117–1125, 2000.
- [56] D. Sangweme, C. Shiff, and N. Kumar, “*Plasmodium yoelii*: adverse outcome of non-lethal *P. yoelii* malaria during co-infection with *Schistosoma mansoni* in BALB/c mouse model,” *Experimental Parasitology*, vol. 122, no. 3, pp. 254–259, 2009.
- [57] S. Wilson, B. J. Vennervald, H. Kadzo et al., “Hepatosplenomegaly in Kenyan schoolchildren: exacerbation by concurrent chronic exposure to malaria and *Schistosoma mansoni* infection,” *Tropical Medicine and International Health*, vol. 12, no. 12, pp. 1442–1449, 2007.
- [58] C. Sokhna, J. Y. Le Hesran, P. A. Mbaye et al., “Increase of malaria attacks among children presenting concomitant infection by *Schistosoma mansoni* in Senegal,” *Malaria Journal*, vol. 3, p. 43, 2004.
- [59] T. O. Diallo, F. Remoue, L. Gaayeb et al., “Schistosomiasis co-infection in children influences acquired immune response against *Plasmodium falciparum* Malaria antigens,” *PLoS ONE*, vol. 5, no. 9, Article ID 12764, pp. 1–7, 2010.
- [60] D. Courtin, A. Djilali-Saïah, J. Milet et al., “*Schistosoma haematobium* infection affects *Plasmodium falciparum*-specific IgG responses associated with protection against malaria,” *Parasite Immunology*, vol. 33, no. 2, pp. 124–131, 2011.

- [61] K. Tetsutani, K. Ishiwata, H. Ishida et al., "Concurrent infection with *Heligmosomoides polygyrus* suppresses anti-*Plasmodium yoelii* protection partially by induction of CD4+CD25+Foxp3+ Treg in mice," *European Journal of Immunology*, vol. 39, no. 10, pp. 2822–2830, 2009.
- [62] N. H. Hunt and G. E. Grau, "Cytokines: accelerators and brakes in the pathogenesis of cerebral malaria," *Trends in Immunology*, vol. 24, no. 9, pp. 491–499, 2003.
- [63] J. H. Waknine-Grinberg, D. Gold, A. Ohayon et al., "*Schistosoma mansoni* infection reduces the incidence of murine cerebral malaria," *Malaria Journal*, vol. 9, no. 1, article 5, 2010.
- [64] K. Bucher, K. Dietz, P. Lackner et al., "*Schistosoma* co-infection protects against brain pathology but does not prevent severe disease and death in a murine model of cerebral malaria," *International Journal for Parasitology*, vol. 41, no. 1, pp. 21–31, 2011.
- [65] M. Legesse, B. Erko, and F. Balcha, "Increased parasitaemia and delayed parasite clearance in *Schistosoma mansoni* and *Plasmodium berghei* co-infected mice," *Acta Tropica*, vol. 91, no. 2, pp. 161–166, 2004.
- [66] R. F. Laranjeiras, L. C. C. Brant, A. C. L. Lima, P. M. Z. Coelho, and E. M. Braga, "Reduced protective effect of *Plasmodium berghei* immunization by concurrent *Schistosoma mansoni* infection," *Memorias do Instituto Oswaldo Cruz*, vol. 103, no. 7, pp. 674–677, 2008.
- [67] A. Yoshida, H. Maruyama, Y. Yabu, T. Amano, T. Kobayakawa, and N. Ohta, "Immune responses against protozoal and nematodal infection in mice with underlying *Schistosoma mansoni* infection," *Parasitology International*, vol. 48, no. 1, pp. 73–79, 1999.
- [68] A. C. La Flamme, P. Scott, and E. J. Pearce, "Schistosomiasis delays lesion resolution during *Leishmania* major infection by impairing parasite killing by macrophages," *Parasite Immunology*, vol. 24, no. 7, pp. 339–345, 2002.
- [69] P. M. Z. W. Coelho Mayrink, M. Dias, and L. H. Pereira, "Susceptibility to *Leishmania mexicana* of mice infected with *Schistosoma mansoni*," *Transactions of the Royal Society of Tropical Medicine and Hygiene*, vol. 74, no. 1, p. 141, 1980.
- [70] M. F. Hassan, Y. Zhang, C. R. Engwerda, P. M. Kaye, H. Sharp, and Q. D. Bickle, "The *Schistosoma mansoni* hepatic egg granuloma provides a favorable microenvironment for sustained growth of *Leishmania donovani*," *American Journal of Pathology*, vol. 169, no. 3, pp. 943–953, 2006.
- [71] K. Kloetzel, J. Faleiros, S. R. Mendes, C. T. Stanley, and H. S. Arias, "Concomitant infection of albino mice by *Trypanosoma cruzi* and *Schistosoma mansoni* parasitological parameters," *Transactions of the Royal Society of Tropical Medicine and Hygiene*, vol. 67, no. 5, pp. 652–658, 1973.
- [72] K. Kloetzel, P. P. Chieffi, J. J. Faleiros, and T. J. Merluzzi Filho, "Mortality and other parameters of concomitant infections in albino mice: the *Schistosoma-Toxoplasma* model," *Tropical and Geographical Medicine*, vol. 29, no. 4, pp. 407–410, 1977.
- [73] J. Andreassen, A. B. Odaibo, and N. O. Christensen, "Antagonistic effects of *Schistosoma mansoni* on superimposed *Hymenolepis diminuta* and *H. microstoma* infections in mice," *Journal of Helminthology*, vol. 64, no. 4, pp. 337–339, 1990.
- [74] A. J. Curry, K. J. Else, F. Jones, A. Bancroft, R. K. Grencis, and D. W. Dunne, "Evidence that cytokine-mediated immune interactions induced by *Schistosoma mansoni* alter disease outcome in mice concurrently infected with *Trichuris muris*," *Journal of Experimental Medicine*, vol. 181, no. 2, pp. 769–774, 1995.
- [75] M. Chamone, C. A. Marques, G. S. Atuncar, A. L. A. Pereira, and L. H. Pereira, "Are there interactions between schistosomes and intestinal nematodes?" *Transactions of the Royal Society of Tropical Medicine and Hygiene*, vol. 84, no. 4, pp. 557–558, 1990.
- [76] H. Maruyama, Y. Osada, A. Yoshida et al., "Protective mechanisms against the intestinal nematode *Strongyloides venezuelensis* in *Schistosoma japonicum*-infected mice," *Parasite Immunology*, vol. 22, no. 6, pp. 279–286, 2000.
- [77] A. L. Njunda and J. P. Oyerinde, "*Salmonella typhi* infection in *Schistosoma mansoni* infected mice," *West African Journal of Medicine*, vol. 15, no. 1, pp. 24–30, 1996.
- [78] I. A. Mikhail, G. I. Higashi, N. S. Mansour et al., "*Salmonella paratyphi* A in hamsters concurrently infected with *Schistosoma mansoni*," *American Journal of Tropical Medicine and Hygiene*, vol. 30, no. 2, pp. 385–393, 1981.
- [79] M. I. Muniz-Junqueira, A. Prata, and C. E. Tosta, "Phagocytic and bactericidal function of mouse macrophages to *Salmonella typhimurium* in schistosomiasis mansoni," *American Journal of Tropical Medicine and Hygiene*, vol. 46, no. 2, pp. 132–136, 1992.
- [80] I. A. Mikhail, G. I. Higashi, D. C. Edman, and S. H. Elwan, "Interaction of *Salmonella paratyphi* A and *Schistosoma mansoni* in hamsters," *American Journal of Tropical Medicine and Hygiene*, vol. 31, no. 2, pp. 328–334, 1982.
- [81] R. F. Melhem and P. T. LoVerde, "Mechanism of interaction of *Salmonella* and *Schistosoma* species," *Infection and Immunity*, vol. 44, no. 2, pp. 274–281, 1984.
- [82] D. Elias, H. Akuffo, A. Pawlowski, M. Haile, T. Schön, and S. Britton, "*Schistosoma mansoni* infection reduces the protective efficacy of BCG vaccination against virulent *Mycobacterium tuberculosis*," *Vaccine*, vol. 23, no. 11, pp. 1326–1334, 2005.
- [83] D. Elias, H. Akuffo, C. Thors, A. Pawlowski, and S. Britton, "Low dose chronic *Schistosoma mansoni* infection increases susceptibility to *Mycobacterium bovis* BCG infection in mice," *Clinical and Experimental Immunology*, vol. 139, no. 3, pp. 398–404, 2005.
- [84] H. A. Hammad, M. M. Bd El Fattah, M. Moris, E. H. Madina, A. A. El Abbasy, and A. T. M. Soliman, "Study on some hepatic functions and prevalence of hepatitis B surface antigenaemia in Egyptian children with schistosomal hepatic fibrosis," *Journal of Tropical Pediatrics*, vol. 36, no. 3, pp. 126–127, 1990.
- [85] H. M. Al-Freihi, "Prevalence of hepatitis B surface antigenemia among patients with *Schistosoma mansoni*," *Annals of Saudi Medicine*, vol. 13, no. 2, pp. 121–125, 1993.
- [86] B. Larouze, M. C. Dazza, C. Gaudebout, M. Habib, M. Elamy, and B. Cline, "Absence of relationship between *Schistosoma mansoni* and hepatitis B virus infection in the Qalyub Governorate, Egypt," *Annals of Tropical Medicine and Parasitology*, vol. 81, no. 4, pp. 373–375, 1987.
- [87] X. P. Ye, Y. L. Fu, R. M. Anderson, and D. J. Nokes, "Absence of relationship between *Schistosoma japonicum* and hepatitis B virus infection in the Dongting lake region, China," *Epidemiology and Infection*, vol. 121, no. 1, pp. 193–195, 1998.
- [88] Z. A. Andrade, P. Berthillon, R. Paraná, J. A. Grimaud, and C. Trépo, "Schistosomiasis mansoni and viral B hepatitis in woodchucks," *Journal of Hepatology*, vol. 34, no. 1, pp. 134–139, 2001.
- [89] H. McClary, R. Koch, F. V. Chisari, and L. G. Guidotti, "Inhibition of hepatitis B virus replication during *Schistosoma mansoni* infection in transgenic mice," *Journal of Experimental Medicine*, vol. 192, no. 2, pp. 289–294, 2000.

- [90] S. Bassily, K. C. Hyams, N. M. El-Ghorab, M. M. Mansour, N. A. El-Masry, and M. A. Dunn, "Immunogenicity of hepatitis B vaccine in patients infected with *Schistosoma mansoni*," *American Journal of Tropical Medicine and Hygiene*, vol. 36, no. 3, pp. 549–553, 1987.
- [91] S. Bassily, K. C. Hyams, N. El-Ghorab, and N. A. El-Masry, "Safety and immunogenicity of a recombinant hepatitis B vaccine in patients infected with *Schistosoma mansoni*," *American Journal of Tropical Medicine and Hygiene*, vol. 42, no. 5, pp. 449–452, 1990.
- [92] S. Bassily, G. T. Strickland, M. F. Abdel-Wahab et al., "Efficacy of hepatitis B vaccination in primary school children from a village endemic for *Schistosoma mansoni*," *Journal of Infectious Diseases*, vol. 166, no. 2, pp. 265–268, 1992.
- [93] S. Kamal, M. Madwar, L. Bianchi et al., "Clinical, virological and histopathological features: long-term follow-up in patients with chronic hepatitis C co-infected with *S. mansoni*," *Liver*, vol. 20, no. 4, pp. 281–289, 2000.
- [94] A. R. el-Zayadi, O. Selim, E. H. Ibrahim et al., "Does schistosomiasis play a role in the high sero prevalence of HCV antibody among Egyptians?" *Tropical Gastroenterology*, vol. 18, no. 3, pp. 98–100, 1997.
- [95] S. M. Kamal, J. W. Rasenack, L. Bianchi et al., "Acute hepatitis C without and with schistosomiasis: correlation with hepatitis C-specific CD4+ T-cell and cytokine response," *Gastroenterology*, vol. 121, no. 3, pp. 646–656, 2001.
- [96] S. M. Kamal, C. S. Graham, Q. He et al., "Kinetics of intrahepatic hepatitis C virus (HCV)-specific CD4+ T cell responses in HCV and *Schistosoma mansoni* coinfection: relation to progression of liver fibrosis," *Journal of Infectious Diseases*, vol. 189, no. 7, pp. 1140–1150, 2004.
- [97] S. M. Kamal, L. Bianchi, A. Al Tawil et al., "Specific cellular immune response and cytokine patterns in patients coinfecting with hepatitis C virus and *Schistosoma mansoni*," *Journal of Infectious Diseases*, vol. 184, no. 8, pp. 972–982, 2001.
- [98] A. Farid, M. Al-Sherbiny, A. Osman et al., "*Schistosoma* infection inhibits cellular immune responses to core HCV peptides," *Parasite Immunology*, vol. 27, no. 5, pp. 189–196, 2005.
- [99] I. M. El-Kady, M. Lotfy, G. Badraf, S. El-Masry, and I. Waked, "Interleukin (IL)-4, IL-10, IL-18 and IFN- $\gamma$  cytokines pattern in patients with combined hepatitis C virus and *Schistosoma mansoni* infections," *Scandinavian Journal of Immunology*, vol. 61, no. 1, pp. 87–91, 2005.
- [100] M. Elrefaei, N. El-Sheikh, K. Kamal, and H. Cao, "HCV-specific CD27- CD28- memory T cells are depleted in hepatitis C virus and *Schistosoma mansoni* co-infection," *Immunology*, vol. 110, no. 4, pp. 513–518, 2003.
- [101] M. M. Bahgat, M. A. E. El-Far, A. A. Mesalam et al., "*Schistosoma mansoni* soluble egg antigens enhance HCV replication in mammalian cells," *Journal of Infection in Developing Countries*, vol. 4, no. 4, pp. 226–234, 2010.
- [102] M. K. El-Awady, S. S. Youssef, M. H. Omran, A. A. Tabll, W. T. El Garf, and A. M. Salem, "Soluble egg antigen of *Schistosoma haematobium* induces HCV replication in PBMC from patients with chronic HCV infection," *BMC Infectious Diseases*, vol. 6, article 91, 2006.
- [103] J. K. Actor, M. Shirai, M. C. Kullberg, R. M. L. Buller, A. Sher, and J. A. Berzofsky, "Helminth infection results in decreased virus-specific CD8+ cytotoxic T- cell and Th1 cytokine responses as well as delayed virus clearance," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 90, no. 3, pp. 948–952, 1993.
- [104] M. A. Marshall, D. Jankovic, V. Ellen Maher, A. Sher, and J. A. Berzofsky, "Mice infected with *Schistosoma mansoni* develop a novel non-T-lymphocyte suppressor population which inhibits virus-specific CTL induction via a soluble factor," *Microbes and Infection*, vol. 3, no. 13, pp. 1051–1061, 2001.
- [105] M. J. Edwards, O. Buchatska, M. Ashton, M. Montoya, Q. D. Bickle, and P. Borrow, "Reciprocal immunomodulation in a schistosome and hepatotropic virus coinfection model," *Journal of Immunology*, vol. 175, no. 10, pp. 6275–6285, 2005.
- [106] A. A. Da'Dara, N. Lautsch, T. Dudek et al., "Helminth infection suppresses T-cell immune response to HIV-DNA-based vaccine in mice," *Vaccine*, vol. 24, no. 24, pp. 5211–5219, 2006.
- [107] J. K. Actor, M. A. Marshall, I. A. Eltoum, R. M. L. Buller, J. A. Berzofsky, and A. Sher, "Increased susceptibility of mice infected with *Schistosoma mansoni* to recombinant vaccinia virus: association of viral persistence with egg granuloma formation," *European Journal of Immunology*, vol. 24, no. 12, pp. 3050–3056, 1994.
- [108] J. F. Bach, "The effect of infections on susceptibility to autoimmune and allergic diseases," *New England Journal of Medicine*, vol. 347, no. 12, pp. 911–920, 2002.
- [109] W. Harnett and M. M. Harnett, "Helminth-derived immunomodulators: can understanding the worm produce the pill?" *Nature Reviews Immunology*, vol. 10, no. 4, pp. 278–284, 2010.
- [110] A. Aumeunier, F. Grela, A. Ramadan et al., "Systemic toll-like receptor stimulation suppresses experimental allergic asthma and autoimmune diabetes in NOD mice," *PLoS ONE*, vol. 5, no. 7, Article ID e11484, 2010.



**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

