Letter to the Editor

Comment on “Emerging Functions of Transcription Factors in Malaria Parasite”

Heather J. Painter and Manuel Llinás

Department of Molecular Biology and Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ 08544-1014, USA

Correspondence should be addressed to Manuel Llinás, manuel@genomics.princeton.edu

Received 19 January 2012; Accepted 18 March 2012

Copyright © 2012 H. J. Painter and M. Llinás. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In a recent Journal of Biomedicine and Biotechnology special issue on Immunology and Cell Biology of Parasitic Disease (2011), Tuteja et al. [1] authored a review summarizing transcription factors in the malaria parasite, *Plasmodium*. It is well known that there are very few characterized transcriptional regulators in the malaria parasite [2–4]. To date, the sole family of transcriptional regulators in *Plasmodium* consists of a conserved group of proteins containing a DNA-interaction domain with high homology to the *Arabidopsis* APETELA2 (AP2) DNA-binding domain [5]. Related AP2-integrase DNA-binding domains are also present in various *Tetrahymena* species, a few viruses, and cyanobacteria (reviewed in [6]). However, there have been no reports of an AP2 expansion in any other eukaryote other than the Apicomplexans. This lineage-specific expansion is now known as the Apicomplexan AP2 (ApiAP2) protein family [5], and since these proteins represent the first family of putative specific transcriptional regulators in the malaria parasite, their characterization has generated a flurry of recent reports [7–10]. The authors of this review, unfortunately, report an incorrect association between the ApiAP2 proteins (pfam PF00847) and the Activator Protein-2 (AP-2) (pfam PF03299) found in higher eukaryotes (reviewed in [11]). Despite the similarity in nomenclature, there is absolutely no evolutionary conservation (homology) or functional relationship between the mammalian AP-2 and the malarial ApiAP2 proteins as the authors suggest. The authors also incorrectly cite a recent in-depth review of the ApiAP2 protein family as a source for this information [12]. As an international journal with a diverse readership, it is pertinent

that this misleading information is corrected so as to prevent further confusion.

References

- [1] R. Tuteja, A. Ansari, and V. S. Chauhan, “Emerging functions of transcription factors in malaria parasite,” *Journal of Biomedicine and Biotechnology*, vol. 2011, Article ID 461979, 6 pages, 2011.
- [2] L. Aravind, L. M. Iyer, T. E. Wellems, and L. H. Miller, “Plasmodium biology: genomic gleanings,” *Cell*, vol. 115, no. 7, pp. 771–785, 2003.
- [3] T. J. Templeton, L. M. Iyer, V. Anantharaman et al., “Comparative analysis of apicomplexa and genomic diversity in eukaryotes,” *Genome Research*, vol. 14, no. 9, pp. 1686–1695, 2004.
- [4] R. M. R. Coulson, N. Hall, and C. A. Ouzounis, “Comparative genomics of transcriptional control in the human malaria parasite *Plasmodium falciparum*,” *Genome Research*, vol. 14, no. 8, pp. 1548–1554, 2004.
- [5] S. Balaji, M. Madan Babu, L. M. Iyer, and L. Aravind, “Discovery of the principal specific transcription factors of Apicomplexa and their implication for the evolution of the AP2-integrase DNA binding domains,” *Nucleic Acids Research*, vol. 33, no. 13, pp. 3994–4006, 2005.
- [6] S. R. Wessler, “Homing into the origin of the AP2 DNA binding domain,” *Trends in Plant Science*, vol. 10, no. 2, pp. 54–56, 2005.
- [7] M. Yuda, S. Iwanaga, S. Shigenobu, T. Kato, and I. Kaneko, “Transcription factor AP2-Sp and its target genes in malarial sporozoites,” *Molecular Microbiology*, vol. 75, no. 4, pp. 854–863, 2010.

- [8] M. Yuda, S. Iwanaga, S. Shigenobu et al., "Identification of a transcription factor in the mosquito-invasive stage of malaria parasites," *Molecular Microbiology*, vol. 71, no. 6, pp. 1402–1414, 2009.
- [9] C. Flueck, R. Bartfai, I. Niederwieser et al., "A major role for the *Plasmodium falciparum* ApiAP2 protein PfSIP2 in chromosome end biology," *PLoS Pathogens*, vol. 6, no. 2, Article ID e1000784, 2010.
- [10] T. L. Campbell, E. K. de Silva, K. L. Olszewski, O. Elemento, and M. Llinás, "Identification and Genome-Wide Prediction of DNA Binding Specificities for the ApiAP2 family of regulators from the malaria parasite," *PLoS Pathogens*, vol. 6, no. 10, Article ID e1001165, 2010.
- [11] D. Eckert, S. Buhl, S. Weber, R. Jäger, and H. Schorle, "The AP-2 family of transcription factors," *Genome Biology*, vol. 6, no. 13, article no. 246, 2005.
- [12] H. J. Painter, T. L. Campbell, and M. Llinás, "The Apicomplexan AP2 family: integral factors regulating *Plasmodium* development," *Molecular and Biochemical Parasitology*, vol. 176, no. 1, pp. 1–7, 2011.