Research Article

Serotype Distribution of *Salmonella* Isolates from Turkey Ground Meat and Meat Parts

Irfan Erol,¹ Muammer Goncuoglu,¹ Naim Deniz Ayaz,¹ Lüppo Ellerbroek,²
Fatma Seda Bilir Ormanci,¹ and Ozlem Iseri Kangal¹

¹ Ankara University, Faculty of Veterinary Medicine, Department of Food Hygiene and Technology, 06110 Diskapi, Ankara, Turkey

² Federal Institute for Risk Assessment (BfR) FGr 42-NRL for *Campylobacter*, Diedersdorfer Weg 1, 12277 Berlin, Germany

Correspondence should be addressed to Irfan Erol; erol@veterinary.ankara.edu.tr

Received 2 April 2013; Revised 10 June 2013; Accepted 16 June 2013

Academic Editor: Stanley Brul

Copyright © 2013 Irfan Erol et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The aim of the study was to find out the serotype distribution of 169 *Salmonella* colonies recovered from 112 *Salmonella* positive ground turkey (115 colonies) and 52 turkey meat parts (54 colonies). Out of 15 *Salmonella* serotypes: *S. Corvallis*, *S. Kentucky*, *S. Bredeney*, *S. Virchow*, *S. Saintpaul* and *S. Agona* were identified as the predominant serovars at the rates of 27%, 13%, 12%, 12%, 11%, and 10%, respectively. Other serotypes were below 6% of the total isolates. All *S. Kentucky* and *S. Virchow* and most of the *S. Corvallis* (39/46) and *S. Heidelberg* (9/9) serotypes were recovered from ground turkey. The results indicate that turkey ground meat and meat parts were contaminated with quite distinct *Salmonella* serotypes. This is the first study reporting *Salmonella* serotype distribution in turkey meat and *S. Corvallis* as predominant serotype in poultry meat in Turkey.

1. Introduction

Salmonella is one of the major food-borne pathogens and has an importance as a leading cause of food-borne bacterial diseases in humans throughout the world [1]. According to the Centers for Disease Control and Prevention (CDC), *Salmonella* spp. are causing 1.4 million food-borne illnesses, 15,000 hospitalizations, and 400 deaths annually in the United States [2]. The total costs of food-borne *Salmonella* infections of humans in the US have been estimated to 3.3 billion dollars per year [3]. According to the European Surveillance System 2010 data, total of 99,020 confirmed *Salmonella* cases were reported through 27 European Union member countries with a notification rate of 21.5 cases per 100,000 population [4]. Different studies indicate that, foods of animal origin, particularly poultry, cattle and pig are the major vehicles of diseases caused by food-borne pathogens. Among them turkey meat and products are attributed to be the important sources of food-borne salmonellosis [4].

Poultry processing plays an important role to increase the contamination rate of *Salmonella* in turkey meat [5, 6]. Scalding, defeathering, evisceration and cooling steps

in slaughtering are the critical points in contamination of carcass [5]. Also contamination of foods with this bacterium can occur at different processing line including distribution, marketing, handling and preparation both in processing plant or home. Therefore, turkey meat can easily be contaminated with *Salmonella* throughout the whole production chain [5, 7]. Nevertheless, *Salmonella* contamination in turkey flocks is generally asymptomatic and detection of the bacterium emerges by the randomly monitoring by the industry [6]. It was pointed out that serotype profiles of *Salmonella* in animal carcasses match with corresponding raw ground products [8] and findings of studies strengthened possibility of transmission of *Salmonella* to humans through the food chain [9].

The prevalence of *Salmonella* in turkey meat has been studied mainly in developed countries. The results of these studies showed that prevalence and serotype distribution of *Salmonella* was varied. The percentage of positive samples varied from zero to 40% in fresh turkey meat and higher than 5% in RTE turkey meat products in EU countries, the US and the Canada [5, 10–14]. The results of these studies showed that serotype profiles of the isolates were different.

Ground meat has high nutritional value and is useful for preparing foods. However, it is a suitable medium for growth of many pathogen and saprophyte microorganisms. Even if ground meat is originally contaminated at a low level with *Salmonella*, growth and/or cross-contamination may occur during storage and handling under poor hygienic conditions [15].

There are no published scientific data on serotype distribution of *Salmonella* spp. in turkey meat in Turkey. Therefore, this study aimed to determine the serotype distribution of *Salmonella* isolated from fresh turkey ground meat and meat parts obtained from retail markets in Turkey, to provide some scientific data for further epidemiological studies.

2. Material and Methods

2.1. Bacterial Strains. A total of 169 *Salmonella* spp. isolated from ground turkey and meat parts (meat cuts, breasts, and legs) purchased from different supermarkets located in Ankara between July 2004 and January 2006 were serotyped. Hundred and fifteen of the colonies were belonging to 112 *Salmonella* positive ground turkey meat samples that have been isolated in a previous study [16]. Additionally 54 colonies were obtained from 52 *Salmonella* spp. detected turkey meat part samples.

2.2. Isolation and Identification of *Salmonella* spp. For the isolation of *Salmonella* spp. standard cultivation technique was performed according to the Bacteriological Analytical Manual of the Food and Drug Administration [16, 17] and the International Organization for Standardization (ISO-6579) [18]. Twenty five grams of each sample was weighed into 225 mL of buffered peptone water (BPW, Oxoid CM0509, Hampshire, UK) for preenriching and shaken for about 2 min, then incubated at 37°C overnight. After incubation, 0.1, 1 and 1 mL of the preenrichment broths were added to 10 mL of Rappaport-Vassiliadis Enrichment broth (RV, Oxoid, CM669), 9 mL of Selenite Cystine broth (SC, Difco 112534 JC, Detroit, USA) and 9 mL of Tetrathionate (TT) broth (FDA, BAM), respectively. For selective enrichment, RV broth, SC broth and TT broth were incubated at 43°C, 37°C and 42°C for 24 h, respectively. After incubation period broths were streaked on to Brilliant-green Phenol-Red Lactose Sucrose agar (BPLS, Merck 1.07237, Darmstadt, Germany) and Xylose Lysine Deoxycholate (XLD) (Merck 1.05287) agar and incubated at 37°C for 18–24 h. Up to five typical grown colonies were picked and inoculated into Triple Sugar Iron Agar (TSIA, Oxoid, CM0277), Lysine Iron Agar (LIA, Oxoid CM0381), and Urea Broth Base (Oxoid, CM0071B) and incubated at 37°C for 24–48 h. TSIA positive, LIA positive, and urease negative colonies were considered as suspect for *Salmonella*. The agglutination test was done with Polyvalent *Salmonella* Antisera (Difco, Cat. No L840114-1, Detroit, USA). Suspect *Salmonella* colonies were mixed with a drop of antiserum on a slide. Agglutination with antiserum was accepted as a positive reaction for *Salmonella* spp. isolates that showed agglutination were stored at 4°C on Tryptone

Soya agar (TSA, Oxoid CM0131) and at –86°C (Sanyo MDF-U5186S, Japan) in cryovials for the PCR verification and serotyping.

2.3. PCR Confirmation of the *Salmonella* Isolates. In order to determine the origin of DNA replication *oriC* gene of *Salmonella* strains for the verification, PCR analysis were performed. For the PCR analysis *Salmonella* Typhimurium ATCC 14028 was used as positive control.

DNA Extraction. isolates stored at 4°C in Tryptone Soy Agar (TSA, Oxoid CM 131) were incubated in Brain Heart Infusion broth (BHI, Oxoid CM0225) at 37°C for 24 h. Then 1 mL of each culture was separately transferred to microcentrifuge tubes. All tubes were centrifuged (Eppendorf Centrifuge 5417R, Hamburg, Germany) for 15 min at 5000 rcf at 10°C. The pellets were resuspended in 1 mL sterile aquabidest. The suspensions were mixed by vortex (IKA MS1 Minishaker, Wilmington, USA). Then all tubes were centrifuged for 5 min at 5000 rcf at 10°C. The pellets were resuspended with 200 µL sterile aquabidest and incubated for 20 min at 95°C in a water bath (Memmert WB/OB 7–45, WBU 45, Schwabach, Germany) then cooled on ice.

PCR Analysis for the Detection of *oriC* Gene. the primers used are specific to the origin of DNA replication (*oriC*) on the *Salmonella* chromosome and produce a 163 bp DNA fragment (Primer 1: 5'-TTA TTA GGA TCG CGC CAG GC-3', Primer 2: 5'-AAA GAA TAA CCG TTG TTC AC-3'). PCR was performed with a final volume of 50 µL reaction mixture containing incomplete 5x PCR Buffer (Promega M7921, Madison USA), 1.5 mM MgCl₂ (Promega A3511), 200 mM each of the deoxynucleoside triphosphates (dNTPs, Promega U1420), 1 U Taq DNA polymerase (Promega M3005), 0.50 mM each of primer and 10 µL DNA. Thermal cycling (Biometra Personal Cycler, Goettingen, Germany) was carried out with the initial denaturation at 94°C for 1 min and then 35 cycles of denaturation at 94°C for 1 min, annealing at 53°C for 1 min, and extension at 72°C for 1 min, 72°C, 10 min for final extension [19–21]. A 10 µL aliquot of each PCR product was subjected to 1.5% agarose gel electrophoresis containing 0.1 mg/mL ethidium bromide for 1 h at 100 V (Biometra, Agagel, B15339). Amplicon visualisation and documentation was performed using gel documentation and analysis system (Syngene Ingenius, Cambridge, UK).

2.4. Serotyping. Serotyping of the *Salmonella* isolates was performed at the National Reference Centre for *Salmonella* and other bacterial enteric pathogens, Robert Koch Institute (RKI-Wernigerode) with the scheme of Kaufmann-White using lam agglutination and serum neutralization tests [22].

3. Results

Conventional cultivation technique was used for the isolation of *Salmonella* spp. from turkey ground meat and meat parts. All 169 colonies (112 from ground turkey and 54 from turkey

meat parts) were confirmed with PCR by detection of *oriC* gene. Using lam agglutination and serum neutralization tests, 15 different serotypes were identified among 169 *Salmonella* isolates. *S. Corvallis* (*n*: 46, 27%), *S. Kentucky* (*n*: 22, 13%), *S. Bredeney* (*n*: 20, 12%), *S. Virchow* (*n*: 20, 12%), *S. Saintpaul* (*n*: 18, 11%), and *S. Agona* (*n*: 17, 10%) were the most frequently isolated serotypes (Table 1). All 22 isolates of *S. Kentucky* and 20 isolates of *S. Virchow* were recovered from ground turkey meat. Likewise *S. Heidelberg*, *S. Stanleyville*, *S. Montevideo*, *S. subsp. I*, *S. group C*, and *S. Newport* were only recovered from ground turkey meat samples. As the predominant serotype 39 of 46 isolates of *S. Corvallis* were recovered from ground turkey meat. However other two major serotypes, 90% of *S. Bredeney* and 67% of *S. Saintpaul* isolates were recovered from turkey meat parts. Also, *S. Hadar*, *S. muenchen* and *S. Typhimurium* were only recovered from turkey meat parts samples (Table 1), although only at low numbers.

The seasonal distribution of *Salmonella* serotypes was determined as follows: 39 (23%), 26 (15%), 53 (31%), and 51 isolates (30%) during winter, spring, summer and autumn, respectively (Table 2). Most of the isolates were determined in warm months. In winter and spring 16 different serotypes, in summer and autumn 19 different serotypes were recovered. As a predominant serotype *S. Corvallis* was detected in spring, summer, autumn and second after *S. Bredeney* in winter months. Also, 86% of *S. Kentucky*, as the second major serotype of the study, was recovered in spring, summer and autumn.

Ground turkey meat samples were collected from nine different companies and no correlation was observed between producing companies and serotype profiles. Unlikely, meat part samples were collected from three different supermarkets (supermarkets A, B and C) and these meat parts were prepared in their own butchery stores. Based on the results of this study a relation as observed between supermarkets and serotype distribution of samples. As shown in Table 1, *S. Bredeney* and *S. Saintpaul* were the mostly recovered serotypes from turkey meat parts. Within the 78% of *S. Bredeney* and 58% of meat parts recoveries of *S. Saintpaul* serotypes were detected in supermarket A.

4. Discussion

In this study, out of 169 *Salmonella* turkey meat isolates 15 different serotypes were identified and showed quite distinct distribution. Among them *S. Corvallis* was found to be the predominant serotype. In a study, *S. Kentucky*, *S. Anatum* and *S. Heidelberg* were reported as the most frequently isolated serotypes from turkeys and their environments [23]. In Great Britain, *S. Kottbus*, *S. Kedougou*, *S. Derby*, *S. Senftenberg*, *S. Newport* and *S. Oslo* were the most common serotypes recovered from different types of turkey flocks [9]. *S. Typhimurium*, *S. Newport*, *S. Derby*, *S. Indiana* and *S. Agona* were the top five reported serovars in British turkey flocks between 1995 and 2006 [24]. According to the EFSA Scientific Report *S. Bredeney*, *S. Hadar*, *S. Derby*, *S. Saintpaul*, *S. Kottbus* and *S. Typhimurium* were the most frequently isolated serotypes in turkey flocks [25]. The contamination

TABLE 1: Number and percentage of *Salmonella* serotypes isolated from ground turkey and turkey meat parts.

Serotypes	Percent	Number of isolates				Total
		Ground turkey	Meat cuts	Breasts	Legs	
<i>S. Corvallis</i>	27	39	5	1	1	46
<i>S. Kentucky</i>	13	22	—	—	—	22
<i>S. Bredeney</i>	12	2	6	2	10	20
<i>S. Virchow</i>	12	20	—	—	—	20
<i>S. Saintpaul</i>	11	6	6	—	6	18
<i>S. Agona</i>	10	9	3	3	2	17
<i>S. Heidelberg</i>	5	9	—	—	—	9
<i>S. Hadar</i>	3	—	2	2	1	5
<i>S. Munchen</i>	2	—	—	2	1	3
<i>S. Stanleyville</i>	1	2	—	—	—	2
<i>S. Montevideo</i>	1	2	—	—	—	2
<i>S. subsp. I</i>	1	2	—	—	—	2
<i>S. group C</i>	1	1	—	—	—	1
<i>S. Typhimurium</i>	1	—	1	—	—	1
<i>S. Newport</i>	1	1	—	—	—	1
Total		115	23	10	21	169

TABLE 2: Seasonal distribution of *Salmonella* serotypes.

Serotypes	Winter	Spring	Summer	Autumn	Total
<i>S. Corvallis</i>	8	11	14	13	46
<i>S. Kentucky</i>	3	9	7	3	22
<i>S. Bredeney</i>	10	—	10	—	20
<i>S. Virchow</i>	—	1	1	18	20
<i>S. Saintpaul</i>	4	3	7	4	18
<i>S. Agona</i>	4	1	1	11	17
<i>S. Heidelberg</i>	2	—	7	—	9
<i>S. Hadar</i>	4	—	—	1	5
<i>S. Munchen</i>	2	—	—	1	3
<i>S. Stanleyville</i>	1	—	1	—	2
<i>S. Montevideo</i>	—	—	2	—	2
<i>S. subsp. I</i>	1	1	—	—	2
<i>S. group C</i>	—	—	1	—	1
<i>S. Typhimurium</i>	—	—	1	—	1
<i>S. Newport</i>	—	—	1	—	1
Total	39	26	53	51	169

of poultry meat by different *Salmonella* serotypes is straightly contact with the flock contamination by the slaughtering process and the contamination of carcasses, and meat mainly during evisceration, also the link between levels of *Salmonella* in the flocks and the slaughterhouse are well defined [8, 26]. Further researches should focus on the controlling the sources of contamination with *Salmonella* spp. and serotypes in turkey meat industry.

There were several reports about distribution of *Salmonella* serotypes in raw turkey meat samples. In turkey

meat in Germany, *S. Saintpaul*, *S. 1,4,(5),12:i-* and *S. Newport* were identified as the major serotypes [27]. A previous study conducted in Canada reported that *S. Heidelberg* and *S. Hadar* were the most common serotypes recovered from 91 turkey meat samples including drumstick, wing or ground turkey [14]. In a different study, *S. Newport*, *S. Hadar*, *S. Heidelberg*, *S. 4:12: nonmotile* and *S. Reading* were recovered from retail turkey meat samples in North Dakota, USA [10]. Khaita et al. [12] reported six *Salmonella* serotypes from 959 turkey products as follows: *Hadar*, *Heidelberg*, *Typhimurium* var. *Copenhagen*, *Newport*, *Saintpaul* and *Agona*. Most of the *Salmonella* serotypes recovered during this study are common isolates from turkey meat samples. Like previous reports, in this study *S. Enteritidis* and *S. Typhimurium* were not considered as the most frequently encountered serotypes in turkey meat samples [28]. However *Beli* et al. [29] reported *S. Enteritidis* as the major serotype isolated from turkey meat samples. Similar results arise about specific serotypes within other studies. A study from the USA has reported that *S. Saintpaul* was almost specific for turkey meat rather than chicken, beef and pork samples [11]. Although the serotype of concern was detected with 11% in turkey meat samples in this study, the results of another work conducted in our country showed that *S. Saintpaul* was not detected in any of broiler carcass and edible offal samples in 70 *Salmonella* positive isolates [30]. The number of turkey farms in our country is relatively low, as well as the number of turkey meat production plants. Therefore presence of a particular serotype in these limited number of facilities may cause a serovar as the most detected one over 2500 *Salmonella* serotypes. In addition serotype variation among these studies is probably due to contaminated feed, infected breeding flocks [24], differences in production system of turkey meat and contamination of meat at slaughterhouse process [8], variability in sampling of isolates from different sources [31], geographical features, socioeconomic and cultural differences between countries, national or international control and surveillance program differences, and also some yearly variations [32, 33].

There have been numerous reports of human salmonellosis due to consumption of different turkey products by different *Salmonella* serotypes including *S. Reading*, *S. Hadar*, *S. Agona*, *S. Saintpaul*, and *S. Typhimurium* [34–38]. All the *Salmonella* serotypes recovered from this study except serotype *Reading* were common in reported international human salmonellosis cases that were caused by consumption of contaminated turkey meat [39–41]. However these serotypes frequently recovered in this study have never been reported as an agent of human salmonellosis in Turkey. In the EU *S. Corvallis* and *S. Kentucky*, which are first two major isolates of this study, were reported from human salmonellosis cases [42, 43].

In conclusion, turkey meat can be contaminated with quite distinct serotypes. According to our results fifteen different *Salmonella* serotype were recovered and among them *S. Corvallis* was detected as a predominant serovar. These results showed that upcoming *Salmonella* monitoring programme should cover turkey meat production chain in Turkey.

References

- [1] J. Y. D'Aoust and J. Maurer, "Salmonella species," in *Food Microbiology: Fundamentals and Frontiers*, M. P. Doyle and L. R. Beuchat, Eds., pp. 187–236, ASM Press, Washington, DC, USA, 3rd edition, 2007.
- [2] A. C. Voetsch, T. J. Van Gilder, F. J. Angulo et al., "FoodNet estimate of the burden of illness caused by nontyphoidal *Salmonella* infections in the United States," *Clinical Infections and Disease*, vol. 38, supplement 3, pp. 127–134, 2004.
- [3] P. D. Frenzen, T. L. Riggs, J. C. Buzby et al., "Salmonella cost estimate updated using Foodnet data," *Food Review*, vol. 22, pp. 10–15, 1999.
- [4] EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control), "The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2010," *The EFSA Journal*, vol. 10, no. 2597, p. 442, 2012.
- [5] C. W. Nde, J. M. McEvoy, J. S. Sherwood, and C. M. Logue, "Cross contamination of turkey carcasses by *Salmonella* species during defeathering," *Poultry Science*, vol. 86, no. 1, pp. 162–167, 2007.
- [6] J. Danguy des Déserts, R. H. Davies, K. Vaughan et al., "A longitudinal study of *Salmonella* infection in different types of Turkey flocks in Great Britain," *Zoonoses and Public Health*, vol. 58, no. 3, pp. 200–208, 2011.
- [7] R. Nayak, P. B. Kenney, J. Keswani, and C. Ritz, "Isolation and characterisation of *Salmonella* in a turkey production facility," *British Poultry Science*, vol. 44, no. 2, pp. 192–202, 2003.
- [8] W. Schlosser, A. Hogue, E. Ebel et al., "Analysis of *Salmonella* serotypes from selected carcasses and raw ground products sampled prior to implementation of the Pathogen Reduction; Hazard Analysis and Critical Control Point Final Rule in the US," *International Journal of Food Microbiology*, vol. 58, no. 1–2, pp. 107–111, 2000.
- [9] S. Zhao, S. Qaiyumi, S. Friedman et al., "Characterization of *Salmonella enterica* serotype Newport isolated from humans and food animals," *Journal of Clinical Microbiology*, vol. 41, no. 12, pp. 5366–5371, 2003.
- [10] M. K. Fakhr, J. S. Sherwood, J. Thorsness, and C. M. Logue, "Molecular characterization and antibiotic resistance profiling of *Salmonella* isolated from retail turkey meat products," *Foodborne Pathogens and Disease*, vol. 3, no. 4, pp. 366–374, 2006.
- [11] S. Zhao, P. F. McDermott, S. Friedman et al., "Antimicrobial resistance and genetic relatedness among *Salmonella* from retail foods of animal origin: NARMS retail meat surveillance," *Foodborne Pathogens and Disease*, vol. 3, no. 1, pp. 106–117, 2006.
- [12] M. L. Khaita, R. B. Kegode, and D. K. Doetkott, "Occurrence of antimicrobial-resistant *Salmonella* species in raw and ready to eat Turkey meat products from retail outlets in the midwestern United States," *Foodborne Pathogens and Disease*, vol. 4, no. 4, pp. 517–525, 2007.
- [13] EFSA, "Scientific Opinion of the Panel on Biological Hazards. A quantitative microbiological risk assessment on *Salmonella* in meat: Source attribution for human salmonellosis from meat," *The EFSA Journal*, vol. 625, pp. 1–32, 2008.
- [14] M. Aslam, S. Checkley, B. Avery et al., "Phenotypic and genetic characterization of antimicrobial resistance in *Salmonella* serovars isolated from retail meats in Alberta, Canada," *Food Microbiology*, vol. 32, pp. 110–117, 2012.

- [15] I. Erol, "Incidence and serotype distribution of *Salmonella* in ground beef in Ankara," *Turkish Journal of Veterinary and Animal Sciences*, vol. 23, no. 4, pp. 321–325, 1999.
- [16] O. Iseri and I. Erol, "Incidence and antibiotic resistance of *Salmonella* spp. in ground turkey meat," *British Poultry Science*, vol. 51, no. 1, pp. 60–66, 2010.
- [17] FDA (Food and Drug Administration), *Salmonella*, chapter 5, Bacteriological Analytical Manual, 2003.
- [18] ISO (International Organization for Standardization), "Microbiology of food and animal feeding stuffs—Horizontal method for the detection of *Salmonella* spp.," Ref. No. ISO 6579:2002/Cor.1:2004(E), 2002.
- [19] M. N. Widjojoatmodjo, A. C. Fluit, R. Torensma, B. H. I. Keller, and J. Verhoef, "Evaluation of the Magnetic Immuno PCR Assay for rapid detection of *Salmonella*," *European Journal of Clinical Microbiology and Infectious Diseases*, vol. 10, no. 11, pp. 935–938, 1991.
- [20] A. C. Fluit, M. N. Widjojoatmodjo, A. T. A. Box, R. Torensma, and J. Verhoef, "Rapid detection of *Salmonella* in poultry with the magnetic immuno-polymerase chain reaction assay," *Applied and Environmental Microbiology*, vol. 59, no. 5, pp. 1342–1346, 1993.
- [21] I. Erol, G. Hildebrandt, J. Kleer, and A. Yurtyeri, "Kopplung von immunomagnetischer separation und polymerasekettenreaktion zum schnellachweis von salmonellen in hackfleisch und geflügelinnereien," *Berliner Und Munchener Tierärztliche Wochenschrift*, vol. 112, pp. 100–103, 1999.
- [22] M. Y. Popoff, "Antigenic formulas of the *Salmonella* serovars," in *WHO Collaborating Centre for Reference and Research on Salmonella*, Institute Pasteur, 2001.
- [23] D. W. Hird, H. Kinde, J. T. Case, B. R. Charlton, R. P. Chin, and R. L. Walker, "Serotypes of *Salmonella* isolated from California turkey flocks and their environment in 1984–89 and comparison with human isolates," *Avian Diseases*, vol. 37, no. 3, pp. 715–719, 1993.
- [24] C. Papadopolou, R. H. Davies, J. J. Carrique-Mas, and S. J. Evans, "*Salmonella* serovars and their antimicrobial resistance in british turkey flocks in 1995 to 2006," *Avian Pathology*, vol. 38, no. 5, pp. 349–357, 2009.
- [25] Anon, "Report of the task force on zoonoses data collection on the analysis of the baseline survey on the prevalence of *Salmonella* in turkey flocks, Part B," *The EFSA Journal*, vol. 198, pp. 1–224, 2008.
- [26] M. H. Rostagno, I. V. Wesley, D. W. Trampel, and H. S. Hurd, "*Salmonella* prevalence in market-age turkeys on-farm and at slaughter," *Poultry Science*, vol. 85, no. 10, pp. 1838–1842, 2006.
- [27] M. Hartung and A. Käsbohrer, "Erreger von zoonosen in deutschland im jahr 2009," *BfR-Wissenschaft*, vol. 1, pp. 1–273, 2012.
- [28] USDA-FSIS, "Serotypes profile of *Salmonella* isolates from meat and poultry products January 1998 through December 2010," 2011, http://www.fsis.usda.gov/PDF/Serotypes_Profile_Salmonella.2010.pdf.
- [29] E. Beli, A. Telo, and E. Duraku, "*Salmonella* serotypes isolated from turkey meat in Albania," *International Journal of Food Microbiology*, vol. 63, no. 1–2, pp. 165–167, 2001.
- [30] I. Erol, G. Hildebrandt, M. Goncuoglu, and J. Kleer, "Serotype distribution of *Salmonella* in broiler carcasses and edible offal in Turkey," *Fleischwirtschaft*, vol. 90, no. 9, pp. 106–109, 2010.
- [31] S. L. Foley, A. M. Lynne, and R. Nayak, "*Salmonella* challenges: prevalence in swine and poultry and potential pathogenicity of such isolates," *Journal of animal science*, vol. 86, no. 14, pp. E149–E162, 2008.
- [32] M. E. Doyle, C. Kaspar, J. Archer, and R. Klos, "White paper on human illness caused by *Salmonella* from all food and non-food vectors. FRIBriefings: *Salmonella* Human illness from food and non-food sources," Food Research Institute, UW Madison, 2008–2009, pp. 1–51, 2009.
- [33] R. S. Hendriksen, A. R. Vieira, S. Karlsmose et al., "Global monitoring of *Salmonella* serovar distribution from the world health organization global foodborne infections network country data bank: Results of quality assured laboratories from 2001 to 2007," *Foodborne Pathogens and Disease*, vol. 8, no. 8, pp. 887–900, 2011.
- [34] Anon, "Foodborne nosocomial outbreak of *Salmonella* reading—connecticut," *MMWR Morbidity and Mortality Weekly Report*, vol. 40, pp. 804–806, 1991.
- [35] S. P. Luby, J. L. Jones, and J. M. Horan, "A large salmonellosis outbreak associated with a frequently penalized restaurant," *Epidemiology and Infection*, vol. 110, no. 1, pp. 31–39, 1993.
- [36] D. L. Baggesen, H. C. Wegener, and J. P. Christensen, "Typing of *Salmonella* enterica serovar Saintpaul: an outbreak investigation," *APMIS*, vol. 104, no. 6, pp. 411–418, 1996.
- [37] T. Grein, D. O'Flanagan, T. McCarthy, and D. Bauer, "An outbreak of multidrug-resistant *Salmonella* Typhimurium food poisoning at a wedding reception," *Irish Medical Journal*, vol. 92, no. 1, pp. 238–241, 1999.
- [38] CDC, "Investigation announcement: multistate outbreak of *Salmonella* Hadar infections associated with turkey burgers," 2011, <http://www.cdc.gov/salmonella/hadar0411/040411/index.html>.
- [39] A. D. Aysev, H. Guriz, and B. Erdem, "Drug resistance of *Salmonella* strains isolated from community infections in Ankara, Turkey, 1993–99," *Scandinavian Journal of Infectious Diseases*, vol. 33, no. 6, pp. 420–422, 2001.
- [40] B. Erdem, S. Ercis, G. Hascelik et al., "Antimicrobial resistance patterns and serotype distribution among *Salmonella enterica* strains in Turkey, 2000–2002," *European Journal of Clinical Microbiology and Infectious Diseases*, vol. 24, no. 3, pp. 220–225, 2005.
- [41] B. Erdem, S. Ercis, G. Hascelik, D. Gur, and A. D. Aysev, "Antimicrobial resistance of *Salmonella enterica* group C strains isolated from humans in Turkey, 2000–2002," *International Journal of Antimicrobial Agents*, vol. 26, no. 1, pp. 33–37, 2005.
- [42] Anon, "International surveillance network for the enteric infections—*Salmonella*, VTEC O157 and *Campylobacter*," *Enter-net quarterly Salmonella report Oct-Dec 2004/4*, 2004.
- [43] Anon, "International surveillance network for the enteric infections—*Salmonella*, VTEC O157 and *Campylobacter*," *Enter-net quarterly Salmonella report Jul-Sept 2005/3*, 2005.