Research Article

Isolation and Molecular Identification of Keratinophilic Fungi from Public Parks Soil in Shiraz, Iran

Keyvan Pakshir, Moosa Rahimi Ghiasi, Kamiar Zomorodian, and Ali Reza Gharavi

Department of Parasitology and Mycology, Basic Sciences in Infectious Diseases Research Center, School of Medicine, Shiraz University of Medical Sciences, Shiraz 71345-45794, Iran

Correspondence should be addressed to Keyvan Pakshir; pakshirk@gmail.com

Received 30 April 2013; Accepted 27 June 2013

Academic Editor: Abdelwahab Omri

Copyright © 2013 Keyvan Pakshir et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction. Keratinophilic fungi are an important group of fungi that live in soil. The aim of this study was to isolate and identify keratinophilic fungi from the soil of different parks in Shiraz. **Materials and Methods.** A total of 196 soil samples from 43 parks were collected. Isolation of the fungi was performed by hair bait technique. The isolated colonies were identified by morphologic feature of macro- and microconidia and molecular method, using DNA sequence analysis. ITS region of ribosomal DNA was amplified and the PCR products were sequenced. **Results.** 411 isolates from 22 genera were identified. *Fusarium* (23.8%), *Chrysosporium* (13.13%), *Acremonium* (12.65%), *Penicillium* (12.39%), *Microsporum gypseum* (1.94%), *Bionectria ochroleuca* (1.21%), *Bipolaris spicifera* (1.21%), *Scedosporium apiospermum* (0.82%), *Phialophora reptans* (0.82%), *Cephalosporium curtipies* (0.49%), *Scedosporium dehoogii* (0.24%), *Ochroconis constricta* (0.24%), *Nectria mauritiicola* (0.49%), *Chaetomium* (0.49%), *Scopulariopsis* (0.24%), *Malbranchea* (0.24%), and *Tritirachium* (0.24%) were the most important isolates. Most of the fungi were isolated from the soils with the PH range of 7 to 8. **Conclusion.** Our study results showed that many keratinophilic fungi isolated from the parks soil are important for public health and children are an important group at a high risk of being exposed to these fungi.

1. Introduction

Keratinophilic fungi are ecologically an important group of fungi which could be found in soil [1]. Some groups of these fungi are causative agents of cutaneous fungal infections named dermatophytosis, and the other saprophyte fungi mainly represent hyalohyphomycosis [2, 3]. The prevalence of these fungi depends on different factors, such as the presence of creatinine in the soil, pH, and geographical location [1]. Some of these fungi, such as dermatophytes, are well known to cause tinea infections which could be transmitted from soil to humans. In general, soil could be considered as a reservoir for human infection. Forests, farmyards, park soils, and sediments of the rivers and oceans containing humus and organic materials are the best candidates for growth of keratinolytic and saprophytic fungi [4].

During the past years, many researchers reported about isolation of keratinophilic fungi around the world [5–14]. Also, a lot of reports were available about the isolation of geophilic dermatophytes and keratinophilic fungi from the soils of many parts of Iran and also dermatophytosis due to

the geophilic fungi during the last decade [15–18]. Nowadays, most people spend their time with their children in the parks for fun and are potentially at risk for direct contact with soil and being exposed to keratinophilic fungi [4]. Shiraz is one of the most populous cities of Iran and the capital of Fars province which is located in the southwest of Iran. It has a moderate to warm climate (–3 to 40°C) with rain falling about 350 mm/year, land area of approximately 225 km², and more than one million population. Up to now, there were no data about keratinophilic fungi in soil of Shiraz and no reports about the molecular identification of these fungi around Iran. Thus, the present study aimed to isolate and identify keratinophilic fungi from soil of the popular parks in Shiraz by molecular and morphological analysis.

2. Materials and Methods

2.1. Sample Collection. In this descriptive study, 196 soil samples were collected from various sites of 43 different parks around Shiraz during spring 2011. The samples were collected

from the superficial layer of the soil whose depth did not exceed 5–10 cm by using an iron spatula. In doing so, 300–400 gram of soil was collected in sterile polyethylene bags and brought to laboratory for further processing.

2.2. Measuring of Soil pH. PH of each soil sample was measured after preparation of soil suspension (one gram of soil to five mL deionized water) using pH meter [17, 19].

2.3. Isolation and Identification of the Isolates. We used Vanbreuseghem's hair bait technique for isolation of keratinophilic fungi [20]. Briefly, each soil sample was thoroughly mixed, and about 70 gram of the soil was packed in a sterile 90 mm Petri dish. Then, several pieces of sterile healthy children hair fragments were dispersed over the surface of the soil samples and moistened with sterile distilled water supplemented with antibiotic solutions, chloramphenicol (0.2 g/mL), and strepto-penicillin (1000 IU/mL). All the baited soil Petri dishes were incubated at room temperature (20–25°C) in the dark for 2 months and got moistened if necessary. After observation of colony growth around the hairs, the colonies were subcultured on Sabouraud's dextrose agar (SDA) with and without chloramphenicol (50 mg/L) and cycloheximide (500 mg/L) and purified. The fungi were identified based on the conventional method (colony morphology and macro- and microconidia characteristics) and DNA-based identification techniques.

2.4. DNA-Based Identification Techniques (DNA Sequence Analysis). Molecular identification of the unknown isolates was achieved by DNA sequence analysis. First, the fungi were grown in flasks containing Sabouraud's dextrose broth and incubated at 25°C for several days using shaker rotator. After colony growth, the culture was filtered (Millipore, USA) and the fungal mass was washed with distilled water for several times and stored in a freezer (–20°C) for further processing.

DNA was extracted by using Lee technique [21] with mild modification. First, the frozen mycelium mass was smashed by mechanical pressure using sterile pounder and liquid nitrogen. The acquired powder was then mixed with lyses buffer and the DNA was extracted. The ITS1–5.8S–ITS 2 rDNA was amplified using ITS1 and ITS4 as forward and reverse primers as described by White et al. [22]. Amplification was performed in 50 µL reaction volumes containing 5 µL of 10× buffer, MgCl₂ (25 mM) 1.5 µL, dNTP (10 mM) 0.5 µL, 0.5 µL of each 0.2 Mm primer (ITS1: 3'-TCC-GTA-GGT-GAA-CCT-GCG-G-5' and ITS4: 3'-TCC-TCC-GCT-TAT-TGA-TAT-GC-5'), Taq Polymerase (1.25 U) 0.5 µL, DNA sample 1 µL, and distilled water 40.5 µL. The PCR reaction was carried out using a Thermal Cycler (R corbet cgl-96) with the following conditions: denaturation at 94°C for 5 min, 34 cycles of (30 s at 94°C, 45 s at 56°C, and 45 s at 72°C) extension at 72°C for 7 min, and storage at 4°C. Negative controls were also used in each set of reactions. The final products were analyzed by electrophoresis on 1.2% agarose gel (Sigma) and stained with 0.5 µg mL⁻¹ ethidium bromide. In addition, the PCR products were sent for sequencing in both directions (Bioneer, Korea). The sequence results

FIGURE 1: Agarose gel electrophoresis and PCR products bands; lines 1 and 2 (marker), lines 3–12 (up), and 2–8 (down) are PCR products of many unknown fungi.

were processed by using the web-based blasting program, basic local alignment search tool (BLAST), at the NCBI site (<http://www.ncbi.nlm.nih.gov/BLAST>), and the data were compared with the NCBI/Genbank database [23].

2.5. Data Analysis. The study data were analyzed through Fisher exact test and Chi-square test.

3. Results

PCR products bands on gel agarose are presented in Figure 1. From the 196 soil samples, a total of 411 colonies of keratinophilic fungi were isolated from 43 parks. The fungal isolates belonged to 22 genera as follows: *Fusarium* (25.30%), *Penicillium* (13.13%), *Chrysosporium* (13.13%), *Acremonium* (12.65%), *Aspergillus* (11.92%), *Mucor* (9.48%), *Paeecilomyces* (4.13%), *Microsporum* (2.42%), *Bipolaris* (1.45%), *Bionectria* (1.21%), *Pseudallescheria* (0.73%), *Phialophora* (0.73%), *Alternaria* (0.73%), *Nectria* (0.48%), *Cephalosporium* (0.48%), *Chaetomium* (0.48%), *Scopulariopsis* (0.24%), *Scedosporium* (0.24%), *Verticillium* (0.24%), *Malbranchea* (0.24%), *Tritirachium* (0.24%), and *Ochroconis* (0.24%). More details about the species are presented in Table 1.

Fusarium spp. was the most common fungal isolate among the parks. Besides, eight species of *Microsporum gypseum* were isolated from four parks with soil pH between seven and nine and two species of *Microsporum fulvum* from one park with the same pH. Most of the fungi were isolated from the soil samples with pH between 7 and 8 (66.42%), and no colony growth was seen in pH > 9 as shown in Table 2. The study results revealed no significant correlation between soil pH and fungal species. More details about the isolates and soil pH are presented in Table 2.

4. Discussion

The keratinolytic activity of keratinophilic fungi is important for ecology and has attracted many researchers' attention around the world. Keratinophilic fungi play an important role in the natural degradation of keratinized residues in the soil [24, 25]. Some types of these fungi, such as geophilic dermatophytes, live in soil and could be transmitted to

TABLE 1: Frequency of keratinophilic fungi isolated from soils with different PH.

Fungal genus	pH							
	6-7		7-8		8-9		>9	
	n	%	n	%	n	%	n	%
<i>Aspergillus</i>	0	0	31	11.35	18	13.43	0	0
<i>Acremonium</i>	0	0	35	12.82	17	12.68	0	0
<i>Alternaria</i>	0	0	2	0.73	1	0.75	0	0
<i>Scopulariopsis</i>	0	0	1	0.36	0	0	0	0
<i>Ochroconis</i>	0	0	1	0.36	0	0	0	0
<i>Bipolaris</i>	0	0	4	1.46	2	1.49	0	0
<i>Bionectria</i>	0	0	4	1.46	1	0.75	0	0
<i>Cephalosporium</i>	0	0	1	0.36	1	0.75	0	0
<i>Paecilomyces</i>	2	50	7	2.56	8	5.97	0	0
<i>Scedosporium</i>	0	0	4	1.45	0	0	0	0
<i>Tritirachium</i>	0	0	1	0.36	0	0	0	0
<i>Penicillium</i>	2	50	36	13.1	16	11.94	0	0
<i>Fusarium</i>	0	0	66	24.1	38	28.35	0	0
<i>Phialophora</i>	0	0	2	0.73	1	0.75	0	0
<i>Chrysosporium</i>	0	0	39	14.28	15	11.19	0	0
<i>Chaetomium</i>	0	0	2	0.73	0	0	0	0
<i>Mucor</i>	0	0	22	8.42	16	11.94	0	0
<i>Microsporum</i>	0	0	8	3.66	0	0	0	0
<i>Malbranchea</i>	0	0	1	0.36	0	0	0	0
<i>Nectria</i>	0	0	2	0.73	0	0	0	0
<i>Verticillium</i>	0	0	1	0.36	0	0	0	0
Total	4	100	273	100	134	100	0	0

humans as well as animals and cause cutaneous fungal infections [26, 27]. Parks are among the popular public places for the people to spend their time and have fun with their family. Our study revealed the presence of keratinophilic fungi, such as dermatophytes, in the soil of Shiraz parks. *M. gypseum* is a frequent geophilic dermatophyte commonly distributed in soil worldwide. Although nondermatophyte fungi isolates were more common than dermatophytes in the present study, *M. gypseum* and *M. fulvum* were the main dermatophytes which were isolated from four parks. These parks potentially have a high risk for transmission of fungal infections. Previously, these fungi were isolated from the soil samples of different parts of Iran [28–31].

Some species of *Aspergillus*, *Fusarium solani*, and *Bipolaris spicifera* are the causative agents of mycotic keratitis [32]. Our study showed that the genus *Fusarium* was the first dominant fungus in the soil of Shiraz, which is in agreement with the other studies conducted on the issue. Moallaei et al. [16, 30] also reported that *Fusarium* was the most prevalent saprophyte in South and Razavi Khorasan Provinces. The second most common species isolates in our study were *Chrysosporium* and *Penicillium*. *Chrysosporium* species have been reported to be the causative agents of disseminated diseases [33]. *Chrysosporium tropicum* was reported from comb lesion in two different breeds of chicken in India [34].

In this study, molecular method was utilized for identification of keratinophilic fungi for the first time in Iran.

TABLE 2: Distribution frequency of keratinophilic fungi isolated from Shiraz parks soil.

Species	Number	Percent
<i>Fusarium</i> spp.	95	23.08
<i>Fusarium chlamydosporum</i>	4	0.97
<i>Fusarium oxysporum</i>	3	0.82
<i>Fusarium solani</i>	2	0.49
<i>Chrysosporium</i> spp.	54	13.13
<i>Acremonium</i> spp.	52	12.65
<i>Penicillium</i> spp.	51	12.39
<i>Penicillium crustosum</i>	2	0.49
<i>Penicillium palmarum</i>	1	0.24
<i>Aspergillus niger</i>	40	9.73
<i>Aspergillus fumigatus</i>	6	1.45
<i>Aspergillus sclerotiorum</i>	2	0.49
<i>Aspergillus flavus</i>	1	0.24
<i>Mucor</i> spp.	39	9.48
<i>Paecilomyces</i> spp.	17	4.13
<i>Microsporum gypseum</i>	8	1.94
<i>Microsporum fulvum</i>	2	0.49
<i>Bipolaris spicifera</i>	5	1.21
<i>Bipolaris</i> sp.	1	0.24
<i>Bionectria ochroleuca</i>	5	1.21
<i>Scedosporium apiospermum</i>	3	0.82
<i>Scedosporium dehoogii</i>	1	0.24
<i>Phialophora reptans</i>	3	0.82
<i>Alternaria solani</i>	2	0.49
<i>Alternaria alternata</i>	1	0.24
<i>Cephalosporium curtipes</i>	2	0.49
<i>Chaetomium</i> spp.	2	0.49
<i>Nectria mauritica</i>	2	0.49
<i>Scopulariopsis</i> sp.	1	0.24
<i>Verticillium</i> sp.	1	0.24
<i>Malbranchea</i> sp.	1	0.24
<i>Tritirachium</i> sp.	1	0.24
<i>Ochroconis constricta</i>	1	0.24
Total	411	100%

We could isolate and identify some genera of fungi, such as *Bionectria* spp., which is important in natural products and medicine. The genus *Bionectria* is endophytic and has great potential for medicinal and agricultural applications [35].

Bionectria species are known as a destructive mycoparasite and grow inside the fungal host hyphae. They are used as a biocontrol agent of plant-pathogenic fungi and are infrequently isolated from dead insects. Besides, they are known as a parasite of living nematodes, ticks, and myxomycetes [35, 36].

Tritirachium species are an insect pathogen whose natural habitats are soil and decaying plant materials. These fungi are occasionally isolated from head and nail infections [37, 38]. Furthermore, *Ochroconis* spp. are dematiaceous fungi that cause deep mycoses, such as chromomycosis, around the

world. There are many reports regarding the fungal diseases caused by this fungus [39–41].

Scedosporium apiospermum is a common soil fungus with a worldwide distribution. Environmental isolations have been made from sewage sludge, polluted streams, and manure of poultry and cattle. In Australia, Cooley et al. [42] reported 59 cases of scedosporiosis caused by *Scedosporium apiospermum* and *S. prolificans* in patients with underlying diseases. Invasive infections in normal patients are usually caused by traumatic implantation. So, fungal infections could acquire or spread through playgrounds in parks.

The current study investigated the relationship between the frequency of fungi and the soils pH. Böhme and Ziegler [19] reported the effect of the soil pH on the presence of keratinophilic fungi for the first time. Many researchers stated that keratinophilic fungi could not be found in the soils with low pH levels (3–4.5). Garg et al. [3] also reported that the acidic soils with pH = 5.9 were free of keratinophilic fungi. Moreover, Asahi et al. [43] demonstrated that keratinolytic enzymes were produced in pH of 6–9, and particularly the extracellular keratinase was active in pH = 9. In the present study, all the 411 keratinophilic fungi were isolated from the soils with pH between 6 and 9. We found 66.42%, 32.6%, and 0.97% keratinophilic fungi from the soil samples with pH of 7.01–8, 8.01–9, and 6–7, respectively. These findings have been confirmed by other studies as well. Overall, in this study, 47% of the places were contaminated with keratinophilic fungi and geophilic dermatophyte species isolated from the soils of four parks. These areas potentially have a high risk for causing cutaneous fungal infections in humans and animals and could be considered as a source of these infections.

Acknowledgments

This study was supported by the Research Vice-Chancellor, Grant no. 6000, of Shiraz University of Medical Sciences, Shiraz, Iran. Hereby, the authors would like to thank the Research Improvement Center of Shiraz University of Medical Sciences and Ms. A. Keivanshekouh for improving the use of English in the paper.

References

- [1] S. K. Deshmukh and S. A. Verekar, "The occurrence of dermatophytes and other keratinophilic fungi from the soils of Himachal Pradesh (India)," *Czech Mycology*, vol. 58, no. 1-2, pp. 117–122, 2006.
- [2] G. Pålsson, "Geophilic dermatophytes in the soil in Sweden. Studies on their occurrence and pathogenic properties," *Acta Veterinaria Scandinavica*, supplement 25, pp. 1–89, 1968.
- [3] A. P. Garg, S. Gandotra, K. G. Mukerji, and G. J. F. Pugh, "Ecology of keratinophilic fungi," *Proceedings: Plant Sciences*, vol. 94, no. 2-3, pp. 149–163, 1985.
- [4] M. S. Ali-Shtayeh and R. M. F. Jamous, "Keratinophilic fungi and related dermatophytes in polluted soil and water habitats," in *Biology of Dermatophytes and Other Keratinophilic Fungi*, R. K. S. Kushawaha and J. Guarro, Eds., pp. 51–59, Revista Iberoamericana de Micología, Bilbao, Spain, 2000.
- [5] S. K. Abdullah and D. A. Hassan, "Isolation of dermatophytes and other keratinophilic fungi from surface sediments of the Shatt Al-Arab river and its creeks at Basrah, Iraq," *Mycoses*, vol. 38, no. 3-4, pp. 163–166, 1995.
- [6] A. K. Gupta, J. E. Ryder, M. Chow, and E. A. Cooper, "Dermatophytosis: the management of fungal infections," *Skinmed*, vol. 4, no. 5, pp. 305–310, 2005.
- [7] S. K. Deshmukh, Q. A. Mandeel, and S. A. Verekar, "Keratinophilic fungi from selected soils of Bahrain," *Mycopathologia*, vol. 165, no. 3, pp. 143–147, 2008.
- [8] S. K. Deshmukh and S. A. Verekar, "Incidence of keratinophilic fungi from the soils of Vedanthangal Water Bird Sanctuary (India)," *Mycoses*, vol. 54, no. 6, pp. 487–490, 2011.
- [9] V. Filipello Marchisio, D. Curetti, C. Cassinelli, and C. Bordese, "Keratinolytic and keratinophilic fungi in the soils of Papua New Guinea," *Mycopathologia*, vol. 115, no. 2, pp. 113–119, 1991.
- [10] R. Papini, F. Mancianti, G. Grassotti, and G. Cardini, "Survey of keratinophilic fungi isolated from city park soils of Pisa, Italy," *Mycopathologia*, vol. 143, no. 1, pp. 17–23, 1998.
- [11] P. Anbu, A. Hilda, and S. C. B. Gopinath, "Keratinophilic fungi of poultry farm and feather dumping soil in Tamil Nadu, India," *Mycopathologia*, vol. 158, no. 3, pp. 303–309, 2004.
- [12] V. M. Ramesh and A. Hilda, "Incidence of keratinophilic fungi in the soil of primary schools and public parks of Madras city, India," *Mycopathologia*, vol. 143, no. 3, pp. 139–145, 1998.
- [13] G. M. Vidyasagar, N. Hosmani, and D. Shivkumar, "Keratinophilic fungi isolated from hospital dust and soils of public places at Gulbarga, India," *Mycopathologia*, vol. 159, no. 1, pp. 13–21, 2005.
- [14] P. A. Volz, M. J. Wlosinski, and S. P. Wasser, "Sparse diversity of potential pathogenic soil micro-fungi in the Ukraine," *Microbios*, vol. 65, no. 264–265, pp. 187–193, 1991.
- [15] M. T. Hedayati and M. Mirzakhani, "Survey of keratinophilic fungi in sewage sludge from wastewater treatment plants of Mazandaran, Islamic Republic of Iran," *Eastern Mediterranean Health Journal*, vol. 15, no. 2, pp. 451–454, 2009.
- [16] H. Moallaei, F. Zaini, M. Pihet, M. Mahmoudi, and J. Hashemi, "Isolation of keratinophilic fungi from soil samples of forests and farm yards," *Iranian Journal of Public Health*, vol. 35, no. 4, pp. 62–69, 2006.
- [17] R. Kachuei, M. Emami, B. Naeimi, and K. Diba, "Isolation of keratinophilic fungi from soil in Isfahan province, Iran," *Journal de Mycologie Medicale*, vol. 22, no. 1, pp. 8–13, 2012.
- [18] K. Pakshir and J. Hashemi, "Dermatophytosis in Karaj, Iran," *Indian Journal of Dermatology*, vol. 51, no. 4, pp. 262–264, 2006.
- [19] H. Böhme and H. Ziegler, "Verbreitung und keratinophilie von *anixiopsis stercoraria* (hansen) hansen," *Archiv für klinische und experimentelle Dermatologie*, vol. 223, no. 4, pp. 422–428, 1965.
- [20] R. Vanbreuseghem, "Technique biologique pour l'isolement des dermatophytes dusol," *Annales de la Société Belge de Médecine Tropicale*, vol. 32, pp. 173–178, 1952.
- [21] S. B. Lee and J. W. Taylor, "Isolation of DNA from fungal mycelium and single cells," in *Protocols: A Guide to Methods and Applications*, M. A. Innis, H. D. Gelfand, J. J. Sninsky, and T. J. White, Eds., pp. 282–287, Academic Press, San Diego, Calif, USA, 1990.
- [22] T. J. White, T. Bruns, S. Lee, and J. Taylor, "Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics," in *PCR Protocols: A Guide to Methods and Applications*, A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White, Eds., pp. 315–322, Academic Press, San Diego, Calif, USA, 1990.

- [23] S. F. Altschul, W. Gish, W. Miller, E. W. Myers, and D. J. Lipman, "Basic local alignment search tool," *Journal of Molecular Biology*, vol. 215, no. 3, pp. 403–410, 1990.
- [24] M. V. Fillipello, "Keratinophilic fungi: their role in nature and degradation of keratinic substrates," in *Biology of Dermatophytes and Other Fungi*, R. K. S. Kushawaha and J. Guarri, Eds., pp. 77–85, Revista Iberoamericana de Micologia, Bilbao, Spain, 2000.
- [25] M. D. Connole, "Review of animal mycoses in Australia," *Mycopathologia*, vol. 111, no. 3, pp. 133–164, 1990.
- [26] V. Filipello Marchisio, L. Preve, and V. Tullio, "Fungi responsible for skin mycoses in Turin (Italy)," *Mycoses*, vol. 39, no. 3-4, pp. 141–150, 1996.
- [27] R. Śpiewak and W. Szostak, "Zoophilic and geophilic dermatophytoses among farmers and non-farmers in eastern Poland," *Annals of Agricultural and Environmental Medicine*, vol. 7, no. 2, pp. 125–129, 2000.
- [28] M. A. Zarei and M. Zarrin, "Isolation of dermatophytes and related keratinophilic fungi from the two public parks in Ahvaz," *Jundishapur Journal of Microbiology*, vol. 1, pp. 20–23, 2008.
- [29] G. Dargahi, *The survey of pathogenic fungi and actinomycetes from soils of Ghazvin [Ph.D. thesis]*, Laboratory Sciences School of Medicine, Tehran University of Medical Science, Tehran, Iran, 1991.
- [30] T. Shokohi, M. T. Hedayati, and H. Bakhshi, "Isolation of fungi and aerobic actinomycetes from surface soil in Sari," *Journal of Kermanshah University of Medical Sciences*, vol. 8, pp. 25–32, 2005.
- [31] S. Shadzi, M. Chadeganipour, and M. Alimoradi, "Isolation of keratinophilic fungi from elementary schools and public parks in Isfahan, Iran," *Mycoses*, vol. 45, no. 11-12, pp. 496–499, 2002.
- [32] M. Eghtedari and K. Pakshir, "Asymptomatic fungal cyst of conjunctiva caused by *Bipolaris spicifera*," *Iranian Journal of Medical Sciences*, vol. 31, no. 1, pp. 56–58, 2006.
- [33] E. Roilides, L. Sigler, E. Bibashi, H. Katsifa, N. Flaris, and C. Panteliadis, "Disseminated infection due to *Chrysosporium zonatum* in a patient with chronic granulomatous disease and review of non-aspergillus fungal infections in patients with this disease," *Journal of Clinical Microbiology*, vol. 37, no. 1, pp. 18–25, 1999.
- [34] S. A. Saidi, S. Bhatt, J. L. Richard, A. Sikdar, and G. R. Ghosh, "Chrysosporium tropicum as a probable cause of mycosis of poultry in India," *Mycopathologia*, vol. 125, no. 3, pp. 143–147, 1994.
- [35] W. Ebrahim, J. Kjer, M. El Amrani et al., "Two new peptides from the endophytic fungus *Bionectria ochroleuca* isolated from the mangrove plant *Sonneratia caseolaris*," *Marine Drugs*, vol. 10, pp. 1081–1091, 2012.
- [36] N. C. Paul, J. X. Deng, J. H. Lee, and S. H. Yu, "New records of endophytic paecilomyces inflatus and bionectria ochroleuca from Chili Pepper Plants in Korea," *Mycobiology*, vol. 41, no. 1, pp. 18–24, 2013.
- [37] R. N. R. Moraes, M. C. T. Ribeiro, M. C. L. Nogueira, K. C. Cunha, M. M. C. N. Soares, and M. T. G. Almeida, "First report of *Tritirachium oryzae* infection of human scalp," *Mycopathologia*, vol. 169, no. 4, pp. 257–259, 2010.
- [38] A. Naseri, A. Fata, and M. J. Najafzadeh, "First case of tritirachium oryzae as agent of onychomycosis and its susceptibility to antifungal drugs," *Mycopathologia*, 2013.
- [39] D. M. Dixon and I. F. Salkin, "Morphologic and physiologic studies of three dematiaceous pathogens," *Journal of Clinical Microbiology*, vol. 24, no. 1, pp. 12–15, 1986.
- [40] N. Singh, F. Y. Chang, T. Gayowski, and I. R. Marino, "Infections due to dematiaceous fungi in organ transplant recipients: case report and review," *Clinical Infectious Diseases*, vol. 24, no. 3, pp. 369–374, 1997.
- [41] K. Singh, J. Flood, R. D. Welsh, J. H. Wyckoff III, T. A. Snider, and D. A. Sutton, "Fatal systemic phaeohyphomycosis caused by *Ochroconis gallopavum* in a dog (*Canis familiaris*)," *Veterinary Pathology*, vol. 43, no. 6, pp. 988–992, 2006.
- [42] L. Cooley, D. Spelman, K. Thursky, and M. Slavin, "Infection with *Scedosporium apiospermum* and *S. prolificans*, Australia," *Emerging Infectious Diseases*, vol. 13, no. 8, pp. 1170–1177, 2007.
- [43] M. Asahi, R. Lindquist, and K. Fukuyama, "Purification and characterization of major extracellular proteinases from *Trichophyton rubrum*," *Biochemical Journal*, vol. 232, no. 1, pp. 139–144, 1985.