

Research Article

Effects of Diphenyl Diselenide on Methylmercury Toxicity in Rats

**Cristiane L. Dalla Corte,¹ Caroline Wagner,² Jéssie H. Sudati,²
Bruna Comparisi,³ Gerlania O. Leite,⁴ Alcindo Busanello,¹ Félix A. A. Soares,¹
Michael Aschner,^{5,6} and João B. T. Rocha¹**

¹ *Biochemistry and Molecular Biology Department, Graduation Program in Biological Sciences: Toxicological Biochemistry, Natural and Exact Sciences Center, Federal University of Santa Maria, 97105-900 Santa Maria, RS, Brazil*

² *Federal University of Pampa—Caçapava do Sul Campus, Avenida Pedro Anunciação, Vila Batista, 96570-000 Caçapava do Sul, RS, Brazil*

³ *Higher Education Cenecista Institute of Santo Ângelo—IESA, Rua Dr. João Augusto Rodrigues 471, 98801-015 Santo Ângelo, RS, Brazil*

⁴ *Regional University of Cariri, Pharmacology and Molecular Chemistry Laboratory, Rua Cel. Antônio Luís 1161, 63100-000 Crato, CE, Brazil*

⁵ *Department of Pharmacology, Vanderbilt University Medical Center, Nashville, TN 37232, USA*

⁶ *Department of Pediatrics, Vanderbilt University Medical Center, Nashville, TN 37232, USA*

Correspondence should be addressed to Cristiane L. Dalla Corte; crisbioq@yahoo.com.br and João B. T. Rocha; jbtrocha@yahoo.com.br

Received 19 September 2013; Revised 25 November 2013; Accepted 25 November 2013

Academic Editor: Fernando Barbosa Jr.

Copyright © 2013 Cristiane L. Dalla Corte et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study investigates the efficacy of diphenyl diselenide [(PhSe)₂] in attenuating methylmercury- (MeHg-)induced toxicity in rats. Adult rats were treated with MeHg [5 mg/kg/day, intragastrically (i.g.)] and/ or (PhSe)₂ [1 mg/kg/day, intraperitoneally (i.p.)] for 21 days. Body weight gain and motor deficits were evaluated prior to treatment, on treatment days 11 and 21. In addition, hepatic and cerebral mitochondrial function (reactive oxygen species (ROS) formation, total and nonprotein thiol levels, membrane potential ($\Delta\Psi_m$), metabolic function, and swelling), hepatic, cerebral, and muscular mercury levels, and hepatic, cerebral, and renal thioredoxin reductase (TrxR) activity were evaluated. MeHg caused hepatic and cerebral mitochondrial dysfunction and inhibited TrxR activity in liver (38,9%), brain (64,3%), and kidney (73,8%). Cotreatment with (PhSe)₂ protected hepatic and cerebral mitochondrial thiols from depletion by MeHg but failed to completely reverse MeHg's effect on hepatic and cerebral mitochondrial dysfunction or hepatic, cerebral, and renal inhibition of TrxR activity. Additionally, the cotreatment with (PhSe)₂ increased Hg accumulation in the liver (50,5%) and brain (49,4%) and increased the MeHg-induced motor deficits and body-weight loss. In conclusion, these results indicate that (PhSe)₂ can increase Hg body burden as well as the neurotoxic effects induced by MeHg exposure in rats.

1. Introduction

MeHg is one of the most poisonous environmental contaminants, causing toxic effects in humans and experimental animals [1, 2]. Environmental MeHg is largely derived from inorganic mercury biomethylation carried out primarily by aquatic microorganisms [3] with subsequent accumulation in

the aquatic food chain and human consumption [4]. MeHg causes acute and chronic damage to multiple organs, most profoundly to the central nervous system (CNS), in particular when exposures occur during the neurodevelopmental period [1, 5, 6].

The events that mediate MeHg toxicity are largely dependent upon its electrophilic properties, which allow for its

interaction with soft nucleophilic groups (mainly thiols and selenols) from either low- or high-molecular-weight biomolecules [7]. The interaction of MeHg with soft nucleophilic groups from biomolecules is responsible, at least in part, for decreased antioxidant capacity and increased ROS generation [7, 8]. Notably, MeHg can disrupt the activity of thiol- and selenol-containing proteins, such as glutathione peroxidase (GPx), thioredoxin (Trx), and TrxR [1, 9–11]. These proteins are important components of the cellular antioxidant system, and their inhibition contributes to the disruption of the normal redox balance of cells [7].

In addition, MeHg can disrupt mitochondrial function by targeting specific thiol-containing proteins, including respiratory chain complexes [12, 13]. The inhibition of these complexes or enzymes can contribute to mitochondrial depolarization and swelling upon MeHg exposure. Mitochondrial targeting by MeHg has also been associated with increased mitochondrial ROS generation, which can further exacerbate the toxicity of MeHg by attacking additional nucleophilic centers in mitochondria and in other subcellular compartments [7, 10, 12–14], leading to a vicious cycle of cell demise.

Several studies demonstrated that organic and inorganic selenium (Se) compounds influence the deposition and toxicity of MeHg [13, 15, 16]. Se is an essential trace element for a wide range of living organisms, including humans [17]. Se is necessary for the expression of approximately 25 Se-dependent proteins, including GPx, TrxR, and several other enzymes and proteins, which can modulate the cellular redox and antioxidant status [17].

In addition to inorganic and naturally occurring organoselenium compounds, synthetic organoselenium compounds can also exhibit protective effects against MeHg. For example, ebselen and (PhSe)₂ have been shown to exert beneficial effects against *in vitro* and *in vivo* MeHg-induced toxicity [18–21]. (PhSe)₂ (which is the simplest of the diaryl diselenides [22]) protected against an array of toxic effects of MeHg and lowered the Hg burden in the brain, liver, and kidneys of adult mice [21]. The molecular mechanism(s) which underlie(s) the protective effects of (PhSe)₂ in mice likely reflect the direct interaction of MeHg with “selenol intermediate” of (PhSe)₂ after its reaction with thiols, or indirectly, by modulating oxidative stress levels [21, 23]. In short, the protective effects of (PhSe)₂ against MeHg-induced toxicity are likely related to its antioxidant properties and its ability to form stable complexes with MeHg, which can increase Hg excretion and decrease the MeHg body burden.

Of particular pharmacological significance, the toxicity and pharmacokinetics of MeHg [24] are different in mice and rat which can be explained by the higher binding affinity of rat hemoglobin, which contains more cysteinyl residues than mice protein, for MeHg when compared to the mice hemoglobin [25]. (PhSe)₂ toxicity and pharmacokinetics differences between mice and rat also exist and could be explained by a faster metabolism of (PhSe)₂ in mice [26–28].

Therefore, the aim of the present study was to investigate the potential protective effects of (PhSe)₂ against MeHg-induced toxicity and mitochondrial dysfunction in rats. To accomplish this goal, the effects of (PhSe)₂ on Hg deposition

in liver and brain and on behavioral and biochemical parameters were studied in rats.

2. Materials and Methods

2.1. Chemicals. Chemicals, including ethylene glycol-bis(β -aminoethylether)-*N,N,N',N'*-tetraacetic acid (EGTA), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), 2,4 dinitrophenol (2,4 DNP), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), glutamic acid, safranin O, 2',7'-dichlorofluorescein diacetate (H₂-DCFDA), and methylmercury chloride were obtained from Sigma Aldrich (St. Louis, MO, USA). (PhSe)₂ was synthesized according to the method by Paulmier [29]. All other chemicals were of analytical reagent grade and purchased from local commercial suppliers.

2.2. Animals. Male Wistar rats, weighing 250–310 g and with age from 3 to 3.5 months, from our own breeding colony were kept in cages (four animals in each). Rats were placed in a room with controlled temperature (22 ± 3°C) on a 12 h light/dark cycle (lights on at 7:00 a.m.) and had continuous access to food and water. All experiments were conducted in accordance with the Committee on Care and Use of Experimental Animal Resources of the Federal University of Santa Maria, Brazil.

2.3. Treatment. Sixteen rats were equally divided into four experimental groups as follows: (1) control (10 mL/Kg of water i.g. and 1 mL/Kg of soybean oil i.p.); (2) (PhSe)₂ (10 mL/Kg of water i.g. and 1 mg/Kg of (PhSe)₂ i.p.); (3) MeHg (5 mg/Kg of MeHg i.g. and 1 mL/Kg of soybean oil i.p.); and (4) (PhSe)₂ + MeHg (5 mg/Kg of MeHg i.g. and 1 mg/Kg of (PhSe)₂ i.p.). Based on previous studies, exposures were performed daily over a 21-day period [21, 30, 31]. Twenty-four hours after the last exposure, the animals were sacrificed and the livers, brains, kidneys, and skeletal muscle were quickly removed, placed on ice and homogenized.

2.4. Determination of Hg Levels. Tissue levels of total Hg were measured in liver, brain, and skeletal muscle collected at the time of euthanasia [32]. Approximately 0.4 g (wet weight) of the tissues was weighed and digested with 5 mL of HNO₃ acid (65%). Digested samples were diluted to 50 mL with ultrapure water before analysis using a Multitype ICP Emission Spectrometer (ICPE-9000, Shimadzu). Calibration standard curve was prepared freshly using mercury stock standard solution.

2.5. Motor Coordination Tests

2.5.1. Open Field Test. General locomotor activity was evaluated as previously described [33]. The number of line crossings (number of segments crossed with the four paws) and rearings was measured over 5 min and taken as an indicator of locomotor activity. The test was carried out at 3 time points: 24 hours prior to treatment (basal), and on treatment days 11 and 21.

2.5.2. Rotarod Test. Motor coordination was tested on the rotarod apparatus as described previously [34, 35]. The latency to fall and the number of falls from the apparatus were recorded for 120 s. The tests were conducted 3 times: 24 hours prior to treatment (basal), and on treatment days 11 and 21.

2.6. TrxR

2.6.1. TrxR Purification. TrxR was partially purified by a modification of the method by Holmgren and Bjornstedt [36]. Tissues were homogenized in buffered saline (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄, and 1.4 mM KH₂PO₄, pH 7.3). Livers, brains, and kidneys (0.5 g) were homogenized in 10, 3, and 5 volumes of buffered saline, respectively. Homogenates were centrifuged at 13,000 g for 30 min. The protein concentration in the supernatant was measured and adjusted to 10 mg/mL. The supernatant was dialyzed against buffered saline for 16 h to remove endogenous glutathione (GSH) and Trx. The dialysate was heated at 55°C for 10 min, cooled, and centrifuged at 13,000 g for 30 min to remove denatured protein.

2.6.2. TrxR Activity. TrxR activity was measured by the method of Holmgren and Bjornstedt [36]. The reaction mixture consisted of the following: 0.24 mM NADPH, 10 mM EDTA, 100 mM potassium phosphate buffer (pH 7.0), 2 mg/mL 5,5'-dithiobis-2-nitrobenzoic acid (DTNB), and 0.2 mg/mL of BSA. The partially purified TrxR was added (to final concentration of 6–8 µg of protein) to the cuvette containing the reaction mixture, and the absorbance was followed at 412 nm for a maximum of 4 min.

2.7. Isolation of Rat Brain and Liver Mitochondria. Rat brain and liver mitochondria were isolated as previously described by Brustovetsky and Dubinsky [37], with some modifications. Brain and liver were rapidly weighed and homogenized in 1:5 (w/v) ice-cold "isolation buffer I" containing 225 mM mannitol, 75 mM sucrose, 1 mM K⁺-EGTA, 0.1% bovine serum albumin (BSA), and 10 mM K⁺-HEPES, pH 7.2. The tissue was then manually homogenized with a potter glass. The resulting suspension was centrifuged for 7 min at 2,000 g. After centrifugation the supernatant was recentrifuged for 10 min at 12,000 g. The pellet was resuspended in "isolation buffer II" containing 225 mM mannitol, 75 mM sucrose, 1 mM K⁺-EGTA, and 10 mM K⁺-HEPES pH 7.2 and recentrifuged at 12,000 g for 10 min. The supernatant was discarded and the final pellet gently washed and resuspended in "isolation buffer II" without EGTA.

2.8. Mitochondrial Nonprotein and Total Thiol Content. Mitochondrial nonprotein and total thiol content were measured according to the method of Ellman [38]. To determine total thiol groups, mitochondria (0.3 mg protein) were added to the reaction medium containing 10 mM Tris-HCl pH 7.2, 1% SDS, and 10 mM DTNB. Nonprotein thiol content was measured by adding 50 µL 10% TCA to 50 µL of the mitochondria (0.3 mg protein). After centrifugation (4,000 ×g at 4°C for 10 min), the protein pellet was discarded and an aliquot of

the clear supernatant, neutralized with 0.1M NaOH, was added to the medium containing 10 mM Tris-HCl pH 7.2 and 10 mM DTNB. The samples absorbance was measured spectrophotometrically at a wavelength of 412 nm.

2.9. Measurements of Mitochondrial ΔΨ_m. Mitochondrial ΔΨ_m was estimated by fluorescence changes in safranin O (3 mM) recorded by RF-5301 Shimadzu spectrofluorometer (Kyoto, Japan) operating at excitation and emission wavelengths of 495 and 535 nm, with slit widths of 1.5 nm [39]. Data on ΔΨ_m in the figures is presented in Arbitrary Fluorescence Units (AFU).

2.10. Estimation of ROS Production. The mitochondrial generation of ROS was determined spectrofluorimetrically, using the membrane permeable fluorescent dye H₂-DCFDA recorded by RF-5301 Shimadzu spectrofluorometer (Kyoto, Japan) operating at excitation and emission wavelengths of 488 and 525 nm, with slit widths of 3 nm [40]. Data of ROS production in the figures is presented as Arbitrary Fluorescence Units (AFU).

2.11. Assessment of Mitochondrial Metabolic Function. The mitochondrial metabolic function was assessed by the conversion of MTT to a dark violet formazan product by mitochondrial dehydrogenases [41]. The rate of MTT reduction was measured spectrophotometrically at a wavelength of 570 nm. Results were expressed as the percentage of MTT reduction relative to control values.

2.12. Assessment of Mitochondrial Swelling. Measurement of mitochondrial swelling was performed in a RF-5301 Shimadzu spectrofluorometer at 600 nm (slit 1.5 nm for excitation and emission) [42]. Data for mitochondrial swelling are expressed as Arbitrary Absorbance Units (AAU). The difference (ΔA) between the initial absorbance reading and the final absorbance reading was used for statistical analysis.

2.13. Protein Measurement. Protein was assayed by the method of Bradford [43] with bovine serum albumin as standard.

2.14. Statistical Analysis. Normality assumption was tested with Kolmogorov-Smirnov test and the distribution of the majority of results is not normal. Data were analyzed statistically by Mann-Whitney or Kruskal-Wallis, followed by Dunn's post-hoc tests when appropriate. The results were considered statistically significant at $P < 0.05$. All statistical analyses were conducted using GraphPad Prism 5 (Version 5.01, GraphPad Software, Inc., USA).

3. Results

3.1. Effects of (PhSe)₂ and MeHg on Body Weight. Treatment with MeHg led to body-weight loss from the second week until the end of the treatment compared to controls ($P < 0.05$, Figure 1). Rats cotreated with (PhSe)₂ and MeHg also




FIGURE 1: Effect of MeHg and/or (PhSe)₂ on the body weight gain in adult rats. Data are expressed as mean \pm S.D., $n = 4$. (*) represents $P < 0.05$ as compared to controls by Mann-Whitney test.

showed a decrease in the body weight when compared to the control group ($P < 0.05$, Figure 1). Rats treated with (PhSe)₂ lost weight after the first week of treatment ($P < 0.05$) but showed a trend towards a recovery and were statistically indistinguishable from the controls at the end of the treatment (Figure 1).

3.2. Effects of (PhSe)₂ and MeHg on Hg Deposition. Treatment with MeHg increased the levels of Hg in liver, brain, and skeletal muscle compared with controls ($P < 0.05$, Figure 2). The cotreatment with (PhSe)₂ caused a greater increase in brain Hg deposition when compared to MeHg alone treatment, both in brain (Figure 2(b)) and liver (Figure 2(a)), and showed a trend towards increased deposition in skeletal muscle (Figure 2(c)).

3.3. Effects of (PhSe)₂ and MeHg on Motor Coordination and Spontaneous Locomotor Activity. The effects of MeHg and/or (PhSe)₂ on locomotion and motor coordination were assessed by the open-field and rotarod tests, respectively. After 11 days, rats treated with MeHg showed increased number of falls on the rotarod and decreased latency to the first fall when compared to controls ($P < 0.05$, Figures 3(a) and 3(b)). Rats treated with (PhSe)₂ did not show statistically significant differences on the rotarod test when compared to controls; however, rats cotreated with (PhSe)₂ and MeHg showed increased loss of motor coordination as evidenced by increased number of falls and reduced latency to the first fall ($P < 0.05$, Figures 3(a) and 3(b)). The rotarod test could not be performed at the end of the treatment in rats receiving MeHg since they were unable to remain in the apparatus due to severe motor impairment caused by MeHg.

Rats treated with MeHg showed a decrease in the number of crossings and rearings in the open-field at the end of the treatment compared to the control rats ($P < 0.05$, Figures 3(c) and 3(d)). Rats cotreated with MeHg and (PhSe)₂ also showed a significant decrease in the number of crossings

after 11 days of treatment and a decrease in the number of rearings at the end of the treatment ($P < 0.05$, Figure 3(d)). Treatment with (PhSe)₂ did not affect the rats' performance in the open-field. The decrease in the number of crossings and rearings observed in all groups on treatment days 11 and 21 was expected given that the animals habituate to the open-field arena [44].

3.4. Effects of (PhSe)₂ and MeHg on Mitochondrial Dysfunction

3.4.1. Mitochondrial Metabolic Function. The hepatic mitochondrial metabolic integrity (MTT reduction) was not affected by MeHg and/or (PhSe)₂ (Figure 4(a)). Treatment with MeHg or cotreatment with MeHg and (PhSe)₂ decreased the capacity of brain mitochondrial dehydrogenases to reduce MTT compared to controls ($P < 0.05$, Figure 4(b)). Treatment with (PhSe)₂ alone did not affect the cerebral mitochondrial metabolic function.

3.4.2. Mitochondrial Total and Nonprotein Thiols. MeHg treatment decreased the total mitochondrial thiol levels in brain and liver when compared to controls ($P < 0.05$, Figure 5). Treatment with (PhSe)₂ alone did not alter the mitochondrial total thiol levels in liver and brain (Figure 5). The cotreatment with (PhSe)₂ blunted the MeHg-induced mitochondrial total thiol level depletion in rats' liver and brain ($P < 0.05$, Figure 5). Rats treated with MeHg showed decreased mitochondrial nonprotein thiol levels in the liver compared to controls, and coadministration of (PhSe)₂ blunted the MeHg-induced decrease in hepatic nonprotein thiol content ($P < 0.05$, Figure 5(a)). Brain mitochondrial nonprotein thiol levels were not affected by any of the treatments (Figure 5(b)).

3.4.3. Mitochondrial Swelling. Treatment with MeHg significantly increased hepatic mitochondrial swelling when compared to controls ($P < 0.05$, Figure 6(a)). Cotreatment with (PhSe)₂ partially prevented the MeHg-induced mitochondrial swelling in liver (Figure 6(a)). Similarly, treatment with MeHg showed a trend towards increased mitochondrial swelling in brain (Figure 6(b)). The cotreatment with MeHg and (PhSe)₂ significantly increased cerebral mitochondrial swelling when compared to controls ($P < 0.05$, Figure 6(b)). Treatment with (PhSe)₂ alone did not alter the mitochondrial swelling in brain or liver compared to the controls (Figures 6(a) and 6(b)).

3.4.4. Mitochondrial ROS Production. Mitochondrial ROS production (DCFH oxidation) was significantly increased in livers of rats treated with MeHg or cotreated with MeHg and (PhSe)₂ ($P < 0.05$, Figure 7(a)). Rats treated with (PhSe)₂ showed hepatic mitochondrial ROS levels indistinguishable from controls. ROS production in cerebral mitochondria was not affected by any of the treatments (Figure 7(b)).

3.4.5. Mitochondrial $\Delta\Psi_m$. Polarization ($\Delta\Psi_m$) of mitochondria from liver of rats cotreated with MeHg and (PhSe)₂




FIGURE 2: Hg content in liver (a), brain (b), and muscle (c) of rats exposed to MeHg and/or (PhSe)₂. Data are expressed as mean \pm S.D., $n = 4$. (*) represents $P < 0.05$ as compared to controls by Mann-Whitney test. (#) represents $P < 0.05$ as compared to MeHg by Mann-Whitney test.

showed only a trend towards decreased (Figures 8(a) and 8(c)). Treatment with (PhSe)₂ and MeHg alone did not cause mitochondrial depolarization in liver of rats (Figures 8(a) and 8(c)). Treatment with (PhSe)₂ and/or MeHg had no effect on mitochondrial $\Delta\Psi_m$ in brain of rats (Figures 8(b) and 8(d)).

3.5. Effects of (PhSe)₂ and MeHg on TrxR Activity. MeHg is known to inhibit TrxR activity both *in vitro* and *in vivo* [1, 9, 11]. (PhSe)₂ treatment significantly increased renal TrxR activities when compared to controls ($P < 0.05$, Figures 9(a) and 9(b)). Hepatic and cerebral TrxR activity showed a trend towards increased in rats treated with (PhSe)₂ (Figure 9(c)). MeHg treatment also led to significant inhibition of TrxR in liver, kidney, and brain compared to controls ($P < 0.05$, Figure 9). Cotreatment with (PhSe)₂ failed to significantly attenuate the MeHg-induced inhibition of TrxR activity in the liver, kidney, or brain (Figure 9).

4. Discussion

The present study investigated the efficacy of (PhSe)₂, an organoselenium compound, in attenuating MeHg-induced

toxicity in rats. Our results established that MeHg decreased body weight (Figure 1) and induced motor deficits (Figure 3) as well hepatic and cerebral mitochondrial dysfunction (Figures 4(b), 5, 6(a), and 7(a)) and inhibited TrxR activity in liver, brain, and kidney (Figure 9) in the rat. The cotreatment with (PhSe)₂ and MeHg increased Hg accumulation in the liver and brain (Figure 2). Furthermore, the cotreatment with (PhSe)₂ protected hepatic and cerebral mitochondrial thiols from depletion by MeHg (Figure 5) but did not prevent hepatic and cerebral mitochondrial dysfunction (Figures 4(b), 6(b), and 7(a)) nor did it reverse the MeHg-induced motor deficits (Figure 3), body-weight loss (Figure 1), and the MeHg-induced inhibition of TrxR activity in liver, brain, and kidney (Figure 9).

Cotreatment with (PhSe)₂ and MeHg increased Hg deposition in the brain and liver of exposed rats (Figure 2). These results differ from those of de Freitas et al. [21] where (PhSe)₂ led to a significant reduction in Hg concentrations in brain, liver, and kidney of MeHg-exposed mice. The discrepancies between the 2 studies may be attributed to metabolic differences between the species and the route of administration. The toxicity and pharmacokinetics of MeHg [24] are different in mice and rat which can be explained




FIGURE 3: Rotarod and open field tests in rats exposed to MeHg and/or (PhSe)₂. The number of falls (a) and latency for the first fall (b) ambulation (crossing) (a) and rearing (b) were recorded. Data are expressed as mean \pm S.D., $n = 4$. (*) represents $P < 0.05$ as compared to controls by Kruskal-Wallis test followed by multiple comparison test. (#) represents $P < 0.05$ as compared to (PhSe)₂ by Kruskal-Wallis test followed by multiple comparison test.

by the higher binding affinity of rat hemoglobin, containing more cysteinyl residues, for MeHg when compared to the mice hemoglobin [25]. (PhSe)₂ toxicity and pharmacokinetics differences between mice and rat also exist and could be explained by a faster metabolism of (PhSe)₂ in mice [26–28]. Notably, herein rats were administered (PhSe)₂ i.p., whilst in the study by de Freitas et al. [21] (PhSe)₂ was subcutaneously (s.c.) administered to the mice. Another difference between the two works is in relation to the dose of MeHg: in our study we used a dose 2.5 times higher than in the study of de Freitas et al. (2 mg/Kg). However, the duration of the treatment was shorter in our study, 21 versus 35 days. On the other hand, the dose of (PhSe)₂ was similar between the two studies. The higher dose of MeHg used in our study may have contributed to the discrepancies since it could generate a more severe toxicity which could not be prevented

by (PhSe)₂. However, we realize that the differences in the pharmacokinetics between rats and mice for the (PhSe)₂ is the major factor involved in the discrepancies found here [28].

In the study by de Freitas et al. [21] the proposed mechanism for the reduction Hg's organ burden by (PhSe)₂ was the formation of a selenol/selenolate (PhSeH/PhSe⁻) intermediate, which could interact with MeHg, generating the readily excretable PhSeHgMe complex. One possible explanation for the increase in hepatic and cerebral Hg deposition (Figures 2(a) and 2(b), resp.) by the cotreatment with (PhSe)₂ observed herein may be the conversion of (PhSe)₂ to inorganic selenium, which is subsequently metabolized to selenidric acid (HSe⁻). HSe⁻ could bind to MeHg to form a less soluble complex [45], which can be degraded to HgSe [46, 47]. In addition, Palmer and Parkin [48] showed




FIGURE 4: MTT reduction in liver (a) and brain (b) mitochondria of rats exposed to MeHg and/or (PhSe)₂. Data are expressed as mean ± S.D., n = 4. (*) represents P < 0.05 as compared to controls by Mann-Whitney test.




FIGURE 5: Total and nonprotein thiol content in liver (a), (c) and brain (b), (d) mitochondria of rats exposed to MeHg and/or (PhSe)₂. Data are expressed as mean ± S.D., n = 4. (*) represents P < 0.05 as compared to controls by Mann-Whitney test.

that organoselenium can also form a complex with mercury. Thus the increase in hepatic and cerebral Hg deposition by the cotreatment with (PhSe)₂ possibly involves Hg:Se interactions and the formation of a less excretable compound that accumulates in these organs [45]. These results are in

agreement with other studies that showed elevated deposition of Hg in key brain regions upon oral Se administration [49, 50]. It has been speculated that the formation of insoluble HgSe salt could reduce the toxicity of MeHg. However, experimental evidence supporting this assumption has yet to be




FIGURE 6: Mitochondrial swelling in liver (a) and brain (b) of rats exposed to MeHg and/or (PhSe)₂. Data are expressed as mean ± S.D., $n = 4$. (*) represents $P < 0.05$ as compared to controls by Mann-Whitney test.




FIGURE 7: ROS production (H₂-DCFH oxidation) in liver (a) and brain (b) mitochondria of rats exposed to MeHg and/or (PhSe)₂. Data are expressed as mean ± S.D., $n = 4$. (*) represents $P < 0.05$ as compared to controls by Mann-Whitney test.

generated. Although the cotreatment with (PhSe)₂ increased Hg levels in brain and liver, these were accompanied by a partial protection against MeHg-induced mitochondrial dysfunction. We suggest that the formation of an insoluble and inert complex between Hg and Se could decrease the availability of MeHg that could react with important cellular components decreasing its toxicity.

Decreased weight gain and weight loss are prominent and readily observed features of severe MeHg toxicity. In this study, rats treated with MeHg showed body-weight loss (Figure 1). Notably, the most severe effect on weight loss occurred in rats cotreated with (PhSe)₂ and MeHg (Figure 1). In addition, rats treated with MeHg showed decreased locomotor activity (Figure 3). Cotreatment with (PhSe)₂ and MeHg increased the severity of motor dysfunction (rotarod test) (Figure 3), likely as a result of increased Hg deposition in the brain (Figure 2(b)). Motor deficits are the most apparent neurological effects following MeHg

exposure [51]. *In vivo* studies in rodents point to impairment in intracellular calcium homeostasis, alteration in glutamate homeostasis, and oxidative stress as critical mediators of MeHg-induced neurotoxicity [52]. The overactivation of N-methyl-D-aspartate- (NMDA-) type glutamate receptors increases Ca²⁺ influx into neurons, thereby leading to cell death [53]. Alternatively, Ca²⁺ taken up by mitochondria may stimulate the generation of ROS [54].

Several studies corroborate MeHg's ability to induce mitochondrial dysfunction and ROS generation [14, 18, 55]. The high affinity binding of MeHg to thiol groups inactivates enzymes, including respiratory chain complexes [7, 13, 55], decreasing mitochondrial dehydrogenases activity. Inhibition of these complexes may contribute to mitochondrial swelling and ROS production after MeHg exposure (Figures 6 and 7). However, in brain, the MeHg-induced decrease in mitochondrial dehydrogenases activity (MTT reduction) was not accompanied by an increase in ROS production. These




FIGURE 8: Mitochondrial depolarization in liver (a), (c) and brain (b), (d) of rats exposed to MeHg and/or (PhSe)₂. Figures (a) and (b) show mitochondrial membrane potential (AFU). Figures (c) and (d) show mitochondrial $\Delta\Psi_m$. $\Delta\Psi_{m1}$ = delta of fluorescence before (time 0) and after addition of mitochondria (time 150 seconds) and $\Delta\Psi_{m2}$ = delta of fluorescence before (time 150 seconds) and after addition of 2,4 DNP (time 300 seconds). Data are expressed as mean \pm S.D., $n = 4$.

results are corroborated by the fact that MeHg affected total thiols but not nonprotein thiol levels in brain mitochondria. MeHg caused a decrease in the total mitochondrial thiol levels in brain, which is related mainly with protein thiols, and is in agreement with the inhibition of mitochondrial dehydrogenases activity in this tissue. On the other hand, MeHg did not affect nonprotein thiol levels (mainly GSH) in brain mitochondria, which can explain the normal ROS production, since GSH is the main antioxidant in brain.

The cotreatment with (PhSe)₂ prevented the MeHg-induced mitochondrial total and nonprotein thiol groups depletion in the brain and liver (Figure 4). The efficacy of (PhSe)₂ in preventing thiol depletion may reside in its ability to form a complex with MeHg, thus effectively reducing MeHg binding to protein and free thiols. Treatment with

(PhSe)₂ also partially protected the liver from mitochondrial MeHg-induced swelling (Figure 6(a)). However, the cotreatment with (PhSe)₂ failed to reverse the MeHg-induced mitochondrial swelling (Figure 6(b)) and decreased mitochondrial metabolic function (Figure 3(b)) in the brain as well as increased mitochondrial ROS production (Figure 7(a)) in the liver. These results indicate that mechanisms other than the interaction with important free and protein thiols are likely involved in the MeHg-induced mitochondrial dysfunction. Thus, the preferential affinity of MeHg for specific, and as of yet unidentified, mitochondrial protein targets may have a critical role in MeHg's toxicity.

Previous studies have demonstrated that MeHg can directly inhibit TrxR activity both *in vitro* and *in vivo* [1, 9, 11]. Mammalian TrxR is a selenoenzyme containing a




FIGURE 9: TrxR activity in liver (a), kidney (b), and brain (c) of rats exposed to MeHg and/or (PhSe)₂. Data are expressed as mean \pm S.D., $n = 4$. (*) represents $P < 0.05$ as compared to controls by Mann-Whitney test. (#) represents $P < 0.05$ as compared to controls by Mann-Whitney test.

unique, catalytically active selenolthiol/selenenylsulfide in the conserved C-terminal sequence (-Gly-Cys-Sec-Gly) [56]. Three mammalian TrxR selenoenzymes have been identified, the cytosolic enzyme TrxR1, the mitochondrial enzyme TrxR2, and a testis-specific enzyme thioredoxin-glutathione reductase (TGR/TrxR3) [56]. Here, we show that MeHg treatment inhibited rat TrxR activity in brain, liver, and kidney (Figure 9). MeHg forms covalent bonds between its Hg moiety and the Se of the selenocysteine of the enzyme, thus directly inhibiting the activity of TrxR [1]. Since TrxR is critical for cellular antioxidant defense system the inhibition of this enzyme likely has a central role in mediating the toxicity of MeHg.

Recently, diphenyl diselenide was demonstrated to be a substrate for cerebral and hepatic rat TrxR, which could account, at least in part, for the antioxidant properties of (PhSe)₂ [23]. Herein, rats treated solely with (PhSe) showed an increase in the activity of renal TrxR (Figures 9(a) and 9(b), resp.). The formation of selenhydric acid from (PhSe)₂ could also explain the increase in TrxR activity, since this inorganic form of Se can be converted to selenocysteine and incorporated to selenoenzymes, such as TrxR [45, 57]. Accordingly, Zhang et al. [58] have demonstrated that organoselenium compounds (including diselenide) increase

the expression of TrxR in white blood cells lines in culture. The cotreatment with (PhSe)₂ and MeHg was ineffective in attenuating the inhibition of MeHg-induced TrxR in liver, kidney, and brain (Figure 9). Similarly, studies *in vitro* and *in vivo* have previously corroborated that selenite was able to recover the activity of HgCl₂-induced TrxR inhibition but not in response to MeHg. The effect of Se (as selenide) was attributed to the displacement of Hg from the active site, giving rise to mercury selenide and regenerating the TrxR selenol [1, 9].

5. Conclusions

In conclusion, the results of this study established that (PhSe)₂ can increase Hg body burden (likely associated with release of inorganic Se from (PhSe)₂) and MeHg neurotoxicity in rats despite the fact that (PhSe)₂ blunted the deleterious effects of MeHg on thiol levels. The results presented herein also reinforce the central role of mitochondrial dysfunction in mediating the aberrant effects of MeHg *in vivo*, as well as the role of TrxR as a molecular target for MeHg in the rat. Further research into MeHg-(PhSe)₂ interactions will be helpful in characterizing the consequences concomitant exposures to these and related compounds.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This work was supported by Fundação de Amparo a Pesquisa do Estado do Rio Grande do Sul (FAPERGS), CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), FINEP (Rede Instituto Brasileiro de Neurociência (IBN-Net) no. 01.06.0842-00), FAPERGS-PRONEX-CNPQ, and INCT-EN (Instituto Nacional de Ciência e Tecnologia em Excitotoxicidade e Neuroproteção). Cristiane L. Dalla Corte is recipient of CAPES fellowship. Félix A. A. Soares and João B. T. Rocha are recipients of CNPq fellowships. Michael Aschner was partially supported by grants from the US PHS, R01 ES007331 and R01 ES200852.

References

- [1] C. M. L. Carvalho, J. Lu, X. Zhang, E. S. J. Arnér, and A. Holmgren, "Effects of selenite and chelating agents on mammalian thioredoxin reductase inhibited by mercury: implications for treatment of mercury poisoning," *The FASEB Journal*, vol. 25, no. 1, pp. 370–381, 2011.
- [2] T. W. Clarkson, L. Magos, and G. J. Myers, "The toxicology of mercury—current exposures and clinical manifestations," *The New England Journal of Medicine*, vol. 349, no. 18, pp. 1731–1737, 2003.
- [3] G. C. Campeau and R. Bartha, "Sulfate-reducing bacteria: principal methylators of mercury in anoxic estuarine sediment," *Applied and Environmental Microbiology*, vol. 50, no. 2, pp. 498–502, 1985.
- [4] J. G. Dórea, "Persistent, bioaccumulative and toxic substances in fish: human health considerations," *Science of the Total Environment*, vol. 400, no. 1–3, pp. 93–114, 2008.
- [5] P. Grandjean and K. T. Herz, "Methylmercury and brain development: imprecision and underestimation of developmental neurotoxicity in humans," *Mount Sinai Journal of Medicine*, vol. 78, no. 1, pp. 107–118, 2011.
- [6] C. Johansson, A. F. Castoldi, N. Onishchenko, L. Manzo, M. Vahter, and S. Ceccatelli, "Neurobehavioural and molecular changes induced by methylmercury exposure during development," *Neurotoxicity Research*, vol. 11, no. 3–4, pp. 241–260, 2007.
- [7] M. Farina, M. Aschner, and J. B. T. Rocha, "Oxidative stress in MeHg-induced neurotoxicity," *Toxicology and Applied Pharmacology*, vol. 256, no. 3, pp. 405–417, 2011.
- [8] J. L. Franco, H. C. Braga, J. Stringari et al., "Mercurial-induced hydrogen peroxide generation in mouse brain mitochondria: protective effects of quercetin," *Chemical Research in Toxicology*, vol. 20, no. 12, pp. 1919–1926, 2007.
- [9] V. Branco, J. Canário, A. Holmgren, and C. Carvalho, "Inhibition of the thioredoxin system in the brain and liver of zebra-seabreams exposed to waterborne methylmercury," *Toxicology and Applied Pharmacology*, vol. 251, no. 2, pp. 95–103, 2011.
- [10] J. L. Franco, T. Posser, P. R. Dunkley et al., "Methylmercury neurotoxicity is associated with inhibition of the antioxidant enzyme glutathione peroxidase," *Free Radical Biology and Medicine*, vol. 47, no. 4, pp. 449–457, 2009.
- [11] C. Wagner, J. H. Sudati, C. W. Nogueira, and J. B. T. Rocha, "In vivo and in vitro inhibition of mice thioredoxin reductase by methylmercury," *BioMetals*, vol. 23, no. 6, pp. 1171–1177, 2010.
- [12] A. Dreiem and R. F. Seegal, "Methylmercury-induced changes in mitochondrial function in striatal synaptosomes are calcium-dependent and ROS-independent," *NeuroToxicology*, vol. 28, no. 4, pp. 720–726, 2007.
- [13] V. Glaser, E. M. Nazari, Y. M. R. Müller et al., "Effects of inorganic selenium administration in methylmercury-induced neurotoxicity in mouse cerebral cortex," *International Journal of Developmental Neuroscience*, vol. 28, no. 7, pp. 631–637, 2010.
- [14] D. H. Roos, R. L. Puntel, M. Farina et al., "Modulation of methylmercury uptake by methionine: prevention of mitochondrial dysfunction in rat liver slices by a mimicry mechanism," *Toxicology and Applied Pharmacology*, vol. 252, no. 1, pp. 28–35, 2011.
- [15] A. L. Choi, E. Budtz-Jørgensen, P. J. Jørgensen et al., "Selenium as a potential protective factor against mercury developmental neurotoxicity," *Environmental Research*, vol. 107, no. 1, pp. 45–52, 2008.
- [16] A. Fredriksson, A. Teiling Gardlund, K. Bergman et al., "Effects of maternal dietary supplementation with selenite on the post-natal development of rat offspring exposed to methyl mercury in utero," *Pharmacology and Toxicology*, vol. 72, no. 6, pp. 377–382, 1993.
- [17] D. L. Hatfield and V. N. Gladyshev, "How selenium has altered our understanding of the genetic code," *Molecular and Cellular Biology*, vol. 22, no. 11, pp. 3565–3576, 2002.
- [18] C. L. Dalla Corte, F. A. A. Soares, M. Aschner, and J. B. T. Rocha, "Diphenyl diselenide prevents methylmercury-induced mitochondrial dysfunction in rat liver slices," *Tetrahedron*, vol. 68, no. 51, pp. 10437–10443, 2012.
- [19] D. H. Roos, R. L. Puntel, M. M. Santos et al., "Guanosine and synthetic organoselenium compounds modulate methylmercury-induced oxidative stress in rat brain cortical slices: involvement of oxidative stress and glutamatergic system," *Toxicology in Vitro*, vol. 23, no. 2, pp. 302–307, 2009.
- [20] Z. Yin, E. Lee, M. Ni et al., "Methylmercury-induced alterations in astrocyte functions are attenuated by ebselen," *NeuroToxicology*, vol. 32, no. 3, pp. 291–299, 2011.
- [21] A. S. de Freitas, V. R. Funck, M. D. S. Rotta et al., "Diphenyl diselenide, a simple organoselenium compound, decreases methylmercury-induced cerebral, hepatic and renal oxidative stress and mercury deposition in adult mice," *Brain Research Bulletin*, vol. 79, no. 1, pp. 77–84, 2009.
- [22] C. W. Nogueira and J. B. T. Rocha, "Diphenyl diselenide a janus-faced molecule," *Journal of the Brazilian Chemical Society*, vol. 21, no. 11, pp. 2055–2071, 2010.
- [23] A. S. de Freitas, A. de Souza Prestes, C. Wagner et al., "Reduction of diphenyl diselenide and analogs by mammalian thioredoxin reductase is independent of their glutathione peroxidase-like activity: a possible novel pathway for their antioxidant activity," *Molecules*, vol. 15, no. 11, pp. 7699–7714, 2010.
- [24] L. Magos, "The absorption, distribution, and excretion of methylmercury," in *The Toxicity of Methylmercury*, C. U. Edclles and Z. Annau, Eds., pp. 24–44, John Hopkins University Press, Baltimore, Md, USA, 1987.
- [25] R. Doi and M. Tagawa, "A study on the biochemical and biological behavior of methylmercury," *Toxicology and Applied Pharmacology*, vol. 69, no. 3, pp. 407–416, 1983.

- [26] V. B. Brito, V. Folmer, G. O. Puntel et al., "Diphenyl diselenide and 2,3-dimercaptopropanol increase the PTZ-induced chemical seizure and mortality in mice," *Brain Research Bulletin*, vol. 68, no. 6, pp. 414–418, 2006.
- [27] C. W. Nogueira and J. B. T. Rocha, "Toxicology and pharmacology of selenium: emphasis on synthetic organoselenium compounds," *Archives of Toxicology*, vol. 85, no. 11, pp. 1313–1359, 2011.
- [28] M. Prigol, R. F. Schumacher, C. WayneNogueira, and G. Zeni, "Convulsant effect of diphenyl diselenide in rats and mice and its relationship to plasma levels," *Toxicology Letters*, vol. 189, no. 1, pp. 35–39, 2009.
- [29] C. Paulmier, "Selenoorganic functional groups," in *Selenium Reagents and Intermediates in Organic Synthesis*, pp. 25–51, Pergamon Press, Oxford, UK, 1 edition, 1986.
- [30] J. Christinal and T. Sumathi, "Effect of *Bacopa monniera* extract on methylmercury-induced behavioral and histopathological changes in rats," *Biological Trace Element Research*, vol. 155, no. 1, pp. 56–64, 2013.
- [31] T. Sumathi, C. Shobana, J. Christinal, and C. Anusha, "Protective effect of *Bacopa monniera* on methyl mercury-induced oxidative stress in cerebellum of rats," *Cellular and Molecular Neurobiology*, vol. 2, no. 6, pp. 979–987, 2012.
- [32] G. R. Bhagure and S. R. Mirgane, "Heavy metal concentrations in groundwaters and soils of Thane Region of Maharashtra, India," *Environmental Monitoring and Assessment*, vol. 173, no. 1–4, pp. 643–652, 2011.
- [33] P. L. Broadhurst, "The place of animal psychology in the development of psychosomatic research," *Fortschritte der Psychosomatischen Medizin*, vol. 1, pp. 63–69, 1960.
- [34] R. Lalonde, T. L. Lewis, C. Strazielle, H. Kim, and K. Fukuchi, "Transgenic mice expressing the β APP695SWE mutation: effects on exploratory activity, anxiety, and motor coordination," *Brain Research*, vol. 977, no. 1, pp. 38–45, 2003.
- [35] R. Lalonde, H. D. Kim, and K. Fukuchi, "Exploratory activity, anxiety, and motor coordination in bigenic APPswe + PS1/ Δ E9 mice," *Neuroscience Letters*, vol. 369, no. 2, pp. 156–161, 2004.
- [36] A. Holmgren and M. Bjornstedt, "Thioredoxin and thioredoxin reductase," *Methods in Enzymology*, vol. 252, pp. 199–208, 1995.
- [37] N. Brustovetsky and J. M. Dubinsky, "Dual responses of CNS mitochondria to elevated calcium," *Journal of Neuroscience*, vol. 20, no. 1, pp. 103–113, 2000.
- [38] G. L. Ellman, "Tissue sulfhydryl groups," *Archives of Biochemistry and Biophysics*, vol. 82, no. 1, pp. 70–77, 1959.
- [39] Q. Guo, S. Christakos, N. Robinson, and M. P. Mattson, "Calbindin D28K blocks the proapoptotic actions of mutant presenilin 1: reduced oxidative stress and preserved mitochondrial function," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 95, no. 6, pp. 3227–3232, 1998.
- [40] C. García-Ruiz, A. Colell, M. Marí, A. Morales, and J. C. Fernández-Checa, "Direct effect of ceramide on the mitochondrial electron transport chain leads to generation of reactive oxygen species: role of mitochondrial glutathione," *Journal of Biological Chemistry*, vol. 272, no. 17, pp. 11369–11377, 1997.
- [41] T. Mosmann, "Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays," *Journal of Immunological Methods*, vol. 65, no. 1-2, pp. 55–63, 1983.
- [42] T. V. Votyakova and I. J. Reynolds, " Ca^{2+} -induced permeabilization promotes free radical release from rat brain mitochondria with partially inhibited complex I," *Journal of Neurochemistry*, vol. 93, no. 3, pp. 526–537, 2005.
- [43] M. M. Bradford, "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding," *Analytical Biochemistry*, vol. 72, no. 1-2, pp. 248–254, 1976.
- [44] P. B. Mello, F. Benetti, M. Cammarota, and I. Izquierdo, "Effects of acute and chronic physical exercise and stress on different types of memory in rats," *Anais da Academia Brasileira de Ciencias*, vol. 80, no. 2, pp. 301–309, 2008.
- [45] C. Pinheiro, J. L. M. Do Nascimento, L. C. L. Silveira, J. B. T. Rocha, and M. Aschner, "Mercury and selenium—a review on aspects related to the health of human populations in the Amazon," *Environmental Bioindicators*, vol. 4, no. 3, pp. 222–245, 2009.
- [46] F. E. Huggins, S. A. Raverty, O. S. Nielsen, N. E. Sharp, J. D. Robertson, and N. V. C. Ralston, "An XAFS investigation of mercury and selenium in Beluga whale tissues," *Environmental Bioindicators*, vol. 4, no. 4, pp. 291–302, 2009.
- [47] H. Iwata, T. Masukawa, H. Kito, and M. Hayashi, "Degradation of methylmercury by selenium," *Life Sciences*, vol. 31, no. 9, pp. 859–866, 1982.
- [48] J. H. Palmer and G. Parkin, "2-Seleno-1-alkylbenzimidazoles and their diselenides: synthesis and structural characterization of a 2-seleno-1-methylbenzimidazole complex of mercury," *Polyhedron*, vol. 52, pp. 658–668, 2013.
- [49] B. Moller-Madsen and G. Danscher, "Localization of mercury in CNS of the rat: IV. The effect of selenium on orally administered organic and inorganic mercury," *Toxicology and Applied Pharmacology*, vol. 108, no. 3, pp. 457–473, 1991.
- [50] J. R. Prohaska and H. E. Ganther, "Interactions between selenium and methylmercury in rat brain," *Chemico-Biological Interactions*, vol. 16, no. 2, pp. 155–167, 1977.
- [51] M. O. Dietrich, C. E. Mantese, G. D. Anjos, D. O. Souza, and M. Farina, "Motor impairment induced by oral exposure to methylmercury in adult mice," *Environmental Toxicology and Pharmacology*, vol. 19, no. 1, pp. 169–175, 2005.
- [52] M. Aschner, T. Syversen, D. O. Souza, J. B. T. Rocha, and M. Farina, "Involvement of glutamate and reactive oxygen species in methylmercury neurotoxicity," *Brazilian Journal of Medical and Biological Research*, vol. 40, no. 3, pp. 285–291, 2007.
- [53] D. W. Choi, "Excitotoxic cell death," *Journal of Neurobiology*, vol. 23, no. 9, pp. 1261–1276, 1992.
- [54] S. Ceccatelli, E. Daré, and M. Moors, "Methylmercury-induced neurotoxicity and apoptosis," *Chemico-Biological Interactions*, vol. 188, no. 2, pp. 301–308, 2010.
- [55] N. Mori, A. Yasutake, and K. Hirayama, "Comparative study of activities in reactive oxygen species production/defense system in mitochondria of rat brain and liver, and their susceptibility to methylmercury toxicity," *Archives of Toxicology*, vol. 81, no. 11, pp. 769–776, 2007.
- [56] L. V. Papp, A. Holmgren, and K. K. Khanna, "Selenium and selenoproteins in health and disease," *Antioxidants and Redox Signaling*, vol. 12, no. 7, pp. 793–795, 2010.
- [57] W. J. Adams Jr., J. J. Kocsis, and R. Snyder, "Acute toxicity and urinary excretion of diphenyldiselenide," *Toxicology Letters*, vol. 48, no. 3, pp. 301–310, 1989.
- [58] G. Zhang, V. Nitteranon, S. Guo et al., "Organoselenium compounds modulate extracellular redox by induction of extracellular cysteine and cell surface thioredoxin reductase," *Chemical Research in Toxicology*, vol. 26, no. 3, pp. 456–464, 2013.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

