Clinical Study

The Role of Hemoclips Reinforcement in the Ligation-Assisted Endoscopic Enucleation for Small GISTs in Gastric Fundus

Ge Nan, Sun Siyu, Wang Sheng, Liu Xiang, and Guo Jintao

The Shengjing Hospital, China Medical University, No. 36 Sanhao Street, Shenyang, Liaoning 110004, China

Correspondence should be addressed to Sun Siyu; sunsy@sj-hospital.org

Received 7 February 2014; Revised 14 April 2014; Accepted 20 April 2014; Published 7 May 2014

Academic Editor: Steven N. Hochwald

Copyright © 2014 Ge Nan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Endoscopic ultrasonography- (EUS-) assisted band ligation has been proven to be a safe and effective procedure for the treatment of small gastrointestinal stromal tumors (GISTs) apart from the relatively high risk of the postligation perforation of the gastric fundus. The aim of this study is to investigate the efficacy of hemoclip reinforcement in treating small GISTs in the gastric fundus. **Method.** During a standard endoscopy, a transparent cap attached to the endoscopic tip was placed over the lesion to exert sustained maximal aspiration before a rubber band was released. Once a definite ligation was confirmed by EUS, the tumor was enucleated. Four to 6 hemoclips were placed on the folds around the ligation band to reduce the tension of the ligation site. **Results.** The small GISTs were resected completely in 192 patients. Two cases of delayed perforation were found 72 hours after the procedure and successfully treated with an ordinary conservative method. **Conclusion.** Hemoclip-reinforced endoscopic band ligation with systematic follow-up using EUS appears to be a simple and effective technique for the resection of small GISTs in the gastric fundus.

1. Introduction

GISTs are a type of neoplasm derived from gastrointestinal mesenchymal tissues and contain spindle cells and positive CD117 expression [1]. Currently, no therapeutic guideline is available for the treatment of asymptomatic small GISTs with a low degree of malignancy, as identified by EUS. The common approach to the management of small GISTs is periodical follow-up examination with EUS after endoscopic resection. Sun et al. [2–4] have reported on the application of EUS-assisted band ligation to treat small GISTs. Although this technique has been proven to be safe and effective, delayed perforation was recently reported as a severe complication, especially for GISTs localized in the gastric fundus [5]. The purpose of this study was to evaluate an improved ligation technique by placing 4 to 5 hemoclips around the ligation band to prevent perforation and complete the tumor enucleation above the band to obtain a pathological diagnosis. The study was approved by the Institutional Review Board of China Medical University.

2. Materials and Methods

2.1. Patients. A total of 192 patients with small tumors originating from the muscularis propria in the gastric fundus treated at Shengjing Hospital between May 2010 and June 2013 were included in this study. All patients gave informed consent for this procedure. Inclusion criteria were as follows. (1) Lesion was localized to the muscularis propria of the gastric fundus as confirmed by EUS (Figure 1). (2) Tumor diameter was no larger than 12 mm measured with EUS (corresponding to the diameter of the transparent hood used for ligation). (3) Patients did not take aspirin or other nonsteroid anti-inflammatory drugs for at least 1 week prior to the procedure. (4) Complete blood counts, prothrombin time, and partial thromboplastin time were normal for all patients.

2.2. Devices. Ligation was carried out with a conventional 2.7 mm channel endoscope (EG2770K, Pentax) and an air-driven band ligator (Sumitbe, Akita, Japan). EUS was performed with a curved linear-array echoendoscope

FIGURE 2: (a) The tumor was ligated completely. (b) A hook knife or triangle knife was used to cut the surrounding mucosa open above the band.

FIGURE 3: The tumor was dissected from the muscular layer and retrieved by the foreign body forceps.

FIGURE 5: Emergent CT scanning showed a small perforation of the ligation site after the band exfoliation.

FIGURE 4: Four to six hemoclips were placed on the folds around the ligation band to reduce tension of the ligation site.

FIGURE 6: Complete healing of gastric mucosa after ligation.

Traditionally, surgical resection is the method of choice for nonmetastatic GISTs. DeMatteo showed that survival is not affected by the microscopic margin of resection. Routine extensive lymphadenectomy is not warranted because lymphatic involvement is extremely rare in GISTs [11, 12]. Therefore, minimal invasive intervention including several endoscopic, laparoscopic, and combined techniques is a more rational choice for GISTs and has resulted in lower mortality and earlier recovery [13]. Endoscopy is the least invasive

technique and has more advantages for elderly patients and those with high surgical risks, especially in treating small tumors.

Various methods including ESD and NOTES for complete endoscopic resection of GISTs have been reported. ESD technique is difficult to master because it requires thorough knowledge and skills in performing endoscopic hemostasis, wound closure, and use of devices under endoscopy, so as

to avoid severe complications. Sometimes, for GISTs tightly adherent to the muscularis propria, lesions have not been able to be resected completely by ESD [14]. Few resections have been done in the gastric fundus when the tumors are localized within muscularis propria. The fundic wall is very thin so that resection is often accompanied by simultaneous perforation.

Sun et al. have reported that small esophageal leiomyomas [2] GISTs, and duodenal stromal tumors could be resected safely using endoscopic band ligation techniques. The band sloughs spontaneously. When aspiration and band ligation were performed using a cap-mounted endoscope, all layers of the gastrointestinal tract and GISTs were ligated. The resultant ischemia and necrosis would cause sloughing of the band and the ligated gastric wall. The healing of the gastric wall is by gradual formation of serosal adhesion outside the band, in response to local inflammation. No perforation or other complications were reported, except for the tumors located in the gastric fundus [5]. As reported by Sun et al. in 2010, two patients with a GIST located at the gastric fundus complained of severe abdominal pain, one at 24 hours and one at 35 hours after the procedure. Upright abdominal pain films confirmed bilateral subphrenic free air, and both patients underwent emergent laparotomy. At operation, circular perforations of 2–2.5 cm diameter were found with gastric mucosa that appeared inverted. Those two cases of severe complications have limited the application of ligation technique in the gastric fundus.

In this study, a novel technique for complete resection of GISTs using hemoclip-reinforced ligation was demonstrated. GISTs were safely resected above rubber bands with 4 to 5 hemoclips placed on the folds around the ligation band. Medical adhesive was sprayed onto the surface to immobilize these clips.

Since all the reported gastric perforations occurred after endoscopic ligation at the fundus, premature sloughing of lesions may account for the perforation, although the exact histopathologic mechanisms of “all-layer” ligation is unknown. Premature sloughing of ligated gastric wall may result from the gastric wall being comparatively thin at this location. The band can often exfoliate prematurely during fundic ligation. Premature exfoliation of the band can cause perforation due to still-incomplete serosal adhesion. The sooner the band has exfoliated, the greater the risk of postprocedural complications is, including bleeding and perforation. We placed 4 to 5 hemoclips on the folds around the ligation band to reduce tension of the gastric wall and hoped this would be sufficient to delay the exfoliation of lesions, since perforation could be avoided when adhesion occurred ahead of exfoliation. In our study, two cases of perforation occurred 72 hours after the procedure, which was far longer than the cases reported in 2010 [5]. Also, the incomplete healing of the serosa revealed in CT scanning was not that large (compared with the 2.5 cm diameter perforation in 2010) and was only a small incompletely-healed defect. In our study, although the risk of perforation could still not be avoided, the patients with perforation did not suffer from complications of severe peritoneal cavity inflammation and require surgical intervention. The patients recovered with conservative treatment with a decompression

tube and extended fasting time. The hemoclips may play an important role in the delayed sloughing time. The exact mechanism should be further tested by the animal study.

Hemoclips, originally developed 3 decades ago in Japan by Jensen and Machicado [15], have been used for endoscopic hemostasis since then and their use has been expanded to the management of many types of perforations and fistulas. However, their application in reinforcing band ligation has not previously been described. The use of small hemoclips can help reduce tension of the gastric wall, a much easier task than repairing a defect in the wall. Hemoclips placed around the ligation band could prevent premature exfoliation of the tumors effectively and sustain a decompressive effect, even after exfoliation of tumors.

It is also the first time that medical adhesive was used in strengthening ligation. Compound medical adhesive was originally developed for hemostasis and wound sealing. The main purpose of using medical adhesive in this procedure was to immobilize the hemoclips firmly and prevent their premature detachment from the gastric wall effectively. In this study, the retention time for these hemoclips ranged from 1 week to 1 year, much longer than the 2 weeks reported by Jensen et al. [16].

In conclusion, hemoclip-reinforced endoscopic band ligation with systematic follow-up using EUS appears to be a simple and effective technique for resection of small GISTs in the fundus by decreasing the risk of perforation. Although it cannot completely prevent perforation during the band exfoliation process, it can minimize a peritoneal cavity inflammation reaction and avoid the following necessary surgical intervention. Patients with GISTs located in gastric fundus should prolong the fasting time after the ligation.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] S. W. Jeon, Y. D. Park, Y. J. Chung et al., “Gastrointestinal stromal tumors of the stomach: endosonographic differentiation in relation to histological risk,” *Journal of Gastroenterology and Hepatology*, vol. 22, no. 12, pp. 2069–2075, 2007.
- [2] S. Sun, N. Ge, C. Wang, M. Wang, and Q. Lü, “Endoscopic band ligation of small gastric stromal tumors and follow-up by endoscopic ultrasonography,” *Surgical Endoscopy and Other Interventional Techniques*, vol. 21, no. 4, pp. 574–578, 2007.
- [3] S. Sun, Y. Jin, G. Chang, C. Wang, X. Li, and Z. Wang, “Endoscopic band ligation without electrosurgery: a new technique for excision of small upper-GI leiomyoma,” *Gastrointestinal Endoscopy*, vol. 60, no. 2, pp. 218–222, 2004.
- [4] S. Sun, N. Ge, S. Wang, X. Liu, and Q. Lü, “EUS-assisted band ligation of small duodenal stromal tumors and follow-up by EUS,” *Gastrointestinal Endoscopy*, vol. 69, no. 3, pp. 492–496, 2009.
- [5] S. Siyu, W. Sheng, W. Guoxin, G. Nan, and L. Jingang, “Gastric perforations after ligation of GI stromal tumors in the gastric fundus,” *Gastrointestinal Endoscopy*, vol. 72, no. 3, pp. 615–616, 2010.

- [6] A. Darnell, E. Dalmau, C. Pericay et al., "Gastrointestinal stromal tumors," *Abdominal Imaging*, vol. 31, no. 4, pp. 387–399, 2006.
- [7] J. Guo, Z. Liu, S. Sun et al., "Endosonography-assisted diagnosis and therapy of gastrointestinal submucosal tumors," *Endoscopic Ultrasound*, vol. 2, pp. 125–133, 2013.
- [8] H. L. Evans, "Smooth muscle tumors of the gastrointestinal tract. A study of 56 cases followed for a minimum of 10 years," *Cancer*, vol. 56, no. 9, pp. 2242–2250, 1985.
- [9] S. J. Kwon, "Surgery and prognostic factors for gastric stromal tumor," *World Journal of Surgery*, vol. 25, no. 3, pp. 290–295, 2001.
- [10] Y. Fujimoto, Y. Nakanishi, K. Yoshimura, and T. Shimoda, "Clinicopathologic study of primary malignant gastrointestinal stromal tumor of the stomach, with special reference to prognostic factors: analysis of results in 140 surgically resected patients," *Gastric Cancer*, vol. 6, no. 1, pp. 39–48, 2003.
- [11] E. M. Connolly, E. Gaffney, and J. V. Reynolds, "Gastrointestinal stromal tumours," *British Journal of Surgery*, vol. 90, no. 10, pp. 1178–1186, 2003.
- [12] B. K. P. Goh, P. K. H. Chow, S. Kesavan, W. Yap, and W. Wong, "Outcome after surgical treatment of suspected gastrointestinal stromal tumors involving the duodenum: is limited resection appropriate?" *Journal of Surgical Oncology*, vol. 97, no. 5, pp. 388–391, 2008.
- [13] C. Dholakia and J. Gould, "Minimally invasive resection of gastrointestinal stromal tumors," *Surgical Clinics of North America*, vol. 88, no. 5, pp. 1009–1018, 2008.
- [14] I.-L. Lee, P. Y. Lin, S.-Y. Tung, C.-H. Shen, K.-L. Wei, and C.-S. Wu, "Endoscopic submucosal dissection for the treatment of intraluminal gastric subepithelial tumors originating from the muscularis propria layer," *Endoscopy*, vol. 38, no. 10, pp. 1024–1028, 2006.
- [15] D. M. Jensen and G. A. Machicado, "Hemoclipping of chronic canine ulcers: a randomized, prospective study of initial deployment success, clip retention rates, and ulcer healing," *Gastrointestinal Endoscopy*, vol. 70, no. 5, pp. 969–975, 2009.
- [16] D. M. Jensen, G. A. Machicado, and K. Hirabayashi, "Randomized controlled study of 3 different types of hemoclips for hemostasis of bleeding canine acute gastric ulcers," *Gastrointestinal Endoscopy*, vol. 64, no. 5, pp. 768–773, 2006.