Research Article

The Renal Effects of Vanadate Exposure: Potential Biomarkers and Oxidative Stress as a Mechanism of Functional Renal Disorders—Preliminary Studies

Agnieszka Ścibior,^{1,2} Dorota Gołębiowska,^{1,2} Agnieszka Adamczyk,¹ Irmina Niedźwiecka,¹ and Emilia Fornal³

¹ Laboratory of Physiology and Animal Biochemistry, Department of Zoology and Invertebrate Ecology, Institute of Environmental Protection, The John Paul II Catholic University of Lublin, 102 Kraśnicka Avenue, 20-718 Lublin, Poland

² Laboratory of Oxidative Stress, Center for Interdisciplinary Research, The John Paul II Catholic University of Lublin, 102 Kraśnicka Avenue, 20-718 Lublin, Poland

³ Laboratory of Separation and Spectroscopic Method Applications, Center for Interdisciplinary Research, The John Paul II Catholic University of Lublin, 102 Kraśnicka Avenue, 20-718 Lublin, Poland

Correspondence should be addressed to Agnieszka Ścibior; cellbiol@kul.lublin.pl

Received 30 April 2013; Revised 22 October 2013; Accepted 28 October 2013; Published 28 January 2014

Academic Editor: Maha Zaki Rizk

Copyright © 2014 Agnieszka Ścibior et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The alterations in the levels/activities of selected biomarkers for detecting kidney toxicity and in the levels of some oxidative stress (OS) markers and elements were studied in male rats to evaluate biochemically the degree of kidney damage, investigate the role of OS in the mechanism of functional renal disorders, reveal potential biomarkers of renal function, and assess the renal mineral changes in the conditions of a 12-week sodium metavanadate (SMV, 0.125 mg V/mL) exposure. The results showed that OS is involved in the mechanism underlying the development of SMV-induced functional renal disturbances. They also suggest that the urinary cystatin C (CysC_u) and kidney injury molecule-1 (KIM-1_u) could be the most appropriate to evaluate renal function at the conditions of SMV intoxication when the fluid intake, excreted urinary volume (EUV), body weight (BW), and the urinary creatinine excretion (Cre_u) decreased. The use of such tests as the urinary lactate dehydrogenase, alkaline phosphatase, γ -glutamyltranspeptidase, and N-acetyl- β -D-glucosaminidase (LDH_u, ALP_u, GGTP_u, and NAG_u) seems not to be valid given their reduced activities. The use of only traditional biomarkers of renal function in these conditions may, in turn, be insufficient because their alterations are greatly influenced by the changes in the fluid intake and/or BW.

1. Introduction

Vanadium (V) is a well-known powerful prooxidant. It may modify oxidative stress (OS) in the cells and be involved in oxidative injury mechanisms [1]. Its prooxidant action has been demonstrated in *in vivo* and *in vitro* conditions [2–7]. It has been shown that the free radical process-lipid peroxidation (LPO), which is a biochemical biomarker of cellular dysfunction and an index of cytotoxicity [8], is enhanced by V in the kidney [3, 8]. It has also been suggested that LPO may be predictive of renal dysfunction [8].

Kidney is particularly vulnerable to deleterious effects of V. The susceptibility of this organ to V may be a reflection of its accumulation in this tissue [8]. It has been reported that V may be present in tubular cells in a readily exchangeable form as well as in low and high molecular mass complexes, and it may be excreted free or bound to proteins after prolonged exposure [9, 10]. It has also been speculated that V may be involved in the pathogenesis of distal renal tubular acidosis (dRTA), renal stone disease, “uremic syndrome,” and acquired cystic kidney disease [11–13]. A suggestion that prolonged intake of high-dose V supplements may cause serious kidney toxicity has been put forward as well [14, 15].

The occupational and environmental toxicological impact of V and the fact that kidneys are critical for its poisoning [16–19], the modest number of biomarkers of possible functional renal disturbances under vanadate exposure examined until now, and the insufficient information about the contribution of OS in the mechanism underlying the vanadate-induced functional kidney disorders prompted us to answer the following questions. (a) To what extent do the effects of 12-week sodium metavanadate (NaVO_3 , SMV, 0.125 mg V/mL) exposure alter the levels/activities of some biomarkers of renal toxicity in rats? We intended to examine traditional biomarkers classified as not very specific or sensitive but routinely used in the diagnosis of kidney function and those that allow distinguishing, to some degree, structural and functional renal disorders and have a potential for determining the site of renal tubular damage [20–22]. (b) To what degree do the effects of SMV exposure change the homeostasis of some micro- and macroelements in the kidney? (c) Which of the biomarkers examined are the most sensitive in our experimental conditions? (d) What value/information do some biomarkers add to the existing data? (e) Is OS involved in the mechanism of the development of kidney function disorders during SMV intoxication? (f) Are there any significant relationships between the measured variables? Since the measurement of the activities of some enzymes in tissues and biological fluids may play a significant role in detection of tissue cellular injury and point to damage long before histological alterations, both cytosolic and lysosomal enzymes as well as those located on the brush-border membrane have been taken into consideration and illustrated in the present report.

2. Materials and Methods

2.1. Reagents. The kits for determination of the plasma concentrations of TP_p , U_p , UA_p , Mg_p , and Ca_p and the urinary levels of U_u , UA_u , and Cre_u were obtained from Emapol (Gdańsk, Poland), whereas the kits for determination of the plasma Cu_p and Zn_p levels were purchased from Sentinel Ch. kits (Milan, Italy). The reagents for determination of the levels/activities of the plasma (Cre_p , ALB_p , LDH_p , GGTP_p , and ALP_p), urinary (ALB_u , TP_u , LDH_u , GGTP_u , and ALP_u), and renal (LDH_k , GGTP_k , and ALP_k) biomarkers as well as the reagent for measurement of the levels of electrolytes in the plasma and urine ($\text{Na}_{p/u}$, $\text{K}_{p/u}$, and $\text{Cl}_{p/u}$) were acquired from Alpha Diagnostics (Warsaw, Poland). The kit for estimation of total antioxidant status in the kidney (TAS_k) was bought from Calbiochem-Novabiochem Corporation (San Diego, CA, USA), whereas the kits for determination of the urinary levels of NAG_u (E90069Ra), CysC_u (E90896Ra), $\beta_2\text{M}_u$ (E90260Ra), and KIM-1_u (E90785Ra) and of the plasma concentrations of CysC_p (E90896Ra) and ($\beta_2\text{M}_p$ E90260Ra) were derived from Uscn Life Science Inc. (Wuhan, China). NaVO_3 , Triton X-100, and the caesium chloride lanthanum chloride buffer (CsClLaCl) were purchased from Sigma Chemical (St. Louis, USA). Nitric acid (HNO_3 , 65% suprapure) and diethyl ether ($\text{C}_4\text{H}_{10}\text{O}$) were acquired from Merck (Darmstadt, Germany), whereas hydrogen peroxide (H_2O_2 ; 30% pure P.A.) and the physiological buffered saline (PBS)

were purchased from POCH (Gliwice, Poland) and from the Serum and Vaccine Factory (Biomed, Lublin, Poland), respectively. Stocks of V, Mg, Zn, and Cu (Inorganic Ventures, Christiansburg, USA) and a stock of Ca (Spectracer, UK) atomic absorption standard solutions as well as a multi-element standard solution for Na and K (Sigma-Aldrich) were used in the element analysis by the Atomic Absorption Spectrometry (AAS) method. Ultrapure water was received from an ultrapure water HLP Spring 5R system¹ (Hydrolab, Gdańsk, Poland). All the chemicals were of the highest quality available.

2.2. Animals and Experimental Design. The experiment was conducted according to the experimental protocol approved by the 1st Local Ethical Committee for Animal Studies in Lublin. Biological material used in this study originated from some of the outbred albino male Wistar rats used in the previous study [23]. The animals were kept in an animal room with controlled conventional conditions (one rat per stainless steel cage) and received, inter alia, deionised water (Group I, Control, 8 rats) or a water solution of NaVO_3 at a concentration of 0.125 mg V/mL (Group II, SMV, 8 rats) to drink in special bottles with the scale every day over a 12-week period. All the rats had *ad libitum* access to fresh deionised water, SMV solution, and a standard rodent diet (Labofeed B, Fodder and Concentrate Factory, Kcynia, Poland) in which the concentration of V had been assessed by Graphite Furnace AAS (GF-AAS) in our laboratory, and it was about 0.17 μg V/g. More details about the chow had been provided by us previously [23]. The intake of food, water, and SMV solution was monitored daily and body weight (BW) was obtained weekly. The water and SMV solution intake was expressed as mL/rat/24 h, whereas the food intake was expressed as g/rat/24 h. The daily V intake in the SMV-exposed rats was estimated based on the 24 h consumption of the drinking SMV solution and expressed as mg V/kg b. wt./24 h. During the whole experiment, the animals were observed in order to assess their general health. The doses of V presented in the study are within the broad dose range that was used for demonstrating the antidiabetic activity of V [19, 24] and for analysing its pharmacokinetic behaviour [25] on an animal model. The concentration of V in rats' urine determined in the present study may reflect exposure to this element occurring especially in persons occupationally exposed to this metal [26, 27].

Every second week in the course of the experiment and in the 11th week, 24-hour urine was collected from each rat placed individually in plastic metabolic cages not equipped with a cooling system (Tecniplast, Italy), which allowed separate collection of urine and faeces. During that time, each rat had access to food and water or the SMV solution. The urine samples from all the control and SMV-exposed rats were used immediately for determination of some biochemical parameters and for measurements of excreted urine volume (EUV) and urine pH. The volumes of excreted 24-hour urine were measured with a measuring cylinder, whereas urine pH and blood in the urine were tested on the H-100 urine analyser (DIRUI, China) using urinary dipsticks. Portions of the urine samples that were not used immediately were frozen

at -80°C in a deep-freezer HFU 486 Basic¹ (Thermo Fisher Scientific, Germany) and stored until further analyses.

All the rats were sectioned at the end of week 12. Whole blood was taken from the jugular vein into plastic tubes with heparin as an anticoagulant under anaesthesia with ketamine/xylazine cocktail (100 mg/mL and 20 mg/mL, resp., i.p.) and centrifuged (5 min, $1500 \times g$, 4°C). Plasma portions were collected for routine analyses of clinical chemistry parameters and for other biological determinations. The kidneys and femurs (right and left) were immediately removed. The kidneys were washed in ice-cold physiological saline (0.9% NaCl) and weighted. The right femurs (after removal of the overlying tissue with stainless steel knives) were also washed in 0.9% NaCl, weighed, and stored frozen at -80°C until the time of bone digestion. Before digestion, all the collected right femurs were cut using a diamond-disk saw (Metkon Micracut 175)¹, cooled with ultrapure water to separate the proximal and distal femoral epiphysis (PFE and DFE) (a region of trabecular bone) from the femoral diaphysis (FD) (a region of cortical bone). Next, bone marrow was removed from all the FDs, which then were first soaked with ether (to remove fat content) and later in H_2O_2 (to remove remaining blood deposits). Afterwards, the cleaned FDs were washed in ice-cold PBS and then in ultrapure water. Next, they were dried at room temperature into constant mass. The cleaned and bone marrow-deprived FD samples (~ 0.355 g, mean) were used for digestion.

2.3. Decomposition of the Kidney, Urine, and FD. In order to determine Mg, Ca, V, Zn, Cu, Na, and K in the kidney and Mg, Ca, V, Zn, and Cu in the urine, 0.5 g of kidney and 1 mL of urine were wet-mineralized with 5 mL of 65% HNO_3 in 12 Teflon Fluor Modified (TFM) closed digestion vessels using a model Speedwave Four microwave digestion system¹ (Berghof, Germany) equipped with a temperature and pressure sensor in each vessel. In turn, ~ 0.355 g of FD was wet-mineralized in the same digestion system but in the presence of 5 mL of 65% HNO_3 and 1 mL of 30% H_2O_2 to determine V, Mg, and Ca. Before the measurements, all the decomposed samples of kidney, FD, and urine were transferred into 25 mL volumetric polypropylene flasks by washing the inner surface of the digestion vessels with ultrapure water three times and filled up to the mark with ultrapure water. Next, determinations of the selected elements were performed by the AAS method.

2.4. Determination of Some Renal Biomarkers. The concentrations of U_p , UA_p and TP_p in the plasma as well as U_u , UA_u , and Cre_u in 24-hour urine were determined colourimetrically by measuring the absorbance with a Thermo Spectronic BioMate5 UV-VIS spectrophotometer (UK). The concentrations of ALB_p , ALB_u , TP_u , and Cre_p and the activities of LDH_p , GGTP_p , ALP_p , LDH_u , GGTP_u , and ALP_u in the plasma and urine as well as the activities of LDH_k , GGTP_k , and ALP_k in the kidney were measured using an automatic biochemical analyser BS-120¹ (Mindray, China). The levels of CysC_p and $\beta_2\text{M}_p$ in the plasma and the levels of CysC_u , $\beta_2\text{M}_u$, KIM-1_u , and NAG_u in the urine were assessed by a

traditional method using rat-specific commercial enzyme-linked immunosorbent assay (ELISA) kits and an ELISA microplate reader Synergy 2¹ equipped with an automated microplate strip washer ELx50¹ and a microplate shaker¹ (BIO-TEK Instruments Inc., USA). All the ELISA tests were performed according to the manufacturer's protocols. Before the measurements, thawed plasma and urine samples were mixed by inversion, centrifuged with cooling (1500 rpm for 10 min, 4°C , or 2000 rpm for 5 min, 4°C , resp.) using a centrifuge Heraeus Megafuge 11R¹ (Thermo Fisher Scientific, Germany), and immediately used for analysis. The optimal factors of dilution for some samples were chosen when necessary and the samples were diluted with ultrapure water or with PBS ($\text{pH } 7.4 \pm 0.2$). In order to evaluate the glomerular filtration rate (GFR), calculation of creatinine clearance (CreC) was performed. The results for all the above-mentioned parameters are expressed as % of the control and presented in Figures 1(a), 2(a), 2(b), 3(a), 3(b), and 4(a).

2.5. Determination of Elements in Biological Fluids and Tissues

2.5.1. Spectrophotometric Determination of Ca, Mg, Zn, and Cu in the Plasma. The measurements of these elements were performed using a BioMate5 spectrophotometer with the direct colourimetric method according to the protocols of the kits, and their concentrations are presented in Figures 1(b) and 1(d).

2.5.2. Determination of Na, K, and Cl in the Plasma and Urine. The electrolytes mentioned were measured using an automatic EasyLyte analyser Na/K/Cl¹ (Medica). Their urinary excretion was first normalized against 24-hour diuresis and against 24-hour urinary Cre_u excretion, and their concentrations are presented in Figures 1(b), 1(c), 2(e), and 3(e).

2.5.3. Atomic Absorption Measurements of Mg, Ca, V, Zn, Cu, Na, and K in Digested Kidney Samples, Mg, Ca, V, Zn, and Cu in Digested Urine Samples, V, Mg, and Ca in Digested FD Samples, and V in Nondigested Plasma Samples. The elements were determined by Flame or Graphite Furnace AAS (F-AAS or GF-AAS, resp.) using a SpectrAA Z-2000 TANDEM atomic absorption spectrometer¹ (Hitachi, Japan) equipped with a Zeeman background corrector. A specific matrix modifier was used for V determination. In order to determine the V concentration in the plasma, the samples were diluted with 0.05% Triton X-100 and in ultrapure water when necessary. All the operating parameters of the instrument and the details of measurements of all the above-mentioned elements together with the values of the detection, quantification limits (LOD and LOQ, resp.), and the coefficient of variations (CV) are shown in Table 1.

The method of standard addition was performed in order to estimate the effect of interferences during Mg assessment. Mg, Ca, V, Zn, Cu, Na, and K were determined by application of a calibration curve using working standard solutions, which were obtained from a stock atomic absorption standard solutions containing 1000 μg Mg, Ca, V, Zn, Cu, Na, and

TABLE 1: Operating parameters of the atomic absorption spectrometer with details of measurement of the levels of elements in biological samples as well as the certified and determined values of elements for the selected Certified Reference Materials (CRMs).

Parameters	Elements						
Technique	V	Cu	Mg	Ca	Zn	Na	K
FT	GF-AAS	GF-AAS			F-AAS Air-acetylene		
FF (L/min)	Argon with the flow rate of 200 (mL/min) in all the steps except the atomization stage when the flow rate was 30 (mL/min)						
					1.8		
GTT	PCGTs (PyroTube CHR)						
SV (μ L)	20 (μ L)						
SM	B-CIA						
LC (mA)	10	7.5	7.5	10	B-CIM 5.0	10.0	10.0
WL (nm)	318.4	324.8	285.2	422.7	213.9	589.0	766.5
SW (nm)	1.3	1.3	1.3	0.2	1.3	0.2	1.3
	K: 0.11 (μ g/L)	K: 0.28 (μ g/L)	K: 1×10^{-5} (mg/L)	K: 0.09 (mg/L)	K: 1×10^{-3} (mg/L)	K: 3.1×10^{-3} (mg/L)	K: 5.5×10^{-3} (mg/L)
	U: 0.23 (μ g/L)	U: 0.14 (μ g/L)	U: 6×10^{-5} (mg/L)	U: 0.36 (mg/L)	U: 0.6 (μ g/L)	U: **	U: **
	P: 0.23 (μ g/L)	P: $\checkmark$	P: $\checkmark$	P: $\checkmark$	P: $\checkmark$	P: **	P: **
DL (LOD)	FD: 0.47 (μ g/L)		FD: 8×10^{-4} (mg/L)	FD: 0.13 (mg/L)			
	K: 0.33 (μ g/L)	K: 0.84 (μ g/L)	K: 3×10^{-5} (mg/L)	K: 0.27 (mg/L)	K: 3×10^{-3} (mg/L)	K: 9.3×10^{-3} (mg/L)	K: 16.5×10^{-3} (mg/L)
	U: 0.69 (μ g/L)	U: 0.42 (μ g/L)	U: 1.8×10^{-4} (mg/L)	U: 1.08 (mg/L)	U: 1.8 (μ g/L)	U: **	U: **
	P: 0.69 (μ g/L)	P: $\checkmark$	P: $\checkmark$	P: $\checkmark$	P: $\checkmark$	P: **	P: **
	FD: 1.41 (μ g/L)		FD: 2.4×10^{-3} (mg/L)	FD: 0.39 (mg/L)			
LOQ							
	0.2-2	0.5-2			0.5-1		
CV (%)							

TABLE 1: Continued.

Elements	Bovine Liver 1577 c		Seronorm Trace Elements Urine 201205		Trace Elements in Natural Water I640 a	
	Certified value	Determined value [§]	Certified value	Determined value [§]	Certified value	Determined value [§]
Mg	620 ± 42 (mg/kg)	657.4 ± 20.2 (mg/kg)	71.1 ± 2.5 (mg/L)	69.7 ± 2.3 (mg/L)	1.058 ± 0.0040 (mg/L)	1.059 ± 0.040 (mg/L)
Ca	131 ± 10 (mg/kg)	133.4 ± 15.2 (mg/kg)	111 ± 2 (mg/L)	111.9 ± 1.2 (mg/L)	5.615 ± 0.021 (mg/L)	5.649 ± 0.028 (mg/L)
V	8.17 ± 0.66 (µg/kg)	10.2 ± 1.6 (µg/kg)	25.2 ± 1.4 (µg/L)	22.43 ± 0.56 (µg/L)	12.99 ± 0.37 (µg/L)	12.33 ± 0.35 (µg/L)
Zn	181.1 ± 1.0 (mg/kg)	180.3 ± 8.7 (mg/kg)	1141 ± 79 (µg/L)	1100.6 ± 24.4 (µg/L)	55.64 ± 0.35 (µg/L)	55.50 ± 0.17 (µg/L)
Cu	275.2 ± 4.6 (mg/kg)	268.6 ± 4.7 (mg/kg)	78 ± 8 (µg/L)	92.4 ± 6.9 (µg/L)	85.75 ± 0.51 (µg/L)	85.45 ± 0.14 (µg/L)
Na	2.033 ± 0.064 (mg/kg)	1.969 ± 0.015 (mg/kg)	2307 ± 56 (mg/L)	2376 ± 45 (mg/L)	3.137 ± 0.031 (mg/L)	3.150 ± 0.025 (mg/L)
K	—	—	1903 ± 42 (mg/L)	1930 ± 40 (mg/L)	0.5799 ± 0.0023 (mg/L)	0.5818 ± 0.0209 (mg/L)
Elements	Bone Ash 1400					
	Certified value				Determined value [§]	
Mg	6.84 ± 0.13 (mg/g)				6.774 ± 0.065 (mg/g)	
Ca	381.8 ± 1.3 (mg/g)				384.8 ± 2.2 (mg/g)	

FT: flame type; FF: fuel flow; GTT: graphite tube type; SV: sample volume; SM: signal mode; LC: lamp current; WL: wavelength; SW: slit width; PCGTs: pyrolytically coated graphite tubes; B-CIA: background-corrected integrated absorbance; B-CIM: background-corrected integral mode.
✓*** Determined colourimetrically or by an EasyLyte analyser, respectively.
DL (LOD): detection limit.
LOQ: quantification limit.
CV: coefficient of variation.
K: kidney; U: urine; P: plasma; FD: femoral diaphysis.
§ Mean ± SD, n = 5.

FIGURE 1: The levels/activities of some biomarkers of renal toxicity (a) and the concentrations of selected elements ((b)–(e)) in rat plasma. ^{1,2,3}Data were tested by Student's *t*-test, Welch's *t*-test, or Mann-Whitney's *U* test, respectively. *Significant differences, compared with the Control (Group I). ^fLogarithmically transformed data. ^{††} $P = 0.05$, $P = 0.08$, respectively, compared with the Control (Group I).

K/mL by dilution with 5% (v/v) HNO₃. The 10% CsCl/LaCl buffer was used to determine Ca. The analytical quality of the measurements was checked with the use of Certified Reference Materials (CRMs) such as Bovine Liver 1577 c (NIST), Seronorm Trace Elements Urine 201205 (SERO), Trace Elements in Natural Water 1640 a (NIST), and Bone Ash 1400 (NIST). The analysis of these CRMs confirmed the reliability of the proposed approach. The certified and determined values of all the elements examined in the above-mentioned CRMs are presented in Table 1. The plasma V level is presented in Figure 1(e). In turn, the urinary excretion of the above-mentioned elements investigated was also first normalized against 24-hour diuresis and against 24-hour urinary Cre_u excretion. This is illustrated in Figures 2(c), 2(d), 2(f), 3(c), 3(d), and 3(f). However, the renal V, Mg, Ca, Zn, Cu, Na, and K concentrations as well as the FD V, Mg, and Ca concentrations are presented in Figures 4(b)–4(e) and in Figures 5(a)–5(c).

2.6. Determination of Renal Lipid Peroxidation (LPO) and Total Antioxidant Status (TAS). All details concerning

the methodology of determination of LPO and the preparation of kidney homogenates and supernatants for the renal MDA_k and TAS_k measurements had already been provided [2, 3]. The results of both the OS markers mentioned are expressed as % of control and illustrated in Figure 4(a).

2.7. Statistical Analysis. The statistical analysis of the data was performed with the Statistica and SPSS, versions 9.0 and 14.0 PL for Windows, respectively. The normal distribution of the data was tested by Shapiro-Wilk's normality test. Grubbs' test was performed to detect the presence of outliers from a normal distribution. The homogeneity of variances was verified employing Levene's test and sometimes additionally Hartley's F_{max}, Cochran's C, and Bartlett's tests. Student's *t*-test and Welch's *t*-test were applied to compare the means of two independent groups when the data met the assumptions of ANOVA and when they had a normal distribution but the variances were not homogenous, respectively. In some cases, logarithmic transformation of the data was used to make them more normal. In turn, the nonparametric Mann-Whitney *U*

FIGURE 2: EUV, urine pH, and the urinary levels/activities of some biomarkers of renal toxicity ((a) and (b)) and the urinary levels of selected elements ((c)–(f)) normalized per 24-hour diuresis in the tested rats.^{1,2,3} Data were tested by Student's *t*-test, Welch's *t*-test, or Mann-Whitney's *U* test, respectively. *Significant differences, compared with the Control (Group I). [†]Logarithmically transformed data. [‡]Correlated with excreted urinary volume (EUV). ^{††}*P* = 0.06, compared with the Control (Group I).

test was applied when the data did not meet the assumptions of ANOVA. The results were presented as mean $\pm$ the standard error of the mean (SEM). A *P* value less than 0.05 was taken as a criterion for a statistically significant difference. Pearson's correlation analyses were applied to assess the relationships between the measured variables. Correlations were considered statistically significant at *P* < 0.05.

3. Results

3.1. General Observation and Changes in Some Basic Parameters. In the animals exposed to SMV, no distinct differences in physical appearance and motor behaviour were observed during the 12-week experimental period, compared with the control. Some of the SMV-intoxicated rats had gastrointestinal disturbances probably caused by the consumption of the V dose. One-day diarrhoea was observed in four animals from this group in the first, second, third, or fourth week of the experiment. Only one rat from this group had three-day diarrhoea in the first week of the experiment. The fluid and

food intake as well as BW (Table 2) and EUV (Figure 2(a)) in the SMV-exposed rats were lower, compared with the control animals. The dose of V consumed by the rats during the 12-week period reached the value of about 13 mg V/kg b. wt./24 h (Table 2).

3.2. Plasma, Urinary, and Renal Levels of the Examined Biomarkers and the Dipstick Urinalysis (Blood and pH). The concentrations of Cre_p and U_p in the SMV-exposed rats increased (Figure 1(a)), whereas CreC (Figure 1(a)) and the levels of urinary excretion of Cre_u (per 24 h) and U_u (per 24 h) decreased significantly (Figure 2(a)), in comparison with the control animals. In turn, the urinary level of U_u (per Cre_u/24 h) (Figure 3(a)) and the concentration of U_k in the kidney (Figure 4(a)) were unaltered. The levels of UA_p (Figure 1(a)) and UA_u (per 24 h and per Cre_u/24 h, Figures 2(a) and 3(a), resp.) also remained unchanged in response to the SMV exposure.

The levels of TP_p and ALB_p in the SMV-exposed rats did not alter significantly, compared with the control (Figure 1(a)), but the urinary excretion of TP_u (per 24 h)

FIGURE 3: The urinary levels/activities of some biomarkers of renal toxicity ((a) and (b)) and the urinary levels of selected elements ((c)–(f)) normalized per 24-hour urinary Cre_u excretion in the tested rats. ^{1,2,3}Data were tested by Student's *t*-test, Welch's *t*-test, or Mann-Whitney's *U* test, respectively. *Significant differences, compared with the Control (Group I). ^fLogarithmically transformed data. ⁱⁱCorrelated with the urinary Cre_u excretion. †,††,† $P = 0.06$, $P = 0.07$, and $P = 0.08$, respectively, compared with the Control (Group I).

TABLE 2: Basic indices in the tested animal groups at week 12.

Parameters	Groups of animals		Percentage decrease (↓) compared with Group I, <i>P</i> value
	(I) Control	(II) SMV	
Fluid intake (mL/rat/24 h) ²	58.53 ± 1.92	39.64 ± 0.73*	↓ 32 < 0.001
Food intake (g/rat/24 h) ¹	37.02 ± 0.86	28.73 ± 0.69*	↓ 22 < 0.001
Body weight (% of initial b. wt.) ¹	264.13 ± 23.59	190.46 ± 15.95*	↓ 28 < 0.05
V dose (mg V/kg b-wt./24 h) [#]	—	13.27 ± 0.33	

^{1,2}Data were tested by Student's *t*-test or Welch's *t*-test, respectively.

*Significant differences ($P < 0.05$), compared with the Control (Group I).

[#]Consumed with drinking water.

and ALB_u (per 24 h) decreased and unchanged, respectively, (Figure 2(a)). On the contrary, the urinary levels of TP_u (per $Cre_u/24$ h) in the same group of rats were unaltered and ALB_u (per $Cre_u/24$ h) increased, in comparison with the control (Figure 3(a)). In turn, the renal concentration of TP_k and ALB_k did not change markedly (Figure 4(a)).

The concentrations of CysC_p and β_2M_p remained unaltered between the groups (Figure 1(a)). However, the urinary excretion of CysC_u (per 24 h, Figure 2(a), and per $Cre_u/24$ h,

Figure 3(a)) increased markedly but the urinary β_2M_u level (per 24 h, Figure 2(a), and per $Cre_u/24$ h, Figure 3(a)) was unchanged. In turn, the urinary level of KIM-1_u (per 24 h, Figure 2(a), and per $Cre_u/24$ h, Figure 3(a)) in the SMV-intoxicated rats did not alter and increased, respectively, compared with the control.

The activity of LDH_p increased, ALP_p decreased, and GGTP_p was unchanged in the rats after the SMV intoxication, in comparison with the control (Figure 1(a)). In turn, the

FIGURE 4: Renal relative weight (RRW), the levels/activities of some OS markers, proteins, and enzymes (a) in the kidney, and the concentrations of selected elements ((b)–(e)) in the same organ of the tested rats.^{1,2,3} Data were tested by Student's *t*-test, Welch's *t*-test, or Mann-Whitney's *U* test, respectively. *Significant differences, compared with the Control (Group I). #, \$ *P* = 0.09, *P* = 0.05, respectively, compared with the Control (Group I).

urinary activities of LDH_u, GGTP_u, and ALP_u (per 24 h) as well as the urinary activities of GGTP_u and ALP_u (per Cre_u/24 h) and NAG_u (per 24 h and per Cre_u/24 h) were lowered in the SMV-exposed rats, compared with the control (Figures 2(b) and 3(b)). However, the urinary activity of LDH_u (per Cre_u/24 h) only showed a visible trend toward an increase (Figure 3(b)). In turn, the renal activity of LDH_k, GGTP_k, and ALP_k in the SMV-intoxicated rats was lowered, in comparison with the control (Figure 4(a)).

Urine dipstick analysis did not reveal presence of blood in the urine of the SMV-intoxicated rats (data not shown). However, the urine pH in the same animals was markedly higher, compared with that in the control rats (Figure 2(a)).

3.3. Element Levels in the Plasma. The exposure to SMV led to a distinct elevation of the Ca_p and K_p concentrations (Figure 1(b)), produced a visible decrease in the plasma Cu_p

level (Figure 1(d)), and increased the plasma V_p concentration (Figure 1(e)). However, no SMV intoxication-related effects on the concentrations of Mg_p, Na_p, Cl_p, and Zn_p (Figures 1(b), 1(c), and 1(d)) in the plasma were detected.

3.4. Excretion of Urinary Elements. In the SMV-intoxicated animals, Mg_u, Ca_u, and V_u were excreted in urine in higher amounts than those found in the control rats (Figures 2(c), 2(f), 3(c), and 3(f)). The urinary Cl_u excretion (per Cre_u/24 h) also tended to be higher, compared with the control (Figure 3(e)). In turn, the urinary excretion of Zn_u (per 24 h and per Cre_u/24 h) distinctly decreased in the SMV-exposed rats, in comparison with the control animals (Figures 2(d) and 3(d)), whereas the urinary K_u excretion (per 24 h, Figure 2(e), and per Cre_u/24 h, Figure 3(e)) dropped and did not change, respectively. However, the urinary levels of Cu_u and Na_u (per 24 h and per Cre_u/24 h), in the SMV-exposed rats, did not turn out to be significantly

FIGURE 5: The concentration of Mg (a), Ca (b), and V (c) in the rat femoral diaphysis (FD). ^{1,2}Data were tested by Student's *t*-test or Welch's *t*-test, respectively. *Significant differences, compared with the Control (Group I).

altered, compared with the control animals (Figures 2(d), 2(e), 3(d), and 3(e)).

3.5. Concentrations of Elements in the Kidney and FD. The exposure to SMV elevated the concentrations of Mg_k, K_k (Figures 4(b) and 4(d)), and V_k (Figure 4(e)) and decreased the concentration of Cu_k (Figure 4(c)) in the kidney, compared with the control. A clear tendency toward decreased and increased concentrations of Ca_k and Zn_k, respectively (Figures 4(b) and 4(c)), was also observed in the SMV-intoxicated animals, in comparison with the control rats. However, the concentration of Na_k in the same organ of the SMV-exposed rats did not change markedly, compared with the control (Figure 4(d)). In turn, the concentrations of Mg_{FD} and Ca_{FD} in FD decreased (Figures 5(a) and 5(b)), whereas the concentration of V_{FD} in the same FD increased (Figure 5(c)) after the SMV intoxication, in comparison with the control.

3.6. Kidney Weight and Renal MDA and TAS Levels. The SMV-exposed rats had a significantly elevated relative renal weight (RRW) and the renal MDA_k level, compared with the control animals (Figure 4(a)). The renal TAS_k level in the same group of rats was also distinctly enhanced, in comparison with the control (Figure 4(a)).

3.7. Correlations between Some Measured Variables. Significant positive and negative correlations and trends toward them were revealed between many different parameters, among others, between the renal V_k concentration, the renal MDA_k, or TAS_k levels and some biomarkers and basic indices as well as between both OS markers (Table 3). They were also found between the fluid intake, BW, EUV, or the urinary Cre_u excretion and some investigated parameters (Table 3) as well as between the renal V_k concentration or the renal MDA_k level and the levels/concentrations of elements in the plasma,

urine, kidney and FD (Table 4). The relationships between the urinary excretion of some elements and Cre_u or EUV were also indicated (Table 4).

4. Discussion

To our knowledge, the present report is the first to describe in a rat model the influence of the 12-week SMV exposure on the levels/activities of a wider panel of biomarkers of renal toxicity and on the concentrations of some elements in the kidney. It reveals a probable mechanism of the development of the SMV-induced functional renal disorders and proposes potential biomarkers of nephrotoxicity for specific conditions of SMV intoxication. It also illustrates for the first time many relationships between some explored indices. Because of different experimental conditions used, comparison of the results presented in the current report with literature data was difficult; therefore, the discussion section relies predominantly on our own results.

The increase in the kidney/body weight ratio observed by us in the present (Figure 4(a)) and previous study [28] in the SMV-intoxicated rats may indicate that V accumulated sufficiently in the kidney to manifest significant alterations in the size of this organ. Taking into account the fact that one of the main excretory routes for V within the body is the urinary system [29], V accumulation in the kidney may result in not only functional but also structural damage. Therefore, the estimation of the structural renal injury by histological examination in the rats after the 12-week SMV exposure is going to be performed in the nearest future. In this report, the degree of kidney damage has been evaluated only biochemically.

The decreased CreC together with the enhanced plasma Cre_p and U_p levels (Figure 1(a)), and the lowered urinary Cre_u and U_u excretion (Figure 2(a)) found in the SMV-exposed rats may indicate a problem with removal of Cre and U

TABLE 3: Correlation coefficients for compared variables.

Variables							
V_k		MDA_k		TAS_k			
Food-I	-0.894 [†]	Food-I	-0.817 [†]	Food-I	-0.676 [†]		
Fluid-I	-0.905 [†]	Fluid-I	-0.754 [†]	Fluid-I	-0.707 [†]		
RRW	0.778 [†]	RRW	0.531*	RRW	0.406 ^f		
BW	-0.601*	BW	-0.672 [†]	BW	-0.780 [†]		
MDA_k	0.889 [†]	TAS_k	0.598*	LDH_k	-0.472 ^a		
TAS_k	0.590*	LDH_k	-0.614*	ALP_k	-0.492 ^{††}		
LDH_k	-0.544*	ALP_k	-0.757 [†]	$GGTP_k$	-0.401 ^g		
ALP_k	-0.669 [†]	$GGTP_k$	-0.577*	EUV	-0.412 ^f		
$GGTP_k$	-0.759 [†]	U_k	-0.445 ^c	pH	0.588*		
EUV	-0.716 [†]	ALB_k	-0.549*	Cre_p	0.624 [†]		
pH	0.917 [†]	EUV	-0.613*	$CreC$	-0.505*		
Cre_u	-0.630 [†]	pH	0.761 [†]	U'_u	-0.429 ^d		
U_p	0.483**	Cre_p	0.757 [†]	$CysC'_u$	0.601*		
Cre_p	0.818 [†]	$CreC$	-0.606*	$\beta_2M'_u$	-0.408 ^h		
$CreC$	-0.652 [†]	TP_p	-0.437 ^d	ALP'_u	-0.552*		
ALP_p	-0.448 ^c	LDH_p	0.539*	NAG'_u	-0.448 ^c		
LDH_p	0.757 [†]	Cre_u	-0.572*				
U'_u	-0.596*	U'_u	-0.574*				
TP'_u	-0.654 [†]	TP'_u	-0.518*				
$CysC'_u$	0.399 ^g	$CysC'_u$	0.496 ^f				
LDH'_u	-0.612*	LDH'_u	-0.523*				
ALP'_u	-0.656 [†]	ALP'_u	-0.514*				
$GGTP'_u$	-0.531 ^a	$GGTP'_u$	-0.425 ⁱ				
NAG'_u	-0.506*	NAG'_u	-0.490 ^{ff}				
Fluid-I		BW		EUV		Cre_u	
EUV	0.759 [†]	Cre_p	-0.773 [†]	Cre_u	0.920 [†]	U''_u	-0.561*
Cre_p	-0.844 [†]	Cre_u	0.595*	U'_u	0.816 [†]	UA''_u	-0.443 ^c
Cre_u	0.668 [†]	$CreC$	0.762 [†]	UA'_u	0.499*	TP''_u	-0.786 [†]
$CreC$	0.755 [†]	U'_u	0.734 [†]	LDH'_u	0.569*	ALB''_u	-0.630 [†]
U_p	-0.470 ^a	$CysC_p$	0.488 ^{‡‡}	ALP'_u	0.424 ^e	$\beta_2M''_u$	-0.697 [†]
U'_u	0.700 [†]	$CysC'_u$	-0.482 ^a	$GGTP'_u$	0.643*	LDH''_u	-0.776 [†]
				NAG'_u	0.780 [†]		
				$KIM-1'_u$	0.447 ^c		

Data are presented as the correlation coefficients (r) and the levels of statistical significance (P).

^{P,u,k}Plasma, urine, and kidney (concentration), respectively.

Fluid-I, Food-I, RRW, BW, and EUV: fluid intake, food intake, renal relative weight, body weight, and excreted urinary volume, respectively.

[†] $P < 0.01$; ^{*} $P < 0.05$; [†] $P = 0.051$; ^{†††} $P = 0.053$; ^{††} $P = 0.054$; ^{‡‡} $P = 0.055$; ^{**} $P = 0.058$; ^{††} $P = 0.06$; [†] $P = 0.08$; [†] $P = 0.09$; [†] $P = 0.10$; [†] $P = 0.11$; [†] $P = 0.12$; [†] $P = 0.13$; [†] $P = 0.14$.

^{1,11}Expressed per 24 h and per Cre₁₁ per 24 h, respectively.

The significant correlations and tendencies toward them are highlighted in normal and italic bold font, respectively.

from the body and implicate glomerular functional disorders reflected by decreased clearance of the above-mentioned compounds. However, the lowered basic parameters such as fluid intake (discussed by us more widely) [30] and BW found in the same group of animals (Table 2) and the significant correlations between them and Cr_u , Cr_p , CrC , U_u , and/or U_p (Table 3) do not allow us to exclude the possibility that dehydration and/or low BW affected the measured standard

renal biomarkers. However, it is difficult to recognize to what extent they could influence the alterations in the biomarkers examined. The levels of Cre and U in the blood/urine are known to be affected not only by renal but also by some nonrenal factors [31–33]. On the other hand, the correlations between the renal V concentration (V_k) and Cre_u , Cre_p , $CreC$, U_u , and U_p (Table 3) confirm the involvement of V in the changes in the basic biomarkers explored. Moreover,

the correlations between V_k and both OS markers such as MDA_k and TAS_k as well as between MDA_k and Cre_u , Cre_p , $CreC$, and U_u (Table 3) also point to the involvement of the SMV-induced OS in their alterations observed. Additionally, the positive correlation between both OS markers allows us to definitively conclude that the changes in the renal TAS_k level (Table 3) are associated with the alterations in LPO in the kidney due to the SMV exposure.

The markedly elevated urinary excretion of $CysC_u$ (a tubular injury biomarker) [20, 32] demonstrated in the SMV-intoxicated rats (Figures 2(a) and 3(a)) without significant alterations in the plasma level of this protein ($CysC_p$) (Figure 1(a)) may point to decreased reabsorption of $CysC$ by injured tubules. Moreover, the positive correlation of $CysC_u$ with TAS_k as well as the clear trends toward the positive correlations of $CysC_u$ with V_k and MDA_k (Table 3) can suggest involvement of the renal V prooxidative action in the proximal tubular dysfunction (PTD), which consequently led to the urinary $CysC$ wasting. It has been reported that when the renal proximal tubule remains intact $CysC$ is completely taken up by the proximal tubular cells (PTC) [32], but in the case of tubular disease and the proximal convoluted tubule (PCT) injury the concentration of this protein in the urine is significantly elevated [34, 35]. In addition, the markedly enhanced urinary KIM-1_u excretion (Figure 3(a)) found in the same group of rats may also point to damage to PTC due to the SMV exposure. KIM-1 is expressed at very high levels in proximal tubule epithelial cells after toxic injury and it is present in urine when the injury of PTC occurs [20].

In turn, the lowered urinary activities of LDH_u , ALP_u , $GGTP_u$, and NAG_u , demonstrated in the SMV-exposed rats (Figure 2(b)), allow us to suggest that the use of these tests for monitoring kidney injury during the SMV intoxication may not be valid, as the activity of these enzymes in the urine usually rises due to release from damaged cells. The lowered urinary ALP_u and GGT_u excretion and the simultaneous increase in the urinary LDH_u and NAG_u excretion were demonstrated by De la Torre et al. [36] in adult rats after vanadate intoxication. The discrepancies between our results concerning the urinary excretion of LDH and NAG and the results obtained by the above-mentioned investigators draw attention to the dependence of the renal effects on the conditions of vanadate exposure. In turn, the significant negative correlations found between the renal V_k concentration and the renal LDH_k , ALP_k , and $GGTP_k$ activities (Table 3) may indicate a direct V impact. The direct inhibitory effect of both vanadate and vanadyl on the ALP activity was described by Cortizo et al. [37]. However, the marked increase in the plasma LDH activity (LDH_p) demonstrated in the SMV-exposed rats (Figure 1(a)) requires further analyses. The measurement of the LDH isoenzyme might be the next important point in a more precise explanation of the significant increase in the activity of this enzyme in the plasma because the total plasma LDH is a highly sensitive test but, simultaneously, it is nonspecific. At the present stage of our study, we may only confirm the involvement of V and the SMV-induced OS in the changes in the activity of LDH in the blood, as evidenced by the significant positive correlations of LDH_p with V_k and MDA_k (Table 3).

TABLE 4: Correlation coefficients for measured variables.

Variables		
	V_k	MDA_k
V_p	0.872[†]	0.710[†]
Cu_p	-0.520[*]	-0.445^b
K_p	0.621[*]	—
V_u	0.977[†]	0.795[†]
Mg_u	0.798[†]	0.678[†]
Zn_u	-0.555[*]	-0.618[*]
K_u	-0.600[*]	-0.535[*]
Cu_k	-0.687[†]	-0.628[†]
Mg_k	0.527[*]	—
K_k	0.599[*]	—
V_k	—	0.889[†]
Ca_k	-0.486[#]	-0.479^a
Mg_{FD}	-0.767[†]	-0.725[†]
Ca_{FD}	-0.492[°]	-0.334^c
V_{FD}		
Mg_{FD}	-0.768[*]	
Ca_{FD}	-0.547[*]	
EUV		
V_u	-0.722[†]	
K_u	0.856[†]	
Cre_u		
V_u	-0.600[*]	
Mg_u	-0.602[*]	
Ca_u	-0.633[†]	
Cu_u	-0.688[†]	
K_u	-0.540[*]	
Cl_u	-0.735[†]	

Data are presented as the correlation coefficients (r) and the levels of statistical significance (P).

p, u, FD, k Plasma, urine, femoral diaphysis, and kidney (concentration), respectively.

EUV: excreted urinary volume.

[†] $P < 0.01$; ^{*} $P < 0.05$; [°] $P = 0.053$; [#] $P = 0.056$; ^a $P = 0.061$; ^b $P = 0.08$; ^c $P = 0.20$.

^{†, °} Expressed per 24 h and per Cre_u per 24 h.

The significant correlations and tendencies toward them are highlighted in normal and italic bold font, respectively.

Additionally, the strong positive correlations of the urinary pH with V_k and with both OS markers (Table 3) and the correlation between the renal V_k and K_k concentration and between the renal V_k concentration and the plasma K_p level (Table 4) also suggest involvement of OS in the mechanism of the rise in urine pH under the conditions of the SMV exposure (Figure 2(a)). They also confirm the dependence between the accumulation of V in the kidney and disorders in K homeostasis (Figures 1(b) and 4(d)). All these results suggest further measurements to elucidate the mechanism(s) of the changes observed in our experimental conditions.

The urinary Mg wasting (Figures 2(c) and 3(c)) and its elevated and diminished concentration in the kidney and FD, respectively (Figures 4(b) and 5(a)), as well as the clearly enhanced plasma and urinary Ca level (Figures 1(b) and

2(c)) and its lowered FD concentration (Figure 5(b)) point to disorders in the homeostasis of both macroelements during the SMV exposure. These changes may be partly explained by the influence of SMV on the bone tissue which is the main site of V uptake and deposition and in which the level of this element has been found to be significantly higher (Figure 5(c)). The presence of V in the bone might lead to mobilization of Mg and Ca from this tissue and cause the alterations observed. The negative correlations between the FD V, Mg, and Ca concentrations (Table 4) confirm the connection between the bone V accumulation and the bone Mg and Ca imbalance. Moreover, the negative correlation of V_k and MDA_k with the Mg_{FD} concentration and the clear trend toward this correlation between V_k and MDA_k with Ca_{FD} concentration (Table 4) also allow us to conclude that disturbances in bone Mg and Ca homeostasis observed in the rats at the SMV-exposure are associated both with the rise in the renal V accumulation and with the SMV-induced OS in the kidney. The changes in the levels of the other elements in biological fluids and in the kidney were also modified in correlation with the renal V_k concentration and/or with the SMV-intensified LPO in the kidney (Table 4).

Studies on the mechanisms of V action have shown that, as a transition metal, it may generate reactive oxygen species (ROS) and/or free radicals (FRs) and stimulate LPO or indirectly modify OS in cells, by releasing FR-generating metals from tissues, modifying enzymatic and antioxidant defence, or interacting with mitochondria [1, 38–40]. The literature data [1, 41] have revealed that OS may be involved in the mechanisms of the toxic action of vanadate. The correlations or clear trends toward them between the basic indices explored, some biomarkers of renal toxicity, or elements and the renal MDA_k and/or TAS_k levels (Table 3) suggest that in our experimental conditions the mechanism of the deleterious action of vanadate (as SMV) was associated with generation of OS in the kidney, as evidenced by the SMV-induced increase in both OS markers in this organ.

Finally, the lowered urinary excretion of Cre_u , U_u , UA_u , $KIM-1_u$, LDH_u , $GGTP_u$, ALP_u , NAG_u , and K_u , when normalized per 24 h (Figure 2), and the unaltered or enhanced levels/activities of U_u , UA_u , TP_u , ALB_u , β_2M_u , LDH_u , NAG_u , and some elements in the urine, when normalized per $Cre_u/24$ h (Figure 3), might be in part a consequence of the reduced EUV or the lowered urinary Cre_u excretion (Figure 2(a)), respectively, as indicated by the correlations of EUV or Cre_u with the above-mentioned indices (Tables 3 and 4).

5. Conclusions

The results of this preliminary study demonstrated that the exposure to SMV led to alterations in the levels/activities of some examined biomarkers of nephrotoxicity and caused renal mineral imbalance. They also pointed to a contribution of SMV-induced OS in the mechanism underlying the changes. Moreover, our findings revealed that the use of standard biomarkers of renal toxicity, such as Cre and U, and calculation of CreC for evaluation of kidney function

at the SMV-induced fall in the fluid intake and BW may have a weak diagnostic value, since CreC and the plasma and urinary levels of Cre and U are greatly influenced by both the indices mentioned. In addition, the results simultaneously showed that the use of other classical enzymatic tests such as LDH_u , ALP_u , $GGTP_u$, and NAG_u might not be valid either, due to their reduced urinary activities. Our results suggest that, from among the biomarkers tested, $CysC_u$ and $KIM-1_u$ might be the most appropriate in monitoring kidney function at the SMV exposure. The elevated urinary levels of both compounds may point to the proximal tubular dysfunction. Finally, for the first time our results also revealed many significant relationships between the parameters explored.

Endnotes

1. They were bought as part of the Project entitled “Building of the Center for Interdisciplinary Research” realized within the frame of the Operating Programme “Development of Eastern Poland” 2007–2013, Priority I: Modern Economy, Action I.3. The Advancement of Innovation, cofinanced by the European Regional Development Fund.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors would like to thank very much the former Vice-Rector of John Paul II Catholic University of Lublin Rev. Professor Stanisław Zięba and Dr. Emilia Fornal from Laboratory of Separation and Spectroscopic Method Applications, Center for Interdisciplinary Research, the John Paul II Catholic University of Lublin for providing partial financial support of the studies. They would also like to thank very much Dr. Jakub Nowak from the Laboratory of Composite and Biomimetic Materials, Center for Interdisciplinary Research, the John Paul II Catholic University of Lublin for providing them with a diamond-disc saw for cutting the bones and Anna Gromba M.S. and Wojciech Gierad M.S. for their help in cutting the rat bones.

References

- [1] J. Z. Byczkowski and A. P. Kulkarni, “Oxidative stress and pro-oxidant biological effects of vanadium,” in *Vanadium in the Environment Part II: Health Effects*, J. O. Nriagu, Ed., pp. 235–264, John Wiley and Sons, New York, NY, USA, 1998.
- [2] A. Ścibior, H. Zaporowska, and I. Niedźwiecka, “Lipid peroxidation in the liver of rats treated with V and/or Mg in drinking water,” *Journal of Applied Toxicology*, vol. 29, no. 7, pp. 619–628, 2009.
- [3] A. Ścibior, H. Zaporowska, and I. Niedźwiecka, “Lipid peroxidation in the kidney of rats treated with V and/or Mg in drinking water,” *Journal of Applied Toxicology*, vol. 30, no. 5, pp. 487–496, 2010.

- [4] A. Ścibior, D. Gołębiowska, and I. Niedźwiecka, "Magnesium can protect against vanadium-induced lipid peroxidation in the hepatic tissue," *Oxidative Medicine and Cellular Longevity*, vol. 2013, Article ID 802734, 11 pages, 2013.
- [5] J. Donaldson and F. LaBella, "Prooxidant properties of vanadate *in vitro* on catecholamines and on lipid peroxidation by mouse and rat tissues," *Journal of Toxicology and Environmental Health*, vol. 12, no. 1, pp. 119–126, 1983.
- [6] M. H. Oster, J. M. Llobet, J. L. Domingo, J. B. German, and C. L. Keen, "Vanadium treatment of diabetic Sprague-Dawley rats results in tissue vanadium accumulation and pro-oxidant effects," *Toxicology*, vol. 83, no. 1–3, pp. 115–130, 1993.
- [7] S. S. Soares, H. Martins, R. O. Duarte et al., "Vanadium distribution, lipid peroxidation and oxidative stress markers upon decavanadate *in vivo* administration," *Journal of Inorganic Biochemistry*, vol. 101, no. 1, pp. 80–88, 2007.
- [8] J. Donaldson, R. Hemming, and F. LaBella, "Vanadium exposure enhances lipid peroxidation in the kidney of rats and mice," *Canadian Journal of Physiology and Pharmacology*, vol. 63, no. 3, pp. 196–199, 1985.
- [9] M. Carmignani, A. R. Volpe, E. Sabbioni, M. Felaco, and P. Boscolo, "Vanadium and the cardiovascular system: regulatory effects and toxicity," in *Vanadium in the Environment Part II: Health Effects*, J. O. Nriagu, Ed., vol. 31, pp. 181–219, John Wiley and Sons, New York, NY, USA, 1998.
- [10] K. De Cremer, M. Van Hulle, C. Chéry et al., "Fractionation of vanadium complexes in serum, packed cells and tissues of Wistar rats by means of gel filtration and anion-exchange chromatography," *Journal of Biological Inorganic Chemistry*, vol. 7, no. 7–8, pp. 884–890, 2002.
- [11] V. Sitprija, K. Tungsanga, P. Tosukhowong et al., "Metabolic problems in northeastern Thailand: possible role of vanadium," *Mineral and Electrolyte Metabolism*, vol. 19, no. 1, pp. 51–56, 1993.
- [12] E. Dafnis and S. Sabatini, "Biochemistry and pathophysiology of vanadium," *Nephron*, vol. 67, no. 2, pp. 133–143, 1994.
- [13] P. Tosukhowong, K. Tungsanga, S. Eiam-Ong, and V. Sitprija, "Environmental distal renal tubular acidosis in Thailand: an enigma," *American Journal of Kidney Diseases*, vol. 33, no. 6, pp. 1180–1186, 1999.
- [14] A. S. Tracey, G. R. Willsky, and E. S. Takeuchi, *Vanadium—Chemistry, Biochemistry, Pharmacology and Practical Applications*, Taylor and Francis, Boca Raton, Fla, USA, 2007.
- [15] S. M. Talbott and K. Hughes, *The Health Professional's Guide to Dietary Supplements*, Lippincott Williams and Wilkins, 2007.
- [16] V. Sitprija and S. Eiam-Ong, "Vanadium and metabolic problems," in *Vanadium in the Environment Part II: Health Effects*, J. O. Nriagu, Ed., pp. 91–120, John Wiley and Sons, New York, NY, USA, 1998.
- [17] Y. Teng, S. Ni, C. Zhang, J. Wang, X. Lin, and Y. Huang, "Environmental geochemistry and ecological risk of vanadium pollution in Panzhihua mining and smelting area, Sichuan, China," *Chinese Journal of Geochemistry*, vol. 25, no. 4, pp. 379–385, 2006.
- [18] Expert group on vitamins and minerals, Review of vanadium. EVM/00/04.REVISED AUG, 2002.
- [19] K. H. Thompson, M. Battell, and J. H. McNeill, "Toxicology of vanadium in mammals," in *Vanadium in the Environment Part II: Health Effects*, J. O. Nriagu, Ed., pp. 21–38, John Wiley and Sons, New York, NY, USA, 1998.
- [20] R. G. Fassett, S. K. Venuthurupalli, G. C. Gobe, J. S. Coombes, M. A. Cooper, and W. E. Hoy, "Biomarkers in chronic kidney disease: a review," *Kidney International*, vol. 80, no. 8, pp. 806–821, 2011.
- [21] G. D'Amico and C. Bazzi, "Urinary protein and enzyme excretion as markers of tubular damage," *Current Opinion in Nephrology and Hypertension*, vol. 12, no. 6, pp. 639–643, 2003.
- [22] V. S. Vaidya, V. Ramirez, T. Ichimura, N. A. Bobadilla, and J. V. Bonventre, "Urinary kidney injury molecule-1: a sensitive quantitative biomarker for early detection of kidney tubular injury," *American Journal of Physiology—Renal Physiology*, vol. 290, no. 2, pp. F517–F529, 2006.
- [23] A. Ścibior, A. Adamczyk, D. Gołębiowska, and I. Niedźwiecka, "Effect of 12-week vanadate and magnesium co-administration on chosen haematological parameters as well as on some indices of iron and copper metabolism and biomarkers of oxidative stress in rats," *Environmental Toxicology and Pharmacology*, vol. 34, no. 2, pp. 235–252, 2012.
- [24] J. L. Domingo, M. Gomez, D. J. Sanchez, J. M. Llobet, and C. L. Keen, "Toxicology of vanadium compounds in diabetic rats: the action of chelating agents on vanadium accumulation," *Molecular and Cellular Biochemistry*, vol. 153, no. 1–2, pp. 233–240, 1995.
- [25] J. Azay, J. Brès, M. Krosniak et al., "Vanadium pharmacokinetics and oral bioavailability upon single-dose administration of vanadyl sulfate to rats," *Fundamental and Clinical Pharmacology*, vol. 15, no. 5, pp. 313–324, 2001.
- [26] World Health Organization, *Vanadium Pentoxide and Other Inorganic Vanadium Compounds*, WHO, Geneva, Switzerland, 2001.
- [27] International Agency for Research on Cancer Monographs on the evaluation of carcinogenic risks to humans, *Cobalt in Hard Metals and Cobalt Sulfate, Gallium Arsenide, Indium Phosphide and Vanadium Pentoxide*, vol. 86, Lyon, France, 2006.
- [28] A. Ścibior and H. Zaporowska, "Effects of vanadium(V) and/or chromium(III) on L-ascorbic acid and glutathione as well as iron, zinc, and copper levels in rat liver and kidney," *Journal of Toxicology and Environmental Health A*, vol. 70, no. 8, pp. 696–704, 2007.
- [29] D. A. Barrio and S. B. Etcheverry, "Vanadium and bone development: putative signaling pathways," *Canadian Journal of Physiology and Pharmacology*, vol. 84, no. 7, pp. 677–686, 2006.
- [30] A. Ścibior, H. Zaporowska, A. Wolińska, and J. Ostrowski, "Antioxidant enzyme activity and lipid peroxidation in the blood of rats co-treated with vanadium (V^{+5}) and chromium (Cr^{+3})," *Cell Biology and Toxicology*, vol. 26, no. 6, pp. 509–526, 2010.
- [31] A. C. Baxmann, M. S. Ahmed, N. C. Marques et al., "Influence of muscle mass and physical activity on serum and urinary creatinine and serum cystatin C," *Clinical Journal of the American Society of Nephrology*, vol. 3, no. 2, pp. 348–354, 2008.
- [32] P. D. Sarkar, G. Rajeshwari, and T. M. Shivaprakash, "Cystatin C—A novel marker of glomerular filtration: a review," *Indian Journal of Clinical Biochemistry*, vol. 20, no. 1, pp. 139–144, 2005.
- [33] G. H. Tesch, "Review: serum and urine biomarkers of kidney disease: a pathophysiological perspective," *Nephrology*, vol. 15, no. 6, pp. 609–616, 2010.
- [34] M. Warwas and A. Piwowar, "Moczowa cystatyna C jako biomarker uszkodzenia kanalików nerkowych," *Postępy Higieny i Medycyny Doświadczalnej*, vol. 65, pp. 562–568, 2011.
- [35] J. Westhuyzen, "Cystatin C: a promising marker and predictor of impaired renal function," *Annals of Clinical and Laboratory Science*, vol. 36, no. 4, pp. 387–394, 2006.

- [36] A. De La Torre, S. Granero, E. Mayayo, J. Corbella, and J. L. Domingo, "Effect of age on vanadium nephrotoxicity in rats," *Toxicology Letters*, vol. 105, no. 1, pp. 75–82, 1999.
- [37] A. M. Cortizo, V. C. Salice, and S. B. Etcheverry, "Vanadium compounds: their action on alkaline phosphatase activity," *Biological Trace Element Research*, vol. 41, no. 3, pp. 331–339, 1994.
- [38] A. M. Evangelou, "Vanadium in cancer treatment," *Critical Reviews in Oncology/Hematology*, vol. 42, no. 3, pp. 249–265, 2002.
- [39] A. Ścibior and H. Zaporowska, "Effects of combined vanadate and magnesium treatment on erythrocyte antioxidant defence system in rats," *Environmental Toxicology and Pharmacology*, vol. 30, no. 2, pp. 153–161, 2010.
- [40] X.-G. Yang, X.-D. Yang, L. Yuan, K. Wang, and D. C. Crans, "The permeability and cytotoxicity of insulin-mimetic vanadium compounds," *Pharmaceutical Research*, vol. 21, no. 6, pp. 1026–1033, 2004.
- [41] J. Liu, H. Cui, X. Liu et al., "Dietary high vanadium causes oxidative damage-induced renal and hepatic toxicity in broilers," *Biological Trace Element Research*, vol. 145, no. 2, pp. 189–200, 2012.