Review Article

Genetic Networks Lead and Follow Tumor Development: MicroRNA Regulation of Cell Cycle and Apoptosis in the p53 Pathways

Kurataka Otsuka^{1,2} and Takahiro Ochiya¹

¹ Division of Molecular and Cellular Medicine, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

² Division of Research and Development, Kewpie Corporation, Sengawa-cho, Chofu-shi, Tokyo 182-0002, Japan

Correspondence should be addressed to Kurataka Otsuka; kurataka_otsuka@kewpie.co.jp and Takahiro Ochiya; tochiya@ncc.go.jp

Received 25 July 2014; Accepted 26 August 2014; Published 11 September 2014

Academic Editor: Chengfeng Yang

Copyright © 2014 K. Otsuka and T. Ochiya. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

During the past ten years, microRNAs (miRNAs) have been shown to play a more significant role in the formation and progression of cancer diseases than previously thought. With an increase in reports about the dysregulation of miRNAs in diverse tumor types, it becomes more obvious that classic tumor-suppressive molecules enter deep into the world of miRNAs. Recently, it has been demonstrated that a typical tumor suppressor p53, known as the guardian of the genome, regulates some kinds of miRNAs to contribute to tumor suppression by the induction of cell-cycle arrest and apoptosis. Meanwhile, miRNAs directly/indirectly control the expression level and activity of p53 to fine-tune its functions or to render p53 inactive, indicating that the interplay between p53 and miRNA is overly complicated. The findings, along with current studies, will underline the continuing importance of understanding this interlocking control system for future therapeutic strategies in cancer treatment and prevention.

1. Introduction

Cancer is commonly an age-related disease triggered by the accumulation of genomic mutations that lead to the dysregulation of tumor-suppressive genes and/or protooncogenes. For example, the functions of *TP53* (tumor-suppressive gene) and *c-MYC* (oncogene) have been extensively investigated, and their critical roles in complexly regulating tumorigenesis, including cell-cycle progression/arrest, apoptosis, senescence, and energy metabolism, have been uncovered [1–4]. Specifically, the significance of tumor suppressor p53 has been suggested by the fact that DNA mutation or loss of *TP53* is observed in many types (over 50%) of human tumors and by the possibility that the dysfunctions affect the p53 signaling network in over 80% of tumors [5, 6]. As a transcriptional activator, the p53 protein induces various kinds of tumor-suppressive genes, such as *p21* (G_1/S -arrest), *14-3-3 σ* (G_2/M -arrest), and *PUMA* (apoptosis) [7–10]. p53 has also been reported to negatively regulate specific proteins: for instance, the p53-mediated repression of the cell-cycle regulators, such

as cyclin-dependent kinase 4 (CDK4) and cyclin E2, may lead to cell-cycle arrest [10, 11]. These prove the pivotal roles of p53 as a cellular gatekeeper.

Recently, it has been realized that small noncoding RNAs known as microRNAs (miRNAs) contribute to many human diseases, including cancers; that a general downregulation of miRNAs is observed in cancers as compared with normal tissues; and that miRNA expression profiles can be used to classify poorly differentiated tumors [12]. In addition, some kinds of miRNAs are shown to be connected to a well-studied tumor-suppressive or oncogenic network [13]. It remains to be investigated how miRNAs are regulated by transcription factors, but it is suggested that p53 enters the miRNA world to control the expression patterns of some miRNAs and promote cell-cycle arrest and apoptosis through the miRNA effector pathway. *miR-34a* is one of the representative miRNAs under the direct control of p53, and this upregulation induces cell-cycle arrest and apoptosis [14–18]. Moreover, there are many studies about miRNA effects on cell proliferation and survival in cancers, with attention

given to the interplay between p53 and the miRNA network. In this review, we will focus on the regulation of the cancer cell cycle and apoptosis by miRNA linked with the p53 axis. We will also summarize the key miRNAs concerned with the cell cycle and apoptosis in cancers.

2. miRNA Discovery, Biogenesis, and Mechanism

The first miRNAs discovered were *lin-4* and *let-7*, both of which are the key regulators in the pathway controlling the timing of postembryonic development in *Caenorhabditis elegans* [19–21]. After this discovery, miRNAs have been identified in diverse organisms, such as worms, flies, mice, humans, and plants. Several miRNAs are conserved among different species, indicating that these miRNAs might have important functions and modulate gene expression. Currently, in humans, over 2,000 microRNAs have been identified or predicted based on the miRBase database (<http://www.mirbase.org/>). Computational analyses suggest that about 5,300 genes contain miRNA target sites: ~30% of human genes might be subject to the translational regulation of miRNAs [22, 23].

miRNAs are initially transcribed by RNA polymerase II/III into primary transcripts (pri-miRNAs) [24, 25], which are processed by the complex of RNase III enzyme, Drosha, and its partner DGCR8 [26]. The pri-miRNAs are converted into ~65 nucleotides (nt) of a stem-loop precursor (pre-miRNA) [27]. These pre-miRNAs are transported to cytoplasm by Exportin-5/Ran-GTP and processed by another RNase III, Dicer, to generate a double-strand RNA of about 19–25 nt in length [28–30]. One strand of miRNA gives rise to the mature miRNA, which is incorporated into the RNA-induced silencing complex (RISC). The miRNAs guide the RISC complex to the 3'-untranslated region (3'-UTR) of the target mRNAs, leading to the translational repression or destabilization of the mRNA [31, 32]. In animal systems, the recognition of target mRNA usually requires the "seed" sequence, which is 2–8 nt from the 5'-end of the miRNA [22, 33]. Unlike with plant systems, because of this imperfect complementarity, there are extensive base-pairings to the sequence of mRNAs, and this makes it more complicated to predict miRNA targets and study miRNA biology. Recently, it has been shown that animal miRNAs can induce the degradation of target mRNAs (mRNA degradation and decay) besides translational repression: inhibition of translation elongation; cotranslational protein degradation; competition for the cap structure; and inhibition of ribosomal subunit joining [34–37]. However, the exact order and impact of these events still need to be investigated further.

3. p53 Transactivation Function in a Relationship with Tumorigenesis

Based on numerous studies at both structural and functional levels, p53 is known as a key player in genome stability and tumor suppression. In an unstressed condition, the expression level of p53 is kept low by the activity of an E3

ubiquitin ligase, mouse double minute 2 (MDM2) [38–40]. Under stressed conditions, p53 is activated in response to diverse intrinsic and extrinsic signals, such as DNA damage, oncogene activation, and hypoxia. As a sequence-specific transcription factor, the activated p53 acts directly on cancer-associated pathways to suppress tumor progression by modulating cell-cycle arrest, senescence, apoptosis, angiogenesis, or invasion and metastasis [41–43]. There are also demonstrations showing that p53 is involved in the regulation of DNA repair, oxidative stress, energy metabolism, and differentiation [44–48]. The approach of genome-wide analyses has identified many p53-binding sites and p53-regulated genes which are related to tumorigenesis and various stress signals [49, 50]. Recent works have highlighted that p53 directly induces some specific miRNAs which function as tumor suppressors through a novel transcriptional mechanism. Now, although unknown aspects of the mechanism still need to be investigated, the cooperative contribution of p53 and miRNAs has been shown to be more important for tumor formation and development.

4. miRNA Network with p53: Cell Cycle and Apoptosis

4.1. miR-34 Family. In 2007, several groups reported that the *miR-34* family members are direct p53 targets and that their expression level is strongly upregulated by genotoxic stress in a p53-dependent manner, inducing cell-cycle arrest and apoptosis [14–16, 51, 52]. In mammals, the *miR-34* family is composed of *miR-34a*, *miR-34b*, and *miR-34c*, which are encoded by two different genes in the *miR-34a* and *miR-34b/c* loci. With the overexpression of the *miR-34* family in certain kinds of cell lines, microarray analyses unveiled hundreds of putative candidate target genes of *miR-34s* [15, 16, 18]. Actually, ectopic expression of *miR-34s* promotes cell-cycle arrest in the G₁ phase, senescence, and apoptosis by directly repressing CDK4, CDK6, cyclin E2, E2F3, MYC, and B-cell CLL/lymphoma 2 (BCL-2) [53]. Note that the triggering event of cell-cycle arrest or apoptosis by *miR-34s* depends on the cell type and context, and the expression level of *miR-34s* would affect the decision to proceed [15, 17, 54]. As seen in the decreased expression of *miR-34s* in several types of malignant cancers, the *miR-34* family powerfully prevents tumorigenesis in general.

In addition to the *miR-34* family, p53 is also engaged in the direct regulation of the transcriptional expression of additional miRNAs, such as *miR-107*, *miR-143/145*, *miR-192/194/215*, *miR-200c/141*, the *let-7* family, and the *miR-17-92* cluster (Figure 1 and Table 1).

4.2. miR-107. *miR-107* is encoded within an intron of pan-tothenate kinase 1 (*PANK1*), and *miR-107* and its host gene are directly activated by p53 under hypoxia condition or with the treatment of DNA damage agents [55, 56]. Hypoxia induces angiogenesis, which is essential for solid tumors to grow in severe environments. *miR-107* inhibits hypoxia signaling and antiangiogenesis by repressing the expression of hypoxia inducible factor-1 β (HIF-1 β), which interacts with HIF-1 α to form the HIF-1 transcription factor complex [55].

TABLE 1: Key microRNAs regulated by p53.

miRNA	Genomic location	Cancer type	Target	Phenotype	References
<i>miR-34s</i>	1p36 and 11q23	Colon cancer, neuroblastoma, pancreatic cancer, CLL, NSCLC, OSCC, breast cancer, bladder cancer, kidney cancer, melanoma	CDK4, CDK6, cyclin E2, E2F3, MYC BCL-2	Cell-cycle arrest apoptosis	[14–16, 18, 51, 53, 54]
<i>miR-107</i>	10q23	Colon cancer, breast cancer	CDK6, P130	Cell-cycle arrest	[55–57]
<i>miR-145</i>	5q23	Colon cancer, breast cancer, MDS, prostate cancer	MYC, E2F3, cyclin D2, CDK4, CDK6	Cell-cycle arrest	[58–61]
<i>miR-192/215</i>	1q41 and 11q13	Colon cancer, lung cancer, multiple myeloma, renal cancer	CDC7, MAD2L1	Cell-cycle arrest	[66–69]
<i>miR-200c</i>	12p13	Breast cancer, ovarian cancer	FAP-1	Apoptosis	[71, 81]
<i>let-7</i>	Multiple locations (11 copies)	Lung cancer, colon cancer, ovarian cancer, breast cancer, lymphoma	CDK6, CDC25A, cyclin D, CDC34, MYC, E2F1, E2F3	Cell-cycle arrest	[61, 83–90]
<i>miR-15a/16-1</i>	13q14	B-CLL, pituitary adenomas, gastric cancer, NSCLC, prostate cancer, ovarian cancer, pancreatic cancer	CDK1, CDK2, CDK6, cyclin D1, D3, E1 BCL-2	Cell-cycle arrest apoptosis	[98–112]

CLL: chronic lymphocytic leukemia; NSCLC: non-small cell lung cancer; OSCC: oral squamous cell carcinoma; MDS: myelodysplastic syndromes; B-CLL: B-cell chronic lymphocytic leukemia.

FIGURE 1: p53-induced miRNAs control cell cycle and cell survival. p53 directly induces many kinds of miRNAs, which repress cell-cycle regulators and/or antiapoptotic proteins and contribute to cell-cycle arrest and apoptosis. The miRNAs regulating apoptosis are shown in the top part of this figure, and the miRNAs regulating the cell cycle are at the bottom.

Furthermore, *miR-107* promotes cell-cycle arrest in the G₁/S phase via targeting the cell-cycle activator CDK6 and the antimitogenic p130 [56]. Nevertheless, *miR-107* has another aspect for directly targeting *DICER1* mRNA and the high level of *miR-107* might affect the production and function of p53-induced miRNAs [57].

4.3. *miR-145*. It has been reported that the expression of *miR-145* is frequently decreased in colon tumors, breast and prostate cancers and that the chromosomal region (chromosome 5 [5q32-33] within a 4.09 kb region) is deleted in myelodysplastic syndrome, suggesting *miR-145* acts as a tumor suppressor [58–61]. The expression of *miR-145* is transcriptionally induced by p53, and *miR-145* downregulates c-MYC, E2F3, cyclin D2, CDK4, and CDK6 and leads to G₁ cell-cycle arrest [62, 63].

Recently, it has been found that *miR-145* contains several CpG sites in its promoter region and that the expression of *miR-145* is affected by epigenetic events such as DNA methylation [60]. The CpG regions are located adjacent to p53 response element upstream of *miR-145*, and DNA hypermethylation inhibits p53 from binding to *miR-145*. In addition to this miRNA, it has been reported that *miR-34a*, *miR-124a*, and *miR-127* are downregulated by DNA methylation [64].

4.4. *miR-192/215*. *miR-192* and *miR-215* share a similar seed sequence and are composed of two clusters: the *miR-215/miR-194-1* cluster on chromosome 1 (1q41) and the *miR-192/miR-194-2* cluster on chromosome 11 (11q13.1) [65]. *miR-192* and *miR-215* are downregulated in colon cancers, lung cancers, multiple myeloma, and renal cancers [66–69]. Some studies have suggested that these miRNAs are also under the control of p53 and can induce p21 expression and cell-cycle arrest in a partially p53-dependent manner [66, 70]. Gene expression analyses indicated that *miR-192* and *miR-215* target a number of transcripts that regulate DNA synthesis and the G₁ and G₂ cell-cycle checkpoints, such as CDC7 and MAD2L1 [70]. Therefore, *miR-192/215* functions as a tumor suppressor contributing to the G₁ and G₂/M cell-cycle arrest.

4.5. *miR-200c*. It is well known that p53 acts as an important regulator in modulating epithelial-mesenchymal transition (EMT) that is implicated in tumor progression, metastasis, and the correlation of poor patient prognosis [71, 72]. The p53-induced *miR-200c* represses EMT by targeting the E-cadherin transcriptional repressors ZEB1 and ZEB2, Krüppel-like factor 4 (KLF4), and the polycomb repressor BMI1, all of which are involved in the maintenance of stemness [73–80]. Moreover, *miR-200c* contributes to the induction of apoptosis in cancer cells via the apoptosis-inducing receptor CD95 by targeting the apoptosis-inhibitor FAS-associated phosphatase 1 (FAP-1) [81].

4.6. *let-7a* and *let-7b*. *let-7* is known to be important for the regulation of development and is evolutionally conserved across bilaterian phylogeny [82]. In humans, some *let-7* gene clusters are located in fragile regions involved in cancers [61]. In lung cancers, it has been reported that the downregulated

expression of *let-7* members is correlated with poor prognosis [83, 84]. Recent works suggested that *let-7a* and *let-7b* expression is dependent on p53 in response to genotoxic stress and *let-7* miRNAs target CDK6, CDC25A, cyclin D, CDC34, and MYC [85–89]. On the other hand, *let-7a-d* and *let-7i* are direct targets of E2F1 and E2F3 during the G₁/S transition and are repressed in E2F1/3-null cells [90]. The *let-7* family plays multiple roles in the regulation of the cell cycle and goes a long way toward suppressing tumor progression.

4.7. *miR-17-92* Cluster. The *miR-17-92* cluster consists of *miR-17-5p*, *miR-17-3p*, *miR-18a*, *miR-19a*, *miR-20a*, *miR-19b*, and *miR-92-1*. Some of these are known to be oncogenic, as suggested in the research showing that the cluster is upregulated in human B-cell lymphoma and amplified in malignant lymphoma [91, 92].

Different from the miRNAs mentioned above, *miR-17-92* miRNAs are more or less repressed transcriptionally by p53 under hypoxia, which leads to the p53-mediated apoptosis [93]. The p53-binding site overlaps with the TATA box of the *miR-17-92* promoter region, and p53 prevents the TATA-binding protein (TBP) transcription factor from binding to the site during hypoxic conditions. Moreover, *miR-17-92* is transcriptionally regulated by c-Myc [94]. Although c-Myc is repressed by p53 activation under some stress conditions, the repression of *miR-17-92* is not dependent on c-MYC but on p53 under hypoxia [93, 95].

Note that some members of *miR-17-92* are likely to function as tumor suppressors in different cancers. For example, in breast cancer, *miR-17-5p* represses the expression of the nuclear receptor coactivator amplified in breast cancer 1 (*AIB1*) that enhances the transcription activity of E2F1 to promote the cell proliferation of breast cancer cells [96]. A recent study showed that *miR-17-3p* reduces tumor growth by targeting MDM2 in glioblastoma cells [97].

4.8. *miR-15a/miR-16-1*. *miR-15a* and *miR-16-1* were identified to be deleted and/or downregulated in approximately 68% of B-cell chronic lymphocytic leukemia (B-CLL) [98], as is the case in pituitary adenomas [99], gastric cancer cells [100], prostate cancer [101–104], non-small cell lung cancer [105, 106], ovarian cancer [107], and pancreatic cancer [108], which indicates their important functions for tumor formation. The miRNAs are encoded by an intron of a long noncoding RNA gene, deleted in lymphocytic leukemia 2 (*DLEU2*), and *DLEU2* (*miR-15a/miR-16-1*) was shown to be transactivated by p53 [109]. In addition, p53 regulates the expression level of precursor and mature *miR-15a* and *miR-16-1* as well as *miR-143* and *miR-145* [110]. It has been reported that *miR-15a/miR-16-1* negatively regulates the antiapoptotic protein BCL-2 and the cell-cycle regulators, such as CDK1, CDK2, and CDK6, and cyclins D1, D3, and E1 [102, 110–112].

5. miRNAs Regulating Negative Regulators of p53

It has been shown that MDM2 negatively controls the stability and transcription activity of p53, which attenuates the tumor-suppressive functions of p53 [40]. Actually, overexpression

Besides MDM2, a NAD-dependent deacetylase, silent information regulator 1 (SIRT1), increases the level of deacetylated p53 and negatively regulates the p53 activity [122, 123]. SIRT1 is targeted by the p53-inducible *miR-34a* and joins the positive feedback loop connecting the miRNA, SIRT1, and p53 (Figure 2) [124]. Additionally, *miR-499* participates in this regulatory circuit as the miRNA possesses a very similar seed sequence of *miR-34* members [125–127]. *miR-449* is upregulated by E2F1, not by p53, and *miR-34* and *miR-449* bring in an asymmetric network to balance the functions between p53 and E2F1.

As is the case in the control of negative regulators of p53 via miRNAs, p53 itself is repressed by several miRNAs through direct interaction with the 3'-UTR of *TP53* mRNA (Figure 3). *miR-125b* is a first-identified p53-repressive miRNA and blocks the p53 expression level to suppress apoptosis in human neuroblastoma and lung fibroblast cells; in contrast, the knockdown of *miR-125b* leads to the opposite results [128]. Plus, *miR-125a*, an isoform of *miR-125*, was suggested to inhibit the translation of *TP53* by binding to a region of the 3'-UTR [129]. The high expression of *miR-125b* is associated with poor prognosis in patients with colorectal cancer [130].

Some studies have shown that *miR-125b* represses factors in the p53 network, including apoptosis regulators like *PUMA*, insulin-like growth factor-binding protein 3 (*IGFBP3*), and BCL2-antagonist/killer 1 (*BAK1*) and cell-cycle regulators like *cyclin C*, *CDC25C*, and cyclin-dependent kinase inhibitor 2C (*CDKN2C*) [131]. These suggest that *miR-125b* modulates and buffers the p53 pathway.

Subsequently, *miR-504* was reported to directly repress the p53 protein level and reduce the p53-mediated apoptosis and cell-cycle arrest in response to stress, and its overexpression promotes the tumorigenicity of colon cancer cells *in vivo* [132]. Additionally, *miR-380-5p*, *miR-33*, and *miR-1285* can downregulate the p53 protein expression by directly binding to the two sites in the 3'-UTR of *TP53*, resulting in the reduction of apoptosis and cell-cycle arrest [133–135]. Indeed, *miR-380-5p* is highly expressed in neuroblastomas with neuroblastoma-derived v-myc myelocytomatosis viral-related oncogene (*MYCN*) amplification, and the high expression level correlates with poor diagnosis [133]. More recently, *miR-30d* and *miR-25* also directly interacted with the 3'-UTR of *TP53* to decrease the p53 level. So then, these miRNAs affect apoptotic cell death, cell-cycle arrest, and cellular senescence in some cell lines, such as multiple myelomas, colon cancer, and lung cancer cells [136–138]. When taken together, the miRNAs targeting *TP53* would hinder p53 from exerting its tumor-suppressive functions (senescence, apoptosis, cell-cycle arrest, etc.) under stressed conditions.

7. Concluding Remarks

For more than a decade, small noncoding RNAs have become increasingly central to the study of tumor biology. The accumulating evidence of cancer-associated miRNAs reveals the missing link between classic tumor-suppressive networks and complex oncogenic pathways. In a stress situation, p53 directly induces various protein-coding genes such as *p21* and *PUMA* to contribute to cell-cycle arrest and apoptosis and, furthermore, utilizes tumor-suppressive miRNAs, such as *miR-34s*, *miR-107*, and *miR-145* (Figure 1 and Table 1). Some of the p53-inducible miRNAs target p53-negative regulators (MDM2 and SIRT1), which creates a positive feedback loop to reinforce p53 stability and activity (Figure 2). However, as expected, miRNAs are not always on p53's side: p53-repressive miRNAs (*miR-125s*, *miR-504*, *miR-380-5p*, etc.) reduce the p53 expression level by binding to a region of the 3'-UTR of *TP53* mRNA and result in the inhibition of cell-cycle arrest and apoptosis (Figure 3). There will be more than one way to arrest the cell-cycle and/or induce apoptosis, and the balance between miRNAs and tumor suppressors might be crucial in deciding which strategy to adapt.

For future diagnostic and therapeutic advances, more extensive studies will be needed to find hidden messages in the tumor-suppressive networks of miRNA. The regulatory mechanism of the p53-miRNA circuit has been excellently shown, but the upstream regulators of almost all miRNAs are unknown at this time. What is more, regardless of computational prediction, the downstream targets of miRNA are hard to identify exactly because of the imperfect complementarity

and the possibility that miRNAs can bind to not only the 3'-UTR but also the 5'-UTR and coding regions.

In recent years, the competitive endogenous RNA (ceRNA) hypothesis has suggested that noncoding pseudogenes and long noncoding RNAs act as miRNA sponges, which is likely to counteract the effect of miRNAs on the target mRNA transcripts [139]. Therefore, we need to move deeper inside the world of noncoding RNAs in order to prevent and treat diverse cancers.

Besides the miRNAs described in this paper, there are many miRNAs related to cell-cycle regulation and apoptosis [140–142]. However, it is unclear how these miRNAs act additively/synergistically on tumor suppression. Even the longest journey to understand the role of miRNA begins with a single experiment. The next ten years will be more exciting in the quest to see cancer conquered.

Conflict of Interests

The authors declare no competing financial interests.

Acknowledgments

The authors thank Fumitaka Takeshita (National Cancer Research Institute), Norimitsu Yamagata, and Ryotaro Fujimura (Kewpie Corporation) for providing valuable comments on this paper. This work was supported in part by a Grant-in-Aid for Scientific Research on Priority Areas, Cancer, from the Ministry of Education, Culture, Sports, Science and Technology, Japan; the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NiBio); Project for Development of Innovative Research on Cancer Therapeutics (P-Direct); and Comprehensive Research and Development of a Surgical Instrument for Early Detection and Rapid Curing of Cancer Project (P10003) of the New Energy and Industrial Technology Development Organization (NEDO).

References

- [1] G. A. Calin and C. M. Croce, "MicroRNA signatures in human cancers," *Nature Reviews Cancer*, vol. 6, no. 11, pp. 857–866, 2006.
- [2] K. H. Vousden and D. P. Lane, "p53 in health and disease," *Nature Reviews Molecular Cell Biology*, vol. 8, no. 4, pp. 275–283, 2007.
- [3] S. Pelengaris, M. Khan, and G. Evan, "c-MYC: more than just a matter of life and death," *Nature Reviews Cancer*, vol. 2, no. 10, pp. 764–776, 2002.
- [4] B. Vogelstein and K. W. Kinzler, "Cancer genes and the pathways they control," *Nature Medicine*, vol. 10, no. 8, pp. 789–799, 2004.
- [5] M. Olivier, S. P. Hussain, C. C. de Fromental, P. Hainaut, and C. C. Harris, "TP53 mutation spectra and load: a tool for generating hypotheses on the etiology of cancer," *IARC Scientific Publications*, vol. 157, no. 6, pp. 247–270, 2004.
- [6] A. J. Levine, W. Hu, and Z. Feng, "The P53 pathway: what questions remain to be explored?" *Cell Death and Differentiation*, vol. 13, no. 6, pp. 1027–1036, 2006.

- [7] W. S. El-Deiry, T. Tokino, V. E. Velculescu et al., "WAF1, a potential mediator of p53 tumor suppression," *Cell*, vol. 75, no. 4, pp. 817–825, 1993.
- [8] J. Yu, L. Zhang, P. M. Hwang, K. W. Kinzler, and B. Vogelstein, "PUMA induces the rapid apoptosis of colorectal cancer cells," *Molecular Cell*, vol. 7, no. 3, pp. 673–682, 2001.
- [9] H. Hermeking, C. Lengauer, K. Polyak et al., "14-3-3 σ is a p53-regulated inhibitor of G2/M progression," *Molecular Cell*, vol. 1, no. 1, pp. 3–11, 1997.
- [10] K. H. Vousden and C. Prives, "Blinded by the light: the growing complexity of p53," *Cell*, vol. 137, no. 3, pp. 413–431, 2009.
- [11] K. B. Spurgers, D. L. Gold, K. R. Coombes et al., "Identification of cell cycle regulatory genes as principal targets of p53-mediated transcriptional repression," *The Journal of Biological Chemistry*, vol. 281, no. 35, pp. 25134–25141, 2006.
- [12] J. Lu, G. Getz, E. A. Miska et al., "MicroRNA expression profiles classify human cancers," *Nature*, vol. 435, no. 7043, pp. 834–838, 2005.
- [13] A. Lujambio and S. W. Lowe, "The microcosmos of cancer," *Nature*, vol. 482, no. 7385, pp. 347–355, 2012.
- [14] N. Raver-Shapira, E. Marciano, E. Meiri et al., "Transcriptional activation of miR-34a contributes to p53-mediated apoptosis," *Molecular Cell*, vol. 26, no. 5, pp. 731–743, 2007.
- [15] T.-C. Chang, E. A. Wentzel, O. A. Kent et al., "Transactivation of miR-34a by p53 broadly influences gene expression and promotes apoptosis," *Molecular Cell*, vol. 26, no. 5, pp. 745–752, 2007.
- [16] L. He, X. He, L. P. Lim et al., "A microRNA component of the p53 tumour suppressor network," *Nature*, vol. 447, no. 7148, pp. 1130–1134, 2007.
- [17] V. Tarasov, P. Jung, B. Verdoodt et al., "Differential regulation of microRNAs by p53 revealed by massively parallel sequencing: miR-34a is a p53 target that induces apoptosis and G₁-arrest," *Cell Cycle*, vol. 6, no. 13, pp. 1586–1593, 2007.
- [18] G. T. Bommer, I. Gerin, Y. Feng et al., "p53-mediated activation of miRNA34 candidate tumor-suppressor genes," *Current Biology*, vol. 17, no. 15, pp. 1298–1307, 2007.
- [19] R. C. Lee, R. L. Feinbaum, and V. Ambros, "The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*," *Cell*, vol. 75, no. 5, pp. 843–854, 1993.
- [20] B. Wightman, I. Ha, and G. Ruvkun, "Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*," *Cell*, vol. 75, no. 5, pp. 855–862, 1993.
- [21] B. J. Reinhart, F. J. Slack, M. Basson et al., "The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*," *Nature*, vol. 403, no. 6772, pp. 901–906, 2000.
- [22] B. P. Lewis, C. B. Burge, and D. P. Bartel, "Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets," *Cell*, vol. 120, no. 1, pp. 15–20, 2005.
- [23] A. Krek, D. Grün, M. N. Poy et al., "Combinatorial microRNA target predictions," *Nature Genetics*, vol. 37, no. 5, pp. 495–500, 2005.
- [24] Y. Lee, M. Kim, J. J. Han et al., "MicroRNA genes are transcribed by RNA polymerase II," *The EMBO Journal*, vol. 23, no. 20, pp. 4051–4060, 2004.
- [25] G. M. Borchert, W. Lanier, and B. L. Davidson, "RNA polymerase III transcribes human microRNAs," *Nature Structural and Molecular Biology*, vol. 13, no. 12, pp. 1097–1101, 2006.
- [26] V. N. Kim, "MicroRNA biogenesis: coordinated cropping and dicing," *Nature Reviews Molecular Cell Biology*, vol. 6, no. 5, pp. 376–385, 2005.
- [27] Y. Lee, K. Jeon, J.-T. Lee, S. Kim, and V. N. Kim, "MicroRNA maturation: stepwise processing and subcellular localization," *The EMBO Journal*, vol. 21, no. 17, pp. 4663–4670, 2002.
- [28] E. Lund, S. Güttinger, A. Calado, J. E. Dahlberg, and U. Kutay, "Nuclear export of microRNA precursors," *Science*, vol. 303, no. 5654, pp. 95–98, 2004.
- [29] D. P. Bartel, "MicroRNAs: genomics, biogenesis, mechanism, and function," *Cell*, vol. 116, no. 2, pp. 281–297, 2004.
- [30] V. N. Kim, "Small RNAs: classification, biogenesis, and function," *Molecules and Cells*, vol. 19, no. 1, pp. 1–15, 2005.
- [31] L. P. Lim, N. C. Lau, P. Garrett-Engle et al., "Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs," *Nature*, vol. 433, no. 7027, pp. 769–773, 2005.
- [32] S. Bagga, J. Bracht, S. Hunter et al., "Regulation by let-7 and lin-4 miRNAs results in target mRNA degradation," *Cell*, vol. 122, no. 4, pp. 553–563, 2005.
- [33] E. C. Lai, "Predicting and validating microRNA targets," *Genome Biology*, vol. 5, no. 9, article 115, 2004.
- [34] L. Wu and J. G. Belasco, "Let me count the ways: mechanisms of gene regulation by miRNAs and siRNAs," *Molecular Cell*, vol. 29, no. 1, pp. 1–7, 2008.
- [35] A. Eulalio, E. Huntzinger, and E. Izaurralde, "Getting to the root of miRNA-mediated gene silencing," *Cell*, vol. 132, no. 1, pp. 9–14, 2008.
- [36] R. W. Carthew and E. J. Sontheimer, "Origins and mechanisms of miRNAs and siRNAs," *Cell*, vol. 136, no. 4, pp. 642–655, 2009.
- [37] L. A. Yates, C. J. Norbury, and R. J. C. Gilbert, "The long and short of microRNA," *Cell*, vol. 153, no. 3, pp. 516–519, 2013.
- [38] Y. Haupt, R. Maya, A. Kazaz, and M. Oren, "Mdm2 promotes the rapid degradation of p53," *Nature*, vol. 387, no. 6630, pp. 296–299, 1997.
- [39] M. H. G. Kubbutat, S. N. Jones, and K. H. Vousden, "Regulation of p53 stability by Mdm2," *Nature*, vol. 387, no. 6630, pp. 299–303, 1997.
- [40] C. L. Brooks and W. Gu, "p53 ubiquitination: mdm2 and beyond," *Molecular Cell*, vol. 21, no. 3, pp. 307–315, 2006.
- [41] L. Roger, G. Gadea, and P. Roux, "Control of cell migration: a tumour suppressor function for p53?" *Biology of the Cell*, vol. 98, no. 3, pp. 141–152, 2006.
- [42] J. G. Teodoro, A. E. Parker, X. Zhu, and M. R. Green, "p53-mediated inhibition of angiogenesis through up-regulation of a collagen prolyl hydroxylase," *Science*, vol. 313, no. 5789, pp. 968–971, 2006.
- [43] M. R. Junttila and G. I. Evan, "P53—a Jack of all trades but master of none," *Nature Reviews Cancer*, vol. 9, no. 11, pp. 821–829, 2009.
- [44] S. A. Gatz and L. Wiesmüller, "p53 in recombination and repair," *Cell Death and Differentiation*, vol. 13, no. 6, pp. 1003–1016, 2006.
- [45] K. Bensaad and K. H. Vousden, "Savior and slayer: the two faces of p53," *Nature Medicine*, vol. 11, no. 12, pp. 1278–1279, 2005.
- [46] K. Bensaad, A. Tsuruta, M. A. Selak et al., "TIGAR, a p53-inducible regulator of glycolysis and apoptosis," *Cell*, vol. 126, no. 1, pp. 107–120, 2006.
- [47] S. Matoba, J.-G. Kang, W. D. Patino et al., "p53 regulates mitochondrial respiration," *Science*, vol. 312, no. 5780, pp. 1650–1653, 2006.

- [48] F. Murray-Zmijewski, D. P. Lane, and J.-C. Bourdon, "p53/p63/p73 isoforms: an orchestra of isoforms to harmonise cell differentiation and response to stress," *Cell Death and Differentiation*, vol. 13, no. 6, pp. 962–972, 2006.
- [49] S. Cawley, S. Bekiranov, H. H. Ng et al., "Unbiased mapping of transcription factor binding sites along human chromosomes 21 and 22 points to widespread regulation of noncoding RNAs," *Cell*, vol. 116, no. 4, pp. 499–509, 2004.
- [50] C.-L. Wei, Q. Wu, V. B. Vega et al., "A global map of p53 transcription-factor binding sites in the human genome," *Cell*, vol. 124, no. 1, pp. 207–219, 2006.
- [51] H. Tazawa, N. Tsuchiya, M. Izumiya, and H. Nakagama, "Tumor-suppressive *miR-34a* induces senescence-like growth arrest through modulation of the E2F pathway in human colon cancer cells," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 39, pp. 15472–15477, 2007.
- [52] D. C. Corney, A. Flesken-Nikitin, A. K. Godwin, W. Wang, and A. Y. Nikitin, "*MicroRNA-34b* and *MicroRNA-34c* are targets of p53 and cooperate in control of cell proliferation and adhesion-independent growth," *Cancer Research*, vol. 67, no. 18, pp. 8433–8438, 2007.
- [53] H. Hermeking, "MicroRNAs in the p53 network: micromanagement of tumour suppression," *Nature Reviews Cancer*, vol. 12, no. 9, pp. 613–626, 2012.
- [54] H. Hermeking, "The *miR-34* family in cancer and apoptosis," *Cell Death & Differentiation*, vol. 17, no. 2, pp. 193–199, 2010.
- [55] M. Yamakuchi, C. D. Lotterman, C. Bao et al., "P53-induced microRNA-107 inhibits HIF-1 and tumor angiogenesis," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 14, pp. 6334–6339, 2010.
- [56] L. Böhlig, M. Friedrich, and K. Engeland, "P53 activates the *PANK1/miRNA-107* gene leading to downregulation of CDK6 and p130 cell cycle proteins," *Nucleic Acids Research*, vol. 39, no. 2, pp. 440–453, 2011.
- [57] G. Martello, A. Rosato, F. Ferrari et al., "A microRNA targeting dicer for metastasis control," *Cell*, vol. 141, no. 7, pp. 1195–1207, 2010.
- [58] M. Z. Michael, S. M. O'Connor, N. G. van Holst Pellekaan, G. P. Young, and R. J. James, "Reduced accumulation of specific microRNAs in colorectal neoplasia," *Molecular Cancer Research*, vol. 1, no. 12, pp. 882–891, 2003.
- [59] M. V. Iorio, M. Ferracin, C.-G. Liu et al., "MicroRNA gene expression deregulation in human breast cancer," *Cancer Research*, vol. 65, no. 16, pp. 7065–7070, 2005.
- [60] S. O. Suh, Y. Chen, M. S. Zaman et al., "MicroRNA-145 is regulated by DNA methylation and p53 gene mutation in prostate cancer," *Carcinogenesis*, vol. 32, no. 5, pp. 772–778, 2011.
- [61] G. A. Calin, C. Sevignani, C. D. Dumitru et al., "Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, no. 9, pp. 2999–3004, 2004.
- [62] M. Sachdeva, S. M. Zhu, F. T. Wu et al., "p53 represses c-Myc through induction of the tumor suppressor *miR-145*," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 106, no. 9, pp. 3207–3212, 2009.
- [63] H. Zhu, U. Dougherty, V. Robinson et al., "EGFR signals down-regulate tumor suppressors *miR-143* and *miR-145* in Western diet-promoted murine colon cancer: role of G₁ regulators," *Molecular Cancer Research*, vol. 9, no. 7, pp. 960–975, 2011.
- [64] A. Lujambio and M. Esteller, "CpG island hypermethylation of tumor suppressor microRNAs in human cancer," *Cell Cycle*, vol. 6, no. 12, pp. 1455–1459, 2007.
- [65] H. W. Khella, M. Bakhet, G. Allo et al., "miR-192, miR-194 and miR-215: a convergent microRNA network suppressing tumor progression in renal cell carcinoma," *Carcinogenesis*, vol. 34, no. 10, pp. 2231–2239, 2013.
- [66] C. J. Braun, X. Zhang, I. Savelyeva et al., "p53-responsive microRNAs 192 and 215 are capable of inducing cell cycle arrest," *Cancer Research*, vol. 68, no. 24, pp. 10094–10104, 2008.
- [67] S. Feng, S. Cong, X. Zhang et al., "MicroRNA-192 targeting retinoblastoma 1 inhibits cell proliferation and induces cell apoptosis in lung cancer cells," *Nucleic Acids Research*, vol. 39, no. 15, pp. 6669–6678, 2011.
- [68] F. Pichiorri, S.-S. Suh, A. Rocci et al., "Downregulation of p53-inducible microRNAs 192, 194, and 215 impairs the p53/MDM2 autoregulatory loop in multiple myeloma development," *Cancer Cell*, vol. 18, no. 4, pp. 367–381, 2010.
- [69] U. Senanayake, S. Das, P. Vesely et al., "miR-192, miR-194, miR-215, miR-200c and miR-141 are downregulated and their common target ACVR2B is strongly expressed in renal childhood neoplasms," *Carcinogenesis*, vol. 33, no. 5, pp. 1014–1021, 2012.
- [70] S. A. Georges, M. C. Biery, S.-Y. Kim et al., "Coordinated regulation of cell cycle transcripts by p53-inducible microRNAs, miR-192 and miR-215," *Cancer Research*, vol. 68, no. 24, pp. 10105–10112, 2008.
- [71] K. Polyak and R. A. Weinberg, "Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits," *Nature Reviews Cancer*, vol. 9, no. 4, pp. 265–273, 2009.
- [72] S. Valastyan and R. A. Weinberg, "Tumor metastasis: molecular insights and evolving paradigms," *Cell*, vol. 147, no. 2, pp. 275–292, 2011.
- [73] C. Chang, C. H. Chao, W. Xia et al., "P53 regulates epithelial-mesenchymal transition and stem cell properties through modulating miRNAs," *Nature Cell Biology*, vol. 13, no. 3, pp. 317–323, 2011.
- [74] T. Kim, A. Veronese, F. Pichiorri et al., "p53 regulates epithelial-mesenchymal transition through microRNAs targeting ZEB1 and ZEB2," *Journal of Experimental Medicine*, vol. 208, no. 5, pp. 875–883, 2011.
- [75] U. Burk, J. Schubert, U. Wellner et al., "A reciprocal repression between ZEB1 and members of the miR-200 family promotes EMT and invasion in cancer cells," *EMBO Reports*, vol. 9, no. 6, pp. 582–589, 2008.
- [76] P. A. Gregory, A. G. Bert, E. L. Paterson et al., "The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1," *Nature Cell Biology*, vol. 10, no. 5, pp. 593–601, 2008.
- [77] M. Korpai, E. S. Lee, G. Hu, and Y. Kang, "The miR-200 family inhibits epithelial-mesenchymal transition and cancer cell migration by direct targeting of E-cadherin transcriptional repressors ZEB1 and ZEB2," *The Journal of Biological Chemistry*, vol. 283, no. 22, pp. 14910–14914, 2008.
- [78] S.-M. Park, A. B. Gaur, E. Lengyel, and M. E. Peter, "The miR-200 family determines the epithelial phenotype of cancer cells by targeting the E-cadherin repressors ZEB1 and ZEB2," *Genes & Development*, vol. 22, no. 7, pp. 894–907, 2008.
- [79] Y. Shimono, M. Zabala, R. W. Cho et al., "Downregulation of miRNA-200c links breast cancer stem cells with normal stem cells," *Cell*, vol. 138, no. 3, pp. 592–603, 2009.

- [80] U. Wellner, J. Schubert, U. C. Burk et al., "The EMT-activator ZEB1 promotes tumorigenicity by repressing stemness-inhibiting microRNAs," *Nature Cell Biology*, vol. 11, no. 12, pp. 1487–1495, 2009.
- [81] R. Schickel, S.-M. Park, A. E. Murmann, and M. E. Peter, "miR-200c regulates induction of apoptosis through CD95 by targeting FAP-1," *Molecular Cell*, vol. 38, no. 6, pp. 908–915, 2010.
- [82] A. E. Pasquinelli, B. J. Reinhart, F. Slack et al., "Conservation of the sequence and temporal expression of let-7 heterochronic regulatory RNA," *Nature*, vol. 408, no. 6808, pp. 86–89, 2000.
- [83] J. Takamizawa, H. Konishi, K. Yanagisawa et al., "Reduced expression of the let-7 microRNAs in human lung cancers in association with shortened postoperative survival," *Cancer Research*, vol. 64, no. 11, pp. 3753–3756, 2004.
- [84] N. Yanaihara, N. Caplen, E. Bowman et al., "Unique microRNA molecular profiles in lung cancer diagnosis and prognosis," *Cancer Cell*, vol. 9, no. 3, pp. 189–198, 2006.
- [85] A. D. Saleh, J. E. Savage, L. Cao et al., "Cellular stress induced alterations in microRNA let-7a and let-7b expression are dependent on p53," *PLoS ONE*, vol. 6, no. 10, Article ID e24429, 2011.
- [86] C. D. Johnson, A. Esquela-Kerscher, G. Stefani et al., "The let-7 microRNA represses cell proliferation pathways in human cells," *Cancer Research*, vol. 67, no. 16, pp. 7713–7722, 2007.
- [87] A. Legesse-Miller, O. Elemento, S. J. Pfau, J. J. Forman, S. Tavazoie, and H. A. Collier, "Let-7 overexpression leads to an increased fraction of cells in G₂/M, direct down-regulation of Cdc34, and stabilization of weel kinase in primary fibroblasts," *The Journal of Biological Chemistry*, vol. 284, no. 11, pp. 6605–6609, 2009.
- [88] V. B. Sampson, N. H. Rong, J. Han et al., "MicroRNA let-7a down-regulates MYC and reverts MYC-induced growth in Burkitt lymphoma cells," *Cancer Research*, vol. 67, no. 20, pp. 9762–9770, 2007.
- [89] M. J. Bueno, M. Gómez de Cedrón, G. Gómez-López et al., "Combinatorial effects of microRNAs to suppress the Myc oncogenic pathway," *Blood*, vol. 117, no. 23, pp. 6255–6266, 2011.
- [90] M. J. Bueno, M. G. de Cedrón, U. Laresgoiti, J. Fernández-Piqueras, A. M. Zubiaga, and M. Malumbres, "Multiple E2F-induced microRNAs prevent replicative stress in response to mitogenic signaling," *Molecular and Cellular Biology*, vol. 30, no. 12, pp. 2983–2995, 2010.
- [91] A. Ota, H. Tagawa, S. Karnan et al., "Identification and characterization of a novel gene, *Clorf25*, as a target for 13q31-q32 amplification in malignant lymphoma," *Cancer Research*, vol. 64, no. 9, pp. 3087–3095, 2004.
- [92] L. He, J. M. Thomson, M. T. Hemann et al., "A microRNA polycistron as a potential human oncogene," *Nature*, vol. 435, no. 7043, pp. 828–833, 2005.
- [93] H.-L. Yan, G. Xue, Q. Mei et al., "Repression of the *miR-17-92* cluster by p53 has an important function in hypoxia-induced apoptosis," *EMBO Journal*, vol. 28, no. 18, pp. 2719–2732, 2009.
- [94] K. A. O'Donnell, E. A. Wentzel, K. I. Zeller, C. V. Dang, and J. T. Mendell, "c-Myc-regulated microRNAs modulate E2F1 expression," *Nature*, vol. 435, no. 7043, pp. 839–843, 2005.
- [95] J. S. L. Ho, W. Ma, D. Y. L. Mao, and S. Benchimol, "p53-dependent transcriptional repression of c-myc is required for G₁ cell cycle arrest," *Molecular and Cellular Biology*, vol. 25, no. 17, pp. 7423–7431, 2005.
- [96] A. Hossain, M. T. Kuo, and G. F. Saunders, "Mir-17-5p regulates breast cancer cell proliferation by inhibiting translation of *AIB1* mRNA," *Molecular and Cellular Biology*, vol. 26, no. 21, pp. 8191–8201, 2006.
- [97] H. Li and B. B. Yang, "Stress response of glioblastoma cells mediated by miR-17-5p targeting PTEN and the passenger strand miR-17-3p targeting MDM2," *Oncotarget*, vol. 3, no. 12, pp. 1653–1668, 2012.
- [98] G. A. Calin, C. D. Dumitru, M. Shimizu et al., "Frequent deletions and down-regulation of micro-RNA genes *miR15* and *miR16* at 13q14 in chronic lymphocytic leukemia," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 24, pp. 15524–15529, 2002.
- [99] A. Bottoni, D. Piccin, F. Tagliati, A. Luchin, M. C. Zatelli, and E. C. D. Uberti, "miR-15a and miR-16-1 down-regulation in pituitary adenomas," *Journal of Cellular Physiology*, vol. 204, no. 1, pp. 280–285, 2005.
- [100] L. Xia, D. Zhang, R. Du et al., "miR-15b and miR-16 modulate multidrug resistance by targeting BCL2 in human gastric cancer cells," *International Journal of Cancer*, vol. 123, no. 2, pp. 372–379, 2008.
- [101] D. Bonci, V. Coppola, M. Musumeci et al., "The *miR-15a-miR-16-1* cluster controls prostate cancer by targeting multiple oncogenic activities," *Nature Medicine*, vol. 14, no. 11, pp. 1271–1277, 2008.
- [102] F. Takeshita, L. Patrawala, M. Osaki et al., "Systemic delivery of synthetic microRNA-16 inhibits the growth of metastatic prostate tumors via downregulation of multiple cell-cycle genes," *Molecular Therapy*, vol. 18, no. 1, pp. 181–187, 2010.
- [103] M. Musumeci, V. Coppola, A. Addario et al., "Control of tumor and microenvironment cross-talk by miR-15a and miR-16 in prostate cancer," *Oncogene*, vol. 30, no. 41, pp. 4231–4242, 2011.
- [104] K. P. Porkka, E.-L. Ogg, O. R. Saramäki et al., "The *miR-15a-miR-16-1* locus is homozygously deleted in a subset of prostate cancers," *Genes, Chromosomes and Cancer*, vol. 50, no. 7, pp. 499–509, 2011.
- [105] N. Bandi, S. Zbinden, M. Gugger et al., "miR-15a and miR-16 are implicated in cell cycle regulation in a Rb-dependent manner and are frequently deleted or down-regulated in non-small cell lung cancer," *Cancer Research*, vol. 69, no. 13, pp. 5553–5559, 2009.
- [106] N. Bandi and E. Vassella, "MiR-34a and miR-15a/16 are co-regulated in non-small cell lung cancer and control cell cycle progression in a synergistic and Rb-dependent manner," *Molecular Cancer*, vol. 10, article 55, 2011.
- [107] R. Bhattacharya, M. Nicoloso, R. Arvizo et al., "MiR-15a and MiR-16 control Bmi-1 expression in ovarian cancer," *Cancer Research*, vol. 69, no. 23, pp. 9090–9095, 2009.
- [108] X. J. Zhang, H. Ye, C. W. Zeng, B. He, H. Zhang, and Y. Q. Chen, "Dysregulation of miR-15a and miR-214 in human pancreatic cancer," *Journal of Hematology & Oncology*, vol. 3, article 46, 2010.
- [109] M. Fabbri, A. Bottoni, M. Shimizu et al., "Association of a microRNA/TP53 feedback circuitry with pathogenesis and outcome of B-cell chronic lymphocytic leukemia," *The Journal of the American Medical Association*, vol. 305, no. 1, pp. 59–67, 2011.
- [110] H. I. Suzuki, K. Yamagata, K. Sugimoto, T. Iwamoto, S. Kato, and K. Miyazono, "Modulation of microRNA processing by p53," *Nature*, vol. 460, no. 7254, pp. 529–533, 2009.
- [111] P. S. Linsley, J. Schelter, J. Burchard et al., "Transcripts targeted by the microRNA-16 family cooperatively regulate cell cycle progression," *Molecular and Cellular Biology*, vol. 27, no. 6, pp. 2240–2252, 2007.

- [112] Q. Liu, H. Fu, F. Sun et al., "miR-16 family induces cell cycle arrest by regulating multiple cell cycle genes," *Nucleic Acids Research*, vol. 36, no. 16, pp. 5391–5404, 2008.
- [113] E. Macias, A. Jin, C. Deisenroth et al., "An ARF-Independent c-MYC-activated tumor suppression pathway mediated by ribosomal protein-Mdm2 interaction," *Cancer Cell*, vol. 18, no. 3, pp. 231–243, 2010.
- [114] J. Zhang, Q. Sun, Z. Zhang, S. Ge, Z.-G. Han, and W.-T. Chen, "Loss of *microRNA-143/145* disturbs cellular growth and apoptosis of human epithelial cancers by impairing the MDM2-p53 feedback loop," *Oncogene*, vol. 32, no. 1, pp. 61–69, 2013.
- [115] J. Xiao, H. Lin, X. Luo, and Z. Wang, "miR-605 joins p53 network to form a p53:miR-605:Mdm2 positive feedback loop in response to stress," *EMBO Journal*, vol. 30, no. 3, pp. 524–532, 2011.
- [116] S.-Y. Park, J. H. Lee, M. Ha, J.-W. Nam, and V. N. Kim, "miR-29 miRNAs activate p53 by targeting p85 α and CDC42," *Nature Structural and Molecular Biology*, vol. 16, no. 1, pp. 23–29, 2009.
- [117] A. P. Ugalde, A. J. Ramsay, J. de la Rosa et al., "Aging and chronic DNA damage response activate a regulatory pathway involving miR-29 and p53," *EMBO Journal*, vol. 30, no. 11, pp. 2219–2232, 2011.
- [118] K. Okamoto, H. Li, M. R. Jensen et al., "Cyclin G recruits PP2A to dephosphorylate Mdm2," *Molecular Cell*, vol. 9, no. 4, pp. 761–771, 2002.
- [119] F. Fornari, L. Gramantieri, C. Giovannini et al., "MiR-122/cyclin G1 interaction modulates p53 activity and affects doxorubicin sensitivity of human hepatocarcinoma cells," *Cancer Research*, vol. 69, no. 14, pp. 5761–5767, 2009.
- [120] S.-S. Suh, J. Y. Yoo, G. J. Nuovo et al., "MicroRNAs/TP53 feedback circuitry in glioblastoma multiforme," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 109, no. 14, pp. 5316–5321, 2012.
- [121] A. A. Dar, S. Majid, C. Rittsteuer et al., "The role of miR-18b in MDM2-p53 pathway signaling and melanoma progression," *Journal of the National Cancer Institute*, vol. 105, no. 6, pp. 433–442, 2013.
- [122] J. Luo, A. Y. Nikolaev, S.-I. Imai et al., "Negative control of p53 by Sir2 α promotes cell survival under stress," *Cell*, vol. 107, no. 2, pp. 137–148, 2001.
- [123] H. Vaziri, S. K. Dessain, E. N. Eaton et al., "*hSIR2^{SIRT1}* functions as an NAD-dependent p53 deacetylase," *Cell*, vol. 107, no. 2, pp. 149–159, 2001.
- [124] M. Yamakuchi, M. Ferlito, and C. J. Lowenstein, "miR-34a repression of SIRT1 regulates apoptosis," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 36, pp. 13421–13426, 2008.
- [125] M. Lizé, S. Pilarski, and M. Dobbstein, "E2F1-inducible microRNA 449a/b suppresses cell proliferation and promotes apoptosis," *Cell Death & Differentiation*, vol. 17, no. 3, pp. 452–458, 2010.
- [126] T. Bou Kheir, E. Futoma-Kazmierczak, A. Jacobsen et al., "miR-449 inhibits cell proliferation and is down-regulated in gastric cancer," *Molecular Cancer*, vol. 10, article 29, 2011.
- [127] M. Lizé, A. Klimke, and M. Dobbstein, "MicroRNA-449 in cell fate determination," *Cell Cycle*, vol. 10, no. 17, pp. 2874–2882, 2011.
- [128] M. T. N. Le, C. Teh., N. Shyh-Chang et al., "MicroRNA-125b is a novel negative regulator of p53," *Genes and Development*, vol. 23, no. 7, pp. 862–876, 2009.
- [129] Y. Zhang, J.-S. Gao, X. Tang et al., "MicroRNA 125a and its regulation of the p53 tumor suppressor gene," *FEBS Letters*, vol. 583, no. 22, pp. 3725–3730, 2009.
- [130] N. Nishida, T. Yokobori, K. Mimori et al., "MicroRNA *miR-125b* is a prognostic marker in human colorectal cancer," *International Journal of Oncology*, vol. 38, no. 5, pp. 1437–1443, 2011.
- [131] M. T. N. Le, N. Shyh-Chang, S. L. Khaw et al., "Conserved regulation of p53 network dosage by microRNA-125b occurs through evolving miRNA-target gene pairs," *PLoS Genetics*, vol. 7, no. 9, Article ID e1002242, 2011.
- [132] W. Hu, C. S. Chan, R. Wu et al., "Negative regulation of tumor suppressor p53 by microRNA miR-504," *Molecular Cell*, vol. 38, no. 5, pp. 689–699, 2010.
- [133] A. Swarbrick, S. L. Woods, A. Shaw et al., "MiR-380-5p represses p53 to control cellular survival and is associated with poor outcome in MYCN-amplified neuroblastoma," *Nature Medicine*, vol. 16, no. 10, pp. 1134–1140, 2010.
- [134] A. Herrera-Merchan, C. Cerrato, G. Luengo et al., "miR-33-mediated downregulation of p53 controls hematopoietic stem cell self-renewal," *Cell Cycle*, vol. 9, no. 16, pp. 3277–3285, 2010.
- [135] S. Tian, S. Huang, S. Wu, W. Guo, J. Li, and X. He, "MicroRNA-1285 inhibits the expression of p53 by directly targeting its 3' untranslated region," *Biochemical and Biophysical Research Communications*, vol. 396, no. 2, pp. 435–439, 2010.
- [136] J. Li, S. Donath, Y. Li, D. Qin, B. S. Prabhakar, and P. Li, "miR-30 regulates mitochondrial fission through targeting p53 and the dynamin-related protein-1 pathway," *PLoS Genetics*, vol. 6, no. 1, Article ID e1000795, 2010.
- [137] M. Kumar, Z. Lu, A. A. L. Takwi et al., "Negative regulation of the tumor suppressor *p53* gene by microRNAs," *Oncogene*, vol. 30, no. 7, pp. 843–853, 2011.
- [138] N. Li, S. Kaur, J. Greshock et al., "A combined array-based comparative genomic hybridization and functional library screening approach identifies mir-30d as an oncomir in cancer," *Cancer Research*, vol. 72, no. 1, pp. 154–164, 2012.
- [139] L. Salmena, L. Poliseno, Y. Tay, L. Kats, and P. P. Pandolfi, "A *ceRNA* hypothesis: the rosetta stone of a hidden RNA language?" *Cell*, vol. 146, no. 3, pp. 353–358, 2011.
- [140] R. Schickel, B. Boyerinas, S.-M. Park, and M. E. Peter, "MicroRNAs: key players in the immune system, differentiation, tumorigenesis and cell death," *Oncogene*, vol. 27, no. 45, pp. 5959–5974, 2008.
- [141] N. Tsuchiya, M. Izumiya, H. Ogata-Kawata et al., "Tumor suppressor *miR-22* determines p53-dependent cellular fate through post-transcriptional regulation of p21," *Cancer Research*, vol. 71, no. 13, pp. 4628–4639, 2011.
- [142] M. J. Bueno and M. Malumbres, "MicroRNAs and the cell cycle," *Biochimica et Biophysica Acta*, vol. 1812, no. 5, pp. 592–601, 2011.