Research Article

Molecular Detection of *Theileria* spp. in Livestock on Five Caribbean Islands

Jilei Zhang,¹ Patrick Kelly,² Jing Li,¹ Chuanling Xu,¹ and Chengming Wang¹

¹Jiangsu Co-Innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonoses, Yangzhou University College of Veterinary Medicine, Yangzhou, Jiangsu 225009, China

²Ross University School of Veterinary Medicine, Basseterre, Saint Kitts and Nevis

Correspondence should be addressed to Chengming Wang; wangcm@yzu.edu.cn

Received 6 October 2015; Revised 19 November 2015; Accepted 23 November 2015

Academic Editor: Carlos E. Almeida

Copyright © 2015 Jilei Zhang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Theileria spp. are tick-transmitted, intracellular apicomplexan protozoan parasites infecting a wide range of animals. As there is very limited information on the prevalence of *Theileria* spp. in the Caribbean we used the recently described genus-specific pan-*Theileria* FRET-qPCR to identify infected animals in the region and a standard 18S rRNA gene PCR and sequencing to determine the species involved. We found *Theileria* spp. in 9% of the convenience samples of animals ($n = 752$) studied from five Caribbean islands. Donkeys (20.0%; 5/25) were most commonly infected, followed by sheep (17.4%; 25/144), cattle (6.8%; 22/325), goats (5.0%; 12/238), and horses (5.0%; 1/20). Six species of *Theileria* were identified: *T. equi* (donkeys, cattle, goats, and sheep), *Theileria* sp. OT3 (sheep and goats), *Theileria* sp. NG-2013a (cattle), *Theileria* sp. YW-2014 (donkeys), *Theileria* sp. B15a (goats), and *Babesia vulpes* or a closely related organism (sheep and goats). Only *T. equi* has been previously reported in the Caribbean. Our findings expand the known host ranges of *Theileria* spp. and the known distribution of the organisms around the world.

1. Background

Theileria spp. are tick-transmitted, intracellular apicomplexan protozoan parasites infecting leukocytes and erythrocytes of a wide range of animals [1, 2]. The organisms have been described in all livestock species and can cause significant economic losses to farmers. They are transmitted by a variety of ixodid ticks of the genera *Rhipicephalus*, *Hyalomma*, *Amblyomma*, and *Haemaphysalis* [3]. Infections with some *Theileria* spp. can result in fever, anemia, hemoglobinuria, and death in severe cases, but many species are benign and cause minor or no signs. Animals that recover from acute or primary infections usually remain persistently infected and may act as reservoirs for tick vectors [4, 5]. Infected animals are found particularly in tropical and subtropical regions in Africa, the Middle East, Southern Europe, and Asia [6–11].

There is little information on infectious agents in livestock in the Caribbean although animal production is an important source of income for many people in the region. In the case of *Theileria* spp., morphological and serological evidence has

been presented that *T. mutans* and *T. velifera*, both benign species transmitted by *Amblyomma* spp., occur in cattle on Guadeloupe [12]. Also, an organism with the morphology of *T. mutans* was seen in a blood smear from a bovine on Martinique [13]. In Trinidad, *T. equi* (previously *Babesia equi*) has been demonstrated in horses with a specific nested 18S rRNA PCR [14, 15] and a serosurvey has provided supporting evidence for its presence [16].

While there are many tests to detect *Theileria* spp. in animals, their specificity varies, as does their usefulness in finding the full spectrum of organisms present in an area. Microscopic detection of parasites can be difficult with low parasitemia and does not readily allow differentiation of species [2]. Serological studies, although sensitive and relatively easy to perform, are not specific as there is cross-reactivity between *Theileria* spp. [17]. Although molecular techniques have been described, many are for specific species which limits their usefulness in surveys. Reverse line blotting (RLB) assays enable the simultaneous identification of multiple species [18, 19], but they are cumbersome and time demanding to perform and identifying stringent

	920	920	930	940	950	960	970	980	990	1000	1010	1020	1030	1040	1050	
<i>Theileria orientalis</i> HM538222	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	612	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	613	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	614	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	615	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria buffeli</i> HQ84967	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	612	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	613	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	614	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	615	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria annulata</i> KF429799	756	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	757	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	758	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	759	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	760	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	761	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	762	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	763	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria sergenti</i> EU083804	740	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	741	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	742	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	743	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	744	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	745	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	746	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	747	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria luwenshuni</i> JX495527	603	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	604	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria velifei</i> AF097993	602	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	603	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	604	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria ovis</i> AY508458	576	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	577	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	578	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	579	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	580	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	581	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	582	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	583	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria parva</i> L02366	602	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	603	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	604	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria ulienbergi</i> JF119835	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	612	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria equi</i> E1 AY534882	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	612	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	613	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	614	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	615	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	616	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	617	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	618	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria cervi</i> AY735119	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	612	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria lestoquardi</i> JQ917458	438	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	439	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	440	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	441	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	442	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	443	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	444	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	445	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria capreoli</i> AY260175	604	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria separata</i> AY726011	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	612	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria bircensis</i> AF499604	598	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	599	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	600	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	601	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	602	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	603	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	604	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria taurotragi</i> L19082	602	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	603	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	604	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria mutans</i> FJ213585	554	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	555	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	556	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	557	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	558	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	559	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	560	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	561	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria annae</i> JX454779	470	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	471	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	472	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	473	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	474	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	475	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	476	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	477	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria sp.</i> Thrivae AB981984	263	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	264	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	265	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	266	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	267	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	268	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	269	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	270	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria sp.</i> OT3 KF470868	604	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria sp.</i> NG2013 KF597076	263	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	264	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	265	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	266	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	267	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	268	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	269	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	270	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria sp.</i> YW2014 AB981984	263	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	264	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	265	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	266	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	267	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	268	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	269	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	270	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>Theileria sp.</i> B15a JN572700	559	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	560	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	561	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	562	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	563	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	564	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	565	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	566	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>B. hongkongensis</i> JQ867356	549	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	550	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	551	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	552	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	553	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	554	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	555	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	556	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>B. divergens</i> AJ439713	596	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	597	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	598	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	599	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	600	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	601	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	602	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	603	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>B. bovis</i> JQ273013	568	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	569	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	570	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	571	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	572	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	573	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	574	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	575	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>B. bigemina</i> JQ23014	585	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	586	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	587	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	588	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	589	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	590	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	591	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	592	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>B. gibsoni</i> EU583386	602	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	603	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	604	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	605	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	606	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	607	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	608	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	609	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>B. microti</i> AB219802	610	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	611	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	612	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	613	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	614	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	615	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	616	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	617	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>B. felis</i> AF244912	586	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	587	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	588	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	589	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	590	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	591	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	592	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	593	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT
<i>B. canis vogeli</i> HM590440	583	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	584	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	585	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	586	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	587	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	588	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	589	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT	590	CGTAGTGAATTTCTGCTGCATTACATATCCTTGTGAGTTTCTTTT

FIGURE 1: Alignment of the sequences in the polymorphic region of the 18S RNA gene of *Theileria* spp. and related organisms that was targeted in the standard PCR used in the study. Nucleotides that are identical for all species are highlighted in blue while those that vary between species and can be used for differentiation are not highlighted.

species-specific oligonucleotide sequences can be challenging [20]. Recently, a sensitive genus-specific pan-*Theileria* FRET-qPCR has been described that detects the recognized *Theileria* spp. of domestic animals in a single reaction (Table 1) [21]. To provide further data on *Theileria* spp. in the Caribbean, we used the pan-*Theileria* FRET-qPCR to screen livestock from five islands for evidence of infection. Further, we used a standard 18S rRNA PCR and gene sequencing on positive reactors to identify the *Theileria* spp. involved. The results of this survey are described below.

2. Materials and Methods

2.1. Samples Collection. Jugular venipuncture was used to collect blood in EDTA from convenience samples of apparently healthy livestock (cattle, goats, sheep, donkeys, and horses) on five Caribbean islands [22]. This study was reviewed and approved by the Institutional Animal Care and Use Committee of the Ross University School of Veterinary Medicine (RUSVM), St. Kitts. Owners of animals gave permission for the blood samples to be collected.

2.2. DNA Extraction. The DNA was extracted from aliquots (200 μ L) of the whole blood samples with the QIAamp DNA Blood Mini Kit (QIAGEN, Valencia, CA, USA) according to the manufacturer's instructions. The DNAs were eluted into 200 μ L Buffer AE and couriered to Yangzhou University College of Veterinary Medicine, China, at room temperature where they were frozen at -80°C until PCRs were performed.

2.3. PCRs for *Theileria* Detection and Species Determination. All the PCRs were performed on a Roche Light-Cycler 480-II platform with the HMBS gene as an endogenous control [23]. Samples found positive in the pan-*Theileria* FRET-qPCR were tested in a conventional PCR with primers targeting a highly polymorphic 584–610-nucleotide region of the 18S

rRNA gene of *Theileria* spp. (Table 2, Figure 1) [21]. The amplicons from positive conventional PCRs were sequenced directly with forward and reverse primers to determine the *Theileria* spp. present (BGI, Shanghai, China) [21] as has been done with 18S rRNA sequences in a number of previous studies [11, 14, 17, 21, 24].

3. Results and Discussion

The sensitive and specific pan-*Theileria* FRET-qPCR [21] we used in our study demonstrated that a substantial proportion of livestock (8.6%; 65/752) on the five Caribbean islands we studied were infected with *Theileria* (Table 3). Each island had positive animals and each livestock species we studied was found to be infected with *Theileria* spp. Sequencing of representative samples of positive 18S rRNA PCR amplicons we obtained ($n = 43$) showed that there was one recognized *Theileria* spp. (*T. equi*) present in the Caribbean, along with five less well characterized *Theileria* sp. (Tables 3 and 4, Figure 2). The average copy number of 18S rRNA per mL whole blood was relatively low at 116.6 ± 440.8 , indicating that the animals we studied were chronically infected.

Theileria equi and the *Theileria* sp. YW-2014 were the species we identified in equids. *Theileria* sp. YW-2014 has been described in a Sika deer (*Cervus nippon*) from Japan but there is little sequence data on the organism with only a 552 bp sequence of the 18S rRNA gene reported in GenBank (AB981984). On the other hand, *T. equi* is a well-recognized cause of equine piroplasmiasis [25], an important disease of horses which has been recognized and studied in the Caribbean [14–16]. The organism has also been found in dogs in Spain [26], South Africa [27], and Nigeria [28] with some having clinical signs that responded to appropriate treatment [29]. Our finding that *T. equi* also occurs in domestic ruminants further expands the host range of the organism. The significance, extent, and consequences of infections with *T. equi* in domestic ruminants require further investigation.

TABLE 1: Alignment of the nucleotides used in the primers and probes of the pan-*Theileria* FRET-qPCR used in this study.

	Forward primer	LCRed-640	6-FAM	Reverse primer
	TACTGACAAGAAATAACAATACGGGGC--TT	GTCTTGTAAATGGGAATGATGGGAATT	AAACCTCTTCCAGAGATATCAATTGG	AGTTAAAGCTCGTAGTTGAATTTCTGCTG
<i>T. orientalis</i>				
<i>T. buffeli</i>				
<i>T. annulata</i>				
<i>T. sergenti</i>				
<i>T. luwenshuni</i>				
<i>T. velifera</i>				
<i>T. ovis</i>				
<i>T. parva</i>				
<i>T. uilenbergi</i>				
<i>T. lestoquardi</i>				
<i>T. equi</i>				
<i>T. separata</i>				
<i>T. capreoli</i>				
<i>T. bicornis</i>				
<i>T. taurotragi</i>				
<i>T. mutans</i>				
<i>Theileria</i> sp. OT3				
<i>Theileria</i> sp. NG				
<i>Theileria</i> sp. YW				
<i>Theileria</i> sp. B15				
<i>B. vulpes</i>				
<i>B. hongkongensis</i>				
<i>B. divergens</i>				
<i>B. bovis</i>				
<i>B. bigemina</i>				
<i>B. gibsoni</i>				
<i>B. microti</i>				
<i>B. felis</i>				
<i>B. canis</i>				
<i>C. felis</i>				
<i>H. americanum</i>				
<i>T. gondii</i>				

Primers and probes are shown at the head of the table. Dots indicate nucleotides identical to primers and probes, and dashes denote absence of the nucleotide. The upstream primer is used as the demonstrated sequences without gaps while the two probes and downstream primer are used as antisense oligonucleotides. The designed oligonucleotides show minimum mismatching with *Theileria* spp. The 6-FAM label is directly attached to the 3-terminal nucleotide of the fluorescent probe, and the LCRed-640 fluorescent label is added via a linker to the 5'-end of the LCRed-640 probe. The 18S rRNA sequences for the available recognized *Theileria* spp. on GenBank and other closely related protozoan species were obtained from GenBank: *T. orientalis* (HM538222), *T. buffeli* (HQ840967), *T. annulata* (KF429799), *T. sergenti* (EU083804), *T. luwenshuni* (JX469527), *T. velifera* (AF097993), *T. ovis* (AY508458), *T. parva* (L02366), *T. uilenbergi* (JF19835), *T. equi* (AY534882), *T. lestoquardi* (JQ917458), *T. separata* (AY260175), *T. capreoli* (AY726011), *T. bicornis* (AF499604), *T. taurotragi* (L19082), *T. mutans* (FJ213585), *Babesia vulpes* (JX454779), *Theileria* sp. OT3 (KF470868), *Theileria* sp. NG-2013a (KF597076), *Theileria* sp. YW-2014 (AB981984), *Theileria* sp. B15a (JN572700), *B. hongkongensis* (JQ867356), *B. divergens* (AJ439713), *B. bovis* (JQ723013), *B. bigemina* (JQ723014), *B. gibsoni* (JQ723014), *B. microti* (AB219802), *B. felis* (AF244912), *B. canis* (HM590440), *Hepatozoon americanum* (AF176836), *Cytauxzoon felis* (AY679105), and *Toxoplasma gondii* (L37415).

TABLE 2: Primers and probes used in this study.

PCR	Primer/probe	Nucleotides sequence	Amplicon
FRET-qPCR	Forward	5'-TAGTGACAAGAAATAACAATACGGGGCTT-3'	178 bp
	Reverse	5'-CAGCAGAGAAATTCAACTACGAGCTTTTAACT-3'	
	6-FAM	5'-CCAATTGATACTCTGGAGAGGGTTT-(6-FAM)-3'	
	LCRed-640	5'-(LCRed640)-AAT'TCCCATCATTCCTCAATTACAAGAC-Phosphate-3'	
Conventional PCR	Upstream	5'-CCTGAGAAACGGGCTACCACATCT-3'	<i>T. orientalis</i> 593 bp; <i>T. buffeli</i> 591 bp; <i>T. annulata</i> 591 bp; <i>T. sergenti</i> 591 bp; <i>T. luwenshumi</i> 594 bp; <i>T. velifera</i> 592 bp; <i>T. ovis</i> 595 bp; <i>T. parva</i> 592 bp; <i>T. uilenbergi</i> 592 bp; <i>T. equi</i> 596 bp; <i>T. cervi</i> 595 bp; <i>T. lestoquardi</i> 591 bp; <i>T. separata</i> 593 bp; <i>T. capreoli</i> 599 bp; <i>T. bicornis</i> 610 bp; <i>T. taurotraghi</i> 587 bp; <i>T. mutans</i> 585 bp; <i>B. vulpes</i> 609 bp; <i>Theileria</i> sp. OT3 600 bp; <i>Theileria</i> sp. NG 597 bp; <i>Theileria</i> sp. YW 593 bp; <i>Theileria</i> sp. BL5 584 bp
	Downstream	5'-GGACTAGGACGGGTATCTGATCG-3'	

TABLE 3: Prevalence of *Theileria* spp. in livestock from five Caribbean islands.

	Bovine	Goat	Sheep	Donkey	Horse	Total	<i>Theileria</i> spp.
Dominica	3/77 (3.9%)	0/70 (0.0%)	1/15 (6.7%)	N/A*	N/A	4/162 (2.5%)	<i>T. equi</i> , <i>Theileria</i> sp. NG-2013a, <i>B. vulpes</i> , or closely related organism
Grenada	N/A	2/31 (6.5%)	N/A	N/A	N/A	2/31 (6.5%)	<i>Theileria</i> sp. B15a
Montserrat	0/12 (0.0%)	8/19 (42.1%)	24/62 (38.7%)	N/A	N/A	32/93 (34.4%)	<i>Theileria</i> sp. OT3, <i>B. vulpes</i> , or closely related organism
Nevis	19/43 (44.2%)	2/114 (1.8%)	0/41 (0.0%)	N/A	N/A	21/198 (10.6%)	<i>T. equi</i> , <i>Theileria</i> sp. NG-2013a
St. Kitts	0/193 (0.0%)	0/4 (0.0%)	0/26 (0.0%)	5/25 (20.0%)	1/20 (5.0%)	6/268 (2.2%)	<i>T. equi</i> , <i>Theileria</i> sp. YW-2014
Total	22/325 (6.8%)	12/238 (5.0%)	25/144 (17.4%)	5/25 (20.0%)	1/20 (5.0%)	65/752 (8.6%)	
<i>Theileria</i> spp.	<i>T. equi</i> , <i>Theileria</i> sp. NG-2013a	<i>T. equi</i> , <i>Theileria</i> sp. OT3, <i>Theileria</i> sp. B15a, <i>B. vulpes</i> , or closely related organism	<i>T. equi</i> , <i>Theileria</i> sp. OT3, <i>B. vulpes</i> , or closely related organism	<i>T. equi</i> , <i>Theileria</i> sp. YW-2014	<i>T. equi</i>		

*No specimen was available.

TABLE 4: The *Theileria* spp. identified in livestock from five Caribbean islands and their similarity with reported organisms on GenBank.

<i>Theileria</i> spp.	Sequences identified in this study		Highly similar sequences in GenBank		Mismatch
	Number	Source	GenBank #	Source	
<i>T. equi</i>	14	8 cattle, 1 goat from Nevis; 1 cow from Dominica; 3 donkeys, 1 sheep from St. Kitts	KF559357	Horse from China	0/550
<i>Theileria</i> sp. OT3	11	8 sheep, 3 goats from Montserrat	KF470868	Sheep from China	0/555
<i>Theileria</i> sp. NG-2013a	7	6 cattle from Nevis; 1 cow from Dominica	KF597076	Waterbuck from Kenya	0/552
<i>Theileria</i> sp. YW-2014	1	1 donkey from St. Kitts	AB981984	Sika deer from Japan	0/549
<i>Theileria</i> sp. B15a	1	1 goat from Grenada	JN572700	Buffalo from South Africa	0/539
<i>B. vulpes</i> or closely related organism	9	4 sheep, 4 goats from Montserrat; 1 goat from Dominica	JX454779	Dog from France	0/178

Numerous species of *Amblyomma*, *Dermacentor*, *Hyalomma*, *Ixodes*, and *Rhipicephalus* are confirmed or suspected vectors of *T. equi* [29]. Of these, only *A. cajennense* [30], *R. microplus* [30], *R. sanguineus* [31], and *R. turanicus* (unpublished observation) occur in the Caribbean. There is conflicting data that *Amblyomma cajennense* is a competent vector of *T. equi* [32] but the tick is localized to Jamaica, Trinidad, and Cuba in the Caribbean [30] and it appears, then, not to be the vector of *T. equi* we found on Nevis, St. Kitts, and Dominica. *Dermacentor nitens*, the tropical horse tick, is very common in the Caribbean and the tropical Americas [33]. Although there is no data on the competence of *D. nitens* as a vector of *T. equi* and there is contradictory epidemiological evidence [32], PCR positive *D. nitens* have been found [32] and Asgarali et al. [16] have suggested that this tick is a vector in Trinidad. They also suggested that *R. microplus* [16], which is very common on cattle throughout the Caribbean, might also be a vector. While there is some

evidence that *R. microplus* is a competent vector [34] and our PCR identified *T. equi* in cattle on two islands, it seems unlikely that *R. microplus* is an important natural vector as it is a one host species and transovarial transmission has not been demonstrated [32]. Although *R. sanguineus* and *R. turanicus* have been implicated as vectors of *T. equi*, more recent studies have failed to confirm their role [29]. Further studies are needed to establish the epidemiology of *T. equi* and its vectors in the Caribbean and the neighboring Americas [32].

Theileria sp. OT3 was first described in sheep, deer, and chamois in Spain [35–37] and later in sheep in Italy [38], China [39], and Turkey [40]. Its pathogenicity and vectors have yet to be determined. Ours is the first report of the organism in the Caribbean and also the first report of the *Theileria* sp. OT3 in goats which further demonstrates that the organism has a wide distribution and host range. Although we used only convenience samples which were

FIGURE 2: Phylogenetic tree of sequences identified in this study and closely related *Theileria* spp. Sequences from GenBank have gene accession numbers in black font; those from this study are in red font. *B. vulpes* or closely related organism DM indicates the organism detected on Dominica and Montserrat; *Theileria* sp. Gre indicates the *Theileria* sp. from Grenada; *T. equi* DNS indicates the sequence of *T. equi* from Dominica, Nevis, and St. Kitts; *Theileria* sp. Mon indicates the *Theileria* sp. detected from Montserrat; and *Theileria* sp. SK is the sequence of *Theileria* sp. from St. Kitts.

not representative of the islands, it is of note that *Theileria* sp. OT3 was the most prevalent species we detected in ruminants. Similar high prevalences of infections have also been reported in the other countries where the organism has been described. The sequences of our *Theileria* sp. OT3 were identical to one another and to that of an isolate from China (KF470868) [39]. A phylogenetic relationship tree established for the Chinese isolate showed that the *Theileria* sp. OT3 forms a separated cluster and that the organism is closely related to *T. uilenbergi*, *T. luwenshuni*, and *T. ovis*. Although the organism has only been found in apparently healthy animals, studies on its pathogenicity seem indicated as, with its high prevalence and wide distribution, it might be causing substantial economic losses for livestock farmers.

Theileria mutans and *T. velifera* are African species that have been reported on Guadeloupe [12] and appear most likely to have been imported with cattle from West Africa in the 18th Century. This might also have been the case with the other African *Theileria* sp. we identified. Seven of the *Theileria* we found had an identical sequence to that of *Theileria* sp. NG-2013a (KF597076) described from waterbuck (*Kobus defassa*) in Kenya [41]. This organism has been found to cluster with *T. equi* but might represent a novel taxon. Its vectors and pathogenicity are unknown. One goat we studied had a *Theileria* spp. with a sequence identical to that of *Theileria* sp. B15a (JN572700) from an African buffalo (*Syncerus caffer*) in South Africa [42]. This organism is close to *T. mutans* which is widespread in Africa where it is transmitted by *Amblyomma* spp. and causes benign theileriosis. Previously *T. mutans* has been reported on Guadeloupe based on serological test results [12] and, since we found no confirmatory molecular evidence for the presence of *T. mutans*, it might be that the cross-reacting antibodies detected in Guadeloupe were against this closely related *Theileria* sp. B15.

The only non-*Theileria* sp., the pan-*Theileria* FRET-qPCR identified in our study, was one that appeared to be *Babesia vulpes* [43] or a closely related species. This organism was previously known as “*T. annae*” or the “*Babesia microti*-like organism” but was reclassified based on 18S rRNA and tubulin-beta gene sequence data [43]. The sequences of the short amplicons we obtained in the pan-*Theileria* FRET-qPCR were identical to those of *B. vulpes* (JX454779; KF773740) and had one mismatch with *B. microti* (AB219802, HQ629933, and LC005772). We were unable, however, to obtain longer amplicons with the standard PCR, probably because there were only very low copy numbers present in the positive animals. We were, then, unable to use this additional sequencing data to confirm that the organism we detected was *B. vulpes* or determine if it was a closely related species or strain. The pan-*Theileria* FRET-qPCR we used in our study was designed to detect seventeen recognized *Theileria* spp., which did not include “*T. annae*” or *B. vulpes* (Table 1) [21]. These seventeen recognized *Theileria* spp. differed from one another by only a maximum of 4 nucleotides in the regions of the primers and the probes used in the pan-*Theileria* FRET-qPCR. It is of note, then, that these primers and probes enabled the multiplication and detection of *B. vulpes*, or a closely related species, with 8 nucleotide differences and it therefore appears that the pan-*Theileria* FRET-qPCR might not be as genus specific as first thought. Further work is currently underway in our laboratory to more clearly characterize the *B. vulpes* or closely related organism found in the Caribbean.

4. Conclusions

Our study has confirmed the sensitivity of the pan-*Theileria* PCR in the rapid detection of a wide range of *Theileria* spp. but has also shown it might detect *B. vulpes* or closely

related organisms. We found livestock infected with *Theileria* spp. on each of the five islands we studied. While we could not confirm previous reports of *T. mutans* and *T. velifera* in cattle, we found that one recognized species, *T. equi*, four poorly characterized *Theileria* spp., and *B. vulpes* or a closely related organism are present in the region. Further studies are indicated to more precisely determine the phylogenetic relationships of these organisms in the Caribbean with closely related organisms from other parts of the world. Also, the prevalences of infections on the different islands should be determined as well as the impact these poorly characterized organisms might have on livestock production, both in the Caribbean and around the world where they are found.

Conflict of Interests

The authors declare that they have no competing interests.

Authors' Contribution

Chengming Wang, Patrick Kelly, and Jilei Zhang designed the study, analyzed the data, and wrote the paper. Jilei Zhang, Jing Li, and Chuanling Xu carried out the experiments. All authors read and approved the final paper.

Acknowledgments

This project was supported by grants from the National Natural Science Foundation of China (no. 31472225) and the Priority Academic Program Development of Jiangsu Higher Education Institutions, Yangzhou, Jiangsu, China and Grant 2006 34135 6930 from the United States Department of Agriculture through its program for Tropical and Subtropical Agricultural Research (T-STAR) and by Ross University School of Veterinary Medicine.

References

- [1] M. K. Shaw, L. G. Tilney, and A. J. Musoke, "The entry of *Theileria parva* sporozoites into bovine lymphocytes: evidence for MHC class I involvement," *The Journal of Cell Biology*, vol. 113, no. 1, pp. 87–101, 1991.
- [2] R. Bishop, A. Musoke, S. Morzaria, M. Gardner, and V. Nene, "*Theileria*: intracellular protozoan parasites of wild and domestic ruminants transmitted by ixodid ticks," *Parasitology*, vol. 129, supplement, pp. S271–S283, 2004.
- [3] M. Florin-Christensen and L. Schnittger, "Piroplasmids and ticks: a long-lasting intimate relationship," *Frontiers in Bioscience*, vol. 14, no. 8, pp. 3064–3073, 2009.
- [4] E. J. Glass, "The balance between protective immunity and pathogenesis in tropical theileriosis: what we need to know to design effective vaccines for the future," *Research in Veterinary Science*, vol. 70, no. 1, pp. 71–75, 2001.
- [5] J. S. Ahmed, E. J. Glass, D. A. Salih, and U. Seitzer, "Innate immunity to tropical theileriosis," *Innate Immunity*, vol. 14, no. 1, pp. 5–12, 2008.
- [6] T. Sivakumar, K. Hayashida, C. Sugimoto, and N. Yokoyama, "Evolution and genetic diversity of *Theileria*," *Infection, Genetics and Evolution*, vol. 27, pp. 250–263, 2014.
- [7] M. R. Rjeibi, M. A. Darghouth, M. Rekik, B. Amor, L. Sassi, and M. Gharbi, "First molecular identification and genetic characterization of *Theileria lestoquardi* in sheep of the Maghreb region," *Transboundary and Emerging Diseases*, 2014.
- [8] S. Bawm, K. Shimizu, J.-I. Hirota et al., "Molecular prevalence and genetic diversity of bovine *Theileria orientalis* in Myanmar," *Parasitology International*, vol. 63, no. 4, pp. 640–645, 2014.
- [9] L. P. Belotindos, J. V. Lazaro, M. A. Villanueva, and C. N. Mingala, "Molecular detection and characterization of *Theileria* species in the Philippines," *Acta Parasitologica*, vol. 59, no. 3, pp. 448–453, 2014.
- [10] M. H. Hussain, M. Saqib, F. Raza et al., "Seroprevalence of *Babesia caballi* and *Theileria equi* in five draught equine populated metropolises of Punjab, Pakistan," *Veterinary Parasitology*, vol. 202, no. 3–4, pp. 248–256, 2014.
- [11] Y. Li, Z. Chen, Z. Liu et al., "Molecular identification of *Theileria* parasites of northwestern Chinese Cervidae," *Parasites and Vectors*, vol. 7, article 225, 2014.
- [12] G. Uilenberg, E. Camus, and N. Barré, "Existence of *Theileria mutans* and *Theileria velifera* (Sporozoa, Theileriidae) in Guadeloupe (French West Indies)," *Revue d'Elevage et de Médecine Vétérinaire des Pays Tropicaux*, vol. 36, no. 3, pp. 261–264, 1983.
- [13] M. Alonso, E. Camus, J. Rodriguez Diego, L. Bertaudière, J. C. Tatareau, and J. M. Liabeuf, "Current status of bovine haemoparasitic diseases in Martinique (French West Indies)," *Revue d'Elevage et de Médecine Vétérinaire des Pays Tropicaux*, vol. 45, no. 1, pp. 9–14, 1992.
- [14] J. Rampersad, E. Cesar, M. D. Campbell, M. Samlal, and D. Ammons, "A field evaluation of PCR for the routine detection of *Babesia equi* in horses," *Veterinary Parasitology*, vol. 114, no. 2, pp. 81–87, 2003.
- [15] K. C. Georges, C. D. Ezeokoli, O. Sparagano et al., "A case of transplacental transmission of *Theileria equi* in a foal in Trinidad," *Veterinary Parasitology*, vol. 175, no. 3–4, pp. 363–366, 2011.
- [16] Z. Asgarali, D. K. Coombs, F. Mohammed, M. D. Campbell, and E. Caesar, "A serological study of *Babesia caballi* and *Theileria equi* in Thoroughbreds in Trinidad," *Veterinary Parasitology*, vol. 144, no. 1–2, pp. 167–171, 2007.
- [17] F. Katzer, S. McKellar, E. Kirvar, and B. Shiels, "Phylogenetic analysis of *Theileria* and *Babesia equi* in relation to the establishment of parasite populations within novel host species and the development of diagnostic tests," *Molecular and Biochemical Parasitology*, vol. 95, no. 1, pp. 33–44, 1998.
- [18] J. M. Gubbels, A. P. de Vos, M. van der Weide et al., "Simultaneous detection of bovine *Theileria* and *Babesia* species by reverse line blot hybridization," *Journal of Clinical Microbiology*, vol. 37, no. 6, pp. 1782–1789, 1999.
- [19] L. Schnittger, H. Yin, B. Qi et al., "Simultaneous detection and differentiation of *Theileria* and *Babesia* parasites infecting small ruminants by reverse line blotting," *Parasitology Research*, vol. 92, no. 3, pp. 189–196, 2004.
- [20] B. J. Mans, R. Pienaar, and A. A. Latif, "A review of *Theileria* diagnostics and epidemiology," *International Journal for Parasitology: Parasites and Wildlife*, vol. 4, no. 1, pp. 104–118, 2015.
- [21] Y. Yang, Y. Mao, P. Kelly et al., "A pan-*Theileria* FRET-qPCR survey for *Theileria* spp. in ruminants from nine provinces of China," *Parasites and Vectors*, vol. 7, article 413, 2014.
- [22] P. Kelly, H. Lucas, L. Beati, C. Yowell, S. Mahan, and J. Dame, "*Rickettsia africae* in *Amblyomma variegatum* and domestic ruminants on eight Caribbean islands," *Journal of Parasitology*, vol. 96, no. 6, pp. 1086–1088, 2010.

- [23] L. Wei, P. Kelly, J. Zhang et al., "Use of a universal hydroxymethylbilane synthase (HMBS)-based PCR as an endogenous internal control and to enable typing of mammalian DNAs," *Applied Microbiology and Biotechnology*, vol. 98, no. 12, pp. 5579–5587, 2014.
- [24] E. Hawkins, R. Kock, D. McKeever et al., "Prevalence of *Theileria equi* and *Babesia caballi* as well as the identification of associated ticks in sympatric grevy's zebras (*Equus grevyi*) and donkeys (*Equus africanus asinus*) in northern Kenya," *Journal of Wildlife Diseases*, vol. 51, no. 1, pp. 137–147, 2015.
- [25] L. N. Wise, L. S. Kappmeyer, R. H. Mealey, and D. P. Knowles, "Review of equine piroplasmosis," *Journal of Veterinary Internal Medicine*, vol. 27, no. 6, pp. 1334–1346, 2013.
- [26] A. Criado-Fornelio, A. Martinez-Marcos, A. Buling-Saraña, and J. C. Barba-Carretero, "Molecular studies on *Babesia*, *Theileria* and *Hepatozoon* in southern Europe. Part II. Phylogenetic analysis and evolutionary history," *Veterinary Parasitology*, vol. 114, no. 3, pp. 173–194, 2003.
- [27] C. T. Rosa, P. Pazzi, S. Nagel et al., "Theileriosis in six dogs in South Africa and its potential clinical significance," *Journal of the South African Veterinary Association*, vol. 85, no. 1, article 1114, 2014.
- [28] M. Adamu, M. Troskie, D. O. Oshadu, D. P. Malatji, B. L. Penzhorn, and P. T. Matjila, "Occurrence of tick-transmitted pathogens in dogs in Jos, Plateau State, Nigeria," *Parasites and Vectors*, vol. 7, no. 1, article 119, 2014.
- [29] G. A. Scoles and M. W. Ueti, "Vector ecology of equine piroplasmosis," *Annual Review of Entomology*, vol. 60, pp. 561–580, 2015.
- [30] E. Camus and N. Barre, "Vector situation of tick-borne diseases in the Caribbean islands," *Veterinary Parasitology*, vol. 57, no. 1–3, pp. 167–176, 1995.
- [31] P. J. Kelly, C. Xu, H. Lucas et al., "Ehrlichiosis, babesiosis, anaplasmosis and hepatozoonosis in dogs from St. Kitts, West Indies," *PLoS ONE*, vol. 8, no. 1, Article ID e53450, 2013.
- [32] M. Peckle, M. S. Pires, T. M. Dos Santos et al., "Molecular epidemiology of *Theileria equi* in horses and their association with possible tick vectors in the state of Rio de Janeiro, Brazil," *Parasitology Research*, vol. 112, no. 5, pp. 2017–2025, 2013.
- [33] G. I. Garriss and K. Scotland, "Ticks on livestock in St. Lucia," *Veterinary Parasitology*, vol. 18, no. 4, pp. 367–373, 1985.
- [34] D. Stiller, W. L. Goff, L. W. Johnson, and D. P. Knowles, "*Dermacentor variabilis* and *Boophilus microplus* (Acari: Ixodidae): experimental vectors of *Babesia equi* to equids," *Journal of Medical Entomology*, vol. 39, no. 4, pp. 667–670, 2002.
- [35] D. Nagore, J. García-Sanmartín, A. L. García-Pérez, R. A. Juste, and A. Hurtado, "Identification, genetic diversity and prevalence of *Theileria* and *Babesia* species in a sheep population from Northern Spain," *International Journal for Parasitology*, vol. 34, no. 9, pp. 1059–1067, 2004.
- [36] J. García-Sanmartín, O. Aurtinetxe, M. Barral et al., "Molecular detection and characterization of piroplasms infecting cervids and chamois in Northern Spain," *Parasitology*, vol. 134, Part 3, pp. 391–398, 2007.
- [37] A. Ros-García, J. F. Barandika, A. L. García-Pérez, R. A. Juste, and A. Hurtado, "Assessment of exposure to piroplasms in sheep grazing in communal mountain pastures by using a multiplex DNA bead-based suspension array," *Parasites and Vectors*, vol. 6, no. 1, article no. 277, 2013.
- [38] A. Giangaspero, M. Marangi, R. Papini, B. Paoletti, M. Wijnveld, and F. Jongejans, "*Theileria* sp. OT3 and other tick-borne pathogens in sheep and ticks in Italy: molecular characterization and phylogeny," *Ticks and Tick-borne Diseases*, vol. 6, no. 1, pp. 75–83, 2015.
- [39] Z. Tian, G. Liu, H. Yin et al., "First report on the occurrence of *Theileria* sp. OT3 in China," *Parasitology International*, vol. 63, no. 2, pp. 403–407, 2014.
- [40] M. F. Aydin, M. Aktas, and N. Dumanli, "Molecular identification of *Theileria* and *Babesia* in sheep and goats in the Black Sea Region in Turkey," *Parasitology Research*, vol. 112, no. 8, pp. 2817–2824, 2013.
- [41] N. Githaka, S. Konnai, R. Bishop et al., "Identification and sequence characterization of novel *Theileria* genotypes from the waterbuck (*Kobus defassa*) in a *Theileria parva*-endemic area in Kenya," *Veterinary Parasitology*, vol. 202, no. 3–4, pp. 180–193, 2014.
- [42] M. E. Chaisi, N. E. Collins, F. T. Potgieter, and M. C. Oosthuizen, "Sequence variation identified in the 18S rRNA gene of *Theileria mutans* and *Theileria velifera* from the African buffalo (*Syncerus caffer*)," *Veterinary Parasitology*, vol. 191, no. 1–2, pp. 132–137, 2013.
- [43] G. Baneth, M. Florin-Christensen, L. Cardoso, and L. Schnitger, "Reclassification of *Theileria annae* as *Babesia vulpes* sp. nov.," *Parasites & Vectors*, vol. 8, no. 1, article 207, 2015.