Clinical Study

The Impact of Vitamin D₃ Supplementation on Mechanisms of Cell Calcium Signaling in Chronic Kidney Disease

Ingrid Lajdova,¹ Viera Spustova,¹ Adrian Oksa,¹ Zuzana Kaderjakova,² Dusan Chorvat Jr.,³ Marcela Morvova Jr.,² Libusa Sikurova,² and Alzbeta Marcek Chorvatova³

¹Department of Clinical and Experimental Pharmacology, Faculty of Medicine, Slovak Medical University, 833 03 Bratislava, Slovakia

²Department of Nuclear Physics and Biophysics, Faculty of Mathematics, Physics and Informatics, Comenius University, 833 03 Bratislava, Slovakia

³Department of Biophotonics, International Laser Centre, 833 03 Bratislava, Slovakia

Correspondence should be addressed to Ingrid Lajdova; ingrid.lajdova@szu.sk

Received 2 October 2014; Accepted 19 December 2014

Academic Editor: Luca De Nicola

Copyright © 2015 Ingrid Lajdova et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Intracellular calcium concentration in peripheral blood mononuclear cells (PBMCs) of patients with chronic kidney disease (CKD) is significantly increased, and the regulatory mechanisms maintaining cellular calcium homeostasis are impaired. The purpose of this study was to examine the effect of vitamin D₃ on predominant regulatory mechanisms of cell calcium homeostasis. The study involved 16 CKD stages 2-3 patients with vitamin D deficiency treated with cholecalciferol 7000–14000 IU/week for 6 months. The regulatory mechanisms of calcium signaling were studied in PBMCs and red blood cells. After vitamin D₃ supplementation, serum concentration of 25(OH)D₃ increased ($P < 0.001$) and $[Ca^{2+}]_i$ decreased ($P < 0.001$). The differences in $[Ca^{2+}]_i$ were inversely related to differences in 25(OH)D₃ concentration ($P < 0.01$). Vitamin D₃ supplementation decreased the calcium entry through calcium release activated calcium (CRAC) channels and purinergic P2X₇ channels. The function of P2X₇ receptors was changed in comparison with their baseline status, and the expression of these receptors was reduced. There was no effect of vitamin D₃ on P2X₇ pores and activity of plasma membrane Ca^{2+} -ATPases. Vitamin D₃ supplementation had a beneficial effect on $[Ca^{2+}]_i$ decreasing calcium entry via CRAC and P2X₇ channels and reducing P2X₇ receptors expression.

1. Introduction

Vitamin D hormonal system has been classically implicated in the regulation of calcium homeostasis and bone metabolism. However, it has also noncalcitropic effects through the activation of tissue vitamin D receptors (VDR) [1]. Vitamin D insufficiency/deficiency is a significant risk factor for the development of various chronic diseases, and the deficiency of calcidiol (25(OH)D₃) as well as calcitriol (1,25(OH)₂D₃) is common in CKD patients [2]. Therefore, the supplementation of native vitamin D (cholecalciferol or ergocalciferol) or active vitamin D (calcitriol and VDR activators) in CKD is well established.

Free cytosolic calcium concentration ($[Ca^{2+}]_i$) is controlled by mechanisms that regulate Ca^{2+} entry from

the extracellular space and Ca^{2+} release from intracellular stores and by the activity of ATP-dependent Ca^{2+} pumps and antiporters that move Ca^{2+} back into stores or out of cells [3]. Already in early stages of chronic kidney disease (CKD), $[Ca^{2+}]_i$ and calcium concentration of intracellular stores were significantly increased in comparison with healthy volunteers, and the regulatory mechanisms of calcium signaling were impaired by the disease [4–7].

Calcium enters into the cells by any of the general classes of calcium/cation channels. In nonexcitable cells like peripheral blood mononuclear cells (PBMCs), the predominant Ca^{2+} entry pathway is the store-operated one, in which the emptying of intracellular Ca^{2+} stores activates the Ca^{2+} influx. This type of the channel is known as the calcium release activated calcium (CRAC) channel in lymphocytes.

The calcium entry through CRAC channels activates certain transcription factors which regulate the gene expression for cytokines responsible for immune responses [8, 9].

Another mechanism of calcium entry into the cell is represented by purinergic P2X receptors. At the present time, purinergic signaling is accepted as a crucial component of diseases and was found to mediate a vast array of biological processes. The P2X₇ receptors are expressed primarily on cells of hemopoietic origin, where they participate in immune responses, cell proliferation, cell death, bone formation, and bone resorption [10]. The P2X₇ receptor is a bifunctional purinoreceptor that opens a nonselective cation channel and consecutively forms a large, cytolytic pore. The key factor of P2X₇-dependent cytotoxicity is the massive intracellular Ca²⁺ increase triggered by its activation. This can lead to membrane blebbing and cell death by apoptosis or necrosis. There is an increasing body of evidence implicating P2X₇ receptors in various pathological conditions [11–14].

The plasma membrane Ca²⁺-ATPases (PMCA) is responsible for removing excessive Ca²⁺ out of the cells to extracellular environment. The decreased PMCA activity increases [Ca²⁺]_i and affects some intracellular processes.

To our knowledge, little information is available regarding the impact of vitamin D₃ supplementation on disturbed cell calcium homeostasis in CKD. Therefore, the aim of the present study was to examine the effect of vitamin D₃ supplementation on essential regulatory mechanisms of cell calcium homeostasis.

2. Materials and Methods

2.1. Patients. The study population consisted of 16 nondiabetic patients with CKD (9 patients CKD stage 2 and 7 patients CKD stage 3). All of them were screened and followed up in the outpatient department of nephrology at the Slovak Medical University. The diagnosis of CKD was based on clinical and laboratory examinations as defined by the K/DOQI criteria [15]. Causes of their renal disease were glomerulonephritis in 9 patients, tubulointerstitial nephritis in 3 cases, hypertensive nephroangiosclerosis in 2 patients, and other in 2 causes. The glomerular filtration rate was estimated by the MDRD study formula [16]. Patients with acute impairment of renal function, nephrotic proteinuria, malignancies, and derangements in mineral metabolism of nonrenal origin were excluded from the study. Concurrent treatments interfering with mineral metabolism were not allowed. Previous therapy with vitamin D₂/D₃, calcitriol, or over-the-counter vitamin D preparations had to be cancelled at least 2 months before enrollment. Hypertension was the most common comorbidity present in all patients and treated with ACE inhibitors or angiotensin II receptor blockers in 14, diuretics in 8, betablockers in 6, and calcium channel blockers in 8 cases. Dihydropyridine calcium channel blockers were allowed as they do not interfere with studied parameters and effects in PBMCs [17]. All patients had vitamin D deficiency (serum 25(OH)D₃ concentration <30 ng/mL) and were supplemented with cholecalciferol 7000–14000 IU/week for 6 months; the dose (approximately 1000–2000 IU/day)

was chosen as a common supplementary dose for the treatment of vitamin D deficiency in general population.

The study was approved by the Ethics Committee of the Slovak Medical University and all participants gave their written informed consent.

2.2. PBMCs Isolation. Human PBMCs were isolated by the Ficoll gradient centrifugation, diluted 1:1 with RPMI-1640 medium, layered onto an equivalent volume of medium LSM-1077, and centrifuged at 700 g for 20 min at 22°C, as previously reported [5]. The PBMC layer was washed in 40 mL RPMI, resuspended in 10 mL RPMI and 10% fetal bovine serum (FBS), and centrifuged at 300 g 10 min at 22°C and the pellet was resuspended in 2 mL aliquots of physiological salt solution. Final concentration of PBMCs was adjusted to 2.5 × 10⁶ cells/mL. Our preparation contained lymphocytes (94–96%), monocytes (3–4%), and natural killer cells (the rest), as determined by flow cytometry (Coulter Epics XL, Ireland). The cell viability was quantified using a 0.8% solution of trypan blue and estimated to be 96–98%.

2.3. Red Blood Cell Membranes Isolation. Isolated red blood cell (RBC) membranes were used to assess the PMCA function. RBC membranes were obtained by hemolytic fragmentation in hypotonic media using standard method of Hanahan and Ekholm [18] modified in our laboratory to achieve a higher quality of the ghosts. RBCs from previous isolation were diluted 1:5 with physiological salt solution and centrifuged at 1270 g for 20 min at 4°C. The supernatant was removed and the procedure was repeated one more time. RBCs were then diluted 1:5 with tris(hydroxymethyl)amino-methane (TRIS) medium (20 mmol/L, pH = 7.4) and centrifuged at 7700 g for 35 min at 4°C. This step was repeated twice for each TRIS medium with decreasing concentrations (20, 10, and 5 mmol/L).

2.4. Intracellular Ca²⁺ Measurements. The population of 2 × 10⁶ PBMCs/mL was loaded with fluorescence dye Fluo-3 AM at a final concentration of 2 μmol/L for 40 min at 22°C in a physiological salt solution. After incubation, the cells were centrifuged at 300 g, washed three times with a physiological salt solution, and kept at room temperature for 10 min before use. The Fluo-3 fluorescence was measured at 37°C in Fluorolog 3–11 spectrofluorometer (HORIBA Jobin Yvon Inc., Edison, NJ, USA) with an excitation at 488 nm (bandpass 3 nm) and an emission at 526 nm (bandpass 5 nm). Each experiment was followed by [Ca²⁺]_i calibration to estimate the actual free cytoplasmic calcium concentration from the measured fluorescence signal (*F*) in each cell population. [Ca²⁺]_i was quantified in nmol/L according to the following equation:

$$[\text{Ca}^{2+}]_i = K_d \cdot \left[\frac{F - F_{\min}}{F_{\max} - F} \right], \quad (1)$$

where *K_d* = 400 nmol/L at 37°C [19]. The maximal fluorescence intensity (*F_{max}*) was assessed by the addition of Triton X-100 (0.1%) with Ca²⁺ (5 mmol/L), and the minimum

fluorescence level ($F_{\min}$) was determined after the addition of 25 mmol/L EGTA (pH = 9). Digitonin (20 μ mol/L) was used to answer for minimal compartmentalization [20]. To assess the role of CRAC channels in Ca^{2+} entry, 2-aminoethyl-diphenyl borate (2APB), a widely used inhibitor of these channels was applied [21]. Although there are already more potent and selective inhibitors of these channels [22], 2APB was used for the possibility of comparing the results with our previous studies [4, 6]. The action of 2APB (50 μ mol/L) was studied in cells where Ca^{2+} entry through these channels was stimulated by thapsigargin (Tg) (1 μ mol/L), a specific inhibitor of endoplasmic reticulum Ca^{2+} -ATPase. To examine the function of P2X_7 receptors, AZ11645373 (50 nmol/L), a highly selective antagonist of human P2X_7 receptors [23], and 2',3'-O-(4-benzoyl)benzoyl ATP (BzATP) (50 μ mol/L), the most potent and selective agonist of these receptors, were used [24].

2.5. Cell Surface P2X_7 Receptors Expression. PBMCs were stained with P2X_7 polyclonal antibody labelled with fluorescein isothiocyanate (FITC) according to the protocol provided by the manufacturer. Briefly, 100 μ L of PBMCs were incubated either with FITC-conjugated anti- P2X_7 (2 μ g/mL) or FITC-conjugated IgG2a (2 μ g/mL, control) for 20 min at 22°C in the dark. PBMCs were simultaneously stained with phycoerythrin- (PE-) conjugated CD14 antibody in order to exclude monocytes from the examined population. After the incubation, cells were washed three times, dissolved in 500 μ L of PBS, and subjected to the analysis on flow cytometer (Cytomics FC 500 cytometer, Beckman Coulter, USA). Values of control samples stained with FITC-conjugated IgG2a were subtracted from the evaluated results.

2.6. P2X_7 Receptors Visualization by Fluorescence Microscopy. To visualize the surface P2X_7 receptors, PBMCs were stained with P2X_7 (extracellular) antibody. Cell imaging was performed by using the Axiovert 200 inverted microscope (Carl Zeiss, Germany) with mercury lamp HBO-100 and FluoArc driver. The 450–490 nm bandpass filter and FT510 dichroic mirror were used for excitation, and emission was detected with a 515 nm long-pass filter. Images were recorded by the PentaMax cooled CCD camera (Roper Scientific) with KODAK KAF-1400 chip. Images were taken with the C-Apochromat 40x/1.2 water immersion objective (Zeiss, Germany).

2.7. Ethidium Bromide Uptake by Flow Cytometry. The ethidium bromide uptake by PBMCs was measured on the Cytomics FC 500 cytometer (Beckman Coulter, USA) and results were processed with the CXP software. PBMCs at concentration of 10^6 cells/100 μ L were kept at room temperature when ethidium bromide at 30 μ mol/L final concentration was added to the sample. After a 5 min incubation, the mean channel ethidium fluorescence was assessed in FL3 sensor after excitation with a 488 nm laser beam. Up to 10000 events were involved in the data analysis. For the investigation of P2X_7 pore function, ethidium fluorescence was measured

after a 10 min incubation with either BzATP (50 μ mol/L) or AZ11645373 (50 nmol/L) at 37°C.

2.8. PMCA Activity Measurement. The RBCs membrane suspension was added to working medium (mmol/L): 100 TRIS, 80 KCl, 3 MgCl_2 , 0.2 ethylenediaminetetraacetic acid (EDTA), and 1 ouabain (pH = 7.4) in the presence or absence of CaCl_2 (5 mmol/L). The reaction was started by the addition of 40 mmol/L of ATP, conducted for 60 min at 37°C and stopped by the addition of a 15% trichloroacetic acid (TCA). The amount of liberated inorganic phosphate was determined using a phosphate colorimetric assay kit (BioVision) on the UV-VIS spectrophotometer Shimadzu UV-1700 (Shimadzu Corp., Japan). The estimated PMCA activity was calculated as the difference between the activities of the enzyme incubated in the presence and absence of CaCl_2 and was expressed as nmol of Pi/mg protein/h. The protein concentration was measured by the method of Lowry [25].

2.9. Reagents. The physiological salt solution contained (mmol/L) 140 NaCl, 5.4 KCl, 1 CaCl_2 , 1 Na_2HPO_4 , 0.5 MgCl_2 , 5 glucose, and 5 HEPES (pH = 7.4). Thapsigargin (Tg) was procured from Calbiochem (San Diego, CA). 2-Aminoethyl-diphenyl borate (2APB), ethidium bromide, AZ11645373, and 2',3'-O-(4-benzoyl) benzoyl ATP (BzATP) were obtained from Sigma-Aldrich (St. Louis, MO, USA). Fluo-3 acetoxymethylester (Fluo-3 AM) was from Molecular Probes (Eugene, OR), lymphocyte separation medium LSM-1077 was from PAA Laboratories GmbH (Pasching, Austria), and fetal bovine serum (FBS) and RPMI-1640 medium were from GIBCO (Grand Island, NY, USA). PE-conjugated anti-CD14, FITC-conjugated IgG2a, and anti- P2X_7 (extracellular) FITC were from Sigma-Aldrich (St. Louis, MO, USA). The phosphate colorimetric assay kit was from BioVision (Hayward, CA, USA). All other chemicals were purchased from Sigma-Aldrich.

2.10. Analytical Procedures. Serum calcium and creatinine were measured by the Vitros 250 Analyzer, Johnson & Johnson, Rochester, NY, USA. Intact parathormone and 25-hydroxyvitamin D_3 were determined by the electrochemoluminescence immunoassays (ECLIA) (Roche-Diagnostics, Mannheim, Germany).

2.11. Statistical Analyses. All values are expressed as means $\pm$ SD. Statistical analysis was carried out by the SPSS 15.0 (SPSS Inc., Chicago, IL, USA). The Shapiro-Wilk test was used to evaluate a sample normality distribution. The statistical significance of differences was tested by the independent 2-population Student's *t*-test for normally distributed data and the Wilcoxon's test for a nonparametric analysis. A *P* value < 0.05 was considered significant. The Spearman's correlations between variables were used as a measure of association.

3. Results

3.1. Effect of Cholecalciferol Treatment on Main Laboratory Variables. Effect of cholecalciferol treatment was evaluated

TABLE 1: Main laboratory variables at baseline and after the cholecalciferol treatment.

Parameters	Baseline	6 months
Total serum calcium (mmol/L)	2.28 ± 0.08	2.34 ± 0.08
[Ca ²⁺] _i (nmol/L)	120 ± 6	105 ± 3***
iPTH (pg/mL)	43 ± 17	43 ± 20
25(OH)D ₃ (ng/mL)	18 ± 2	36 ± 9***
Serum creatinine (μmol/L)	102 ± 21	101 ± 28
eGFR (mL/min)	64.8 ± 5.4	63 ± 4.2

Values are expressed as mean ± SD, *** $P < 0.001$, $n = 16$ for comparison with baseline.

FIGURE 1: Relationships between absolute changes from baseline in [Ca²⁺]_i and 25(OH)D₃ concentrations after a 6-month vitamin D₃ supplementation ($P < 0.01$, $n = 16$).

in 16 CKD patients (mean age 55 ± 4 , range 32–80 years, 9 males and 7 females). Table 1 shows laboratory data of CKD patients at baseline and after the vitamin D₃ supplementation. Initial 25(OH)D₃ concentrations were low (18 ± 8 ng/mL) and significantly increased after the vitamin D₃ treatment (36 ± 9 ng/mL). Eleven patients reached the recommended 25(OH)D₃ level above 30 ng/mL. The [Ca²⁺]_i in PBMCs was increased in CKD patients (120 ± 6 nmol/L). After the 6-month vitamin D₃ supplementation, [Ca²⁺]_i significantly decreased (105 ± 3 nmol/L, $P < 0.001$, $n = 16$) to values comparable with those in healthy subjects. The absolute changes from baseline in [Ca²⁺]_i were related to the absolute changes in 25(OH)D₃ concentrations ($P < 0.01$, $n = 16$) (Figure 1). The PTH concentrations were in reference range (43 ± 17 pg/mL) at basal conditions and vitamin D₃ supplementation had no effect on their values. The correlation between [Ca²⁺]_i and PTH concentrations has not been found.

3.2. CRAC Channels. CRAC channels were activated indirectly by intracellular Ca²⁺ store depletion using Tg ($1 \mu\text{mol/L}$). The 2APB ($50 \mu\text{mol/L}$), a reliable inhibitor of these channels, was applied during the sustained phase of

Tg effect. It evoked a decrease in [Ca²⁺]_i, which represented particularly the Ca²⁺ influx through CRAC channels. After the treatment with vitamin D₃, the enhanced Ca²⁺ entry through these types of channels (68 ± 27 nmol/L) was significantly decreased (43 ± 16 nmol/L; $P < 0.01$, $n = 16$) in CKD patients (Figures 2(a) and 7(b)).

3.3. P2X₇ Receptors

3.3.1. P2X₇ Channels. The application of P2X₇ receptors antagonist AZ11645373 (50 nmol/L) led to reduction in [Ca²⁺]_i in CKD patients from 120 ± 6 to 112 ± 8 nmol/L ($P < 0.001$, $n = 16$). On the other hand, after the vitamin D₃ supplementation, AZ11645373 had no effect on [Ca²⁺]_i (106 ± 4 versus 104 ± 6 nmol/L; ns, $n = 16$) (Figure 3(a)). Differences in [Ca²⁺]_i after the inhibition of P2X₇ receptors were significantly decreased after the vitamin D₃ supplementation ($P < 0.001$, $n = 16$) (Figure 3(b)). In BzATP stimulated cells, AZ11645373 (50 nmol/L) decreased the calcium influx before and also after the vitamin D₃ supplementation (Figures 3(c) and 3(d)), but the effect of an inhibitor was attenuated ($P < 0.01$, $n = 16$) (Figure 3(e)). The 6-month vitamin D₃ supplementation had an inhibitory effect on function of P2X₇ channels and thereby decreased the Ca²⁺ entry. The agonist of purinergic P2X₇ receptors, BzATP ($50 \mu\text{mol/L}$), caused a sustained increase in [Ca²⁺]_i in CKD patients at baseline 120 ± 6 to 155 ± 12 nmol/L ($P < 0.001$, $n = 16$) and also after the vitamin D₃ supplementation 106 ± 4 to 136 ± 13 nmol/L ($P < 0.001$, $n = 16$). However, the P2X₇ receptors activation after the vitamin D₃ supplementation did not reach the values of [Ca²⁺]_i before supplementation (Figure 4(a)). Furthermore, the effect of BzATP ($50 \mu\text{mol/L}$) on AZ11645373-inhibited calcium influx in PBMCs was evaluated. Under these conditions, a rising calcium influx through P2X₇ channels was found at baseline and after supplementation (Figures 4(b) and 4(c)). All these results demonstrate the inhibitory effect of vitamin D₃ supplementation on calcium entry through P2X₇ channels.

3.3.2. P2X₇ Pores. The uptake of ethidium bromide into PBMCs was measured by flow cytometry at basal conditions and with BzATP ($50 \mu\text{mol/L}$) stimulation or AZ11645373 (50 nmol/L) inhibition. The permeability of ethidium bromide through P2X₇ pores in PBMCs of CKD patients was significantly increased in comparison with healthy volunteers [4, 5] and remained unchanged after the vitamin D₃ supplementation. The treatment did not change the permeability of P2X₇ pores after the application of either BzATP ($50 \mu\text{mol/L}$) or AZ11645373 (50 nmol/L) (Figures 5(a) and 5(b)).

3.3.3. Expression of Cell Surface P2X₇ Receptors. The expression of cell surface P2X₇ receptors was 1.5-fold greater on PBMCs from CKD patients compared to healthy donors [4]. We assessed a decreased expression of these receptors after vitamin D₃ supplementation in the whole population of PBMCs ($P < 0.001$) (Figures 6(a), 6(b), and 6(c)).

FIGURE 2: The calcium entry through CRAC channels. (a) A typical experiment where CRAC channels were activated by intracellular Ca^{2+} depletion by Tg ($1 \mu\text{mol/L}$) and subsequently inhibited by 2APB ($50 \mu\text{mol/L}$). The representative trace was obtained from a single experiment from PBMCs of a CKD patient. (b) The comparison of the calcium entry through CRAC channels in PBMCs of CKD patients at baseline and after a 6-month vitamin D_3 supplementation (** $P < 0.01$, $n = 16$).

3.4. Plasma Membrane Ca^{2+} -ATPases. The PMCA activity of RBCs membranes is decreased by 25% in patients with early stages of CKD when compared to healthy subjects [4]. Vitamin D_3 supplementation did not increase the PMCA activity (Figures 7(a) and 7(b)). The concentrations of total plasma membrane proteins were significantly enhanced (7.7 ± 1.6 versus $11.8 \pm 2.7 \text{ mg/mL}$, $P < 0.001$) which may indicate increased expression of PMCA.

4. Discussion

Disturbances in intracellular calcium homeostasis in patients with CKD represent a complex process which aggravates with CKD progression. The mechanisms of cell calcium influx and efflux are impaired in renal disease. Already in early stages of CKD, cytosolic calcium concentration ($[\text{Ca}^{2+}]_i$) and calcium concentration of intracellular stores are increased [6, 7]. The elevated calcium entry through CRAC channels and P2X_7 receptors, increased expression of P2X_7 receptors, and decreased PMCA activity contribute to this state [4]. We have previously shown that vitamin D_3 supplementation in CKD patients led to a decline in $[\text{Ca}^{2+}]_i$ to values comparable with healthy people [6]. The aim of this study was to examine the effect of vitamin D_3 supplementation on predominant regulation mechanisms of cell calcium homeostasis in nonexcitable cells from patients with early stages of CKD. The principal finding of the present study is that vitamin D_3 supplementation affected the mechanisms of intracellular calcium homeostasis as follows: the Ca^{2+} entry through CRAC and P2X_7 channels was decreased, while no effect was found on the permeability and functionality of P2X_7 pores, and the expression of P2X_7 receptors was reduced. Finally, the activity of PMCA was not increased after treatment.

4.1. Vitamin D_3 Supplementation. All clinical studies of vitamin D supplementation in CKD patients reported

a significant improvement in $25(\text{OH})\text{D}_3$ concentrations, although several studies did not reach a mean $25(\text{OH})\text{D}_3$ concentration in optimal range ($\geq 30 \text{ ng/mL}$). In our study, vitamin D_3 supplementation with weekly cholecalciferol dosing significantly increased $25(\text{OH})\text{D}_3$ to the recommended levels. PTH values were in normal range and did not change after treatment, and no correlation was observed between $25(\text{OH})\text{D}_3$ and PTH concentrations. Total serum calcium was not changed but $[\text{Ca}^{2+}]_i$ significantly decreased after supplementation. Linear regression analysis demonstrated that changes in $25(\text{OH})\text{D}_3$ were significantly and inversely correlated with that of $[\text{Ca}^{2+}]_i$ ($R = 0.617$, $P < 0.01$). Besides our previous work [6], no studies have investigated the relationship between vitamin D status and $[\text{Ca}^{2+}]_i$ in CKD.

4.2. Ca^{2+} Entry through CRAC Channels. Dysregulation of Ca^{2+} homeostasis involving the endoplasmic reticulum (ER) and store-operated calcium channels has been manifested in patients with neurodegenerative disorders, immunodeficiency, acute pancreatitis, polycystic kidney disease, and cardiac hypertrophy [9, 26–29]. In nonexcitable cells, CRAC channels are the main pathway of Ca^{2+} entry. These channels are composed of two proteins Orail and STIM1 (stromal interaction molecule). Orail protein is located in the plasma membrane and forms the channel pore. It is activated by STIM1 located in the membrane of ER. STIM1 has a dual function of sensing the Ca^{2+} concentration in ER and activating CRAC channels. A decrease in the ER Ca^{2+} concentration induces STIM1 translocation close to the plasma membrane where it binds to and activates the Orail channel. Alterations in STIM1/Orail system may contribute to several pathophysiological conditions including cardiovascular [30, 31] and pulmonary diseases [32], hypertension [33], immunodeficiency, and autoimmune and lymphoproliferative diseases

FIGURE 3: The effect of vitamin D₃ supplementation on an inhibitory effect of AZ11645373. (a) The comparison of the AZ11645373 (50 nmol/L) effect on $[Ca^{2+}]_i$ in CKD patients at baseline and after vitamin D₃ supplementation (** $P < 0.01$, *** $P < 0.001$, $n = 16$). After vitamin D₃ supplementation, AZ11645373 (50 nmol/L) had no effect on $[Ca^{2+}]_i$. (b) Differences in $[Ca^{2+}]_i$ after an inhibition with AZ11645373 (50 nmol/L) at baseline and after vitamin D₃ supplementation (*** $P < 0.001$, $n = 16$). (c) The representative trace where AZ11645373 (50 nmol/L) inhibited the $[Ca^{2+}]_i$ rise induced by BzATP (50 μ mol/L). (d) The comparison of the AZ11645373 (50 nmol/L) effect on BzATP- (50 μ mol/L) activated calcium influx before and after supplementation (*** $P < 0.001$, $n = 16$). (e) Differences in $[Ca^{2+}]_i$ in BzATP- (50 μ mol/L) activated PBMCs after the AZ11645373 (50 nmol/L) inhibition at baseline and after vitamin D₃ supplementation (** $P < 0.01$, $n = 16$).

FIGURE 4: The effect of vitamin D₃ supplementation on activation by BzATP. (a) The comparison of the BzATP (50 μ mol/L) effect on [Ca²⁺]_i in CKD patients at baseline and after vitamin D₃ supplementation (***) $P < 0.001$, $n = 16$). (b) A typical experiment where PBMCs were pretreated with AZ11645373 (50 nmol/L) prior to the application of BzATP (50 μ mol/L). (c) The comparison of the BzATP (50 μ mol/L) effect on AZ11645373 (50 nmol/L)-inhibited calcium influx in PBMCs before and after supplementation (***) $P < 0.001$, $n = 16$).

[8, 34, 35]. In our previous study, we have demonstrated that the function of CRAC channels is altered in PBMCs of patients in early stages of CKD [6]. It is not known what changes in the STIM1/Orai1 system develop over the course of CKD. In the current study, a 6-month vitamin D₃ supplementation significantly reduced the increased Ca²⁺ entry through CRAC channels which contributed to the decrease in [Ca²⁺]_i. In our previous study, a 12-month vitamin D₃ supplementation decreased the Ca²⁺ entry through

CRAC channels insignificantly, which could be due to less rigorous patient selection (patients with diabetes mellitus and polycystic kidney disease were included). To our best knowledge, the effects either 25(OH)D₃ or 1,25(OH)₂D₃ on CRAC channels were not studied in CKD.

4.3. Ca²⁺ Entry via P2X₇ Channels. The most recent advances provide compelling evidence for P2X receptors playing a key role in regulating physiological and pathophysiological

FIGURE 5: The effect of vitamin D₃ supplementation on permeability of P2X₇ pores. (a) Representative flow cytometry histograms of ethidium bromide entry into PBMCs of a CKD patient at basal conditions (white peak) and after a stimulation by BzATP (50 μ mol/L) (grey peak). (b) The comparison of the ethidium bromide permeability under the basal conditions, in BzATP- (50 μ mol/L) stimulated and AZ11645373- (50 nmol/L) inhibited PBMCs before and after vitamin D₃ supplementation.

processes in the kidney [36]. In our study, P2X₇ receptors were involved in the disrupted calcium homeostasis in PBMCs of CKD patients. We have shown that the Ca²⁺ entry via P2X₇ channels and pores was increased and also the permeability of P2X₇ pores was higher. In addition, the function of P2X₇ channels and pores was altered [5]. It is known that 1,25(OH)₂D₃, an active metabolite of vitamin D, prevents Ca²⁺ increase through P2X₇ channels and reduces the plasma membrane permeability through P2X₇ pores in human PBMCs of healthy subjects [17]. 1,25(OH)₂D₃ can also up- or downregulate the expression of several genes in many cell types. To our knowledge, the effect of 1,25(OH)₂D₃ on P2X₇ receptor expression has not been studied. The data of this study disclosed that vitamin D₃ supplementation reduced Ca²⁺ influx through P2X₇ channels and affected their functionality. Moreover, the supplementation had no effect on permeability of P2X₇ pores, and the differences in responses to stimulation or inhibition were not found. Different effects of vitamin D₃ on P2X₇ channels and pores in CKD patients may have been made due to different sensitivity to these channels and pores. It is known that various agonists and antagonists of P2X₇ receptors may have a distinct effect on the function of P2X₇ channels and/or pores [37]. Not only the function but also the expression may be altered in some pathological conditions [38–41]. In our recent study, we found a 1.5-fold increase in the expression of surface P2X₇ receptors on PBMCs from CKD patients compared to healthy subjects [4].

In the current study, the flow cytometric measurement revealed that vitamin D₃ decreased the expression of P2X₇ receptors by 45%.

4.4. Activity of PMCA. The PMCA is critical for the maintenance of resting [Ca²⁺]_i in nonexcitable cells and may be the last gatekeeper for the control of low [Ca²⁺]_i. We have observed a decreased PMCA activity in early CKD patients [4]. This finding is consistent with other studies and points out that the PMCA activity decreases with kidney disease progression [42]. The decline in PMCA activity may be caused by numerous factors such as calmodulin deficiency, an activity of other endogenous protein regulators, and inhibition of mitochondrial or glycolytic metabolism [43]. The Ca²⁺ influx through CRAC channels is also an important regulator of PMCA activity, and an altered communication between them may be another cause of PMCA malfunction [44]. 1,25(OH)₂D₃ is known to increase PMCA expression and activity in Ca²⁺ transporting tissues such as the intestine, as well as in osteoblasts and Madin-Darby bovine kidney epithelial cells [45–47]. Several experimental studies have observed direct and/or indirect effects of 1,25(OH)₂D₃ and/or 24,25(OH)₂D₃ on activity and expression of PMCA in different cells, but the effects of 25(OH)D₃ itself have not been studied. The studies targeting at the effects of native vitamin D₃ supplementation on PMCA activity in patients with early stages of CKD are also missing. In the present

FIGURE 6: The expression of surface P2X₇ receptors. (a) Representative flow cytometry histograms of PBMCs immunostained with primary antibody for the extracellular domain of the P2X₇ receptor (gray peak) and an isotype-matched control (IgG2a, white peak). (b) Immunofluorescence staining of P2X₇ receptors in plasma membrane using anti-P2X₇ (extracellular) antibody of lymphocytes visualised by fluorescence microscopy. Images represent surface P2X₇ receptors in four different samples. (c) Decreased expression of surface P2X₇ receptors after vitamin D₃ supplementation in a whole population of PBMCs (***P* < 0.001, *n* = 16).

study, we did not find changes in activity of PMCA after treatment; however, the concentration of plasma membrane proteins was increased by 47%. Therefore, we cannot preclude the possibility of increased expression of PMCA.

In conclusion, we have shown that vitamin D₃ supplementation reduces elevated $[Ca^{2+}]_i$ via CRAC and P2X₇ channels and decreases the expression of cell surface P2X₇ receptors in early CKD. We did not find the effect on P2X₇

pores and PMCA activity. Thus, vitamin D₃ supplementation had a beneficial effect on disturbed cell calcium homeostasis in early CKD.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

FIGURE 7: The assessment of PMCA activity. (a) The representative absorbance spectrum of a CKD patient PMCA in the medium with (black line) and without (grey line) Ca^{2+} . (b) The comparison of PMCA activity in CKD patients at baseline and after a 6-month vitamin D_3 treatment.

Acknowledgments

The authors would like to thank Mrs M. Karolyova, Mrs D. Nagyova, M.Sc. D. Lakatos-Hanicova, and Mrs J. Hullmanova for excellent technical assistance. This paper was created by the realisation of the project "Center of Excellence of Environmental Health," ITMS no. 26240120033, based on the Supporting Operational Research and Development Program financed from the European Regional Development Fund, and was supported by the Grants from the VG SZU 19-90-07, "NanoNet 2" (ITMS 26240120018), SK-FR-2013-0020, APVV-0134-12, and UK/361/2014.

References

- [1] A. S. Dusso, A. J. Brown, and E. Slatopolsky, "Vitamin D," *American Journal of Physiology: Renal Physiology*, vol. 289, no. 1, pp. F8–F28, 2005.
- [2] S. Z. Fadem and S. M. Moe, "Management of chronic kidney disease mineral-bone disorder," *Advances in Chronic Kidney Disease*, vol. 14, no. 1, pp. 44–53, 2007.
- [3] M. J. Berridge, "Elementary and global aspects of calcium signalling," *Journal of Experimental Biology*, vol. 200, no. 2, pp. 315–319, 1997.
- [4] Z. Kaderjakova, I. Lajdova, M. Horvathova, M. Morvova, and L. Sikurova, "Effects of chronic kidney disease on blood cells membrane properties," *Bioelectrochemistry*, vol. 87, pp. 226–229, 2012.
- [5] I. Lajdova, A. Oksa, D. Chorvat Jr., P. Topor, and V. Spustova, "Purinergic P2X_7 receptors participate in disturbed intracellular calcium homeostasis in Peripheral Blood Mononuclear Cells of Patients with Chronic Kidney Disease," *Kidney and Blood Pressure Research*, vol. 35, no. 1, pp. 48–57, 2012.
- [6] I. Lajdova, V. Spustova, A. Oksa, A. Chorvatova, D. Chorvat, and R. Dzurik, "Intracellular calcium homeostasis in patients with early stages of chronic kidney disease: effects of vitamin D_3 supplementation," *Nephrology Dialysis Transplantation*, vol. 24, no. 11, pp. 3376–3381, 2009.
- [7] D. Polak-Jonkisz, L. Purzyc, K. Laszki-Szczchor, K. Musiał, and D. Zwolińska, "The endogenous modulators of Ca^{2+} - Mg^{2+} -dependent ATPase in children with chronic kidney disease (CKD)," *Nephrology Dialysis Transplantation*, vol. 25, no. 2, pp. 438–444, 2010.
- [8] S. Feske, C. Picard, and A. Fischer, "Immunodeficiency due to mutations in ORAI1 and STIM1," *Clinical Immunology*, vol. 135, no. 2, pp. 169–182, 2010.
- [9] M. Partiseti, F. Le Deist, C. Hivroz, A. Fischer, H. Korn, and D. Choquet, "The calcium current activated by T cell receptor and store depletion in human lymphocytes is absent in a primary immunodeficiency," *The Journal of Biological Chemistry*, vol. 269, no. 51, pp. 32327–32335, 1994.
- [10] V. Ralevic and G. Burnstock, "Receptors for purines and pyrimidines," *Pharmacological Reviews*, vol. 50, no. 3, pp. 413–492, 1998.
- [11] F. Di Virgilio and A. Solini, "P2 receptors: new potential players in atherosclerosis," *British Journal of Pharmacology*, vol. 135, no. 6, pp. 831–842, 2002.
- [12] N. R. Jørgensen, Z. Henriksen, O. H. Sørensen, E. F. Eriksen, R. Civitelli, and T. H. Steinberg, "Intercellular calcium signaling occurs between human osteoblasts and osteoclasts and requires activation of osteoclast P2X_7 receptors," *Journal of Biological Chemistry*, vol. 277, no. 9, pp. 7574–7580, 2002.
- [13] R. A. North, "Molecular physiology of P2X receptors," *Physiological Reviews*, vol. 82, no. 4, pp. 1013–1067, 2002.
- [14] O. Vonend, C. M. Turner, C. M. Chan et al., "Glomerular expression of the ATP-sensitive P2X_7 receptor in diabetic and hypertensive rat models," *Kidney International*, vol. 66, no. 1, pp. 157–166, 2004.

- [15] National Kidney Foundation, "K/DOQI clinical practice guidelines for bone metabolism and disease in chronic kidney disease," *The American Journal of Kidney Disease*, vol. 42, supplement 3, pp. S1–S201, 2013.
- [16] A. S. Levey, J. Coresh, T. Greene et al., "Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate," *Annals of Internal Medicine*, vol. 145, no. 4, pp. 247–254, 2006.
- [17] I. Lajdova, D. Chorvat Jr., and A. Chorvatova, "Rapid effects of $1\alpha,25(\text{OH})_2\text{D}_3$ in resting human peripheral blood mononuclear cells," *European Journal of Pharmacology*, vol. 586, no. 1–3, pp. 14–23, 2008.
- [18] D. J. Hanahan and J. E. Ekholm, "The preparation of red cell ghosts (membranes)," *Methods in Enzymology*, vol. 31, pp. 168–172, 1974.
- [19] R. Y. Tsien, "Fluorescence measurement and photochemical manipulation of cytosolic free calcium," *Trends in Neurosciences*, vol. 11, no. 10, pp. 419–424, 1988.
- [20] J. P. Kao, "Practical aspects of measuring $[\text{Ca}^{2+}]$ with fluorescent indicators," *Methods in Cell Biology*, vol. 40, pp. 155–181, 1994.
- [21] M. D. Bootman, T. J. Collins, L. Mackenzie, H. Llewellyn Roderick, M. J. Berridge, and C. M. Peppiatt, "2-Aminoethoxydiphenyl borate (2-APB) is a reliable blocker of store-operated Ca^{2+} entry but an inconsistent inhibitor of InsP_3 -induced Ca^{2+} release," *FASEB Journal*, vol. 16, no. 10, pp. 1145–1150, 2002.
- [22] J.-I. Goto, A. Z. Suzuki, S. Ozaki et al., "Two novel 2-aminoethyl diphenylborinate (2-APB) analogues differentially activate and inhibit store-operated Ca^{2+} entry via STIM proteins," *Cell Calcium*, vol. 47, no. 1, pp. 1–10, 2010.
- [23] L. Stokes, L.-H. Jiang, L. Alcaraz et al., "Characterization of a selective and potent antagonist of human P_2X_7 receptors, AZ11645373," *British Journal of Pharmacology*, vol. 149, no. 7, pp. 880–887, 2006.
- [24] J. S. Wiley, C. E. Gargett, W. Zhang, M. B. Snook, and G. P. Jamieson, "Partial agonist and antagonists reveal a second permeability state of human lymphocyte $\text{P}_2\text{Z}/\text{P}_2\text{X}_7$ channel," *The American Journal of Physiology—Cell Physiology*, vol. 275, no. 5, pp. C1224–C1231, 1998.
- [25] O. H. Lowry, N. J. Rosebrough, A. L. Farr, and R. J. Randall, "Protein measurement with the folin phenol reagent," *The Journal of Biological Chemistry*, vol. 193, no. 1, pp. 265–275, 1951.
- [26] X. Luo, B. Hojayeve, N. Jiang et al., "STIM1-dependent store-operated Ca^{2+} entry is required for pathological cardiac hypertrophy," *Journal of Molecular and Cellular Cardiology*, vol. 52, no. 1, pp. 136–147, 2012.
- [27] M. P. Mattson and S. L. Chan, "Neuronal and glial calcium signaling in Alzheimer's disease," *Cell Calcium*, vol. 34, no. 4–5, pp. 385–397, 2003.
- [28] M. Raraty, J. Ward, G. Erdemli et al., "Calcium-dependent enzyme activation and vacuole formation in the apical granular region of pancreatic acinar cells," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 24, pp. 13126–13131, 2000.
- [29] S. Somlo and B. Ehrlich, "Human disease: calcium signaling in polycystic kidney disease," *Current Biology*, vol. 11, no. 9, pp. R356–R360, 2001.
- [30] S. G. Baryshnikov, M. V. Pulina, A. Zulian, C. I. Linde, and V. A. Golovina, "Modulation of plasma membrane calcium-ATPase activity by local calcium microdomains near CRAC channels in human T cells," *The American Journal of Physiology—Cell Physiology*, vol. 297, no. 5, pp. C1103–C1112, 2009.
- [31] M. Potier, J. C. Gonzalez, R. K. Motiani et al., "Evidence for STIM1- and Orai1-dependent store-operated calcium influx through ICRC in vascular smooth muscle cells: role in proliferation and migration," *FASEB Journal*, vol. 23, no. 8, pp. 2425–2437, 2009.
- [32] W. Lu, J. Wang, L. A. Shimoda, and J. T. Sylvester, "Differences in STIM1 and TRPC expression in proximal and distal pulmonary arterial smooth muscle are associated with differences in Ca^{2+} responses to hypoxia," *The American Journal of Physiology—Lung Cellular and Molecular Physiology*, vol. 295, no. 1, pp. L104–L113, 2008.
- [33] F. R. C. Giachini, C.-W. Chiao, F. S. Carneiro et al., "Increased activation of stromal interaction molecule-1/Orai-1 in aorta from hypertensive rats: a novel insight into vascular dysfunction," *Hypertension*, vol. 53, no. 2, pp. 409–416, 2009.
- [34] C.-A. McCarl, C. Picard, S. Khalil et al., "ORAI1 deficiency and lack of store-operated Ca^{2+} entry cause immunodeficiency, myopathy, and ectodermal dysplasia," *Journal of Allergy and Clinical Immunology*, vol. 124, no. 6, pp. 1311–1318.e7, 2009.
- [35] C. Picard, C.-A. McCarl, A. Papolos et al., "STIM1 mutation associated with a syndrome of immunodeficiency and autoimmunity," *The New England Journal of Medicine*, vol. 360, no. 19, pp. 1971–1980, 2009.
- [36] R. E. Birch, E. M. Schwiebert, C. M. Peppiatt-Wildman, and S. S. Wildman, "Emerging key roles for P_2X receptors in the kidney," *Frontiers in Physiology*, vol. 4, article 262, 2013.
- [37] L. G. B. Ferreira, R. A. M. Reis, L. A. Alves, and R. X. Faria, "Intracellular signaling pathways integrating the pore associated with $\text{P}_2\text{X}_7\text{R}$ receptor with other large pores," in *Patch Clamp Technique*, F. S. Kanez, Ed., pp. 37–54, InTech Open Access Publishing, Vienna, Austria, 2012.
- [38] S. Franco-Martínez, P. Niño-Moreno, S. Bernal-Silva et al., "Expression and function of the purinergic receptor P_2X_7 in patients with pulmonary tuberculosis," *Clinical & Experimental Immunology*, vol. 146, no. 2, pp. 253–261, 2006.
- [39] M. H. García-Hernández, L. Portales-Cervantes, N. Cortez-Espinosa et al., "Expression and function of P_2X_7 receptor and CD39/Entpd1 in patients with type 2 diabetes and their association with biochemical parameters," *Cellular Immunology*, vol. 269, no. 2, pp. 135–143, 2011.
- [40] B. J. Gu, W. Y. Zhang, L. J. Bendall, I. P. Chessell, G. N. Buell, and J. S. Wiley, "Expression of P_2X_7 purinoceptors on human lymphocytes and monocytes: evidence for nonfunctional P_2X_7 receptors," *American Journal of Physiology—Cell Physiology*, vol. 279, no. 4, pp. C1189–C1197, 2000.
- [41] S. Madec, C. Rossi, M. Chiarugi et al., "Adipocyte P_2X_7 receptors expression: a role in modulating inflammatory response in subjects with metabolic syndrome?" *Atherosclerosis*, vol. 219, no. 2, pp. 552–558, 2011.
- [42] U. Gafter, T. Malachi, H. Barak, M. Djaldetti, and J. Levi, "Red blood cell calcium homeostasis in patients with end-stage renal disease," *Journal of Laboratory and Clinical Medicine*, vol. 114, no. 3, pp. 222–231, 1989.
- [43] M. L. Holton, W. Wang, M. Emerson, L. Neyses, and A. L. Armesilla, "Plasma membrane calcium ATPase proteins as novel regulators of signal transduction pathways," *World Journal of Biological Chemistry*, vol. 1, no. 6, pp. 201–208, 2010.
- [44] D. M. Bautista and R. S. Lewis, "Modulation of plasma membrane calcium-ATPase activity by local calcium microdomains near CRAC channels in human T cells," *Journal of Physiology*, vol. 556, no. 3, pp. 805–817, 2004.

- [45] P. Glendenning, T. Ratajczak, I. M. Dick, and R. L. Prince, "Calcitriol upregulates expression and activity of the 1b isoform of the plasma membrane calcium pump in immortalized distal kidney tubular cells," *Archives of Biochemistry and Biophysics*, vol. 380, no. 1, pp. 126–132, 2000.
- [46] P. Glendenning, T. Ratajczak, I. M. Dick, and R. L. Prince, "Regulation of the 1b isoform of the plasma membrane calcium pump by 1,25-dihydroxyvitamin D₃ in rat osteoblast-like cells," *Journal of Bone and Mineral Research*, vol. 16, no. 3, pp. 525–534, 2001.
- [47] S. N. Kip and E. E. Strehler, "Vitamin D₃ upregulates plasma membrane Ca²⁺-ATPase expression and potentiates apico-basal Ca²⁺ flux in MDCK cells," *The American Journal of Physiology—Renal Physiology*, vol. 286, no. 2, pp. F363–F369, 2004.