

Research Article

Effects of Chronic REM Sleep Restriction on D₁ Receptor and Related Signal Pathways in Rat Prefrontal Cortex

Yan Han,¹ Xiaosa Wen,^{2,3} Fei Rong,² Xinmin Chen,² Ruying Ouyang,⁴ Shuai Wu,¹ Hua Nian,⁵ and Wenling Ma²

¹Department of Neurology, Changhai Hospital, Second Military Medical University, Shanghai 200433, China

²Department of Environmental Hygiene, Second Military Medical University, Shanghai 200433, China

³Minhang District Center for Disease Control and Prevention, Shanghai 201101, China

⁴Shanghai Sanatorium for Naval Retired Cadres, Shanghai 200434, China

⁵Pharmaceutical Center of Yueyang Hospital, Shanghai University of Traditional Chinese Medicine, Shanghai 200433, China

Correspondence should be addressed to Wenling Ma; wenlingma@163.com

Received 3 July 2014; Accepted 17 November 2014

Academic Editor: John J. Gildea

Copyright © 2015 Yan Han et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The prefrontal cortex (PFC) mediates cognitive function that is sensitive to disruption by sleep loss, and molecular mechanisms regulating neural dysfunction induced by chronic sleep restriction (CSR), particularly in the PFC, have yet to be completely understood. The aim of the present study was to investigate the effect of chronic REM sleep restriction (REM-CSR) on the D₁ receptor (D₁R) and key molecules in D₁R signal pathways in PFC. We employed the modified multiple platform method to create the REM-CSR rat model. The ultrastructure of PFC was observed by electron microscopy. HPLC was performed to measure the DA level in PFC. The expressions of genes and proteins of related molecules were assayed by real-time PCR and Western blot, respectively. The general state and morphology of PFC in rats were changed by CSR, and DA level and the expression of D₁R in PFC were markedly decreased ($P < 0.01$, $P < 0.05$); the expression of phosphor-PKAc α was significantly lowered in CSR rats ($P < 0.05$). The present results suggested that the alteration of neuropathology and D₁R expression in PFC may be associated with CSR induced cognitive dysfunction, and the PKA pathway of D₁R may play an important role in the impairment of advanced neural function.

1. Introduction

As one of the most striking problems in modern society, sleep disorder greatly shortens the time of sleep [1]. Sleep disorder can lead to sleep restriction/sleep deprivation, further resulting in hazardous physiological effects [2]. Sleep deprivation is particularly serious among people who are in some special professions, such as air transport, seafaring, emergency medicine, military contingency, and social security. Sleep deprivation can reduce job performance and cause cognitive impairments in humans including deficits in attention, learning, and memory, executive function, and alertness as well as emotional reactivity [3]. Compared with acute sleep restriction (ASR), chronic sleep restriction (CSR) which is induced by long-term intermittent sleep deprivation

or sleep fragmentation is more prevalent and can lead to more serious consequences [4]. However, the exact neurochemical mechanism underlying CSR-induced advanced neuronal dysfunction has not been fully understood.

The prefrontal cortex (PFC) plays an important role in many higher-order executive tasks. It is involved in learning [5], memory [6], cognitive flexibility [7], and emotional processing. The PFC mediates cognitive responses that are sensitive to disruption by sleep loss [8]; therefore, PFC-mediated cognitive abilities might be impaired by sleep restriction. Previous studies have found reduced cognitive function with reduced neuronal activation following sleep loss in the PFC [9] and indicated that sleep loss induces alteration in synaptic plasticity and membrane excitability in the cortex [10], which might be modulated by dopaminergic

system. PFC receives dopaminergic projections from both the substantia nigra and the ventral tegmental area [11]. Dopaminergic innervation constitutes a major regulating system that modulates higher cognitive function. In addition to modulating neural activity within PFC, dopamine (DA) also alters activity-dependent neural plasticity; these changes might play a pivotal role in learning, memory, and cognitive function [12, 13]. Therefore, DA and its receptors might be associated with the changes of synaptic excitability and plasticity in PFC during sleep restriction.

The diverse physiological actions of DA are mediated by five distinct subtypes, which are commonly divided into two classes: the D_1 -like receptors (D_1 Rs, comprised of D_1 and D_5 receptors) and the D_2 -like receptors (D_2 , D_3 , and D_4 receptors) [14]. Within cortex, the D_1 Rs are expressed in prefrontal V-VI layer pyramidal neurons and I-III layer interneurons. The expression of D_1 Rs is enriched in the PFC of both primates and rodents, suggesting an important role in specifically circuit functions in PFC. D_1 receptor (D_1 R) is a principal member of D_1 Rs, and, in PFC, D_1 R has been shown to modulate many neural activities [15, 16]. As a G-protein-coupled receptor, D_1 R can be coupled to the G protein Gs and activate adenylyl cyclase and regulate the expression of cAMP response element binding (CREB) protein which is a nuclear transcription factor, modulating the functions of neurons. This regulation can be performed through several pathways such as phosphate kinase A (PKA) and mitogen activated protein kinase (MAPK) pathways [17]. However, less is known about how CSR induces the changes of neuronal ultrastructure in PFC and which pathway of D_1 R plays the major role in PFC during the cognitive dysfunction. In this study, we employed transmission electron microscope to observe the ultrastructure of PFC and used PCR and Western blot to measure the expression of related signal molecules. The design of the study aimed at exploring preliminarily the modulatory mechanism of cognition underlying D_1 R and its signal pathways in PFC during CSR.

2. Materials and Methods

2.1. Animals and Groups. Sixty male Sprague-Dawley rats, weighing 300 ± 20 g, were purchased from the Animal Center of Second Military Medical University (SMMU), Shanghai, China. All treatments and experimental procedures were approved by the Animal Ethics Committee of SMMU in accordance with the international guidelines for animal experiments. After a week of adaptive feeding and 3 days of training, 16 rats were rejected due to escape latency exceeding 90 seconds in each trial in water maze or failure to stand on the platform constantly in the sleep deprivation tank. The screening regimen and requirements were described in the previous study [18]. The remaining 44 animals were randomly assigned to the two groups: the treatment control (TC) group and the chronic sleep restriction (CSR) group, each consisting of 22 animals.

2.2. Experimental Procedures. The animals were housed in a room ($22 \pm 1^\circ\text{C}$; 12 h light-dark cycle, light on at 8:00 a.m., light off at 8:00 p.m.) with normal feed and water. The CSR

model was created by modified multiple platform method as stated earlier [19]. Eight narrow platforms (diameter 6.5 cm) or a grid floor was placed inside a water tank. Six rats in the CSR group were forced to stand on the platforms surrounded by water for 18 h (beginning at 16:00) daily; at least two platforms in each tank are empty, thus allowing animals to move around. Animals in the TC group were placed on the grid, thus allowing them to sleep and move freely in a similar water environment. This process continued for 21 days to create the chronic REM sleep restriction (REM-CSR) model. Tufik and his colleagues employed the method of EEG monitoring to discover that this sleep restriction led to a complete suppression of REM sleep during the 18 h sleep deprivation period each day, which persisted throughout the whole restriction period [1, 20]. Food and water were available through a grid placed on top of the water cage during the time of sleep restriction.

2.3. Preparation of PFC. After 21 days of CSR, the animals were perfused with 4°C saline through the ascending aorta under chloral hydrate anesthesia (400 mg/Kg, i.p.). PFC from 18 animals in each group (including prelimbic medial cortex and infralimbic medial cortex) was dissected under microscope (Figure 1(a)). The samples were immediately frozen in liquid nitrogen and stored at -80°C for subsequent experiment. Meanwhile, samples of the last 4 rats in each group were fixed with 2.5% glutaraldehyde and 4% paraformaldehyde for the observation of ultrastructure.

2.4. Observation of Electron Microscope. The isolated PFC tissues were fixed with 1% osmium tetroxide and then dehydrated and embedded in resin. The 80 nm sections were cut and double stained with uranyl acetate and lead citrate. The ultrastructure of PFC was observed under transmission electron microscope (H-7650, Hitachi, Japan) and images were recorded.

2.5. The DA Concentration in PFC. The DA concentration was detected by High-Performance Liquid Chromatography with electrochemical detection (ESA, USA, $n = 8$ for each group). The PFC tissues were sonicated in 0.5 mL of 0.1 M perchloric acid and DA extraction was performed. The supernatant was filtered through a 0.22 Millex-GV filter and transferred to an autosampler tube prior to injection. Detection was performed with an ESA Coulochem III electrochemical detector (ESA, USA). The area under the curve (AUC) of standard solutions was used for the comparison of dopamine concentrations in the samples.

2.6. RNA Isolation and Quantitative Real-Time PCR. The mRNA for 7 signal transduction molecules was examined by real-time PCR ($n = 6$ for each group). Total RNA was extracted from PFC tissue with Trizol reagent (Invitrogen), according to the instructions of the manufacturer. RNA yield and quality were determined spectrophotometrically by A260/A280 ratio. After reverse transcription, the real-time PCRs were done in a 20 μL reaction mixture system. Amplification was performed by the Applied Biosystems




FIGURE 1: Transmission electron microscopic results of PFC ($n = 4$). (a) The shadow area in the middle of cross-section is the objective PFC observed by electron microscope; compared to the TC group (b), the CSR group (c) showed that the number of mitochondria (thin arrows) was decreased and the structure of mitochondrial cristae was fuzzy and ruptured; the postsynaptic density (arrowheads) became much thinner or disappeared; the rough endoplasmic reticulum (thick arrows) structure(s) became fuzzy.

7500 Real-Time PCR System (ABI, USA) with an initial denaturation of 5 min at 95°C, followed by 45 cycles of 95°C for 15 s, 60°C for 30 s, and 72°C for 30 s. At the end of the amplification phase, a melting curve analysis was performed to ensure the amplification of a single PCR product. Primers used in the amplification were designed and synthesized by Sangon Biotech (Shanghai) Co., Ltd. The sequences of 7 gene primers were shown in Table 1.

2.7. Protein Extraction and Western Blotting. The frozen PFC samples were dropped into SDS extraction buffer and homogenized with an ultrasonic cell disrupter and then boiled and centrifuged. The supernatant was transferred to a new tube, and, after determination of the protein concentrations, the samples were then diluted to SDS extraction buffer and stored at -20°C for later experiments. Western blotting ($n = 4$ for each group) was used to measure the protein expressions of D₁R (1:500, Sigma, D2994), phosphorylated PKA α (1:1000, CST, #5661), PKA α (1:1000, abcam, ab76238), phosphorylated ERK1/2 (1:1000, Bioworld, BS5016), ERK1/2 (1:1000, Bioworld, BS1112), phosphorylated CREB (1:1000, CST, #9198), CREB (1:1000, CST, #9197), and β -actin (1:1000, Santa Cruz, sc-1616r). The sample proteins were separated by 10% SDS-polyacrylamide gel electrophoresis and transferred to nitrocellulose membrane (Millipore, USA). The membranes were blocked with 5% nonfat dry milk in Tris buffered saline (TBS) for 2 h at room temperature and incubated with the primary antibodies in TBS with 5% dry milk overnight at 4°C. The membranes were washed with TBS plus 0.1% Tween-20 (TBS-T) three times (10 min each) and incubated for 2 h at room temperature with different horseradish peroxidase-conjugated secondary IgG antibodies (1:3000, KPL, anti-rabbit or anti-mouse) and washed again in TBS-T (10 min \times 3). Blots were revealed by ECL advanced kit (Thermo Fisher, USA). To normalize protein content, blots were probed with corresponding β -actin, and the ratio to β -actin was calculated.

2.8. Statistical Analysis. All the assessment data were analyzed by a two-tailed Student's t -test for the comparison

TABLE 1: The sequences of 7 primers in real-time PCR.

Primer name		Sequence (5' to 3')
Drd1a	Forward	GCGAGCACCACAAAGCC
	Reverse	GGAAACTAGATTCAAACCCACAGA
Creb1	Forward	TTGTTGTTCAAGCTGCCTCTG
	Reverse	TGCTGTGCGAATCTGGTATGT
Adcy5	Forward	CAACTACCTGAACGGGGACTAT
	Reverse	CGATGCTGTGCTCCTTGA
Prkaca	Forward	CCCACTCCCTAAATCCATTCTG
	Reverse	AAGCCCAGTTCCCTTCCTTGAC
Darpp32	Forward	CTCCCCAGAAACCCACTCT
	Reverse	GCAAACACAGACCAGCCTTAGT
Erk1	Forward	CTGGAATGGAAGGGCTATGAC
	Reverse	CAACAGGATGAGTAGGGCAGAG
Erk2	Forward	GCGTTCAGATGTCCGGTGTC
	Reverse	CAAAGGAGTCAAGAGTGGGTAAG
Rat-gapdh	Forward	CAGTGCCAGCCTCGTCAT
	Reverse	AGGGGCATCCACAGTCTTC

of mean values between the two groups, using GraphPad Prism 5.01 (GraphPad Software, Inc., USA). Values were expressed as mean \pm standard deviation and differences were considered to be significant when $P < 0.05$.

3. Results

3.1. The Change of Behavioral Ability and General State of CSR Rats. In this study, we found that the learning and memory ability, stamina, and locomotor activity in rats were significantly reduced in the CSR group, as compared with the TC group. This result repeats our previous reports [18]. In addition, during the course of sleep restriction, we also found

that CSR rats displayed increased excitability and augmented alertness to environmental stimulation in the early phase (CSR 1st–7th d), and the animals showed irritability and bite and attack enhancements in the middle stage (CSR 8th–14th d). In the late phase (CSR 15th–22th d), the rats demonstrated emaciation, response slowness, arched back, and trembling and fell into water more frequently.

3.2. The Pathological Changes and DA Levels in PFC. To assess the morphological changes, the ultrastructures of PFC in the two groups were observed under the transmission electron microscope according to the previous method [21]. We randomly examined 5 fields in each group. The mitochondrial impairment was identified by observing the vague or disappeared cristae and the fragmentation of membrane. We found that only 18.3% (11/60) of the mitochondria in the TC group was damaged, while the damage rate in the CSR group was 51.0% (26/51). The damages of postsynaptic density were identified when the structure of synapse became much thinner or disappeared. Both the number and the rate of damaged synapses in the CSR group (44.4%, 12/27) were obviously increased, compared with those of the control group (24.1%, 7/29). Moreover, the structure of rough endoplasmic reticulum became fuzzy (Figure 1). Meanwhile, when compared with that of the TC group, the DA concentration of PFC in the CSR group was notably decreased ($P < 0.01$, Figure 2(a)).

3.3. Effects of CSR on the Expression of Key Molecules mRNAs in D_1R Pathway. In addition to D_1R , we also chose 6 other key molecules in the D_1R signal pathways and detected their mRNA expression. *Drd1a* was the gene of D_1R ; *Acly5*, *Prkaca*, and *Darpp32* are the crucial regulators of PKA pathway; *Erk1* and *Erk2* play an essential role in the MAPK pathway; *Creb1* is a transcription factor and is involved in the regulation of learning and memory. The results showed that only *Drd1a* was significantly reduced by CSR ($P < 0.05$, Figure 2(B1)), and the expressions of other genes showed no differences between the two groups during CSR (Figure 2(B2)–(B7)).

3.4. Effects of CSR on the Expression of Key Molecules Proteins in D_1R Pathway. The D_1R protein expression in the CSR group was significantly decreased, as compared to that in the TC group ($P < 0.05$, Figure 3(a)). The phosphor-PKAc α was also notably reduced in the CSR rats, as compared to the corresponding control ($P < 0.05$, Figure 3(B1)). Both the phosphor-CREB and CREB in the CSR group were obviously decreased as compared with CREB in the TC group ($P < 0.05$, Figure 3(D1)/(D2)). There were no significant changes in PKAc α (Figure 3(B2)), phosphor-Erk1/2 (Figure 3(C1)), or total Erk1/2 (Figure 3(C2)) between the CSR and the control group.

4. Discussion

According to the results of our preliminary experiment, there was only one control group in the study. The blank control

(cage control) was rejected due to its similar outcome to the treatment control. The modified multiple platform method used to create the CSR model had been repeatedly tested by previous researches [1]. This method had been further modified in an attempt to eliminate the social isolation and the restriction of movement. The treatment control group for the deprivation environment consisted of animals lying on a grid placed inside the water tank, which is commonly used as a control for the confounding factors inherent to this technique [19].

Sleep deprivation has a broad variety of effects on performance and neural function that manifest themselves at different levels of description [22]. Our previous study demonstrated that REM-CSR can significantly drop the learning and memory ability, stamina, and locomotor activity. And the results also showed that, after one week of CSR, the ability of learning and memory began to decline and gradually reduced in the second and third weeks [18]. Those results are consistent with many other studies [23, 24]. This research also observed that the general state and mental condition of rats were affected by REM-CSR and changed with the prolonged restriction.

Our latest study pointed out that the cognitive dysfunction of the CSR rat was related to the pathological change of ultrastructure of hippocampus [18]. This research is the first to observe the PFC neuronal degeneration after REM-CSR, and we also found that the mitochondria and endoplasmic reticulum in the PFC were destructed. The rate of mitochondrial damage caused by REM-CSR in PFC was 51%, which was much more than those of hippocampus. Furthermore, the same destruction of endoplasmic reticulum and postsynaptic density in PFC demonstrated that REM-CSR can cause similar damage of PFC and hippocampus. Different exposure or stress can all lead to the destruction of mitochondria, such as swelling and the disappearance or fuzziness of crista. And the damage of mitochondria was associated with the degeneration, apoptosis, and cell death of neurons, which can reduce the discharge activity of neurons. The results indicated that, through altering the structure of neurons, REM-CSR might reduce the neuronal excitability and then affect the function of PFC. PFC is involved in working memory, decision making, and vigilance maintenance [25, 26]. It could be therefore inferred that PFC structural destruction might play a crucial role in the CSR-induced advanced neuronal dysfunction.

The neuronal degeneration of PFC was associated with the reduction of neuronal activity [8, 27]. Moreover, previous studies have indicated that the activation of D_1R increases neuronal activity, suggesting that the D_1R may be involved in the modulation of neuronal activity [28]. The present study displayed that DA concentration and the expression of D_1R mRNA and protein were significantly reduced after REM-CSR, suggesting that the CSR-induced changes of PFC structure and activity may be associated with the alteration of D_1R expression and its signal pathways. Therefore, the DA and D_1R in the PFC might play an important role in the CSR-induced advanced neural dysfunction, and elevation of




FIGURE 2: Continued.




FIGURE 2: (a) The results of DA content in the two groups ($n = 8$). Compared to that of the TC group, the DA content in the CSR group was notably reduced. (b) Effects of CSR on the expressions of some genes in PFC ($n = 6$). CSR notably reduced the *Drda1a* in the CSR rats (B1). The expressions of *Creb1*, *Acly5*, *Prkaca*, *Darpp32*, *Erk1*, and *Erk2* in PFC showed no significant differences between the CSR and TC groups (B2–B7). TC: treatment control; CSR: chronic sleep restriction. * $P < 0.05$, and ** $P < 0.01$.

DA concentration or stimulation of D_1R in the PFC might improve the cognitive dysfunction during the CSR process.

D_1R regulates the nuclear transcription factor via two major pathways: PKA and MAPK pathways, and PKA and extracellular signal-regulated kinase (ERK1/2) are the key regulatory molecules in PKA and MAPK pathways, respectively (Figure 4). Studies have shown that the ability of D_1R to increase both the persistence and early magnitude of long-term potentiation (LTP) was associated with activation of PKA pathway [29], and other researchers found that the decreased phosphorylation of DARPP-32 and CREB, which were also the key signal molecules in PKA pathway, was linked to the notable impairment of delayed spatial memory retrieval caused by selective D_1R antagonist in the prefrontal cortex [28]. In many cases, ERK1/2 was involved in memory consolidation. Guan et al. [30] indicated that decreased ERK1/2 activation in the hippocampus was associated with sleep deprivation-induced spatial memory impairment, but the alterations of ERK1/2 were not observed in the cortex. In this study, we demonstrated that REM-CSR decreased the phosphorylation of PKA and CREB but did not alter the level of phosphor-ERK1/2 (Figure 4), suggesting that the reduction of DA concentration and D_1R expression induced by REM-CSR might impair the function of PFC by repressing PKA pathway activity but not the MAPK pathway. These results provide new insights in the molecular mechanisms involved in D_1R -dependent function modulation of PFC, where PKA pathway plays a pivotal role in the synthesis of CREB during REM-CSR.

However, Chen et al. [27] once discovered that the form of $D_{1/5}$ receptor-mediated LTP of the intrinsic excitability relied on activation of Ca^{2+} -dependent intracellular signal in PFC, indicating that the $D_{1/5}$ receptor-mediated phosphoinositol (PI) pathway, including the activation of PLC, intracellular

Ca^{2+} elevation, and the activation of Ca^{2+} -dependent CaMKs and PKC, might be associated with the regulation of overall excitability of PFC neurons. In our future studies, we plan to further demonstrate the regulatory mechanism of PI pathway in PFC dysfunction induced by CSR.

5. Conclusions

This study demonstrated that PFC is associated with the changes in behavioral performance and the impairments of cognitive function induced by REM-CSR, and the DA and its D_1R in PFC might be involved in the regulation of this process. A key finding in this research is the reduction of the level of PKA and CREB protein phosphorylation, which indicates that PKA pathway in PFC seems to play a major role in the cognitive impairment of REM-CSR. However, mechanism for the modulation of PFC under CSR needs to be further clarified in the future.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of the paper.

Authors' Contribution

Xiaosa Wen and Yan Han contributed equally to this work.

Acknowledgments

This study was supported by National Natural Science Foundation of China (Grant nos. 81172638 and 81100847). The authors would like to thank professor Xuting Ye (Department of Biophysics, SMMU, China) for the observation




FIGURE 3: Effects of CSR on the expressions of some proteins in PFC. The D₁R (a) and p-PKAα (B1) in the CSR rats were significantly decreased. CSR did not influence PKAα (B2), P-Erk1/2 (C1), and Erk1/2 (C2) in PFC. Both the p-CREB (D1) and CREB (D2) in PFC were notably reduced by CSR. **P* < 0.05, *n* = 4 per group.




FIGURE 4: The scheme of signal pathways of D₁R. This simplified view of the signal pathways shows major downstream mediators between the two major pathways of D₁R. D₁R regulates CREB through PKA pathway (expressed by purple molecules) and MAPK pathway (yellow members), and the key molecules are marked by red. CREB was significantly decreased by CSR, while PKA showed a notable reduction (expressed by downward arrow).

and analysis of electron microscopy. Special thanks go to Professor Guochuan Gong (Naval Medical Research Institute, Shanghai, China) for his helpful review and revision of the paper.

References

- [1] S. Tufik, M. L. Andersen, L. R. A. Bittencourt, and M. T. de Mello, "Paradoxical sleep deprivation: neurochemical, hormonal and behavioral alterations. Evidence from 30 years of research," *Anais da Academia Brasileira de Ciencias*, vol. 81, no. 3, pp. 521–538, 2009.
- [2] M. Basner, H. Rao, N. Goel, and D. F. Dinges, "Sleep deprivation and neurobehavioral dynamics," *Current Opinion in Neurobiology*, vol. 23, no. 5, pp. 854–863, 2013.
- [3] J. G. McCoy and R. E. Strecker, "The cognitive cost of sleep lost," *Neurobiology of Learning and Memory*, vol. 96, no. 4, pp. 564–582, 2011.
- [4] J. Orzel-Gryglewska, "Consequences of sleep deprivation," *International Journal of Occupational Medicine and Environmental Health*, vol. 23, no. 1, pp. 95–114, 2010.
- [5] E. G. Antzoulatos and E. K. Miller, "Differences between neural activity in prefrontal cortex and striatum during learning of novel abstract categories," *Neuron*, vol. 71, no. 2, pp. 243–249, 2011.
- [6] M. R. Warden and E. K. Miller, "Task-dependent changes in short-term memory in the prefrontal cortex," *The Journal of Neuroscience*, vol. 30, no. 47, pp. 15801–15810, 2010.
- [7] A. J. Gruber, G. G. Calhoun, I. Shusterman, G. Schoenbaum, M. R. Roesch, and P. O'Donnell, "More is less: a disinhibited prefrontal cortex impairs cognitive flexibility," *The Journal of Neuroscience*, vol. 30, no. 50, pp. 17102–17110, 2010.
- [8] B. D. Winters, Y. H. Huang, Y. Dong, and J. M. Krueger, "Sleep loss alters synaptic and intrinsic neuronal properties in mouse prefrontal cortex," *Brain Research*, vol. 1420, pp. 1–7, 2011.
- [9] B. S. McKenna and L. T. Eyler, "Overlapping prefrontal systems involved in cognitive and emotional processing in euthymic bipolar disorder and following sleep deprivation: a review of functional neuroimaging studies," *Clinical Psychology Review*, vol. 32, no. 7, pp. 650–663, 2012.
- [10] J. Yan, J.-C. Li, M.-L. Xie et al., "Short-term sleep deprivation increases intrinsic excitability of prefrontal cortical neurons," *Brain Research*, vol. 1401, pp. 52–58, 2011.
- [11] K. L. Clark and B. Noudoost, "The role of prefrontal catecholamines in attention and working memory," *Frontiers in Neural Circuits*, vol. 8, no. 33, pp. 1–19, 2014.
- [12] T. Aubele and M. F. Kritzer, "Gonadectomy and hormone replacement affects in vivo basal extracellular dopamine levels in the prefrontal cortex but not motor cortex of adult male rats," *Cerebral Cortex*, vol. 21, no. 1, pp. 222–232, 2011.
- [13] V. Pawlak and J. N. D. Kerr, "Dopamine receptor activation is required for corticostriatal spike-timing-dependent plasticity," *The Journal of Neuroscience*, vol. 28, no. 10, pp. 2435–2446, 2008.
- [14] J. K. Seamans and C. R. Yang, "The principal features and mechanisms of dopamine modulation in the prefrontal cortex," *Progress in Neurobiology*, vol. 74, no. 1, pp. 1–58, 2004.
- [15] J. R. Glausier, Z. U. Khan, and E. C. Muly, "Dopamine D₁ and D₅ receptors are localized to discrete populations of interneurons in primate prefrontal cortex," *Cerebral Cortex*, vol. 19, no. 8, pp. 1820–1834, 2009.
- [16] M. V. Puig and E. K. Miller, "The role of prefrontal dopamine D1 receptors in the neural mechanisms of associative learning," *Neuron*, vol. 74, no. 5, pp. 874–886, 2012.
- [17] A. S. Undieh, "Pharmacology of signaling induced by dopamine D₁-like receptor activation," *Pharmacology and Therapeutics*, vol. 128, no. 1, pp. 37–60, 2010.
- [18] X.-S. Wen, X.-M. Chen, F. Rong, T. Jing, S. Chen, and W.-L. Ma, "The regulation of SKF38393 on the dopamine and D₁ receptor expression in hippocampus during chronic rem sleep restriction," *CNS Neuroscience and Therapeutics*, vol. 19, no. 9, pp. 730–733, 2013.
- [19] D. Suchecki and S. Tufik, "Social stability attenuates the stress in the modified multiple platform method for paradoxical sleep deprivation in the rat," *Physiology and Behavior*, vol. 68, no. 3, pp. 309–316, 2000.

- [20] R. B. Machado, D. Suchecki, and S. Tufik, "Sleep homeostasis in rats assessed by a long-term intermittent paradoxical sleep deprivation protocol," *Behavioural Brain Research*, vol. 160, no. 2, pp. 356–364, 2005.
- [21] W. Xie, Q. Xie, M. Jin et al., "The β -SiC nanowires (~100 nm) induce apoptosis via oxidative stress in mouse osteoblastic cell line MC3T3-E1," *BioMed Research International*, vol. 2014, Article ID 312901, 9 pages, 2014.
- [22] T. W. Boonstra, J. F. Stins, A. Daffertshofer, and P. J. Beek, "Effects of sleep deprivation on neural functioning: an integrative review," *Cellular and Molecular Life Sciences*, vol. 64, no. 7-8, pp. 934–946, 2007.
- [23] Z. Zhao, Y. Li, H. Chen et al., "Xylaria nigripes mitigates spatial memory impairment induced by rapid eye movement sleep deprivation," *International Journal of Clinical and Experimental Medicine*, vol. 7, no. 2, pp. 356–362, 2014.
- [24] L. Chen, S. Tian, and J. Ke, "Rapid eye movement sleep deprivation disrupts consolidation but not reconsolidation of novel object recognition memory in rats," *Neuroscience Letters*, vol. 563, pp. 12–16, 2014.
- [25] K. L. Clark and B. Noudoost, "The role of prefrontal catecholamines in attention and working memory," *Frontiers in Neural Circuits*, vol. 8, article 33, 2014.
- [26] M. L. Dixon and K. Christoff, "The lateral prefrontal cortex and complex value-based learning and decision making," *Neuroscience and Biobehavioral Reviews*, vol. 45, pp. 9–18, 2014.
- [27] L. Chen, J. D. Bohanick, M. Nishihara, J. K. Seamans, and C. R. Yang, "Dopamine D1/5 receptor-mediated long-term potentiation of intrinsic excitability in rat prefrontal cortical neurons: Ca^{2+} -dependent intracellular signaling," *Journal of Neurophysiology*, vol. 97, no. 3, pp. 2448–2464, 2007.
- [28] M. Hotte, S. Thuault, F. Lachaise et al., "D₁ receptor modulation of memory retrieval performance is associated with changes in pCREB and pDARPP-32 in rat prefrontal cortex," *Behavioural Brain Research*, vol. 171, no. 1, pp. 127–133, 2006.
- [29] M. Stramiello and J. J. Wagner, "D_{1/5} receptor-mediated enhancement of LTP requires PKA, Src family kinases, and NR2B-containing NMDARs," *Neuropharmacology*, vol. 55, no. 5, pp. 871–877, 2008.
- [30] Z. Guan, X. Peng, and J. Fang, "Sleep deprivation impairs spatial memory and decreases extracellular signal-regulated kinase phosphorylation in the hippocampus," *Brain Research*, vol. 1018, no. 1, pp. 38–47, 2004.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

