Corrigendum

Corrigendum to “Transmission Model of Hepatitis B Virus with the Migration Effect”

Muhammad Altaf Khan,¹ Saeed Islam,¹ Muhammad Arif,¹ and Zahoor ul Haq²

¹Department of Mathematics, Abdul Wali Khan University, Mardan, Khyber Pakhtunkhwa, Pakistan

²Department of Management Sciences, Abdul Wali Khan University, Mardan, Khyber Pakhtunkhwa, Pakistan

Correspondence should be addressed to Muhammad Altaf Khan; altafdir@gmail.com

Received 13 March 2016; Accepted 3 October 2016

Copyright © 2016 Muhammad Altaf Khan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Hepatitis B is one of the major public health problems in the world. It is an infection that causes the liver disease. The aim of this note is to provide corrections to the basic reproduction number $\mathcal{R}_0$ in [1] as described in the letter to the editor [2, 3]. In view of [2, 3], the model is reconstructed and all their mathematical results are computed.

2. Model Correction

In view of [2, 3], we decided that the product of F and V^{-1} , that is, FV^{-1} , is incorrect. If the product FV^{-1} is incorrect, then the obtained basic reproduction number in [1] is incorrect. The incorrectness of the basic reproduction number affects the stability (local and global), which is to be addressed again. Further, in light of [2], the model is valid biologically. By observing all these assumptions in [2, 3], we decide to reformulate the model by making the following changes to the model revised; we add the parameter α , a rate from migrated class to susceptible, and ξ is the rate of flow from exposed to migrated classes to the model published in [1]. The complete flow diagram of our new improved model can be seen in Figure 1. For complete details of each class with their parameters description, we refer the reader to [1], except α and ξ . Adding the above parameters we obtain the following improved model:

$$\begin{aligned}\frac{dS(t)}{dt} &= \delta\pi(1 - \eta C(t)) - (\delta + p)S(t) \\ &\quad - \beta(A(t) + \kappa C(t))S(t) + \delta_o V(t) \\ &\quad + \alpha M(t),\end{aligned}$$

$$\begin{aligned}\frac{dE(t)}{dt} &= \beta(A(t) + \kappa C(t))S(t) - (\delta + \gamma_1 + \xi)E(t) \\ &\quad + \delta\pi\eta C(t) + \mu_1 M(t),\end{aligned}$$

$$\frac{dA(t)}{dt} = \gamma_1 E(t) - (\delta + \gamma_2)A(t) + \mu_2 M(t),$$

$$\frac{dC(t)}{dt} = q\gamma_2 A(t) - (\delta + \gamma_3)C(t),$$

$$\begin{aligned}\frac{dV(t)}{dt} &= \gamma_3 C(t) + (1 - q)\gamma_2 A(t) - (\delta_o + \delta)V(t) \\ &\quad + \delta(1 - \pi) + pS(t),\end{aligned}$$

$$\frac{dM(t)}{dt} = \xi E(t) - (\mu_1 + \mu_2 + \delta + \alpha)M(t),$$

(1)

being subject to the initial conditions

$$S(0) \geq 0,$$

$$E(0) \geq 0,$$

$$A(0) \geq 0,$$

$$C(0) \geq 0,$$

$$V(0) \geq 0,$$

$$M(0) \geq 0.$$

(2)

FIGURE 1: The complete flow diagram of hepatitis B virus transmission model.

The model published in [1] should be replaced by (1) as well as the results. Assume $S(t) + E(t) + A(t) + C(t) + V(t) + M(t) = 1$; we obtain the following reduced model for (1):

$$\begin{aligned} \frac{dS(t)}{dt} = & \delta\pi + \delta_o - (\delta\pi\eta + \delta_o)C(t) \\ & - (\delta + p + \delta_o)S(t) \\ & - \beta(A(t) + \kappa C(t))S(t) \\ & - \delta_o(E(t) + A(t) + M(t)) + \alpha M(t), \end{aligned}$$

$$\begin{aligned} \frac{dE(t)}{dt} = & \beta(A(t) + \kappa C(t))S(t) - (\delta + \gamma_1 + \xi)E(t) \\ & + \delta\pi\eta C(t) + \mu_1 M(t), \\ \frac{dA(t)}{dt} = & \gamma_1 E(t) - (\delta + \gamma_2)A(t) + \mu_2 M(t), \\ \frac{dC(t)}{dt} = & q\gamma_2 A(t) - (\delta + \gamma_3)C(t), \\ \frac{dM(t)}{dt} = & \xi E(t) - (\mu_1 + \mu_2 + \delta + \alpha)M(t). \end{aligned} \quad (3)$$

Let

$$\begin{aligned} \Gamma = & \left\{ (S, E, A, C, M) \in \mathfrak{R}_+^5 \mid S(t) \leq \frac{\delta\pi + \delta_o}{\delta + \delta_o + p}, S + E \right. \\ & \left. + A + C + M \leq \frac{\delta\pi + \delta_o}{\delta + \delta_o} \right\}. \end{aligned} \quad (4)$$

Here Γ is a positively invariant set. All the solutions inside Γ are our main focus.

3. Equilibria and Basic Reproduction Number

System (3) has the disease-free equilibrium at $D_o = (S^o, 0, 0, 0, 0)$; that is, $S^o = (\delta\pi + \delta_o)/(\delta + \delta_o + p)$. At $D_1 = (S^*, E^*, A^*, C^*, M^*)$, the endemic equilibria of system (3) are given by

$$\begin{aligned} S^* = & \frac{(\delta + \gamma_1 + \xi)(\delta + \gamma_2)(\delta + \gamma_3)(\alpha + \delta + \mu_1 + \mu_2) - [\delta\pi\eta q\gamma_2(\xi\mu_2 + \gamma_1(\alpha + \delta + \mu_1 + \mu_2)) + \xi\mu_1(\delta + \gamma_2)(\delta + \gamma_3)]}{\beta((\delta + \gamma_3) + q\kappa\gamma_2)(\xi\mu_2 + \gamma_1(\alpha + \delta + \mu_1 + \mu_2))}, \\ E^* = & \frac{(\delta + \gamma_2)(\alpha + \delta + \mu_1 + \mu_2)A^*}{(\gamma_1(\alpha + \delta + \mu_1 + \mu_2) + \mu_2\xi)}, \\ C^* = & \frac{q\gamma_2 A^*}{(\delta + \gamma_3)}, \\ M^* = & \frac{\xi(\delta + \gamma_2)A^*}{(\xi\mu_2 + \gamma_1(\alpha + \delta + \mu_1 + \mu_2))}. \end{aligned} \quad (5)$$

The basic reproduction number $\mathcal{R}_0$ for system (3) can be obtained by using the method in [4]. We obtain the following matrices:

$$F = \begin{pmatrix} 0 & \beta S^o & \beta \kappa S^o & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\begin{aligned} V = & \begin{pmatrix} (\delta + \xi + \gamma_1) & 0 & -\delta\pi\eta & -\mu_1 \\ -\gamma_1 & (\delta + \gamma_2) & 0 & -\mu_2 \\ 0 & -q\gamma_2 & (\delta + \gamma_3) & 0 \\ -\xi & 0 & 0 & (\alpha + \delta + \mu_1 + \mu_2) \end{pmatrix}. \end{aligned} \quad (6)$$

The basic reproduction number $\mathcal{R}_0$ for (3) is given by

$$\mathcal{R}_0 = \frac{[\beta S^o(\delta + \gamma_3 + \kappa q\gamma_2) + \delta\pi\eta q\gamma_2](\mu_2\xi + \gamma_1(\alpha + \delta + \mu_1 + \mu_2)) + \mu_1\xi(\delta + \gamma_2)(\delta + \gamma_3)}{(\delta + \gamma_1 + \xi)(\delta + \gamma_2)(\delta + \gamma_3)(\alpha + \delta + \mu_1 + \mu_2)}. \quad (7)$$

Define

$$\begin{aligned}\mathcal{R}_1 &= \frac{[\beta(\delta\pi + \delta_o)(\delta + \gamma_3 + \kappa q\gamma_2) + \delta\pi q\gamma_2(\delta + \delta_o + p)]\gamma_1}{(\delta + \gamma_1 + \xi)(\delta + \gamma_2)(\delta + \gamma_3)(\delta + \delta_o + p)}, \\ \mathcal{R}_2 &= \frac{\beta(\delta\pi + \delta_o)(\delta + \gamma_3 + \kappa q\gamma_2)\xi\mu_2 + \xi(\delta + \delta_o + p)[\delta\pi q\gamma_2\mu_2 + \mu_1(\delta + \gamma_2)(\delta + \gamma_3)]}{(\delta + \gamma_1 + \xi)(\delta + \gamma_2)(\delta + \gamma_3)(\delta + \delta_o + p)(\alpha + \delta + \mu_1 + \mu_2)},\end{aligned}\quad (8)$$

where $S^o = (\delta\pi + \delta_o)/(\delta + \delta_o + p)$ and $\mathcal{R}_0 = \mathcal{R}_1 + \mathcal{R}_2$.

4. Stability of Disease-Free Equilibrium (DFE)

In this section, we discuss the local and global stability of system (3) at disease-free equilibrium. We have the following results.

Theorem 1. *The disease-free equilibrium D_o of system (3) is locally asymptotically stable if $\mathcal{R}_0 < 1$ and the conditions $(\gamma_2 + \delta)(\gamma_3 + \delta) > \beta S^o \gamma_1$ and $(\gamma_2 + \delta) > \beta S^o$ are satisfied; otherwise, they become unstable.*

Proof. At the disease-free equilibrium D_o the corresponding Jacobian matrix $J_o(\zeta)$ of system (3) is computed as follows:

$$J_o(\zeta) = \begin{pmatrix} -(\delta + \delta_o + p) & -\delta_o & -(\beta S^o + \delta_o) & -T_2 & -\delta_o + \alpha \\ 0 & -(\delta + \gamma_1 + \xi) & \beta S^o & \beta\kappa S^o + \delta\pi\eta & \mu_1 \\ 0 & \gamma_1 & -(\delta + \gamma_2) & 0 & \mu_2 \\ 0 & 0 & q\gamma_2 & -(\delta + \gamma_3) & 0 \\ 0 & \xi & 0 & 0 & -(\mu_1 + \mu_2 + \delta + \alpha) \end{pmatrix}, \quad (9)$$

where $T_2 = (\delta\pi\eta + \beta\kappa S^o + \delta_o)$. The root $-(\delta + \delta_o + p)$ of $J_o(\zeta)$ is clearly negative; the other roots can be obtained from the following equation:

$$\lambda^4 + c_1\lambda^3 + c_2\lambda^2 + c_3\lambda + c_4 = 0, \quad (10)$$

where

$$\begin{aligned}c_1 &= (\gamma_2 + \delta) + (\gamma_3 + \delta) + (\alpha + \delta + \mu_1 + \mu_2) \\ &\quad + (\gamma_1 + \delta + \xi), \\ c_2 &= (\alpha + \delta + \mu_2)(\gamma_1 + \delta + \xi) + \mu_1(\gamma_1 + \delta) \\ &\quad + ((\gamma_2 + \delta)(\gamma_3 + \delta) - \beta\gamma_1 S^o) \\ &\quad + ((\gamma_2 + \delta) + (\gamma_3 + \delta)) \\ &\quad \cdot [(\alpha + \delta + \mu_1 + \mu_2) + (\gamma_1 + \delta + \xi)], \\ c_3 &= [(\gamma_2 + \delta) + (\gamma_3 + \delta)]\mu_1(\gamma_1 + \delta) \\ &\quad + (\alpha + \delta + \mu_1 + \mu_2)((\gamma_2 + \delta)(\gamma_3 + \delta) - \beta\gamma_1 S^o) \\ &\quad + (\gamma_2 + \delta)((\alpha + \delta)(\gamma_1 + \delta + \xi) + \mu_2(\gamma_1 + \delta)) \\ &\quad + \mu_2\xi((\gamma_2 + \delta) - \beta S^o) + (\gamma_1 + \delta + \xi)(\gamma_3 + \delta) \\ &\quad \cdot [(\alpha + \delta + \mu_2) + (\gamma_2 + \delta)(1 - \mathcal{R}_1)],\end{aligned}$$

$$\begin{aligned}c_4 &= (\gamma_2 + \delta)(\gamma_3 + \delta)(\alpha + \delta + \mu_1 + \mu_2)(\gamma_1 + \delta + \xi) \\ &\quad \cdot (1 - \mathcal{R}_0).\end{aligned}\quad (11)$$

The eigenvalues of the characteristics equation have negative real part if the Routh-Hurwitz condition is satisfied; that is, $c_i, i = 1, 2, 3, 4$, with $c_1c_2c_3 - c_3^2 - c_1^2c_4 > 0$. Therefore, if $\mathcal{R}_0 < 1$ and $(\gamma_2 + \delta)(\gamma_3 + \delta) > \beta S^o \gamma_1$ and $(\gamma_2 + \delta) > \beta S^o$, we get $c_i > 0$ for $i = 1, 2, 3, 4$. Thus, at D_o the disease-free equilibrium of system (3) is locally asymptotically stable if the above conditions are satisfied. $\square$

Next, we show the global stability of DFE by using the method in [5].

Theorem 2. *The disease-free equilibrium of system (3) is globally asymptotically stable if $\mathcal{R}_0 < 1$.*

Proof. Consider $Y_1 = S, Y_2 = (E, A, C, M)^T$, and $Y = (Y_1, Y_2)$. The invariant domain Γ is clearly a compact positive set. We can present the subsystem $Y_1 = G_1(Y_1, 0)(Y_1 - Y_1^*)$ as

$$\dot{S} = -(\delta + \delta_o + p)\left(S - \frac{\delta\pi + \delta_o}{(\delta + \delta_o + p)}\right). \quad (12)$$

System (3) represents a linear system which is globally asymptotically stable at $Y_1^* = (\delta\pi + \delta_o)/(\delta + \delta_o + p)$.

The hypotheses in [5] are satisfied. The matrix $G_2(Y)$ is given by

$$G_2(Y) = \begin{pmatrix} -(\delta + \gamma_1 + \xi) & \beta S & \beta \kappa S + \delta \pi \eta & \mu_1 \\ \gamma_1 & -(\delta + \gamma_2) & 0 & \mu_2 \\ 0 & q\gamma_2 & -(\delta + \gamma_3) & 0 \\ \xi & 0 & 0 & -(\mu_1 + \mu_2 + \delta + \alpha) \end{pmatrix}. \quad (13)$$

Following the hypothesis H_3 in [5], for any $Y \in \Gamma$ the above matrix $G_2(Y)$ is Metzler and irreducible. Further, we check the fourth condition H_4 in [5]. There is a maximum in Γ if

$S = (\delta\pi + \delta_o)/(\delta + \delta_o + p)$. This corresponds to disease-free equilibrium and the maximum $G_2(\bar{Y})$ is given by

$$G_2(\bar{Y}) = \begin{pmatrix} -(\delta + \gamma_1 + \xi) & \beta S^o & \beta \kappa S^o + \delta \pi \eta & \mu_1 \\ \gamma_1 & -(\delta + \gamma_2) & 0 & \mu_2 \\ 0 & q\gamma_2 & -(\delta + \gamma_3) & 0 \\ \xi & 0 & 0 & -(\mu_1 + \mu_2 + \delta + \alpha) \end{pmatrix}. \quad (14)$$

The last hypothesis H_5 requires that $\alpha(G_2(\bar{Y})) \leq 0$. We write $G_2(\bar{Y})$ in block form as

$$G_2(\bar{Y}) = \begin{pmatrix} B_1 & B_2 \\ B_3 & B_4 \end{pmatrix}, \quad (15)$$

where

$$B_1 = -(\delta + \gamma_1 + \xi),$$

$$B_2 = (\beta S^o, \beta \kappa S^o + \delta \pi \eta, \mu_1),$$

$$B_3 = \begin{pmatrix} \gamma_1 \\ 0 \\ \xi \end{pmatrix},$$

$$B_4 = \begin{pmatrix} -(\delta + \gamma_2) & 0 & \mu_2 \\ q\gamma_2 & -(\delta + \gamma_3) & 0 \\ 0 & 0 & -(\mu_1 + \mu_2 + \delta + \alpha) \end{pmatrix}. \quad (16)$$

Clearly B_1 is stable Metzler matrix, so we can write $\alpha(B_4 - B_3 B_1^{-1} B_2) \leq 0$. Let $K = B_4 - B_3 B_1^{-1} B_2$.

$$K = \begin{pmatrix} -\left((\delta + \gamma_2) - \frac{\gamma_1 \beta S^o}{(\delta + \gamma_1 + \xi)}\right) & \frac{\gamma_1 (\beta \kappa S^o + \delta \pi \eta)}{(\delta + \gamma_1 + \xi)} & \mu_2 - \frac{\gamma_1 \mu_1}{(\delta + \gamma_1 + \xi)} \\ q\gamma_2 & -(\delta + \gamma_3) & 0 \\ \frac{\xi \beta S^o}{(\delta + \gamma_1 + \xi)} & \frac{\xi (\beta \kappa S^o + \delta \pi \eta)}{(\delta + \gamma_1 + \xi)} & -\left((\mu_1 + \mu_2 + \delta + \alpha) - \frac{\xi \mu_1}{(\delta + \gamma_1 + \xi)}\right) \end{pmatrix}. \quad (17)$$

The characteristics equation of K are given by

$$\lambda^3 + l_1 \lambda^2 + l_2 \lambda + l_3 = 0, \quad (18)$$

where

$$l_1 = (\gamma_1 + \delta + \xi) [(\gamma_2 + \delta) + (\gamma_3 + \delta) + (\alpha + \delta + \mu_2) + \mu_1 (\gamma_1 + \delta) - \beta \gamma_1 S^o],$$

$$l_2 = (\gamma_2 + \delta) (\gamma_3 + \delta) (1 - \mathcal{R}_1) + ((\gamma_2 + \delta) + (\gamma_3 + \delta)) \left((\alpha + \delta + \mu_2) + \frac{\mu_1 (\gamma_1 + \delta)}{(\gamma_1 + \delta + \xi)} \right) + \frac{\beta S^o}{(\gamma_1 + \delta + \xi)^2} (2\gamma_1 \mu_1 \xi - (\gamma_1 + \delta + \xi) \cdot (\xi \mu_2 + \gamma_1 (\alpha + \delta + \mu_1 + \mu_2) (\gamma_1 + \delta + \xi))),$$

$$l_3 = \frac{2\mu_1\xi\gamma_1}{(\gamma_1 + \delta + \xi)^2} [\eta\gamma_2 (\delta\pi\eta + \beta\kappa S^o) + (\gamma_3 + \delta) \beta S^o] + (\gamma_2 + \delta) (\gamma_3 + \delta) \times (\alpha + \delta + \mu_1 + \mu_2) (1 - \mathcal{R}_0). \quad (19)$$

The Routh-Hurwitz criteria ensure that the above characteristics equation has three negative eigenvalues if $\mathcal{R}_0 < 1$ and the condition $l_1 l_2 - l_3 > 0$. All the conditions in [5] are satisfied. Thus, we conclude that system (3) at disease-free equilibrium is globally asymptotically stable. $\square$

5. Stability EE

In this section, we determine the local and global stability of (3) at endemic equilibrium.

Theorem 3. *If $\mathcal{R}_0 > 1$, then model (3) at endemic equilibrium is locally asymptotically stable.*

Proof. The Jacobian matrix J^* computed at D_1 is given by

$$J^* = \begin{pmatrix} -z_1 & -\delta_o & -\beta S^* - \delta_o & -\delta\pi\eta - \beta\kappa S^* - \delta_o & -\delta_o + \alpha \\ z_2 & -(\delta + \gamma_1 + \xi) & \beta S^* & \delta\pi\eta + \beta\kappa S^* & \mu_1 \\ 0 & \gamma_1 & -(\delta + \gamma_2) & 0 & \mu_2 \\ 0 & 0 & \eta\gamma_2 & -(\delta + \gamma_3) & 0 \\ 0 & \xi & 0 & 0 & -(\alpha + \delta + \mu_1 + \mu_2) \end{pmatrix}. \quad (20)$$

The characteristics equation at D_1 is

$$\lambda^5 + g_1\lambda^4 + g_2\lambda^3 + g_3\lambda^2 + g_4\lambda + g_5 = 0, \quad (21)$$

where

$$\begin{aligned} g_1 &= 4\delta + z_1 + \alpha + \gamma_1 + \gamma_2 + \gamma_3 + \mu_1 + \mu_2 + \xi, \\ g_2 &= (\alpha + \delta + \mu_1 + \mu_2) [(\gamma_1 + \delta) + (\gamma_2 + \delta) \\ &\quad + (\gamma_3 + \delta)] + \xi(\alpha + \delta + \mu_2) + [(\alpha + \delta + \mu_1 + \mu_2) \\ &\quad + (\gamma_1 + \delta + \xi) + (\gamma_2 + \delta) + (\gamma_3 + \delta)] z_1 + (\gamma_2 + \delta) \\ &\quad \cdot [(\gamma_1 + \delta + \xi) + (\gamma_3 + \delta)] + (\gamma_3 + \delta) (\gamma_1 + \delta + \xi) \\ &\quad - (z_3 + \delta_o z_2), \\ g_3 &= (\alpha + \delta + \mu_1 + \mu_2) \\ &\quad \cdot [(\gamma_1 + \delta + \xi) ((\gamma_3 + \delta) + (\gamma_2 + \delta))] \\ &\quad + (\gamma_2 + \delta) (\gamma_3 + \delta) - \delta_o z_2] + z_2 \xi [(\alpha - \delta_o) \\ &\quad - (\gamma_3 + \delta) \delta_o - (\gamma_2 + \delta) \delta_o] + (\gamma_2 + \delta) \\ &\quad \cdot z_1 [(\gamma_1 + \delta + \xi) + (\gamma_3 + \delta)] + (\alpha + \delta + \mu_1 + \mu_2) \\ &\quad \cdot z_1 [(\gamma_1 + \delta + \xi) + (\gamma_3 + \delta) + (\gamma_2 + \delta)] + (\gamma_3 + \delta) \\ &\quad \cdot (\gamma_1 + \delta + \xi) z_1 - z_1 (\mu_1 \xi + z_3) - \gamma_1 z_2 z_4 \\ &\quad - \mu_1 \xi [(\gamma_3 + \delta) + (\gamma_2 + \delta)] + (\gamma_2 + \delta) (\gamma_3 + \delta) (\gamma_1 \\ &\quad + \delta + \xi) - z_5 - z_3 (\alpha + \delta + \mu_1 + \mu_2) - z_3 (\gamma_3 + \delta) \\ &\quad - z_6, \end{aligned}$$

$$\begin{aligned} g_4 &= (\gamma_2 + \delta) (\gamma_3 + \delta) (\alpha + \delta + \mu_1 + \mu_2) (\gamma_1 + \delta + \xi) \\ &\quad - (\gamma_2 + \delta) \delta_o (\alpha + \delta + \mu_1 + \mu_2) z_2 - (\gamma_3 + \delta) \delta_o (\alpha \\ &\quad + \delta + \mu_1 + \mu_2) z_2 + \xi (\gamma_2 + \delta) z_2 (\alpha - \delta_o) + z_2 \xi (\gamma_3 \\ &\quad + \delta) (\alpha - \delta_o) - \mu_1 \xi (\gamma_2 + \delta) z_1 - \mu_1 \xi (\gamma_3 + \delta) z_1 \\ &\quad - (\gamma_2 + \delta) (\gamma_3 + \delta) \delta_o z_2 - z_3 (\alpha + \delta + \mu_1 + \mu_2) z_1 \\ &\quad + [(\gamma_2 + \delta) + (\gamma_3 + \delta)] (\alpha + \delta + \mu_1 + \mu_2) (\gamma_1 + \delta \\ &\quad + \xi) z_1 - \gamma_1 (\gamma_3 + \delta) z_2 z_4 - z_1 z_5 + (\gamma_2 + \delta) (\gamma_3 + \delta) \\ &\quad \cdot (\alpha + \delta + \mu_1 + \mu_2) z_1 + (\gamma_2 + \delta) (\gamma_3 + \delta) (\gamma_1 + \delta \\ &\quad + \xi) z_1 - \mu_2 \xi z_2 z_4 - z_3 (\gamma_3 + \delta) z_1 - z_1 z_6 - z_1 z_5 \\ &\quad - \gamma_1 (\alpha + \delta + \mu_1 + \mu_2) z_2 z_4 + \mu_1 \xi (\gamma_2 + \delta) (\gamma_3 + \delta) \\ &\quad - (\alpha + \delta + \mu_1 + \mu_2) z_5 - z_7 - z_3 (\gamma_3 + \delta) (\alpha + \delta \\ &\quad + \mu_1 + \mu_2) - z_6 (\gamma_3 + \delta), \\ g_5 &= -(\gamma_2 + \delta) (\gamma_3 + \delta) \delta_o (\alpha + \delta + \mu_1 + \mu_2) z_2 \\ &\quad + \xi (\gamma_2 + \delta) (\gamma_3 + \delta) z_2 (\alpha - \delta_o) + (\gamma_2 + \delta) (\gamma_3 + \delta) \\ &\quad \cdot (\alpha + \delta + \mu_1 + \mu_2) (\gamma_1 + \delta + \xi) z_1 - \mu_1 \xi (\gamma_2 + \delta) \\ &\quad \cdot (\gamma_3 + \delta) z_1 - (\alpha + \delta + \mu_1 + \mu_2) z_5 z_1 - z_1 z_7 \\ &\quad - z_3 (\gamma_3 + \delta) (\alpha + \delta + \mu_1 + \mu_2) z_1 - z_6 z_1 \\ &\quad - \gamma_1 z_2 \eta \gamma_2 (\alpha + \delta + \mu_1 + \mu_2) z_8 - \mu_2 \xi \eta \gamma_2 z_2 z_8 \\ &\quad - \gamma_1 (\gamma_3 + \delta) (\alpha + \delta + \mu_1 + \mu_2) z_2 z_4 - \mu_2 \xi (\gamma_3 + \delta) \\ &\quad \cdot z_2 z_4, \end{aligned} \quad (22)$$

where $z_1 = (\delta + \delta_o + p) + \beta(A^* + \kappa C^*)$, $z_2 = \beta(A^* + \kappa C^*)$, $z_3 = \beta\gamma_1 S^*$, $z_4 = (\delta_o + \beta S^*)$, $z_5 = \gamma_1 q\gamma_2 (\pi\delta\eta + \beta\kappa S^*)$, $z_6 = \beta\mu_2 \xi S^*$, $z_7 = \mu_2 \xi q\gamma_2 (\pi\delta\eta + \beta\kappa S^*)$, and $z_8 = (\pi\delta\eta + \delta_o + \beta\kappa S^*)$. The characteristics equation will give negative roots if the following conditions are satisfied:

- (i) $g_1, g_2, g_3, g_4, g_5 > 0$,
- (ii) $g_1 g_2 - g_3 > 0$,
- (iii) $g_4 g_5 (g_3 (g_1 g_2 - g_3) - g_1^2 g_4) > 0$,
- (iv) $g_5^2 (2g_1 g_4 - g_2 (g_1 g_2 - g_3) - g_5) > 0$.

Conditions (i) and (ii) are easy to satisfy. If conditions (iii) and (iv) are satisfied; then the characteristics equation given above will give negative eigenvalues. Thus, it follows from Routh-Hurwitz criteria that system (3) is locally asymptotically stable at D_1 . $\square$

Next result shows the global stability of the endemic equilibrium D_1 of system (3) for special case when $\mu_1 = 0$ and $\alpha = 0$. We have the following result.

Theorem 4. *The endemic equilibrium D_1 of system (3) is globally asymptotically stable if condition (30) holds.*

Proof. We define the Lyapunov function in the following form:

$$\begin{aligned} \mathcal{V} = & w_1 \left(S - S^* - S^* \log \frac{S}{S^*} \right) \\ & + w_2 \left(E - E^* - E^* \log \frac{E}{E^*} \right) + w_3 A + w_4 C \\ & + w_5 M. \end{aligned} \quad (23)$$

The time durative of $\mathcal{V}$ is

$$\begin{aligned} \mathcal{V}' = & w_1 \left(1 - \frac{S^*}{S} \right) S' + w_2 \left(1 - \frac{E^*}{E} \right) E' + w_3 A' \\ & + w_4 C' + w_5 M'. \end{aligned} \quad (24)$$

The coefficients $w_1, \dots, w_5$ are positive and will be determined later. At endemic steady state the first equation of (3) is given by

$$\begin{aligned} \delta\pi + \delta_o = & \delta\pi\eta C^* + (\delta + p) S^* + \beta(A^* + \kappa C^*) S^* \\ & + \delta_o X^*. \end{aligned} \quad (25)$$

Using (25) and the S equation (3) we obtain

$$\begin{aligned} w_1 \left(1 - \frac{S^*}{S} \right) S' = & w_1 \left(1 - \frac{S^*}{S} \right) [\delta\pi\eta C^* + (\delta + p) S^* \\ & + \beta(A^* + \kappa C^*) S^* + \delta_o X^*] - w_1 [\delta\pi\eta C \\ & + (\delta + p) S + \beta(A + \kappa C) S + \delta_o X] + w_1 \\ & \cdot \frac{S^*}{S} [\delta\pi\eta C(t) + (\delta + p) S(t) \\ & + \beta(A(t) + \kappa C(t)) S(t) + \delta_o X]. \end{aligned} \quad (26)$$

Similarly,

$$\begin{aligned} w_2 \left(1 - \frac{E^*}{E} \right) E' = & w_2 [\beta(A(t) + \kappa C(t)) S(t) \\ & - (\delta + \gamma_1 + \xi) E(t) + \delta\pi\eta C(t)] - w_2 \frac{E^*}{E} \beta(A(t) \\ & + \kappa C(t)) S(t) + w_2 \beta(A^* + \kappa C^*) S^* + w_2 \delta\pi\eta C^* \\ & + \delta\pi\eta C(t) \frac{E^*}{E}. \end{aligned} \quad (27)$$

Using the values of (26) and (27) and the last three equations of system (3) and substituting in (24) we obtain

$$\begin{aligned} \mathcal{V}' = & w_1 \left(1 - \frac{S^*}{S} \right) [\delta\pi\eta C^* + (\delta + p) S^* \\ & + \beta(A^* + \kappa C^*) S^* + \delta_o X^*] - w_1 \delta\pi\eta C - w_1 (\delta \\ & + p) S - w_1 \beta(A + \kappa C) S - w_1 \delta_o X + w_1 \frac{S^*}{S} \\ & \cdot \delta\pi\eta C(t) + w_1 (\delta + p) S + w_1 \beta(A + \kappa C) S \\ & + w_1 \delta_o X + w_2 [\beta(A + \kappa C) S - (\delta + \gamma_1 + \xi) E \\ & + \delta\pi\eta C] - w_2 \frac{E^*}{E} \beta(A + \kappa C) S + w_2 \beta(A^* + \kappa C^*) \\ & \cdot S^* + w_2 \delta\pi\eta C^* + w_2 \delta\pi\eta C \frac{E^*}{E}. \end{aligned} \quad (28)$$

After implication we obtain

$$\begin{aligned} \mathcal{V}' = & -\frac{q\gamma_2 (\mu_2 \xi + \gamma_1 (\mu_1 + \mu_2 + \delta + \alpha))}{(\mu_1 + \mu_2 + \delta + \alpha) (\delta + \gamma_1 + \xi)} (\delta + p) \\ & \cdot \frac{(S - S^*)^2}{S} - \frac{q\gamma_2 (\mu_2 \xi + \gamma_1 (\mu_1 + \mu_2 + \delta + \alpha))}{(\mu_1 + \mu_2 + \delta + \alpha) (\delta + \gamma_1 + \xi)} \\ & \cdot \beta S^* A^* \times \left(-2 + \frac{S^*}{S} + \frac{A S E^*}{A^* S^* E} \right) \\ & - \frac{q\gamma_2 (\mu_2 \xi + \gamma_1 (\mu_1 + \mu_2 + \delta + \alpha))}{(\mu_1 + \mu_2 + \delta + \alpha) (\delta + \gamma_1 + \xi)} \beta S^* \kappa C^* \left(-2 \right. \\ & \left. + \frac{S^*}{S} + \frac{C S E^*}{C^* S^* E} \right) \\ & - \frac{q\gamma_2 (\mu_2 \xi + \gamma_1 (\mu_1 + \mu_2 + \delta + \alpha))}{(\mu_1 + \mu_2 + \delta + \alpha) (\delta + \gamma_1 + \xi)} \left(-2 + \frac{S^*}{S} \right. \\ & \left. + \frac{C}{C^*} \frac{E^*}{E} - \frac{C}{C^*} \frac{S^*}{S} \right) \delta\pi\eta C^* - (\delta + \gamma_2) (\delta + \gamma_3) C \\ & - \frac{q\gamma_2 (\mu_2 \xi + \gamma_1 (\mu_1 + \mu_2 + \delta + \alpha))}{(\mu_1 + \mu_2 + \delta + \alpha) (\delta + \gamma_1 + \xi)} \delta_o X^* \left(-1 + \frac{S^*}{S} \right. \\ & \left. + \frac{X}{X^*} - \frac{X}{X^*} \frac{S^*}{S} \right), \end{aligned} \quad (29)$$

FIGURE 2: Population behavior of individuals when $\mu_1 = 0.1, \mu_2 = 0.1$.

FIGURE 5: Population behavior of individuals when $\mu_1 = 0.1, \mu_2 = 0.2$.

FIGURE 3: Population behavior of individuals when $\mu_1 = 0.2, \mu_2 = 0.2$.

FIGURE 6: Population behavior of carriers individuals: solid line, $\pi = 1, p = 0$; dashed $\pi = 0.1, p = 0.1$.

FIGURE 4: Population behavior of individuals when $\mu_1 = 0.01, \mu_2 = 0.02$.

FIGURE 7: Population behavior of carriers individuals: solid line $p = 0, \mu_1 = 0.1, \mu_2 = 0.1$; dashed $p = 0.1, \mu_1 = 0.01, \mu_2 = 0.01$.

FIGURE 8: Population behavior of susceptible, exposed, acute, and carriers individuals: solid line $\mu_1 = 0.2, \mu_2 = 0.2$; dashed $\mu_1 = 0.002, \mu_2 = 0.002$.

where the constants $w_1, \dots, w_5$ are chosen as $w_4 = (\delta + \gamma_2)$, $w_3 = q\gamma_2$, $w_5 = q\gamma_2\mu_2/(\mu_1 + \mu_2 + \delta + \alpha)$, $w_2 = w_1 = q\gamma_2(\mu_2\xi + \gamma_1(\mu_1 + \mu_2 + \delta + \alpha))/(\mu_1 + \mu_2 + \delta + \alpha)(\delta + \gamma_1 + \xi)$, and $X = S + E + A + C + M$, $X^* = S^* + E^* + A^* + C^* + M^*$.

Equation (29) $\mathcal{V}' \leq 0$ if the following inequalities are satisfied:

$$\begin{aligned}
 &\left(-2 + \frac{S^*}{S} + \frac{C}{C^*} \frac{E^*}{E} - \frac{C}{C^*} \frac{S^*}{S}\right) \geq 0, \\
 &\left(-2 + \frac{S^*}{S} + \frac{CSE^*}{C^*S^*E}\right) \geq 0, \\
 &\left(-2 + \frac{S^*}{S} + \frac{C}{C^*} \frac{E^*}{E} - \frac{C}{C^*} \frac{S^*}{S}\right) \geq 0, \\
 &\left(-1 + \frac{S^*}{S} + \frac{X}{X^*} - \frac{X}{X^*} \frac{S^*}{S}\right) \geq 0.
 \end{aligned} \tag{30}$$

The endemic equilibrium D_1 of system (3) is said to be globally asymptotically stable if condition (30) holds. $\square$

6. Numerical Simulation

This section deals with the numerical solution of model (3). The numerical results for model (3) are presented in Figures 2–8. In this paper, the value for α and ξ is taken between 0 and 1. The rest of the parameters values are taken from [1], except those mentioned in the figures. Figure 1 shows the population behavior of incidentals when $\mu_1 = 0.1$ and $\mu_2 = 0.1$. Figure 3 represents the population of incisiveness when $\mu_1 = 0.2$ and $\mu_2 = 0.2$. In Figures 2 and 3, the individuals of carriers decrease sharply. For $\mu_1 = 0.001$ and $\mu_2 = 0.02$, we present Figure 4. Figure 5 is population behavior of individuals when $\mu_1 = 0.1$ and $\mu_2 = 0.2$. Figures 6 and 7, respectively, represent the population of carriers individuals for different values of parameters. The population behavior of susceptible, exposed, acute, and carriers individuals for different parameters is presented in Figure 8. The numerical results from Figures 2 to 8 show when there is a decrease in the value of suggested parameters, and the population of individuals in the host decreases sharply.

7. Conclusion

In this corrigendum, we make all the necessary changes to the published paper [1], which are highlighted in the comment papers [2, 3]. We added the parameter α , a rate from migrated class to susceptible, and ξ is the rate of flow from exposed to migrated classes. Further, the basic reproduction number has been investigated. The mathematical results for the revised model are presented successfully.

References

- [1] M. A. Khan, S. Islam, M. Arif, and Z. Ul Haq, "Transmission model of hepatitis b virus with the migration effect," *BioMed Research International*, vol. 2013, Article ID 150681, 10 pages, 2013.
- [2] A. Zeb and G. Zaman, "Comment on 'transmission model of hepatitis B virus with migration effect,'" *BioMed Research International*, vol. 2015, Article ID 492513, 4 pages, 2015.
- [3] A. A. Lashari, "Comment on 'transmission model of Hepatitis B virus with the migration effect,'" *BioMed Research International*, vol. 2015, Article ID 469240, 2 pages, 2015.
- [4] P. van den Driessche and J. Watmough, "Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission," *Mathematical Biosciences*, vol. 180, pp. 29–48, 2002.
- [5] J. C. Kamgang and G. Sallet, "Computation of threshold conditions for epidemiological models and global stability of the disease-free equilibrium (DFE)," *Mathematical Biosciences*, vol. 213, no. 1, pp. 1–12, 2008.