Research Article

Overexpression of *CaAPX* Induces Orchestrated Reactive Oxygen Scavenging and Enhances Cold and Heat Tolerances in Tobacco

Jiangying Wang,^{1,2} Bin Wu,¹ Hengfu Yin,¹ Zhengqi Fan,¹ Xinlei Li,¹ Sui Ni,³ Libo He,^{1,4} and Jiyuan Li¹

¹Research Institute of Subtropical Forestry, Chinese Academy of Forestry, Zhejiang Provincial Key Laboratory of Tree Breeding, Hangzhou, Zhejiang 311400, China

²Lianyungang Academy of Agricultural Sciences, Flower Research Center, Lianyungang, Jiangsu 222000, China

³School of Marine Sciences, Ningbo University, Ningbo, Zhejiang 315211, China

⁴College of Horticulture and Landscape Architecture, Hunan Agricultural University, Changsha, Hunan 410128, China

Correspondence should be addressed to Hengfu Yin; hfyin@sibs.ac.cn, Libo He; helibo@hunau.net, and Jiyuan Li; jiyuan.li@126.com

Received 7 September 2016; Accepted 6 February 2017; Published 13 March 2017

Academic Editor: Atanas Atanasov

Copyright © 2017 Jiangying Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ascorbate peroxidase (APX) acts indispensably in synthesizing L-ascorbate (AsA) which is pivotal to plant stress tolerance by detoxifying reactive oxygen species (ROS). Enhanced activity of APX has been shown to be a key step for genetic engineering of improving plant tolerance. However it needs a deeper understanding on the maintenance of cellular ROS homeostasis in response to stress. In this study, we identified and characterized an APX (*CaAPX*) gene from *Camellia azalea*. Quantitative real-time PCR (qRT-PCR) analysis showed that *CaAPX* was expressed in all tissues and peaked in immature green fruits; the expression levels were significantly upregulated upon cold and hot stresses. Transgenic plants displayed marked enhancements of tolerance under both cold and heat treatments, and plant growth was correlated with *CaAPX* expression levels. Furthermore, we monitored the activities of several ROS-scavenging enzymes including Cu/Zn-SOD, CAT, DHAR, and MDHAR, and we showed that stress tolerance was synchronized with elevated activities of ROS-scavenging. Moreover, gene expression analysis of ROS-scavenging enzymes revealed a role of *CaAPX* to orchestrate ROS signaling in response to temperature stresses. Overall, this study presents a comprehensive characterization of cellular response related to *CaAPX* expression and provides insights to breed crops with high temperature tolerances.

1. Introduction

Temperature stress, such as heat, cold, or freezing, causes an increase in reactive oxygen species (ROS) levels and induces the oxidative stress in plant cells [1]. The elevation of reactive ROS levels can cause deleterious damage to organelle function and cellular metabolism or leads to programmed cell death [2]. To quickly remove excessive ROS, plants evolved a complex system containing nonenzymatic antioxidants, such as ascorbate (AsA), glutathione (GSH), flavonoids and carotenoids, and ROS-scavenging enzymes including superoxide dismutase (SOD), peroxidase (POD),

catalase (CAT), and enzymes in the water-water cycle [3]. The antioxidants and enzymes need to act synergistically to keep the homeostasis of cellular redox state. For example, the water-water cycle, also known as Halliwell-Asada or AsA-GSH cycle, includes the activity of APX, DHAR, GR, and MDHAR and was a pathway involved in scavenging superoxide radicals and H₂O₂ [4].

There is an effective regulatory network containing multiple pathways in plant cell to deal with cellular ROS efficiently. The regulation of antioxidants and enzymatic activities was involved in a signaling transduction leading to changes of gene expression levels. Recently the ROS were revealed as

central signaling components to orchestrate the multiple scavenging genes expressions under various stressed conditions [5, 6]. Upon temperature stress, the cellular ROS levels were upregulated through the “temperature sensor” which amplified the signal to activate the ROS-scavenging enzymes [2, 7]. Heat shock proteins (HSPs) and heat shock transcription factors (HSFs) were thought to be responsible for temperature stress sensing [8, 9]. On the other hand, the NADPH oxidases (respiratory burst oxidase homologous, RBOHs) played a key role in generating ROS upon stress stimuli, which were involved in different signaling pathways [10, 11]. In Arabidopsis, functional analyses of HSF3 and HSF21 revealed that ROS-scavenging enzymes APX1 and APX2 were imminent targets which were required to protect against heat induced oxidative damage [12, 13]. In addition, APX could act interactively with RBOHD to generate systematic signals of ROS tolerance [10, 13].

Ascorbate peroxidase (APX) was found to be a key enzyme in the ascorbate-glutathione pathway to scavenge cellular H_2O_2 produced in various stressful conditions [5, 14–19]. When APX reduces hydrogen peroxide to water by utilizing AsA as an electron donor, monodehydroascorbate (MDA) accumulates. Meanwhile MDA is somewhat unstable and likely to be rereduced disproportionately into AsA and dehydroascorbate (DHA), and both MDA and DHA can be reduced back to AsA through the activity of reductases [20–22]. Many studies over the years have demonstrated that APX played a specific role in improving plant’s tolerance response to certain abiotic stresses [19, 23–25]. Enhanced expression of APX in transgenic plants could intensify their resistance to multiple environmental stresses through eliminating H_2O_2 [19, 23–27]. For example, APX overexpression improved chilling tolerances in rice and sweet potato [28, 29]; the transgenic potato plants with APX expression were more tolerant to high temperature stress [30, 31]. In addition, after exposure to some environmental stresses, the APX knockdown plants exhibited much more severe cellular injuries [32–34].

Although the genetic manipulation of ROS has been practiced in many plant species, our understanding of ROS removing and signaling remains to be expanded to better improve the plant performance by the genetic manipulation approach. Here, in order to investigate the potential improvements of temperature stresses in *Camellia* species, we cloned APX gene from *Camellia azalea* and investigated its expression pattern. We showed that *CaAPX* was expressed in all examined *Camellia* tissues and was quickly induced by the temperature stresses. Functional analysis in transgenic tobacco proved that overexpression of *CaAPX* enhanced plant performances in both cold and heat conditions. Furthermore, we demonstrated that overexpression of *CaAPX* altered the cellular content of AsA, MDA, and H_2O_2 concentrations and led to lower electrolyte leakage rate. Enzymatic activity analysis of APX, SOD, POD, and CAT and gene expression analysis of tobacco endogenous Cu/Zn-SOD, CAT, DHAR, and MDHAR genes revealed that *CaAPX* may have a central role in synchronizing cellular ROS-scavenging network. This work may provide useful information for engineering plants of high resistance to oxidative damage under temperature stresses.

2. Materials and Methods

2.1. Plant Materials and Treatments. *C. azalea* materials selected for present study including four-year-old grafted seedlings and two-year-old cutting seedlings were collected from a greenhouse of the Research Institute of Subtropical Forestry located in Fuyang (119°57' E, 30°04' N; Fuyang city, Zhejiang, China), growing under natural light condition with regular irrigation. And subculture aseptic tobacco seedlings with six true leaves (*Nicotiana tabacum* L., cv. SR1) were grown in a growth chamber at 25°C and 1200 Lux light (16 h light/8 h dark). For treatments, plants were exposed to 4°C low and 38°C high temperatures for 0, 2, 4, 8, and 12 h, respectively, in the growth chamber at 25°C and 1200 Lux light (16 h day/8 h night). The samples treated for 0 h were set as controls. Plant tissues of *C. azalea* seedlings and the forth leaves of treatment tobacco plants collected were immediately put into liquid nitrogen and stored at –80°C for later use.

2.2. Cloning of *CaAPX* Gene. Total RNA was extracted using the plant total RNA isolation Kit (Aidlab Biotech, China) for reverse transcription polymerase chain reaction (RT-PCR) (Takara, China). To isolate the APX gene, the 3',5'-RACE amplification Kit (Takara, China) was used to construct libraries from *C. azalea* young leaves. Two amplified fragments (gene-specific primers were listed in Table S1 (see Table S1 in Supplementary Material available online at <https://doi.org/10.1155/2017/4049534>)) were cloned into pGEM-Teasy vector (Promega, USA) for sequencing, and full-length cDNA sequence was assembled. The cDNA sequence was designated as *CaAPX* and submitted to GenBank (accession number KP635267) (<https://www.ncbi.nlm.nih.gov/genbank/>).

2.3. Quantitative Real-Time PCR. DNA traces of the total RNA were removed by DNase I (RNase-free) (TransGen Biotech, China). The cDNAs from different organs of *C. azalea* cutting seedlings on, respectively, different temperature stress levels were obtained by employing the PrimeScript™ RT reagent Kit (Perfect Real Time) (Takara, China). The *18S* RNA gene of *Camellias* was used as an internal control to normalize the amount of mRNA present in each sample. And all samples were amplified in four independent replicates by an Applied Biosystems 7300 Fast Real-Time PCR system using SYBR-Premix Ex Taq™ II (Tli RNaseH Plus) (Takara, China) for fluorescence detection and the expression data was analyzed using $2^{-\Delta\Delta C_t}$ method.

2.4. Construction of the Plant Expression Vector and Transformation of Tobacco. Full-length cDNA sequence of *CaAPX* gene was obtained by amplification of primers containing *Xba*I and *Bam*HI restriction sites (Table S1). The PCR product was integrated into pGEM-Teasy vector (Promega) and then cloned into vector pBI121 by *Xba*I and *Bam*HI double digestion. Tobacco transformation through *Agrobacterium tumefaciens*-mediated was performed as the leaf disk method described by Horsch et al. [35], and transgenic plants were screened by 50 mg/L kanamycin.

2.5. Southern Blot Analysis. Genomic DNAs (about 25 µg) were obtained from three independent T1 transgenic plant lines and double digested with restriction endonuclease *EcoRI* and *BamHI* (TaKaRa) at 37°C for 20 h and electrophoretically separated on a 0.8% agarose gel and transferred to positively charged nylon membrane. *HindIII*-digested lambda DNA with digoxigenin labeling (Cat. 11218590910, Roche) was employed as marker. The DNA was fixed on the membrane by baking at 120°C for 30 min. According to the protocol of DIG-High Prime DNA Labeling and Detection Starter Kit (Roche, USA), all procedures including the preparation of probe, prehybridization, hybridization, and immunological detection were carried out in proper sequence.

2.6. Cold and Heat Stress Tolerance Assays. Three independent T1 transgenic plant lines and WT tobacco plants were incubated on tissue culture medium under a 16 h light/8 h dark cycle at 25°C for 30 d and then transferred into two different growth chambers separately at 4°C and 38°C, 1200 Lux light (16 h day/8 h dark), for 12 h treatment. The treated plants were photographed every two hours, and the forth functional leaves taken at the same time were exploited to conduct physiological and gene expression experiments.

2.7. Determination of APX, SOD, POD, and CAT Activity and AsA Concentration. APX activity was determined according to the method of Nakano and Asada [36]. 0.4 g ground frozen fresh samples were homogenized in 4 mL of 50 mM extraction PBS buffer (pH 7.8) containing 0.2 mM EDTA and 2 mM AsA. The homogenate was transferred into a tube and centrifuged at 12,000 ×g for 20 min (4°C) and the supernatant was collected for APXase activity analysis. AsA content was determined according to the method of Arakawa et al. [37]. The crude extracts homogenized through grinding 0.3 g fresh samples macerated with liquid nitrogen in presence of 3 mL 5% (w/v) TCA into a fine powder were centrifuged at 4,000 ×g for 10 min at room temperature and the supernatant was collected for AsA content analysis.

0.2 g ground frozen fresh samples were homogenized in 10 mL precooling PBS buffer (pH 7.8), followed by being centrifuged at 12,000 ×g for 15 min (4°C), and the supernatant was extracted for enzyme activity analysis. SOD, POD, and CAT activities measurement was performed as described by Lee [38].

2.8. Biochemical and Physiological Analyses. MDA content was determined by the thiobarbituric acid- (TBA-) based colorimetric method as described by Heath and Packer [39]. H₂O₂ content was assayed by modified method according to Shi et al. [40]. 0.5 g fine grinding fresh samples were homogenized in an ice bath with 0.1% (w/v) TCA, and the homogenate was centrifuged at 12,000 ×g for 15 min. Then, 1 mL supernatant was mixed together with 1 mL potassium phosphate buffer (100 mM, pH 7.0) and 2 mL KI (1M), and the absorbance was measured at 390 nm using a standard curve plotted with known concentration of H₂O₂. Relative electric conductivity was measured referring to the method

of Zhang [41]. 0.5 g fresh leaf discs (0.5 × 0.5 cm) were cut into several conical flasks containing 25 mL distilled water and vacuumed for 8 min, 6 conical flasks totally, and the conical flasks were, respectively, treated under different temperatures lasting 20 min to determine the initial electric conductivity (*R*) using Conductivity Meter (HI-9033). Then, the whole sample mixtures were heated with boiling water for 15 min to kill the cells and final electric conductivity (*R*₀) was determined after 2 h later, with electric conductivity of plant leaf disks at 25°C as a control group (CK). Relative electric conductivity (REC) was evaluated as $REC (\%) = [(R - CK) / (R_0 - CK)] \times 100\%$.

For photosynthetic analysis, a Chlorophyll Fluorescence Analyzer (Imaging-PAM-Maxi, Walz, Germany) was used to determine all of the parameters according to the manufacturer's instructions.

3. Results

3.1. Molecular Cloning and Characterization of CaAPX. A fragment (566-bp) was obtained by amplification of primers (Table S1) designed based on conserved regions, and sequence analysis showed that it was homologous to APX genes (not shown). The full-length sequence of CaAPX (GenBank Accession ID: KP635267) was retrieved by 3' and 5'-RACE amplification (primers listed in Table S1). CaAPX cDNA consisted of 1,017-bp nucleotides and encoded an open reading frame of 250 amino acid residues (Figure S1).

The deduced amino acid sequence of CaAPX was aligned with homologous sequences from various species, including *Arabidopsis thaliana*, *Gossypium hirsutum*, *Glycine max*, *Nicotiana tomentosiformis*, *Oryza sativa*, *Phoenix dactylifera*, and *Solanum tuberosum*, using BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). The predicted structure of CaAPX dominantly contained alpha-helix and random coil, in which a POD domain containing the highly conserved amino acid sequences, namely, "XANX," "LPDAX," and "(E)RSGF/W" (Figure 1(b)), was found. These conserved motifs were shown to be critical to react with substrate AsA [42]. A neighbor-joining bootstrap tree was constructed based on the homologous APX proteins sequences from various plant species (Figure 1(a)). We revealed that they were divided into four groups and CaAPX was categorized into a cytosolic-type APX protein (Figure 1(a)).

3.2. Expression Pattern of CaAPX in C. azalea. We studied the expression patterns of CaAPX by qRT-PCR analysis in various tissues and different temperature stresses (Figure 1(c)). It was found that CaAPX was expressed in all tested tissues and the expression level of CaAPX in immature green fruit was the highest, with nearly 2.68 times higher than in mature leaf (Figure 1(c)). Notably, the expression levels in immature green fruit and mature leaf were higher than other tissue types (leaf bud, flower bud, flower, seed, and petal) (Figure 1(c)). To investigate the expression of CaAPX under temperature stresses, we examined the transcription levels in leaves after exposing to 4°C and 38°C, respectively, with 5 time points.

We showed that the expression levels of CaAPX in leaves were both upregulated under the low and high temperature

FIGURE 1: Sequence analysis and gene expression profiles of CaAPX. (a) Phylogenetic tree analysis of CaAPX and related proteins. (b) Alignment of amino acid sequences. Ca, *C. azalea* (KP635267); At, *Arabidopsis thaliana* (NP_172267); Gh, *Gossypium hirsutum* (ABR18607); Gm, *Glycine max* (NP_001237785); Nt, *Nicotiana tomentosiformis* (XP_009597491); Os, *Oryza sativa* (japonica cultivar group) (P93404); Pd, *Phoenix dactylifera* (XP_008783664); St, *Solanum tuberosum* (NP_001275066). Red boxes indicate the highly conserved APX active-site signature. (c) Relative expression of CaAPX in different tissues of *C. azalea*. (d) Relative expression of CaAPX in leaves of *C. azalea* during different times under 4°C cold stress. (e) Relative expression of CaAPX in leaves of *C. azalea* during different times under 38°C hot stress. The data are the means of four replicates and were compared by Duncan's multiple range tests.

FIGURE 2: Analysis of transgenic plants. (a) Phenotypes of transgenic and WT plants under cold and hot stress treatments after 8 h. Upper panel, 4 degrees; bottom panel, 38 degrees. (b) Southern blot analysis of transgenic lines. (c) Detection of *CaAPX* transgenic tobacco by RT-PCR. Expression level of *CaAPX* gene in different lines (L1, L2, and L3) of transgenic and wild type (WT) plants. M2: λ -Hind III digested DNA marker; L1, L2, and L3: 3 lines of *CaAPX* transgenic tobacco; +: positive control of pBI121-*CaAPX* combined vector; -: water as the negative control; WT: wild type. The data are the means of four replicates and were compared by Duncan's multiple range tests.

treatment, reaching peaks at 8 h treatment. Interestingly, the expression levels both sharply declined in 12 h temperature stresses (Figures 1(d) and 1(e)).

3.3. Heterogeneous Expression of *CaAPX* Altered the Temperature Stress Tolerance and Displayed Elevated APX Activity and AsA Contents. To investigate the molecular functions of *CaAPX*, we generated transgenic tobacco plants with heterogeneous expression. We identified 20 positive transgenic lines by kanamycin selection and gene-specific amplification (Table S1, Figure S2). Three lines were randomly selected for southern blotting analysis, which all confirmed the single insertion of target construct (Figure 2(a)). We quantified the expression levels of *CaAPX* in these three transgenic lines using qRT-PCR. The results showed that expression

level in L2 line was the highest, approximately 3.83 times higher than in L1 line, and L3 line had a higher expression level of around 1.23-fold compared to L1 line (Figure 2(b)). To study the tolerance to temperature stresses, transgenic plants were transferred into growth chamber at 4°C and 38°C, respectively. And phenotypes of treated plants were recorded every 15 min until significant wilting was emerged. After 8 h treatment under both 4°C and 38°C, the WT plants displayed obvious wilting symptoms, while transgenic lines displayed significantly enhanced performance (Figure 2(c)).

The activity of APX at different temperatures was examined in transgenic and WT plants. It was found that APX activity was much higher in three transgenic lines than in untreated WT plants (at 0 h in Figures 3(a) and 3(b)). The APX activities showed a similar increasing pattern in all

FIGURE 3: AsA content and APX activity in plants under temperature stress. (a) APX activities of transgenic and WT plants under 4°C temperature stress. (b) APX activities of transgenic and WT plants under 38°C temperature stress. (c) AsA contents of transgenic and WT plants under 4°C temperature stress. (d) AsA contents of transgenic and WT plants under 38°C temperature stress. L1, L2, and L3: 3 lines of *CaAPX* transgenic tobacco. WT: wild type. FW: fresh weight.

tested transgenic lines as well as WT plants, upon cold and hot treatments (Figures 3(a) and 3(b)); meanwhile the L2 line displayed higher activity than the rest of assayed plants at each time point (Figures 3(a) and 3(b)). We further measured the contents of AsA in WT and transgenic plants under treatments. The results showed that, before treatments (at 0 h), all transgenic lines had significantly higher levels of AsA than WT (Figures 3(c) and 3(d)). Upon both 38°C and 4°C treatments, the AsA levels were increasing along with treatment durations in all transgenic plants and WT (Figures 3(c) and 3(d)). Also the L2 line had the highest AsA levels than other assayed plants at each time point (Figures 3(c)

and 3(d)). Their results indicated that ectopic expression of *CaAPX* caused increased APX activity and AsA levels and enhanced the plant tolerances to cold and hot stresses. Line L2 displayed highest APX activity and AsA content, which is consistent with the accumulation patterns of *CaAPX* transcripts in transgenic plants (Figure 2(c)).

3.4. Overexpression of *CaAPX* Triggered Upregulation of Antioxidant Enzyme Activities. To study how the temperature stresses and *CaAPX* expression affected the ROS removing enzymes, the activities of SOD, POD, and CAT were measured. We showed that the SOD activities were

upregulated before treatments (at 0 h in Figures 4(a) and 4(b)) in transgenic lines; there was a gradually increasing trend of SOD activity for each assayed plant during 38°C and 4°C treatments (Figures 4(a) and 4(b)); transgenic lines had higher SOD activities than WT at each timepoint of treatments (Figures 4(a) and 4(b)). The POD activities were upregulated in transgenic lines comparing to WT before treatments (at 0 h in Figures 4(c) and 4(d)), while the patterns of POD activities during treatments were different from SOD, which reached peaks at 8 h and declined at 12 h in transgenic plants (Figures 4(c) and 4(d)); in WT, the accumulation of POD activities was gradually increasing during 8 hours of treatments and decreased at 12 h (Figures 4(c) and 4(d)). The results of CAT analysis were similar to POD activities, except that CAT activities reached peaks at 8 h in all transgenic lines under both 38°C and 4°C treatments, and there was a sudden drop of activities at 12 h (Figures 4(e) and 4(f)). The activities of CAT in WT did not show strong induction under temperature treatments (Figures 4(e) and 4(f)).

3.5. Overexpression of *CaAPX* Reduced Accumulation of MDA, H_2O_2 , and Electrolyte Leakage under Stresses. To investigate the effects of altered ROS-scavenging activities (APX, CAT, and SOD) in transgenic plants, we measured the contents of MDA and H_2O_2 and the rate of electrolyte leakage. We found that MDA levels in WT plants were immediately induced when exposed to the abnormal temperatures (4°C and 38°C) and accumulated more dramatically compared with transgenic lines. The MDA contents in transgenic lines were significantly decreased, accounting for 71.09% to 82.91% in comparison to WT plants (0 h at Figures 5(a) and 5(b)).

Although there was no significant difference in H_2O_2 concentration between WT and transgenic plants incubated at room temperature (25°C), H_2O_2 concentration in WT plants increased markedly under 4°C and 38°C temperature stresses (Figures 5(c) and 5(d)). In transgenic plants, the H_2O_2 contents kept at the baseline level after 8 h treatments of temperature stresses (Figures 5(c) and 5(d)); but an increase at 12 h under 38°C was observed (Figure 5(d)). In addition, the rate of electrolyte leakage was higher in WT plants than in transgenic lines under the temperature ranges from 15°C to -20°C (Figure 5(e)). There was a slight rise in electrolyte leakage in all plants between 30 and 40°C (Figure 5(f)), and the conductivity rate increased dramatically to the highest level at 60°C and 70°C (Figure 5(f)), which suggested that the cells death of leaves had occurred since 60°C.

3.6. Parameters Associated with Chlorophyll Fluorescence in Leaves under Temperature Stresses. The photosynthetic electron transport (PET) chain is a sensitive process associated with cellular ROS homeostasis. To understand the regulatory roles of *CaAPX* under stresses, we measured several parameters such as *Fv/Fm* and *Y(II)* associated with chlorophyll fluorescence. Before the stress treatments (0 h at Figures 6(a) and 6(b)), the *Fv/Fm* was nearly the same between WT and transgenic plants. Upon temperature stresses, *Fv/Fm* decreased markedly under both 4°C and 38°C temperature stresses; the decline in WT plants was much accentuated

(Figures 6(a) and 6(b)). The *Y(II)* was slightly higher in the transgenic lines compared to WT plants under control conditions, and the trend for *Y(II)* was similar to *Fv/Fm* under temperature stresses (Figures 6(c) and 6(d)). After 12 h treatment, the figures for transgenic lines were approximately 1.32- and 1.12-fold for average in comparison to WT sampling, respectively, at 4 and 38°C. Therefore, it was suggested that low and high temperatures significantly descended the parameters associated with chlorophyll fluorescence in both genotypes, in which the effects of stresses were more noticeable on WT plants than on the transgenic lines.

3.7. Divergent Induction Patterns of ROS-Scavenging Genes upon Cold and Heat Stresses. The elevated levels of ROS by temperature stresses are important signals to control the expression of ROS-scavenging enzymes [2, 5, 43]. To understand the ROS signal transduction in transgenic *CaAPX* plants under stresses, we analyzed the transcripts levels of tobacco ROS-scavenging genes, *NtCu/Zn-SOD*, *NtCAT*, *NtDHAR*, and *NtMDHAR*, in response to temperature stresses. We showed that, before the temperature stress treatments, the expression of *NtCAT* was obviously upregulated in transgenic lines compared with WT plants (Figure 7(c)), while expression of *NtCu/Zn-SOD*, *NtDHAR*, and *NtMDHAR* was not significantly induced in *CaAPX* transgene plants. Upon cold and hot stresses treatments, the expression levels of all assayed genes were upregulated at some time points, but the patterns of induction between cold and hot stresses were mostly different (Figures 7(a)–7(h)). For instance, remarkable induction of *NtCu/Zn-SOD* was detected only at 2 h treatment of 38°C and expression dropped thereafter in *CaAPX* transgenic plants; under 4°C treatments moderate inductions were observed in both WT and transgenic plants (Figures 7(a) and 7(b)). In WT, the expression of *NtCAT* was induced significantly after 2 h of hot treatment (Figure 7(d)), while, in *CaAPX* transgenic plants, the expression of *NtCAT* did not change after 2 h and 4 h and greatly reduced at 8 h and 12 h of hot treatment. Under cold treatment, the expression of *NtCAT* was upregulated in both WT and transgenic plants (Figure 7(c)). These results suggested the *CaAPX* overexpression affected the ROS mediated regulation of SOD and CAT expression, and the hot and cold stresses had different responsive mechanisms of signaling. Meanwhile, the expression levels of *NtDHAR* and *NtMDHAR* displayed similar induction patterns between WT and transgenic plants under both cold and hot stresses (Figures 7(e)–7(h)), suggesting a different regulatory pathway was involved.

3.8. *CaAPX* Overexpression Plants Displayed Greater Thermotolerance in Seed Germination Assay. To further investigate the thermotolerance effects of *CaAPX* in transgenic plants, we tested the germination of transgenic lines. The assayed lines were confirmed by T2 segregation analysis through PCR amplification of progenies (Figure S3). No segregation was found for all three lines. Then we tested the germination rates of seeds on the MS medium containing antibiotics selection at 25°C, 4°C, and 38°C, respectively. We found that the germination rates of transgenic lines were a little higher

FIGURE 4: Activities of antioxidant enzyme in plants under temperature stress. (a) SOD activities of transgenic and WT plants under 4°C temperature stress. (b) SOD activities of transgenic and WT plants under 38°C temperature stress. (c) POD activities of transgenic and WT plants under 4°C temperature stress. (d) POD activities of transgenic and WT plants under 38°C temperature stress. (e) CAT activities of transgenic and WT plants under 4°C temperature stress. (f) CAT activities of transgenic and WT plants under 38°C temperature stress. L1, L2, and L3: 3 lines of *CaAPX* transgenic tobacco. WT: wild type. FW: fresh weight.

FIGURE 5: The contents of MDA, H_2O_2 , and electrolyte leakage rates in plants under temperature stress. (a) MDA concentration of transgenic and WT plants under 4°C temperature stress. (b) MDA concentration of transgenic and WT plants under 38°C temperature stress. (c) H_2O_2 concentration of transgenic and WT plants under 4°C temperature stress. (d) H_2O_2 concentration of transgenic and WT plants under 38°C temperature stress. (e) Electrolyte leakage rates of transgenic and WT plants under low temperature stress. (f) Electrolyte leakage rate of transgenic and WT plants under high temperature stress. L1, L2, and L3: 3 lines of *CaAPX* transgenic tobacco. WT: wild type. FW: fresh weight.

FIGURE 6: Photosynthetic parameters in plants under temperature stress. (a) F_v/F_m of transgenic and WT plants under 4°C temperature stress. (b) F_v/F_m of transgenic and WT plants under 38°C temperature stress. (c) $Y(II)$ of transgenic and WT plants under 4°C temperature stress. (d) $Y(II)$ of transgenic and WT plants under 38°C temperature stress. L1, L2, and L3: 3 lines of *CaAPX* transgenic tobacco. WT: wild type.

than from WT seeds at 25°C treatment but not significant (Figures 8(a), 8(b), and 8(c)).

Under 4°C and 38°C temperature stresses, the rates of transgenic lines were significantly higher than WT seeds (Figures 8(b), 8(d) and 8(c)). Particularly, under 4°C, almost no germination was found for WT plants, but the strong line (L2) showed a comparable rate to WT under normal condition (Figure 8(d)). The results indicated that overexpressing *CaAPX* significantly enhanced temperature stresses tolerance and improved the seed germination rate in transgenic lines under temperature stresses.

4. Discussions

Temperature stress, namely, low and high temperatures, often caused negative impacts on plant metabolism, cellular

homeostasis, and major physiological processes [2]. Previous studies demonstrated that ROS is one of the main factors causing oxidative damage to plants exposed to environmental stresses [29]. Antioxidant enzymes play a significant role in eliminating toxic levels of ROS, generated during stress from living cells [44]. In stressful conditions, plants generally increased the APX activity and the elevated activity level usually correlates with increased stress tolerance [3]. During high and low temperature treatments, many physiological changes occur that result in increased expression of antioxidant enzyme genes to protect from oxidative stress. APX utilizes AsA and its specific electron to reduce H_2O_2 to H_2O , and overexpressing APX might play a critical role in growth and when subjected to high and low temperature. Previous research indicated that APX was highly responsive to temperature stresses and played an important role in

FIGURE 7: Expressions of *CaAPX* and defense-related genes in plants exposed to temperature stress. (a) Expression level of *NtCu/Zn-SOD* in transgenic and WT plants under 4°C temperature stress. (b) Expression level of *NtCu/Zn-SOD* in transgenic and WT plants under 38°C temperature stress. (c) Expression level of *NtCAT* in transgenic and WT plants under 4°C temperature stress. (d) Expression level of *NtCAT* in transgenic and WT plants under 38°C temperature stress. (e) Expression level of *NtDHAR* in transgenic and WT plants under 4°C temperature stress. (f) Expression level of *NtDHAR* in transgenic and WT plants under 38°C temperature stress. (g) Expression level of *NtMDHAR* in transgenic and WT plants under 38°C temperature stress. (h) Expression level of *NtMDHAR* in transgenic and WT plants under 4°C temperature stress. L1, L2, and L3: 3 lines of *CaAPX* transgenic tobacco; WT: wild type. Different lowercase letters represent significant differences among the *CaAPX* gene transgenic and WT plants, whereas different capital letters represent significant differences among the stress treatments, at a confidence level of 0.05. The data are the means of four replicates and were compared by Duncan's multiple range tests.

the scavenging of ROS in plants. For example, the *APX1* expression of *A. thaliana* significantly increased after being treated with heat stress [45]; Padaria et al. [46] found that *TapAPX* of a thermotolerant wheat cv. Raj3765 displayed up to 203-fold level of expression at 42°C heat stress exposure. The expression level of *tAPX* during cold acclimation of

wheat was upregulated [47]; the expression of *A. andraeanum* *APX* mRNA was evidently elevated by cold stress [48]; *OsAPX2* expression of *Oryza sativa* was also developmental- and spatial-regulated and was induced by cold stress [49]. In this study, the expression levels of *CaAPX* were both remarkably upregulated under low and high temperature treatment

FIGURE 8: Germination rates of T1 generation seeds under temperature stress. (a) The state of seeds germination under 25°C. (b) The state of seeds germination under 4°C. (c) The state of seeds germination under 38°C. (d) Rate of seeds germination of T1 generation transgenic and WT seeds under different temperature treatments. L1, L2, and L3: 3 lines of *CaAPX* transgenic tobacco; WT: wild type. Different lowercase letters represent significant differences among the *CaAPX* transgenic and WT plants, whereas different capital letters represent significant differences among the stress treatments, at a confidence level of 0.05. The data are the means of four replicates and were compared by Duncan's multiple range tests.

(Figures 1(d) and 1(e)). It demonstrated that *CaAPX* was induced by heat and cold. We found that overexpression of *CaAPX* increased tolerance to temperature stresses, and transgenic tobacco grew better than wild type under 4°C and 38°C temperature stresses (Figure 2(a)). We showed that the activity of APX and the content of AsA were indeed remarkably enhanced in transgenic plants than those in wild type plants (Figures 3(a)–3(d)), suggesting their roles in scavenging H_2O_2 (Figures 5(c) and 5(d)).

Previous studies indicated that antioxidant enzyme activities were tightly associated with the expression of antioxidant enzyme genes, which could be upregulated after the plants were treated with temperature stresses. For instance, Lin et al. [50] reported that CAT and APX expression profiles were well matched with the data for CAT and APX enzyme activities in the broccoli and Chinese cabbage plants, respectively. And overexpressing chloroplast APX in cotton has conferred higher photochemical activity of photosystem II (PSII) and

stronger antioxidant capacity to transgenic plants [51]. Overexpression of *CaAPX* also enhanced the activities of SOD, POD, and CAT in transgenic tobacco plants than that in the wild type plants under 4°C and 38°C temperature stresses (Figures 4(a)–4(f)). Furthermore, it was demonstrated that the high activities of SOD, POD, and CAT could result in producing the high level of AsA, which could alleviate photoinhibition to protect the plants [52]. Moreover, the high level of AsA could detoxify ROS effectively, further reduce the damage to membrane lipids, enhance the repair of PSII, and protect photosystem. Zhou et al. [53] concluded that *Fv/Fm* is an early indicator of temperature stress tolerance in tomato and that this provides effective and reliable information about leaf photosynthetic performance. Hence, several parameters such as *Fv/Fm* (Figures 6(a) and 6(b)) and *Y(II)* (Figures 6(c) and 6(d)) associated with chlorophyll fluorescence in this study were decreased under extreme temperature stresses, but these effects were more accentuated in wild type plants. These

results manifested that *CaAPX* is a key component to balance the cellular ROS homeostasis and could be an ideal target for improving tolerance to temperature stresses.

The transcript levels of *NtCu/Zn-SOD*, *NtCAT*, *NtDHAR*, and *NtMDHAR* genes significantly increased on exposure to stress in transgenic tobacco plants (Figure 7). The critical role of APX in stress tolerance and upregulation of *Cu/Zn-SOD*, *CAT*, *DHAR*, and *MDHAR* transcripts have been reported [54] under stress. Variations in physiological parameters potentially contribute to the complex phenotype of cold and heat tolerance; however, a clear relation between stress and coexpression of antioxidant genes has so far not been evident. In this study, the expression pattern of *NtCu/Zn-SOD* and *NtCAT* in times was specific to two genotypes and showed no direct correlation with the corresponding enzymes activity. This result was likely because of the complex regulation mechanisms of gene expression; gene expression cannot be directly correlated with enzyme activity [55].

CaAPX was expressed in all tested tissues (leaf, leaf bud, flower, flower bud, petal, immature green fruit, and seed embryo) but was much more abundant in the immature green fruit relative to other tissues (Figure 1(c)). Wang et al. [56] showed that a wide range of *MaAPX1* transcript expression levels among the different tissues in banana, with relatively strong expression, was observed in leaves and roots. Our results indicated that *CaAPX* might have different functions in different tissues during the growth and development of *C. azalea*. To test the thermotolerance and cold-tolerance of transgenic plants, we employed a seed germination assay. Sun et al. [57] reported that overexpression of *StAPX* improved seed germination, and Faize et al. [58] showed that overexpression of *cytapx* also promoted seed germination and germination rates. In our conditions, we found that overexpressing *CaAPX* significantly enhanced temperature stresses tolerance (Figures 8(a), 8(b), and 8(c)) and improved the seed germination rate in transgenic plants (Figure 8(d)) under hot and cold stresses, which was consistent with the other previous plants. Overall, this work characterized the functions of *CaAPX* in response to heat and cold stresses. The results provided comprehensive information about the orchestration of cell ROS homeostasis mediated by *CaAPX*, which could give some insights into rational engineering of plants toward temperature tolerances.

Additional Points

Key Message. *CaAPX* encodes a cytosolic-type APX. Ectopic expression of *CaAPX* enhanced heat and cold tolerances and displayed elevated ROS-scavenging activities. *CaAPX* is central to ROS homeostasis and temperature tolerance.

Competing Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contributions

Hengfu Yin, Jiyuan Li, and Libo He conceived the study. Jiangying Wang performed gene cloning and transgenic

analysis in tobacco. Bin Wu, Zhengqi Fan, and Xinlei Li contributed to sample collection and gene expression analysis. Sui Ni helped on preparing plant materials. Jiangying Wang and Hengfu Yin wrote the draft and all authors contributed to the writing and revision of manuscript.

Acknowledgments

The authors thank Professor Zhongchi Liu from the University of Maryland (College Park) for help with experimental design and technical assistance. This work was supported by funds from Zhejiang Science and Technology Major Program on Agricultural New Variety Breeding (2016C02056-12). They also acknowledge SFA Introduction Program on International Advanced Forestry Sci. & Tech. (2014-4-16), MOST International Sci. & Tech. Cooperation Program (2016YFE0126100), CAF Special Grant on International Cooperation & Innovation (CAFYBB2017ZF001), MOST Nonprofit Sector Special Research Funds on Forestry (201504707), and Ningbo Special Sci. & Tech. Program on Agriculture (2014C11002).

References

- [1] S. S. Karpinski, G. Wingsle, B. Karpinska, and J. Hallgren, "Low-temperature stress and antioxidant defense mechanisms in higher plants," in *Oxidative Stress in Plants*, pp. 69–103, Taylor & Francis, 2002.
- [2] N. Suzuki and R. Mittler, "Reactive oxygen species and temperature stresses: a delicate balance between signaling and destruction," *Physiologia Plantarum*, vol. 126, no. 1, pp. 45–51, 2006.
- [3] R. D. Allen, "Dissection of oxidative stress tolerance using transgenic plants," *Plant Physiology*, vol. 107, no. 4, pp. 1049–1054, 1995.
- [4] K. Asada, "The water-water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess photons," *Annual Review of Plant Biology*, vol. 50, pp. 601–639, 1999.
- [5] R. Mittler, S. Vanderauwera, M. Gollery, and F. Van Breusegem, "Reactive oxygen gene network of plants," *Trends in Plant Science*, vol. 9, no. 10, pp. 490–498, 2004.
- [6] R. Mittler, "Abiotic stress, the field environment and stress combination," *Trends in Plant Science*, vol. 11, no. 1, pp. 15–19, 2006.
- [7] S. Azarabadi, H. Abdollahi, M. Torabi, Z. Salehi, and J. Nasiri, "ROS generation, oxidative burst and dynamic expression profiles of ROS-scavenging enzymes of superoxide dismutase (SOD), catalase (CAT) and ascorbate peroxidase (APX) in response to *Erwinia amylovora* in pear (*Pyrus communis* L)," *European Journal of Plant Pathology*, vol. 147, no. 2, pp. 279–294, 2017.
- [8] L. Pnueli, H. Liang, M. Rozenberg, and R. Mittler, "Growth suppression, altered stomatal responses, and augmented induction of heat shock proteins in cytosolic ascorbate peroxidase (Apx1)-deficient Arabidopsis plants," *Plant Journal*, vol. 34, no. 2, pp. 187–203, 2003.
- [9] W. R. Swindell, M. Huebner, and A. P. Weber, "Transcriptional profiling of Arabidopsis heat shock proteins and transcription factors reveals extensive overlap between heat and non-heat stress response pathways," *BMC Genomics*, vol. 8, article no. 125, 2007.

- [10] A. Baxter, R. Mittler, and N. Suzuki, "ROS as key players in plant stress signalling," *Journal of Experimental Botany*, vol. 65, no. 5, pp. 1229–1240, 2014.
- [11] D. Marino, C. Dunand, A. Puppo, and N. Pauly, "A burst of plant NADPH oxidases," *Trends in Plant Science*, vol. 17, no. 1, pp. 9–15, 2012.
- [12] I. I. Panchuk, R. A. Volkov, and F. Schöfl, "Heat stress- and heat shock transcription factor-dependent expression and activity of ascorbate peroxidase in Arabidopsis," *Plant Physiology*, vol. 129, no. 2, pp. 838–853, 2002.
- [13] S. Davletova, L. Rizhsky, H. Liang et al., "Cytosolic ascorbate peroxidase 1 is a central component of the reactive oxygen gene network of Arabidopsis," *Plant Cell*, vol. 17, no. 1, pp. 268–281, 2005.
- [14] K. Asada, "Ascorbate peroxidase—a hydrogen peroxide-scavenging enzyme in plants," *Physiologia Plantarum*, vol. 85, no. 2, pp. 235–241, 1992.
- [15] G. Noctor and C. H. Foyer, "Ascorbate and glutathione: keeping active oxygen under control," *Annual Review of Plant Biology*, vol. 49, pp. 249–279, 1998.
- [16] S. Shigeoka, T. Ishikawa, M. Tamoi et al., "Regulation and function of ascorbate peroxidase isoenzymes," *Journal of Experimental Botany*, vol. 53, no. 372, pp. 1305–1319, 2002.
- [17] Z. Lu, D. Liu, and S. Liu, "Two rice cytosolic ascorbate peroxidases differentially improve salt tolerance in transgenic Arabidopsis," *Plant Cell Reports*, vol. 26, no. 10, pp. 1909–1917, 2007.
- [18] S. S. Gill and N. Tuteja, "Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants," *Plant Physiology and Biochemistry*, vol. 48, no. 12, pp. 909–930, 2010.
- [19] Q. Guan, Z. Wang, X. Wang, T. Takano, and S. Liu, "A peroxisomal APX from *Puccinellia tenuiflora* improves the abiotic stress tolerance of transgenic *Arabidopsis thaliana* through decreasing of H_2O_2 accumulation," *Journal of Plant Physiology*, vol. 175, pp. 183–191, 2015.
- [20] C. H. Foyer and B. Halliwell, "The presence of glutathione and glutathione reductase in chloroplasts: a proposed role in ascorbic acid metabolism," *Planta*, vol. 133, no. 1, pp. 21–25, 1976.
- [21] P. P. Jablonski and J. W. Anderson, "Light-dependent reduction of hydrogen peroxide by ruptured pea chloroplasts," *Plant Physiology*, vol. 69, no. 6, pp. 1407–1413, 1982.
- [22] M. A. Hossain and K. Asada, "Purification of dehydroascorbate reductase from spinach and its characterization as a thiol enzyme," *Plant and Cell Physiology*, vol. 25, pp. 85–92, 1984.
- [23] S.-H. Lee, N. Ahsan, K.-W. Lee et al., "Simultaneous overexpression of both CuZn superoxide dismutase and ascorbate peroxidase in transgenic tall fescue plants confers increased tolerance to a wide range of abiotic stresses," *Journal of Plant Physiology*, vol. 164, no. 12, pp. 1626–1638, 2007.
- [24] K. Kavitha, G. Venkataraman, and A. Parida, "An oxidative and salinity stress induced peroxisomal ascorbate peroxidase from *Avicennia marina*: molecular and functional characterization," *Plant Physiology and Biochemistry*, vol. 46, no. 8–9, pp. 794–804, 2008.
- [25] R. Ahmad, Y.-H. Kim, M.-D. Kim et al., "Simultaneous expression of choline oxidase, superoxide dismutase and ascorbate peroxidase in potato plant chloroplasts provides synergistically enhanced protection against various abiotic stresses," *Physiologia Plantarum*, vol. 138, no. 4, pp. 520–533, 2010.
- [26] H. Moon, B. Lee, G. Choi et al., "NDP kinase 2 interacts with two oxidative stress-activated MAPKs to regulate cellular redox state and enhances multiple stress tolerance in transgenic plants," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 1, pp. 358–363, 2003.
- [27] Y.-H. Kim, S. Lim, K.-S. Yang et al., "Expression of arabidopsis NDPK2 increases antioxidant enzyme activities and enhances tolerance to multiple environmental stresses in transgenic sweetpotato plants," *Molecular Breeding*, vol. 24, no. 3, pp. 233–244, 2009.
- [28] Y. Sato, T. Murakami, H. Funatsuki, S. Matsuba, H. Saruyama, and M. Tanida, "Heat shock-mediated APX gene expression and protection against chilling injury in rice seedlings," *Journal of Experimental Botany*, vol. 52, no. 354, pp. 145–151, 2001.
- [29] S. Lim, Y.-H. Kim, S.-H. Kim et al., "Enhanced tolerance of transgenic sweetpotato plants that express both CuZnSOD and APX in chloroplasts to methyl viologen-mediated oxidative stress and chilling," *Molecular Breeding*, vol. 19, no. 3, pp. 227–239, 2007.
- [30] L. Tang, S.-Y. Kwon, S.-H. Kim et al., "Enhanced tolerance of transgenic potato plants expressing both superoxide dismutase and ascorbate peroxidase in chloroplasts against oxidative stress and high temperature," *Plant Cell Reports*, vol. 25, no. 12, pp. 1380–1386, 2006.
- [31] M. D. Kim, Y.-H. Kim, S.-Y. Kwon, D.-J. Yun, S.-S. Kwak, and H.-S. Lee, "Enhanced tolerance to methyl viologen-induced oxidative stress and high temperature in transgenic potato plants overexpressing the CuZnSOD, APX and NDPK2 genes," *Physiologia Plantarum*, vol. 140, no. 2, pp. 153–162, 2010.
- [32] S. B. Rosa, A. Caverzan, F. K. Teixeira et al., "Cytosolic APx knockdown indicates an ambiguous redox responses in rice," *Phytochemistry*, vol. 71, no. 5–6, pp. 548–558, 2010.
- [33] A. Bonifacio, M. O. Martins, C. W. Ribeiro et al., "Role of peroxidases in the compensation of cytosolic ascorbate peroxidase knockdown in rice plants under abiotic stress," *Plant, Cell and Environment*, vol. 34, no. 10, pp. 1705–1722, 2011.
- [34] A. Caverzan, A. Bonifacio, F. E. L. Carvalho et al., "The knockdown of chloroplastic ascorbate peroxidases reveals its regulatory role in the photosynthesis and protection under photo-oxidative stress in rice," *Plant Science*, vol. 214, pp. 74–87, 2014.
- [35] R. B. Horsch, J. E. Fry, N. L. Hoffmann, D. Eichholtz, S. G. Rogers, and R. T. Fraley, "A simple and general method for transferring genes into plants," *Science*, vol. 227, no. 4691, pp. 1229–1231, 1985.
- [36] Y. Nakano and K. Asada, "Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts," *Plant and Cell Physiology*, vol. 22, no. 5, pp. 867–880, 1981.
- [37] N. Arakawa, K. Tsutsumi, N. G. Sanceda, T. Kurata, and C. Inagaki, "A rapid and sensitive method for the determination of ascorbic acid using 4,7-diphenyl-1,10-phenanthroline," *Agricultural and Biological Chemistry*, vol. 45, no. 5, pp. 1289–1290, 1981.
- [38] H. S. Lee, *The Principles and Techniques of the Plant Physiological and Biochemical*, Higher Education Press, Beijing, China, 2000.
- [39] R. L. Heath and L. Packer, "Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation," *Archives of Biochemistry and Biophysics*, vol. 125, no. 1, pp. 189–198, 1968.
- [40] X. Shi, D. C. Flynn, D. W. Porter, S. S. Leonard, V. Vallyathan, and V. Castranova, "Efficacy of taurine based compounds as hydroxyl radical scavengers in silica induced peroxidation," *Annals of Clinical and Laboratory Science*, vol. 27, no. 5, pp. 365–374, 1997.

- [41] X. Z. Zhang, *The Research Method of Physiology in the Crop*, Agriculture Press, Beijing, China, 1992.
- [42] H. M. Jespersen, I. V. H. Kjærsgård, L. Østergaard, and K. G. Welinder, "From sequence analysis of three novel ascorbate peroxidases from *Arabidopsis thaliana* to structure, function and evolution of seven types of ascorbate peroxidase," *Biochemical Journal*, vol. 326, no. 2, pp. 305–310, 1997.
- [43] R. Mittler, "Oxidative stress, antioxidants and stress tolerance," *Trends in Plant Science*, vol. 7, no. 9, pp. 405–410, 2002.
- [44] A. Shafi, V. Dogra, T. Gill, P. S. Ahuja, and Y. Sreenivasulu, "Simultaneous over-expression of *PaSOD* and *RaAPX* in transgenic *Arabidopsis thaliana* confers cold stress tolerance through increase in vascular lignifications," *PLoS ONE*, vol. 9, no. 10, Article ID e110302, 14 pages, 2014.
- [45] S. Koussevitzky, N. Suzuki, S. Huntington et al., "Ascorbate peroxidase 1 plays a key role in the response of *Arabidopsis thaliana* to stress combination," *Journal of Biological Chemistry*, vol. 283, no. 49, pp. 34197–34203, 2008.
- [46] J. C. H. Padaria, H. Vishwakarma, K. Biswas, R. S. I. Jasrotia, and G. P. R. Singh, "Molecular cloning and in-silico characterization of high temperature stress responsive *pAPX* gene isolated from heat tolerant Indian wheat cv. Raj 3765," *BMC Research Notes*, vol. 7, pp. 713–714, 2014.
- [47] K.-H. Baek and D. Z. Skinner, "Alteration of antioxidant enzyme gene expression during cold acclimation of near-isogenic wheat lines," *Plant Science*, vol. 165, no. 6, pp. 1221–1227, 2003.
- [48] H.-C. Liu, D.-Q. Tian, J.-X. Liu, G.-Y. Ma, Q.-C. Zou, and Z.-J. Zhu, "Cloning and functional analysis of a novel ascorbate peroxidase (APX) gene from *Anthurium andraeanum*," *Journal of Zhejiang University: Science B*, vol. 14, no. 12, pp. 1110–1120, 2013.
- [49] Z. Zhang, Q. Zhang, J. Wu et al., "Gene knockout study reveals that cytosolic ascorbate peroxidase 2 (*OsAPX2*) plays a critical role in growth and reproduction in rice under drought, salt and cold stresses," *PLOS ONE*, vol. 8, no. 2, Article ID e57472, pp. 1–13, 2013.
- [50] K.-H. Lin, H.-C. Huang, and C.-Y. Lin, "Cloning, expression and physiological analysis of broccoli catalase gene and Chinese cabbage ascorbate peroxidase gene under heat stress," *Plant Cell Reports*, vol. 29, no. 6, pp. 575–593, 2010.
- [51] D. Korniyev, A. L. Barry, P. Payton, R. D. Allen, and A. Scott Holaday, "Enhanced photochemical light utilization and decreased chilling-induced photoinhibition of photosystem II in cotton overexpressing genes encoding chloroplast-targeted antioxidant enzymes," *Physiologia Plantarum*, vol. 113, no. 3, pp. 323–331, 2001.
- [52] K. Asada, "Mechanisms for scavenging reactive molecules generated in chloroplasts under light stress," in *Photoinhibition of Photosynthesis. From Molecular Mechanisms to the Field*, N. R. Baker and J. R. Bowyer, Eds., pp. 129–142, Bios Scientific Publishers, Oxford, UK, 1994.
- [53] R. Zhou, X. Yu, K. H. Kjær, E. Rosenqvist, C.-O. Ottosen, and Z. Wu, "Screening and validation of tomato genotypes under heat stress using *Fv/Fm* to reveal the physiological mechanism of heat tolerance," *Environmental and Experimental Botany*, vol. 118, pp. 1–11, 2015.
- [54] A. Bartwal and S. Arora, "Drought stress-induced enzyme activity and *mdar* and *apx* gene expression in tolerant and susceptible genotypes of *Eleusine coracana* (L.)," *In Vitro Cellular & Developmental Biology—Plant*, pp. 1–9, 2016.
- [55] A. Harb, D. Awad, and N. Samarah, "Gene expression and activity of antioxidant enzymes in barley (*Hordeum vulgare* L.) under controlled severe drought," *Journal of Plant Interactions*, vol. 10, no. 1, pp. 109–116, 2015.
- [56] Z. Wang, Z. Q. Jin, J. S. Wang et al., "Cloning and characterization of an ascorbate peroxidase gene regulated by ethylene and abscisic acid during banana fruit ripening," *African Journal of Biotechnology*, vol. 11, no. 43, pp. 10089–10097, 2012.
- [57] W.-H. Sun, M. Duan, D.-F. Shu, S. Yang, and Q.-W. Meng, "Over-expression of *StAPX* in tobacco improves seed germination and increases early seedling tolerance to salinity and osmotic stresses," *Plant Cell Reports*, vol. 29, no. 8, pp. 917–926, 2010.
- [58] M. Faize, E. Nicolás, L. Faize, P. Díaz-Vivancos, L. Burgos, and J. A. Hernández, "Cytosolic ascorbate peroxidase and Cu, Zn-superoxide dismutase improve seed germination, plant growth, nutrient uptake and drought tolerance in tobacco," *Theoretical and Experimental Plant Physiology*, vol. 27, no. 3-4, pp. 215–226, 2015.