Research Article

Esculetin Neutralises Cytotoxicity of *t*-BHP but Not of H₂O₂ on Human Leukaemia NB4 Cells

Virginia Rubio, Ana I. García-Pérez, M. Cristina Tejedor, Angel Herráez, and José C. Diez

Unidad de Bioquímica y Biología Molecular, Departamento de Biología de Sistemas, Universidad de Alcalá, Alcalá de Henares, 28871 Madrid, Spain

Correspondence should be addressed to Angel Herráez; angel.herraez@uah.es

Received 20 December 2016; Accepted 9 February 2017; Published 7 March 2017

Academic Editor: Swaran J. S. Flora

Copyright © 2017 Virginia Rubio et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The coumarin esculetin shows antioxidant action on some cell types, both by scavenging ROS and by decreasing ROS production. We have previously demonstrated the induction of apoptosis by esculetin on NB4 human leukaemia cells by an ill-defined mechanism related to ROS levels. In this work, we analyze the effect of the simultaneous treatment with esculetin and two oxidants to observe the early events in the mechanism of esculetin-induced apoptosis. Our results show that, from the early time of 15 min, esculetin acts synergistically with H₂O₂ to decrease cell viability and metabolic activity and to increase apoptosis in NB4 cells. In contrast, the early oxidative effects of *t*-BHP are neutralised by esculetin, protecting human leukaemia NB4 cells from apoptosis. Esculetin seems to restrict the increase in peroxides caused by H₂O₂ or *t*-BHP in the time interval analyzed. These results contribute to a better understanding of the cytotoxic effect caused by esculetin on NB4 cells. At the same time, the early neutralisation of exogenous oxidants could be of interest to prevent diseases related to oxidative stress imbalance.

1. Introduction

Intrinsic and microenvironmental oxidative stress has a critical role in cancer development. Reactive oxygen species (ROS) may support survival of tumour cells but also may kill them. The therapeutic use of the cytotoxic action of ROS is exemplified by therapies based on oxidative stress, such as radiotherapy and photodynamic therapy [1]. The imbalance between the production of ROS and antioxidant defence systems in the cells seems to be also implicated in the pathogenesis of a variety of other diseases like cardiovascular diseases, Alzheimer's disease, Parkinson's disease, human aging, and progression of cancer [2–4]. ROS are also molecules modulating many cellular signalling pathways [1]. Disturbance of intracellular redox balance by both oxidants and antioxidants appears to affect cell fate [5]. Therefore, both pro- and antioxidant therapies have been proposed for treating cancer [6].

It has long been proven that cancer cells display a prooxidative state caused by an enhanced aerobic metabolism [7]. Human NB4 cell line, derived from acute promyelocytic leukaemia, possesses increased ROS levels compared to

noncancerous cells [8, 9]. In this cell line ROS promote cellular proliferation by protecting from apoptosis and contribute to cancer development [9]. Thus, a good strategy against this leukaemia might be the use of antioxidant compounds to reduce oxidative stress and promote cell death [6]. A similar approach has been the use of phenolic compounds with antioxidant activity to decrease cell proliferation, by inducing apoptosis, in human erythromegakaryocytic leukaemia cells, characterised by very high intracellular ROS levels [10].

Coumarins are natural benzopyrones that bind weakly to a wide variety of enzymes and receptors in the organism, so exhibiting therapeutic potential as anticoagulant, anticancer, antioxidant, anti-inflammatory, antibacterial, and antiviral activities, among others [11, 12]. Their antioxidant properties with the ability to inhibit lipid peroxidation and to act as ROS scavengers [13, 14] seem to be related to the position and number of hydroxyl groups. Thus esculetin (6,7-dihydroxycoumarin) shows higher antioxidant activity than other coumarins [14]. Esculetin is present in many plants traditionally used in natural medicine, displaying cytoprotective properties against oxidative stress-induced cell damage [14–16], whereas it induces apoptosis in other cell

lines such as adipocyte 3T3-L1 cells [17], human leukaemia U937 cells [18, 19], and HL-60 cells [20]. Esculetin shows other biological activities such as inhibition of lipoxygenase and tyrosinase [21, 22].

In a previous study, we demonstrated a proapoptotic activity of esculetin on NB4 cells that produced significant cell death by apoptosis, involving the mitochondrial pathway and modulating ROS [23]. We also studied whether changes in ROS levels could be related to esculetin-induced apoptosis. Pretreatment of NB4 cells with esculetin prevented the oxidative effect of exogenous ROS generated from *tert*-butyl hydroperoxide (*t*-BHP). On the contrary, esculetin potentiated superoxide anion production from H_2O_2 as well as apoptosis induced by H_2O_2 on NB4 cells. Both oxidants, *t*-BHP and H_2O_2 , increased ROS in NB4 cells via different mechanisms. *t*-BHP is a short chain analogue of lipid hydroperoxides, similar to peroxidised fatty acids whereas H_2O_2 is also generated under oxidative stress conditions modulating fundamental intracellular processes [13]. Therefore, the objective of the current study was to analyze the effect of esculetin acting simultaneously with such oxidants and to observe the early events in the mechanism of esculetin-induced apoptosis. This approach may yield a better understanding of the mechanism of cytotoxicity exerted by esculetin on NB4 cells.

2. Material and Methods

2.1. Reagents. Esculetin (6,7-dihydroxycoumarin, 98% purity) was obtained from Sigma-Aldrich (Steinheim, Germany), prepared as 196 mM stock solution in dimethyl sulfoxide and stored at $-20^{\circ}C$. H_2O_2 (hydrogen peroxide) and *t*-BHP (*tert*-butyl hydroperoxide) were also purchased from Sigma-Aldrich and prepared as 880 mM and 78 mM stock solutions, respectively, in distilled water at the time of use. Fluorescent probes hydroethidine (DHE) and 2',7'-dichlorodihydrofluorescein diacetate (DCFDA) were obtained from Molecular Probes (Eugene, Oregon, USA).

2.2. Cell Culture. The NB4 leukaemic cell line was obtained from the American Type Culture Collection (Manassas, VA, USA). Cells were maintained in culture at a density of 3×10^5 cells/ml in RPMI medium (Gibco-Life Technologies) supplemented with 10% foetal bovine serum (FBS), 1% penicillin/streptomycin, and 0.02 mg/ml gentamicin at $37^{\circ}C$ in a humidified 5% CO_2 atmosphere.

2.3. Cell Treatments for Flow Cytometric and MTT Analyses. NB4 cells (5×10^5 /ml) were treated with 100 μM esculetin and either 1 mM H_2O_2 or 100, 250, or 500 μM *t*-BHP. These treatments were carried out independently or in combination, for different times (15 min, 30 min, 1 h, and 2 h). Appropriate controls were run in parallel.

2.4. Cell Viability. Viability of NB4 cells treated with esculetin was determined by measuring the level of impermeability to propidium iodide (PI, Calbiochem) by flow cytometry. Fluorescence increase will be due to necrotic cells which are permeable to PI. After the different treatments, 2.5×10^5 cells

were washed with 500 μl phosphate buffered saline (PBS) and resuspended in 300 μl PBS; then 15 μl of propidium iodide was added and the fluorescence was measured using a Becton Dickinson FACScalibur flow cytometer (San José, CA, USA) and processed using the WinMDI 2.8 software.

2.5. Cell Metabolic Activity. Cell metabolic activity was determined by the colorimetric MTT assay kit (Roche Diagnostics, SL) in which tetrazolium salts (MTT) are reduced to formazan by the mitochondrial succinate-reductase system. Cells were seeded in 96-well microplates and after treatment were incubated with 10 μl of MTT labelling reagent for 4 h, and then the formazan dye formed was dissolved with 100 μl of solubilisation solution. The absorbance was measured using a multiwell spectrophotometer, to estimate the number of viable cells.

2.6. Analysis of Apoptosis by Annexin-FITC Cytometry Assay. Apoptosis in NB4 cells was quantified by the presence of phosphatidylserine on the outer side of the membrane using the Annexin V-FITC Apoptosis Detection Kit (BioVision). After treatments with esculetin, 2.5×10^5 cells were sedimented at 1200 rpm for 5 minutes and incubated with 500 μl 1x annexin Binding Buffer and 1 μl annexin-FITC for 5 min at room temperature in the dark. Then, 10 μl of PI was added and apoptotic cells were quantified by fluorescence by flow cytometry as indicated before. The results were analyzed using the WinMDI 2.8 software.

Lower left quadrant (FITC⁻/PI⁻) corresponds to living cells, lower right quadrant (FITC⁺/PI⁻) corresponds to early apoptotic cells, upper right quadrant (FITC⁺/PI⁺) corresponds to late apoptotic cells, and upper left quadrant (FITC⁻/PI⁺) corresponds to necrotic cells. Quadrants were defined based on the major population of live control cells (low PI; low annexin), and their position kept constant for all other samples.

2.7. Measurement of Intracellular ROS Levels. To measure intracellular superoxide anion ($\cdot O_2^-$), cells were incubated with 2 μM DHE (dihydroethidine) during the last 15 min of the different treatments and fluorescence detected by flow cytometry. Intracellular ROS levels (mainly peroxides) were detected using the fluorescent probe DCFDA (2',7'-dichlorodihydrofluorescein diacetate), a fluorogenic dye that measures hydroxyl, peroxy, and other reactive oxygen species (ROS) activity within the cell. After diffusion into the cell, DCFDA is deacetylated by cellular esterases and later oxidised by ROS into 2',7'-dichlorofluorescein, a highly fluorescent compound. After treatment, cells were incubated with 10 μM DCFDA for 30 min at $37^{\circ}C$ and then washed with PBS and fluorescence intensity was measured by flow cytometry.

2.8. Statistical Analysis. Data are expressed as the mean $\pm$ standard error of the results of at least three independent experiments. Significance of differences was determined using the Tukey-Kramer test. Comparisons between treated and untreated cells are labelled with * in the figures. Comparisons of cells treated with esculetin plus oxidant versus esculetin alone are labelled with +. Comparisons of cells

FIGURE 1: Cell viability, measured by flow cytometry as impermeability to PI, of NB4 cells treated with esculetin alone or combined with oxidants. (a) Viability of NB4 cells treated with esculetin alone or combined with H₂O₂. Cells (5×10^5 /ml) were treated with 100 μM esculetin alone or in combination with 1 mM H₂O₂ for different times. (b) Viability of NB4 cells treated with esculetin alone or combined with *t*-BHP. Cells (5×10^5 /ml) were treated with 100 μM esculetin, 100, 250, and 500 μM *t*-BHP, or both compounds, for the indicated times. Data are expressed as mean ± SEM of three independent experiments. *Treated versus untreated cells; * $p < 0.05$; ** $p < 0.01$. +Treated with esculetin plus oxidant versus esculetin alone; + $p < 0.05$; ++ $p < 0.01$. #Treated with esculetin plus oxidant versus oxidant alone; # $p < 0.05$; ## $p < 0.01$.

treated with esculetin plus oxidant versus the same oxidant alone are labelled with #. In all cases, one and two symbols indicate $p < 0.05$ and $p < 0.01$, respectively.

3. Results

3.1. Viability of NB4 Cells Treated with Esculetin, H₂O₂, and *t*-BHP. Previous studies showed that the lowest concentration of esculetin that decreased both cell viability (to 75%) and metabolic activity (to 65%) of human leukaemia NB4 cells was 100 μM applied for 19 h [23]. Such a concentration was hence chosen to study the early influence of ROS on the esculetin-induced apoptosis of NB4 cells. To this purpose, cells were treated with esculetin and either H₂O₂ or *t*-BHP, alone or in combination with esculetin, for different time intervals ranging from 15 min to 2 h.

As it can be observed in Figure 1(a), 1 mM H₂O₂ produced a decrease in cell viability from 15 min to 2 h of treatment. In this interval, esculetin did not produce any changes in cell viability. However, when cells were simultaneously treated with both H₂O₂ and esculetin a significant decrease of viability was observed.

Correspondingly, *t*-BHP (100, 250, or 500 μM) reduced cell viability from 30 min to 2 hours, and such reduction was abolished by esculetin (Figure 1(b)).

Therefore, esculetin is potentiating the effect of H₂O₂ while it protects cells from *t*-BHP.

3.2. Metabolic Activity of NB4 Cells Treated with Esculetin, H₂O₂, and *t*-BHP. Metabolic activity was measured by the

MTT assay. Treatment of NB4 cells with 1 mM H₂O₂ reduced dramatically the metabolic activity (Figure 2(a)). Esculetin reduced cell metabolic activity only marginally, even after 2 hours (Figure 2(a)). The simultaneous cotreatment with these two compounds showed a reduction similar to that produced by H₂O₂ alone.

Treatment with *t*-BHP also reduced significantly metabolic activity in a concentration- and time-dependent manner (Figure 2(b)). Again, the effect of esculetin alone was weak even at 2 hours. However, when applied simultaneously with *t*-BHP an early moderate reduction of metabolic activity was observed, somewhat reverted a long time.

Hence, esculetin does not prevent the loss of metabolic activity caused by H₂O₂ but it does protect cells from *t*-BHP.

3.3. Induction of Apoptosis on NB4 Cells Treated with Esculetin and H₂O₂. Figure 3 shows that 1 mM H₂O₂ produced a time-dependent increase in apoptosis as measured by double labelling with propidium iodide and annexin-FITC (Figure 3(a)). H₂O₂ treatment increased apoptosis already at 15 min. Simultaneous treatment with esculetin strengthened apoptosis induced by H₂O₂ (Figure 3(b)), mainly early apoptosis. Therefore, esculetin potentiates apoptosis induced by H₂O₂ in a time-dependent way.

3.4. ROS Production in NB4 Cells Treated with Esculetin and H₂O₂. Levels of superoxide anion and peroxides were measured using the probes DHE and DCFDA, respectively. Figure 4(a) shows superoxide anion levels after treatment

FIGURE 2: Metabolic activity, measured by MTT assay, of NB4 cells treated with esculetin alone or combined with oxidants. (a) Metabolic activity of NB4 cells treated with esculetin and H₂O₂. Cells (5×10^5 /ml) were exposed to 100 μM esculetin, 1 mM H₂O₂, or both, for different times. (b) Metabolic activity of NB4 cells treated with esculetin and *t*-BHP. Cells (5×10^5 /ml) were exposed to 100 μM esculetin or 100, 250, and 500 μM *t*-BHP separately or in combination, for the indicated times. Data are expressed as mean $\pm$ SEM of four independent experiments. *Treated versus untreated cells; ** $p < 0.01$. †Treated with esculetin plus oxidant versus esculetin alone; †† $p < 0.01$. #Treated with esculetin plus oxidant versus oxidant alone; ## $p < 0.01$.

FIGURE 3: Apoptosis and necrosis of NB4 cells treated with esculetin and H₂O₂. Cells (5×10^5 /ml) were treated with 100 μM esculetin, 1 mM H₂O₂, or the combination of both, for the indicated times. Cells were then stained with FITC-conjugated annexin and PI and measured by flow cytometry. The results show one representative experiment out of three. Vertical axes: propidium iodide intensity (log scale); horizontal axes: annexin-FITC intensity (log scale).

FIGURE 4: Intracellular ROS levels in NB4 cells treated with esculetin and H₂O₂. Cells (5×10^5 /ml) were treated with 100 μ M esculetin, 1 mM H₂O₂, or the combination of both, for different times, and then analyzed by flow cytometry using a specific probe. (a) To measure intracellular superoxide levels, 2 μ M DHE was added during the last 15 min of the esculetin treatment. (b) To analyze intracellular peroxides, 10 μ M DCFDA was added for 30 minutes at 37°C after the treatments. One representative experiment is shown out of five. Vertical axes: number of events (linear scale); horizontal axes: fluorescence intensity (log scale). F.I. represents mean fluorescence intensity.

FIGURE 5: Apoptosis and necrosis of NB4 cells treated with esculetin and *t*-BHP. Cells (5×10^5 /ml) were treated with 100 μ M esculetin or 100 or 500 μ M *t*-BHP separately or in combination, for different times. Cells were then stained with FITC-conjugated annexin and PI and measured by flow cytometry. One representative experiment is shown out of three. Vertical axes: propidium iodide intensity (log scale); horizontal axes: annexin-FITC intensity (log scale).

with esculetin and/or H_2O_2 . As it can be observed, H_2O_2 produced an early and definite increase in superoxide between 15 min and 1 hour. Esculetin did not change the fluorescence profile obtained with DHE in comparison to control cells. Simultaneous cotreatments with esculetin and H_2O_2 seem not to have any effects over that produced by H_2O_2 alone at any time of treatment.

When using DCFDA as a probe for testing peroxide levels, an early increase is seen at 15 min that later decreases at 2 hours. Esculetin alone does not produce peroxides but in combined treatments with H_2O_2 it decreases peroxide levels in comparison to the control cells (Figure 4(b)).

Therefore, esculetin does not neutralise superoxide anions produced by H_2O_2 in the time interval analyzed, while it prevents the increase of peroxides.

3.5. Induction of Apoptosis on NB4 Cells Treated with Esculetin and *t*-BHP. Figure 5 shows apoptosis induction by treatments with *t*-BHP. As it can be observed, *t*-BHP at both 100 and 500 μ M produced a time-dependent and concentration-dependent increase in apoptosis of NB4 cells. However, simultaneous treatment with esculetin did not display apoptosis even for the highest time and concentration of *t*-BHP. Therefore, esculetin is protecting cells against *t*-BHP-induced apoptosis.

3.6. Free Radical Production in NB4 Cells Treated with Esculetin and *t*-BHP. Levels of superoxide anion and peroxides after incubation with esculetin and *t*-BHP were measured (Figure 6). Treatments with *t*-BHP increased both superoxide radical and peroxides in a concentration- and time-dependent way. Simultaneous treatment of NB4 cells with *t*-BHP and esculetin neutralised both superoxide anion and peroxides at any *t*-BHP concentration and time assayed. Esculetin seems hence to compensate the oxidative action of *t*-BHP by neutralising the peroxides generated from it.

4. Discussion

We had previously demonstrated that esculetin is able to prevent ROS production in NB4 human leukaemia cells. The reduction of ROS was temporally followed by cell growth arrest and caspase-dependent apoptosis via the intrinsic mitochondrial pathway. Apoptosis was preceded by an early (6 h) loss of mitochondrial membrane potential with a release of cytochrome c to the cytosol [23]. Esculetin also reduced Bcl2/Bax ratio, in correlation with the increased apoptosis. To know whether the antioxidant activity of esculetin can modulate death of NB4 cells, we increased the oxidative stress by adding *t*-BHP or H_2O_2 to cells that were pretreated with esculetin. Thus, we have studied the early events (from 30 min to 2 h) of the cotreatment with esculetin and these two exogenous oxidants in order to understand the cytotoxic mechanism of esculetin.

Our results show that, from the earliest time of 15 min of cotreatment, esculetin acts synergistically with H_2O_2 decreasing cell viability and metabolic activity (Figures 1(a) and 2(a)) and increasing apoptosis (Figure 3). This suggests a possible implication of ROS in esculetin-induced apoptosis of NB4 cells. H_2O_2 increased superoxide anion and peroxides at early times of treatment. That increase of peroxides, but not of superoxide, is neutralised by esculetin, potentiating apoptosis. The increase may be due to production of superoxide anion in the presence of esculetin (Figure 3) in comparison to H_2O_2 -independent treatment. On the contrary, in human hepatoma HepG2 cells esculetin has been shown to suppress cell damage induced by H_2O_2 [24].

It is well known that H_2O_2 , like some of the most common chemotherapeutic agents, induces ROS generation and cytotoxicity in cancer cells by producing DNA breaks, lipid peroxidation, PARP activation, glutathione depletion, alteration of the energy metabolism, and so on [11, 25]. However, the dose of H_2O_2 determines the type of cell death

FIGURE 6: Intracellular ROS levels in NB4 cells treated with esculetin and *t*-BHP. Cells ($5 \times 10^5/\text{ml}$) were treated with 100 μM esculetin or 100 or 500 μM *t*-BHP or the combination of both, for the indicated times, and then analyzed by flow cytometry using a specific probe. (a) To measure intracellular superoxide levels, 2 μM DHE was added during the last 15 min of the esculetin treatment. (b) To quantify intracellular peroxides, 10 μM DCFDA was added for 30 minutes at 37°C after the treatments. One representative experiment is shown out of three. Vertical axes: number of events (linear scale); horizontal axes: fluorescence intensity (log scale). F.I. represents mean fluorescence intensity.

and while low doses trigger apoptosis, higher doses cause severe oxidative modifications which can be irreversible and lead to necrosis [26]. For example, in HeLa cells a low H_2O_2 concentration induces caspase-dependent cell death [24].

Although the redox potentials of H_2O_2 and, particularly, *t*-BHP are hard to estimate accurately, peroxy radicals are commonly described as weakly oxidising, with a reduction potential ca. 0.43 V or 0.71 V for *t*-BHP* [27, 28]. Hydrogen peroxide is, in contrast, a stronger oxidant (1.78 V for the $\text{H}_2\text{O}_2/\text{H}_2\text{O}$ pair). This may account for the different ability of esculetin to neutralise the production of ROS and superoxide

from both agents. In addition, there is wide evidence that the actions of both oxidants towards cell damage are exerted in different manner, and H_2O_2 affects a wider range of mechanisms, resulting in higher toxicity in some cases [29].

Our results indicate that esculetin potentiates apoptosis of NB4 cells in H_2O_2 presence, correlating it with an increase in superoxide anion between 15 min and 1 h (Figure 4). It could be suggested that superoxide increase and therefore oxidative stress are an early event in the apoptotic mechanism of esculetin. On the other hand, *t*-BHP slightly increased the levels of superoxide anion along the 2 h of treatment and

peroxides from 30 min to 2 h of treatment. These increases were neutralised by esculetin from the earlier time of 15 min at both low and high *t*-BHP concentrations (Figure 6). Esculetin also neutralised the apoptosis (Figure 5) caused by *t*-BHP. This behaviour differs from that observed for H_2O_2 .

tert-Butyl hydroperoxide is a short chain analogue of lipid hydroperoxides that forms a stable radical species, and it has been widely used to induce oxidative stress in a variety of cells. In hepatocytes *t*-BHP is metabolised by cytochrome P450 generating free radical intermediates such as alkoxyl or peroxy radicals and superoxide anion [30]. *t*-BHP is also metabolised by glutathione peroxidase, leading to glutathione depletion [31]. In erythrocytes, *t*-BHP reacts with haemoglobin forming *t*-butoxyl radical which causes membrane lipid peroxidation and haemolysis [32]. In these and other cells, *t*-BHP also produces alteration of calcium homeostasis and DNA damage [33] leading to cell death via apoptosis or necrosis [16].

Esculetin seems to neutralise these radical species. According to our results, esculetin suppresses cell damage induced by *t*-BHP, as also reported in rat liver [30] and DNA damage by 1,2-dimethylhydrazine in rat colon [16]. The protective effect of esculetin against oxidation by *t*-BHP could be related to prevention of early membrane lipid peroxidation and the intracellular oxidation.

5. Conclusions

Combined treatments with esculetin and one of the exogenous oxidants H_2O_2 or *t*-BHP show that esculetin neutralises the early oxidative effects of *t*-BHP, protecting human leukaemia NB4 cells from apoptosis, whereas esculetin potentiates the oxidative stress and cell death induced by H_2O_2 . This suggests a different cytotoxic mechanism for these two oxidants. Any other mechanism independent of ROS scavenging could also be involved in the cytotoxic H_2O_2 effects potentiation by esculetin. These results could be relevant to design therapies against diseases related to oxidative stress imbalance.

Abbreviations

Annexin-FITC:	Fluorescein isothiocyanate coupled to annexin V
DCFDA:	2',7'-Dichlorodihydrofluorescein diacetate
DHE:	Hydroethidine
PI:	Propidium iodide
MTT:	3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
ROS:	Reactive oxygen species
<i>t</i> -BHP:	<i>tert</i> -Butyl hydroperoxide.

Competing Interests

The authors declare that they have no competing interests.

Acknowledgments

This work was supported in part by Grants F.I.S.-PI060119, CCG10-UAH/SAL-5966, and UAH2011/BIO-006. The authors also want to thank Isabel Trabado for her technical assistance in cytometric analyses (C.A.I. Medicina-Biología, Unidad de Cultivos, Universidad de Alcalá).

References

- [1] D. H. J. Forman, M. Maiorino, and F. Ursini, "Signaling functions of reactive oxygen species," *Biochemistry*, vol. 49, no. 5, pp. 835–842, 2010.
- [2] V. M. Labunskyy and V. N. Gladyshev, "Role of reactive oxygen species-mediated signaling in aging," *Antioxidants and Redox Signaling*, vol. 19, no. 12, pp. 1362–1372, 2013.
- [3] J. M. Matés, J. A. Segura, F. J. Alonso, and J. Márquez, "Roles of dioxins and heavy metals in cancer and neurological diseases using ROS-mediated mechanisms," *Free Radical Biology and Medicine*, vol. 49, no. 9, pp. 1328–1341, 2010.
- [4] I. Juránek, D. Nikitovic, D. Kouretas, A. W. Hayes, and A. M. Tsatsakis, "Biological importance of reactive oxygen species in relation to difficulties of treating pathologies involving oxidative stress by exogenous antioxidants," *Food and Chemical Toxicology*, vol. 61, pp. 240–247, 2013.
- [5] J. Fiedor and K. Burda, "Potential role of carotenoids as antioxidants in human health and disease," *Nutrients*, vol. 6, no. 2, pp. 466–488, 2014.
- [6] J. Wang and J. Yi, "Cancer cell killing via ROS: to increase or decrease, that is a question," *Cancer Biology and Therapy*, vol. 7, no. 12, pp. 1875–1884, 2008.
- [7] J. Zheng, "Energy metabolism of cancer: glycolysis versus oxidative phosphorylation (review)," *Oncology Letters*, vol. 4, no. 6, pp. 1151–1157, 2012.
- [8] M. Lanotte, V. Martin-Thouvenin, S. Najman, P. Balerini, F. Valensi, and R. Berger, "NB4, a maturation inducible cell line with t(15;17) marker isolated from a human acute promyelocytic leukemia (M3)," *Blood*, vol. 77, no. 5, pp. 1080–1086, 1991.
- [9] P. S. Hole, J. Zabkiewicz, C. Munje et al., "Overproduction of NOX-derived ROS in AML promotes proliferation and is associated with defective oxidative stress signaling," *Blood*, vol. 122, no. 19, pp. 3322–3330, 2013.
- [10] L. Zamboni, C. Caliceti, F. V. D. Segal et al., "Dietary phenolic acids act as effective antioxidants in membrane models and in cultured cells, exhibiting proapoptotic effects in leukaemia cells," *Oxidative Medicine and Cellular Longevity*, vol. 2012, Article ID 839298, 12 pages, 2012.
- [11] M. Liu, Y. Xu, X. Han et al., "Potent effects of flavonoid-rich extract from *Rosa laevigata* Michx fruit against hydrogen peroxide-induced damage in PC12 cells via attenuation of oxidative stress, inflammation and apoptosis," *Molecules*, vol. 19, no. 8, pp. 11816–11832, 2014.
- [12] K. N. Venugopala, V. Rashmi, and B. Odhav, "Review on natural coumarin lead compounds for their pharmacological activity," *BioMed Research International*, vol. 2013, Article ID 963248, 14 pages, 2013.
- [13] V. Rincheval, M. Bergeaud, L. Mathieu et al., "Differential effects of Bcl-2 and caspases on mitochondrial permeabilization during endogenous or exogenous reactive oxygen species-induced cell death: a comparative study of H_2O_2 , paraquat, *t*-BHP, etoposide and TNF- α -induced cell death," *Cell Biology and Toxicology*, vol. 28, no. 4, pp. 239–253, 2012.

- [14] S.-H. Kim, K.-A. Kang, R. Zhang et al., "Protective effect of esculetin against oxidative stress-induced cell damage via scavenging reactive oxygen species," *Acta Pharmacologica Sinica*, vol. 29, no. 11, pp. 1319–1326, 2008.
- [15] T. Kaneko, S. Tahara, and F. Takabayashi, "Suppression of lipid hydroperoxide-induced oxidative damage to cellular DNA by esculetin," *Biological and Pharmaceutical Bulletin*, vol. 26, no. 6, pp. 840–844, 2003.
- [16] T. Kaneko, S. Tahara, and F. Takabayashi, "Inhibitory effect of natural coumarin compounds, esculetin and esculin, on oxidative DNA damage and formation of aberrant crypt foci and tumors induced by 1,2-dimethylhydrazine in rat colons," *Biological and Pharmaceutical Bulletin*, vol. 30, no. 11, pp. 2052–2057, 2007.
- [17] J.-Y. Yang, M. A. Della-Fera, D. L. Hartzell, C. Nelson-Dooley, D. B. Hausman, and C. A. Baile, "Esculetin induces apoptosis and inhibits adipogenesis in 3T3-L1 cells," *Obesity*, vol. 14, no. 10, pp. 1691–1699, 2006.
- [18] C. Park, C.-Y. Jin, G.-Y. Kim et al., "Induction of apoptosis by esculetin in human leukemia U937 cells through activation of JNK and ERK," *Toxicology and Applied Pharmacology*, vol. 227, no. 2, pp. 219–228, 2008.
- [19] C. Park, C.-Y. Jin, H. J. Kwon et al., "Induction of apoptosis by esculetin in human leukemia U937 cells: roles of Bcl-2 and extracellular-regulated kinase signaling," *Toxicology in Vitro*, vol. 24, no. 2, pp. 486–494, 2010.
- [20] C.-Y. Chu, Y.-Y. Tsai, C.-J. Wang, W.-L. Lin, and T.-H. Tseng, "Induction of apoptosis by esculetin in human leukemia cells," *European Journal of Pharmacology*, vol. 416, no. 1-2, pp. 25–32, 2001.
- [21] K. Sekiya, H. Okuda, and S. Arichi, "Selective inhibition of platelet lipoxygenase by esculetin," *Biochimica et Biophysica Acta*, vol. 713, no. 1, pp. 68–72, 1982.
- [22] Y. Masamoto, H. Ando, Y. Murata, Y. Shimoishi, M. Tada, and K. Takahata, "Mushroom tyrosinase inhibitory activity of esculetin isolated from seeds of *Euphorbia lathyris* L," *Bioscience, Biotechnology and Biochemistry*, vol. 67, no. 3, pp. 631–634, 2003.
- [23] V. Rubio, E. Calviño, A. García-Pérez, A. Herráez, and J. C. Díez, "Human acute promyelocytic leukemia NB4 cells are sensitive to esculetin through induction of an apoptotic mechanism," *Chemico-Biological Interactions*, vol. 220, pp. 129–139, 2014.
- [24] S. R. Subramaniam and E. M. Ellis, "Esculetin-induced protection of human hepatoma HepG2 cells against hydrogen peroxide is associated with the Nrf2-dependent induction of the NAD(P)H: quinone oxidoreductase 1 gene," *Toxicology and Applied Pharmacology*, vol. 250, no. 2, pp. 130–136, 2011.
- [25] M. Tochigi, T. Inoue, M. Suzuki-Karasaki, T. Ochiai, C. Ra, and Y. Suzuki-Karasaki, "Hydrogen peroxide induces cell death in human TRAIL-resistant melanoma through intracellular superoxide generation," *International Journal of Oncology*, vol. 42, no. 3, pp. 863–872, 2013.
- [26] G.-Y. Li, B. Fan, and Y.-C. Zheng, "Calcium overload is a critical step in programmed necrosis of ARPE-19 cells induced by high-concentration H_2O_2 ," *Biomedical and Environmental Sciences*, vol. 23, no. 5, pp. 371–377, 2010.
- [27] K. Jomová, O. Kysel, J. C. Madden et al., "Electron transfer from all-trans β -carotene to the t-butyl peroxy radical at low oxygen pressure (an EPR spectroscopy and computational study)," *Chemical Physics Letters*, vol. 478, no. 4-6, pp. 266–270, 2009.
- [28] T. N. Das, T. Dhanasekaran, Z. B. Alfassi, and P. Neta, "Reduction potential of the *tert*-butylperoxy radical in aqueous solutions," *Journal of Physical Chemistry A*, vol. 102, no. 1, pp. 280–284, 1998.
- [29] A. Spector, W. Ma, F. Sun, D. Li, and N. J. Kleiman, "The effect of H_2O_2 and tertiary butyl hydroperoxide upon a murine immortal lens epithelial cell line, α TN4-1," *Experimental Eye Research*, vol. 75, no. 5, pp. 573–582, 2002.
- [30] W.-L. Lin, C.-J. Wang, Y.-Y. Tsai, C.-L. Liu, J.-M. Hwang, and T.-H. Tseng, "Inhibitory effect of esculetin on oxidative damage induced by t-butyl hydroperoxide in rat liver," *Archives of Toxicology*, vol. 74, no. 8, pp. 467–472, 2000.
- [31] O. Kučera, R. Endlicher, T. Roušar et al., "The effect of *tert*-butyl hydroperoxide-induced oxidative stress on lean and steatotic rat hepatocytes in vitro," *Oxidative Medicine and Cellular Longevity*, vol. 2014, Article ID 752506, 12 pages, 2014.
- [32] S. Singh, R. Agarwal, F. Jamal, S. Mehrotra, and R. Singh, "Effects of *tert*-butyl hydroperoxide on Ca^{+2} ATPase activity in isolated rat hepatocytes and its reversal by antioxidants," *Journal of Environmental Biology*, vol. 33, no. 5, pp. 867–870, 2012.
- [33] V. A. Sardão, P. J. Oliveira, J. Holy, C. R. Oliveira, and K. B. Wallace, "Vital imaging of H9c2 myoblasts exposed to *tert*-butylhydroperoxide—characterization of morphological features of cell death," *BMC Cell Biology*, vol. 8, article 11, 2007.