Corrigendum

Corrigendum to “Natural Scaffolds for Regenerative Medicine: Direct Determination of Detergents Entrapped in Decellularized Heart Valves”

Monica Dettin ¹, **Annj Zamuner**,¹ **Filippo Naso**,² **Antonella Monteleone**,² **Michele Spina** ³, and **Gino Gerosa**²

¹Department of Industrial Engineering, University of Padova, Padova, Italy

²Department of Cardiac, Thoracic and Vascular Science, University of Padova, Padova, Italy

³Department of Biomedical Sciences, University of Padova, Padova, Italy

Correspondence should be addressed to Monica Dettin; monica.dettin@unipd.it

Received 14 January 2018; Accepted 16 January 2018; Published 20 February 2018

Copyright © 2018 Monica Dettin et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Natural Scaffolds for Regenerative Medicine: Direct Determination of Detergents Entrapped in Decellularized Heart Valves” [1], there were errors in the Discussion section which should be corrected as follows:

- (i) “To the best of the present authors’ knowledge, the only report of bile acid surfactant determination within cell-depleted natural scaffolds concerns the evaluation of DOC in decellularized human saphenous vein by the dimethylmethylene blue (DMMB) assay for anionic detergents [30]” should be “To the best of the present authors’ knowledge, the only report of bile acid surfactant determination within cell-depleted natural scaffolds concerns the evaluation of DOC in decellularized human saphenous vein by the methylene blue (MB) assay for anionic detergents [30].”
- (ii) “The analytical evidence of sizeable amounts of DOC and other anionic surfactants in ECM preparations (as assessed here) would also provide information about possible positive interferences, as in the case of SDS [39], where DMMB is widely used in assays for GAG evaluation [40, 41]” should be “The analytical evidence of sizeable amounts of DOC and other anionic surfactants in ECM preparations (as assessed here) would also provide information about possible positive interferences of SDS [39] with a reagent

similar to MB, the dimethylmethylene blue (DMMB) widely used in assays for GAG evaluation [40, 41].”

- (iii) “On the other hand, the use of DMMB-based assays might present some advantages in nondestructive investigations and in terms of convenience when simple washing fluid mixtures of expected composition are to be analyzed [43]” should be “On the other hand, the use of MB-based assays might present some advantages in nondestructive investigations and in terms of convenience when simple washing fluid mixtures of expected composition are to be analyzed [43].”

References

- [1] M. Dettin, A. Zamuner, F. Naso, A. Monteleone, M. Spina, and G. Gerosa, “Natural scaffolds for regenerative medicine: Direct determination of detergents entrapped in decellularized heart valves,” *BioMed Research International*, vol. 2017, Article ID 9274135, 9 pages, 2017.