

Research Article

Characterization of a Microalgal UV Mutant for CO₂ Biofixation and Biomass Production

Feng Qi ^{1,2,3,4} Daoji Wu ¹ Ruimin Mu,^{1,4} Shuo Zhang,^{2,5} and Xinyi Xu⁶

¹School of Municipal and Environmental Engineering, Shandong Jianzhu University, Jinan 250101, China

²Shandong Provincial Engineering Centre on Environmental Science and Technology, Jinan 250061, China

³College of Environmental Science and Engineering, Ocean University of China, Qingdao 266100, China

⁴Shandong Co-Innovation Center of Green Building, Jinan 250101, China

⁵Shandong Geomineral Engineering Group Co. Ltd., Jinan 250200, China

⁶School of Environmental Science and Engineering, Shandong University, Jinan 250100, China

Correspondence should be addressed to Feng Qi; qifeng@sdjzu.edu.cn and Daoji Wu; wdj@sdjzu.edu.cn

Received 19 July 2018; Revised 19 October 2018; Accepted 25 November 2018; Published 23 December 2018

Academic Editor: Kok Tat Tan

Copyright © 2018 Feng Qi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The mutagenesis is an emerging strategy for screening microalgal candidates for CO₂ biofixation and biomass production. In this study, by 96-well microplates-UV mutagenesis, a mutant stemmed from *Scenedesmus obliquus* was screened and named as SDEC-1M. To characterize SDEC-1M, it was cultivated under air and high level CO₂ (15% v/v), and its parental strain (PS) was considered as control. Growth characterizations showed that SDEC-1M grew best in high level CO₂. It indicated that the mutant had high CO₂ tolerance (HCT) and growth potential under high level CO₂. Richer total carbohydrate content (37.26%) and lipid content (24.80%) demonstrated that, compared to its parental strain, SDEC-1M was apt to synthesize energy storage materials, especially under high CO₂ level. Meanwhile, the highest light conversion efficiency (approximately 18 %) was also obtained. Thus, the highest overall biomass productivities were achieved in SDEC-1M under high level CO₂, largely attributed to that the highest productivities of total lipid, total carbohydrate, and crude protein were also achieved in the meantime. By modified UV, therefore, mutagenized SDEC-1M was the better candidate for CO₂ biofixation and biofuel production than its parental strain.

1. Introduction

Energy shortage and climatic change have been greatly focused [1–3]. To solve the problems, the techniques of CO₂ capture and sequestration and clean energy alternative to fossil fuel have been widely applied, among which microalgae is one of the most interesting strategies due to being sustainable, environment-friendly, and noncompetitive with other edible feedstocks [1, 4–10].

Microalgae cultivation via autotrophic system with high level CO₂ is considered as a more viable approach for commercialization due to lower cultivation costs cut by free sunlight and carbon source, high contents of lipid or carbohydrate, acquiring high cell density and sequestering CO₂ [1, 7, 8, 11]. However, the growth of microalgae can also be inhibited by high level CO₂, due to acidified medium [12]. Thus, an ideal candidate should have tolerance for high

level CO₂. In previous reports, in addition to the isolation of microalgae from the region affected by atmospheric pollution source [1] and acclimation by bubbled gas with CO₂ gradually increasing [13], screening microalgae with high CO₂ tolerance (HCT) by genetic strategy coupled to high-throughput screening were emerging [3, 4, 13–17]. In consideration of the economic and technological feasibility of genetic manipulations [3, 13, 15], random mutagenesis was more employed to provide microalgae candidates, rather than targeted genetic manipulation [17]. For instance, some mutants with HCT were obtained by chemical [4], nuclear irradiation [13], plasmas [18], and UV mutagenesis [14, 15]. Following mutagenesis, general, surviving mutants should be isolated [4, 15], which means enormous work and risks from invasion of infectious bacteria and algae. According to novel 96-well microplates-UV mutagenesis, each cell was isolated and then mutagenized in a well of 96-well microplates [14].

Since each surviving colony in the closed well was pure strain, luckily, invasion and reisolation were avoided. The novel method can shorten operating time, simplify operation process, and maintain purity of isolated strains [14].

Mutants can access some characteristics which their parental strains do not have, for instance, high biomass productivities [17], high abilities to capture CO₂ [4], high light conversion efficiencies (*LCE*) [14, 17, 19], high CO₂ requiring (HCR) [14, 20], and high contents of lipid or carbohydrate [14–16]. These genetic characteristics affecting their ability to biofix CO₂ and product biomass are key to commercialize microalgae [16]. In view of randomness of mutagenesis, the above-mentioned characteristics might synchronize. Therefore, it is necessary that the mutants are characterized to test their commercialization potential of the CO₂ biofixation and biomass production.

In this study, we aimed to have microalgae strains for CO₂ biofixation and biomass production; *Scenedesmus obliquus* was mutagenized by 96-well microplates-UV mutagenesis. One competitive potential mutant was screened out. To characterize its growth, biochemical components, and *LCE*, moreover, the mutant and its parental strain were cultivated under air and high level CO₂ (15% v/v). After continuous subculture for five generations under 15% CO₂, its genetic stability was tested. According to the experimental results, finally, the mutant's suitability for CO₂ fixation and application of its biomass were evaluated.

2. Material and Methods

2.1. Microalgae Strains and Culture Medium. In this study, as a parental strain (PS), one freshwater microalgae *Scenedesmus obliquus* observed dominant in freshwater water systems [27] was obtained from FACHB-Collection of Institute of Hydrobiology, Chinese Academy of Sciences. By 96-well microplates-UV mutagenesis [14], several *S. obliquus* mutants survived in wells with low pH (4.5) BG11 culture medium, and the survival ratio was approximately 3%. The growth status and genetic stabilities of survivors cultivated under 15% (v/v) CO₂ were compared and the best-growing and good genetically stable mutant was selected and named as SDEC-1M and employed in this study. The modified SE media (Bristol's solutions) with 1000 mg L⁻¹ of NaNO₃ nitrate concentrations [28] were used, which can supply more nitrogen source during the cultivation period. In addition to NaNO₃, there are 75 mg of K₂HPO₄·3H₂O, 75 mg of MgSO₄·7H₂O, 25 mg of CaCl₂·2H₂O, 175 mg of KH₂PO₄, 25 mg of NaCl, 5 mg of FeCl₃·6H₂O, 1 mL of A₅ solution, 1 mL of Fe-EDTA, and 40 mL of soil extract in 958 mL of deionized water. 1L A₅ solution contained 2.86 g of H₃BO₃, 1.81 g of MnCl₂·4H₂O, 0.22 g of ZnSO₄·4H₂O, 79 mg of CuSO₄·5H₂O, and 39 mg of (NH)₆Mo₇O₂₄·4H₂O. 1L Fe-EDTA contained 10g of Na₂EDTA, 0.81 g of FeCl₃·6H₂O, and 500 mL of 0.1 M HCl. The soil extract was the filtered supernatant from boiled soil solution.

2.2. Culture Conditions. The parental strain cells or mutant cells were cultured in photobioreactors (inside diameter (ID), 120 mm; working height (H_w), 221 mm; working volume

(V_w, 2.5L)) [22] containing fresh medium. Then, they were kept in a phytotron at 25 ± 1°C. A continuous illumination was provided by a row of fluorescent lamps which were horizontally fixed on the wall at one side of the bottles. The illumination intensities on the photobioreactor surface toward and back to the light source were 47.25 and 2.7 μmol m⁻²s⁻¹, respectively, read by a photometer. The initial OD₆₈₆ (optical density at 686 nm) was 0.3 (approximately 73 mg L⁻¹ of biomass concentration). During cultivation, deionized water was added to keep working volume. Aeration was carried out by air or high level CO₂. Simulating flue gas with high level CO₂ (up to 15%, v/v) [10], 15% (v/v) CO₂ mixed by air and pure CO₂ that were prearranged in industrial cylinders were employed. Using gas flow meters (Sevenstar, Beijing, China), the flow rates were adjusted to 0.2 vvm (volume gas per volume culture per min) of air and 0.04 vvm of mixture, respectively.

2.3. Measurement Methods. Between 680 and 690 nm, Chang and Yang [29] and Akkerman [30] found absorption peaks of microalgal broth and a linear correlation between the biomass concentrations and the optical densities. Thus, the indirect estimation of biomass concentration by optical density (OD) between 680 and 690 nm has been widely used in the microalgal research [2, 14, 28]. The absorption peaks of PS and SDEC-1M were located at 686nm by spectral scanning with a UV-2450 spectrophotometer (SHIMADZU, Japan). OD₆₈₆ and the biomass concentration of PS and SDEC-1M broth diluted in different proportions were determined by a UV-2450 spectrophotometer (SHIMADZU, Japan) and weighing the dry mass, respectively. Then, (1) relating OD₆₈₆ to biomass concentration was established by the linear regression.

$$X = \text{OD}_{686} \times 254.1 - 3.182 \quad R^2 = 0.9927 \quad (1)$$

Sample was taken once every day. After proper dilution, the biomass concentrations (X, mg L⁻¹) of sample were indirectly calculated via (1). pH of the sample was determined using PHS-3C pH meter (Leici, Shanghai, China).

At the end of cultivation (day 7), the microalgae were harvested and the contents of their primary biochemical components were measured. Microalgal pellets were formed by centrifuging microalgae culture at 4000 rpm at -3°C for 10 min and were washed twice with 0.5 M of ammonium formate to desalinate. Then, the microalgal pellets were dried and ground into powder. Higher heating values (HHV) of microalgal powder were determined by an isotherm oxygen bomb calorimeter [31]. Lipids of the microalgae were extracted by solvent, and the total lipid contents were estimated gravimetrically using a modified method [27]. The total carbohydrate contents of the microalgae were measured by the phenol-sulfuric acid method [32]. The crude protein content in the biomass was calculated via (2).

$$\text{Protein content} = \text{Nitrogen content} \times 6.25 \quad (2)$$

where nitrogen content was measured according to the Kjeldahl method [33], and the factor 6.25 is the correlation

TABLE 1: Maximum biomass concentrations (X_{max}), maximum specific growth rates (μ_{max}), and maximum biomass productivities for *S. obliquus* parental strain (PS) and mutant (SDEC-1M) under air and 15% (v/v) CO₂ for 7 days. Each data indicates the mean \pm SD, which was measured from three independent cultures. Each value in parentheses indicates the time (d) in which the maximum value of the parameter occurs. Data in the same parameter followed by different letters are significantly different by Duncan's test ($p < 0.05$).

Strains	X_{max} (mg L ⁻¹)		μ_{max} (d ⁻¹)		$P_{overall}$ (mg L ⁻¹ d ⁻¹)	
	Air	15% (v/v) CO ₂	Air	15% (v/v) CO ₂	Air	15% (v/v) CO ₂
PS	182.52 \pm 5.85 (7) ^a	240.62 \pm 10.2 (7) ^b	0.21 \pm 0.05 (2) ^a	0.21 \pm 0.02 (4) ^a	15.96 \pm 1.29 (6) ^a	24.01 \pm 3.2 (7) ^b
SDEC-1M	169.13 \pm 10.74 (7) ^a	320.28 \pm 13.9 (7) ^c	0.2 \pm 0.02 (3) ^a	0.23 \pm 0.03 (2) ^a	13.73 \pm 1.89 (7) ^a	35.32 \pm 1.81 (7) ^c

between protein content and nitrogen content reported by previous studies [34, 35].

2.4. Calculations on Important Properties. The maximum biomass concentration (mg L⁻¹) was designated as X_{max} .

The overall biomass productivity ($P_{overall}$, mg L⁻¹ d⁻¹) was calculated via (3).

$$P_{overall} = \frac{(X_7 - X_0)}{7} \quad (3)$$

where X_7 is the biomass concentration on day 7; X_0 is the initial biomass concentration in mg L⁻¹; 7 is the cultured time (d).

Specific growth rate (μ_t , d⁻¹) in a day was calculated via (4). The maximum specific growth rate was designated as μ_{max} (d⁻¹).

$$\mu_t = \frac{(\ln X_t - \ln X_{t-1})}{1} \quad (4)$$

where X_t and X_{t-1} were the biomass concentration (mg L⁻¹) on day t and day $t-1$, respectively; 1 was the time (d) gone through from day $t-1$ to day t .

The light conversion efficiency (LCE, %) based photosynthetic active radiation was estimated via (5).

$$LCE = \frac{(HHV \times P_{overall} \times V_w \times 100)}{[(I^+ + I^-) \times k \times PAR \times A \times t]} \quad (5)$$

$$= 2.08 \times 10^{-2} \times HHV \times P_{overall}$$

where the units of HHV , $P_{overall}$ and V_w are J mg⁻¹, mg L⁻¹ d⁻¹ and L; I^+ and I^- are illumination intensities (mol m⁻² s⁻¹) on the photobioreactor surface toward and back to the light source, respectively; the constant k that converts illumination intensity to light energy density (W m⁻²) is 218800 J mol⁻¹ photons [36]; the coefficient of photosynthetic active radiation (PAR) is 48% [37]; A is the irradiated area (m²); t is 86400 seconds in a day.

Total lipid productivity (P_L , mg L⁻¹ d⁻¹), total carbohydrate productivity (P_C , mg L⁻¹ d⁻¹), and crude protein productivity (P_P , mg L⁻¹ d⁻¹) used to test potential of lipid, carbohydrate, and protein productions were calculated via (6), (7), and (8).

$$P_L = P_{overall} \times \text{total lipid content} \quad (6)$$

$$P_C = P_{overall} \times \text{total carbohydrate content} \quad (7)$$

$$P_P = P_{overall} \times \text{crude protein content} \quad (8)$$




FIGURE 1: Growth curves of *S. obliquus* parental strain (PS) and mutant (SDEC-1M) under air and 15% (v/v) CO₂ for 7 days. Each data indicates the mean \pm SD, which was measured from three independent cultures.

2.5. Statistical Analysis. The differences between parameters of PS and SDEC-1M determined under air and 15% CO₂ were assessed using one-way analysis of variance (ANOVA). A difference was considered statistically significant when $p < 0.05$. Duncan's test was performed to detect the statistical significance of differences ($p > 0.05$).

3. Results and Discussion

3.1. Growth Characterizations

3.1.1. Growth Characterization under Air. As shown in Figure 1, PS grew slightly better than SDEC-1M under air. More details were seen in Table 1; no significant differences on X_{max} , μ_{max} and $P_{overall}$ were observed between PS and SDEC-1M under air. Thus, there were no significant changes in growth characteristics of SDEC-1M under air by mutagenesis. Due to randomness of mutagenesis, obviously, SDEC-1M did not attain the high CO₂ requiring (HCR) characteristic closely related to defectiveness in CO₂ concentrating mechanisms (CCMs) [20]. Although it was confirmed that a UV mutant, *Chlorella vulgaris* SDEC-3M obtained by 96-well




FIGURE 2: Contents of total lipid, total carbohydrate, and crude protein in cell (a) and productivities (b, mg L⁻¹ d⁻¹) of overall biomass, total lipid, total carbohydrate, and crude protein in *S. obliquus* parental strain (PS) and mutant (SDEC-1M) under air and 15% (v/v) CO₂ for 7 days. Each data indicates the mean \pm SD, which was measured from three independent cultures. Data of the same component followed by different letters are significantly different by Duncan's test ($p < 0.05$).

microplates-UV mutagenesis, was a HCR mutant [14], this method is not an efficient method for screening HCR algae strains.

3.1.2. Growth Characterization under High Level CO₂. Some species in *Scenedesmus* screened for CO₂ fixation were reported [1, 24, 25, 38]. These species generally have two characteristics: survival in low pH medium caused by high level CO₂ (high CO₂ tolerance) and high CO₂ fixation efficiency that exhibits good growing ability [7, 12, 13]. As shown in Figure 1, PS and SDEC-1M both exhibited higher growth rates under 15% (v/v) CO₂, and the latter adapted to high level CO₂ faster and attained the linear growth phase earlier. More details were seen in Table 1; thus, although there was no significant difference in μ_{max} between SDEC-1M and PS under 15% CO₂, SDEC-1M maintained a longer logarithmic phase (from day 1 to day 6), and its X_{max} and $P_{overall}$ were significantly 33.10% and 47.09% higher than those of PS, respectively. These results demonstrate that longer logarithmic phase can increase the biomass productivity.

3.1.3. Growth Distinction under Air and High Level CO₂. As shown in Figure 1, PS or SDEC-1M grew better under 15% CO₂ than under air. More details were seen in Table 1; there were no significant differences on their μ_{max} between under air and under 15% CO₂, while μ_{max} occurred at different times. Comparing their performances under air, the X_{max} and $P_{overall}$ of PS and SDEC-1M were both significantly higher. It implies that their growth potentials are similar, but the performances under air are poorer. Insufficient carbon source should be the main reason to decrease the metabolism of microalgae under air [7].

3.2. Contents and Productivities of Biochemical Components. Due to their plenty compounds as lipids, carbohydrates,

and protein, microalgae have enormous potential for the sustainable production of food, fuels, and other biochemicals [17]. Compounds contents and productivities are essential parameters of evaluation on applied purposes of microalgae biomass [23]. As shown in Figure 2(a), whether under air or 15% CO₂, the protein contents in PS were significantly higher than their lipid and carbohydrate contents, which agreed with recent reports, representing 32-58% of protein contents in *Scenedesmus* sp. [5, 23, 26, 39, 40]. However, in mutant SDEC-1M, the contents of lipid and carbohydrate were significantly higher than in PS, while the protein contents were just the opposite. Thus, carbohydrates became the chief biochemical component in SDEC-1M. Meanwhile, the lipid contents in PS and SDEC-1M under 15% CO₂ were both significantly higher than those under air. It agrees with the previous findings that high CO₂ stress enhanced lipid production [7, 41]. These results imply that compared to its parental strain, SDEC-1M was apt to synthesize energy storage materials, especially under high CO₂ level.

Largely attributed to significant higher overall biomass productivities, the highest productivities of total lipid (8.76 ± 0.47 mg L⁻¹ d⁻¹), total carbohydrate (13.16 ± 0.82 mg L⁻¹ d⁻¹), and crude protein (10.14 ± 0.30 mg L⁻¹ d⁻¹) all occurred in SDEC-1M under 15% CO₂, even though crude protein content was lower. Considering that two strains were harvested in logarithmic phase or linear growth phase, the results imply that biomass productivity plays a leading role in the primary biochemical components production until stationary phase, which agreed with previous reports [2, 6, 42].

3.3. Genetic Stabilities under High Level CO₂. In view of the instability of mutagenized strains, especially in continuous subculture, their genetic stability should be tested [18]. Thus, to confirm its stability of biomass product, SDEC-1M, was continuously subcultured under 15% CO₂ for five

TABLE 2: Higher heating values (*HHV*) and light conversion efficiencies (*LCE*) for *S. obliquus* parental strain (PS) and mutant (SDEC-1M) under air and 15% (v/v) CO₂ for 7 days. Each data indicates the mean ± SD, which was measured from three independent cultures. Data in the same parameter followed by different letters are significantly different by Duncan's test ($p < 0.05$).

Strains	<i>HHV</i> (kJ g ⁻¹)		<i>LEC</i> (%)	
	Air	15% (v/v) CO ₂	Air	15% (v/v) CO ₂
PS	22.03 ± 0.47 ^a	23.1 ± 0.26 ^{bc}	7.2 ± 0.29 ^a	11.54 ± 0.74 ^b
SDEC-1M	22.53 ± 0.35 ^{ab}	24.43 ± 0.59 ^c	6.43 ± 0.62 ^a	17.93 ± 0.6 ^c




FIGURE 3: Contents of total lipid, total carbohydrate, and crude protein in cell (a) and productivities (b, mg L⁻¹ d⁻¹) of overall biomass, total lipid, total carbohydrate, and crude protein in SDEC-1M of the first generation and the fifth generation under 15% (v/v) CO₂ for 7 days. Each data indicates the mean ± SD, which was measured from three independent cultures.

generations. The characterized results in SDEC-1M of the first generation and the fifth generation were shown in Figure 3.

As shown in Figure 3, no significant changes were observed on parameters, such as contents of total lipid, total carbohydrate, and crude protein and productivities of overall biomass, total lipid, total carbohydrate, and crude protein in SDEC-1M between the first generation and the fifth generation. The results suggest SDEC-1M is genetically stable from the aspect of biomass production.

3.4. Light Conversion Efficiency (*LEC*). As shown in Table 2, there was no significant difference in *HHV* between PS and SDEC-1M under air, and similar results under 15% CO₂. Furthermore, *HHV* of SDEC-1M was significantly higher than that of PS whether under 15% CO₂ or under air. The results were similar to their growth characterizations. Thus, the highest overall biomass productivity (35.32 ± 1.99 mg L⁻¹ d⁻¹) (Figure 2) or the highest *HHV* (24.43 ± 0.59 KJ g⁻¹) were both obtained by SDEC-1M under 15% CO₂. As a result, the highest calculated value of *LCE* (17.93 ± 0.6 %) occurred in SDEC-1M under 15% CO₂.

The higher *LCE* means more light energy transferred into biomass under same light conditions. Thus, the high overall biomass productivity was obtained in SDEC-1M under 15% CO₂ (Section 3.2). Only photosynthetic active radiation (PAR) is usable radiation fraction (400–700 nm) wavelengths between 400 and 700 nm [43], accounting for 42.3%–45.8% of the total energy from the solar spectrum [44, 45]. Based on PAR, *LCEs* of most microalgae varied between 4–9% [45],

just in which *LCEs* of PS and SDEC-1M were under air. A few recorded higher *LCEs* of 21.6% [19] and 14.52% [11] were comparable with those under 15% CO₂ in this study. This result is close to the theoretical upper limit of conversion efficiencies from solar energy to chemical energy (26.7%–29.8%) previously estimated by Brennan and Owende [45] and Robertson et al. [37]. Interestingly, the higher *LCEs* all were obtained in the cases with additional CO₂ aeration. It shows that CO₂ level is the crucial factor for enhancing biomass production of algal strains with HCT characteristic.

3.5. Characteristics Comparison with Other Algae Strains. The main characteristics represented the growth potential and conversion efficiency of SDEC-1M and other algae strains were shown in Table 3. By comparison, it is found that the contents of lipid, carbohydrate, and protein of SDEC-1M are all moderate, representing the normal level of primary biochemical components in algae, especially *Scenedesmus* sp. Optimization and stress aiming at production enhancement of specific component, such as lipid [2, 22], carbohydrate [23], and protein [1, 26] were not implemented, and the cultured time is not long enough to attain the stable phase in this study. It implies that SDEC-1M has yet great potential to enhance production of lipid, carbohydrate, and protein.

4. Conclusions

S. obliquus SDEC-1M with genetic stability was obtained after UV mutagenesis and was cultivated under air and

TABLE 3: Comparison between characterizations of SDEC-1M and other algae strains.

Strain	Lipid content (%)	Carbohydrate content (%)	Protein content (%)	HHV (kJ g ⁻¹)	LEC (%)	Reference
SDEC-1M	24.80	37.26	28.74	24.43	17.93	This study
<i>Chlorella protothecoides</i>	-	-	-	21.1	-	[21]
<i>Chlorella</i> sp.	33.6	-	-	-	-	[22]
<i>Chlorella vulgaris</i>	32.7	-	-	-	-	[2]
<i>Chlorella vulgaris</i> SDEC-3M	19.15	42.48	-	22.1	14.52	[14]
<i>Cladophora fracta</i>	-	-	-	25.1	-	[21]
<i>Phaeodactylum tricornutum</i> UTEX 640	-	--	-	-	21.6	[19]
<i>Scenedesmus bajacalifornicus</i> BBKLP-07	25.81	26.19	32.89	-	-	[1] ^a
<i>Scenedesmus dimorphus</i>	12	53.7	17.4	-	-	[23]
<i>Scenedesmus obliquus</i>	19.80	25.39	31.26	-	-	[2]
<i>Scenedesmus obliquus</i>	27.5	-	-	-	-	[24]
<i>Scenedesmus obliquus</i>	-	-	-	22.9	-	[25]
<i>Scenedesmus</i> sp.	21	38	23	-	-	[5]
<i>Scenedesmus</i> sp.	25.4	-	49.97	-	-	[26] ^a

a: the optimal value under different conditions.

b: unit is 10⁶ cell/mL.

15% CO₂, with its parental strain used as control. SDEC-1M got the best growth performance and the highest LCE (17.93 %) under 15% CO₂, which confirms its high CO₂ tolerance and high CO₂ fixation efficiency. Meanwhile, the highest total carbohydrate and lipid contents (37.26% and 24.80 %, respectively) and productivity (13.16 and 8.76 mg L⁻¹d⁻¹, respectively) were obtained. These results confirmed SDEC-1M's ability to convert more efficiently light into energy storage materials. Compared to its parental strain, therefore, SDEC-1M is a more suitable candidate for CO₂ fixation and biomass production, especially biofuel production, which mitigates the global warming and energy shortage.

Abbreviations List

UV:	Ultraviolet
PS:	Parental strain
HCT:	High CO ₂ tolerance
LCE:	Light conversion efficiencies
HCR:	High CO ₂ requiring
OD:	Optical density
HHV:	Higher heating values
PAR:	Photosynthetic active radiation
ANOVA:	Analysis of variance.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

The financial assistance for this investigation was gratefully provided by National Key Research and Development Planning of China (2017YFF0209903), Key Research and Development Planning of Shandong Province (2016GSF117014 and 2016GSF117018), Shandong Province Water Conservancy Research and Technology Extension Project (SDSLKY201812), Science and Technology Plans of Ministry of Housing and Urban-Rural Development of the People's Republic of China (UDC2017031512), and the Innovation Team of the Co-Innovation Center for Green Building of Shandong Province in Shandong Jianzhu University.

References

- [1] L. Patil and B. Kaliwal, "Effect of CO₂ Concentration on Growth and Biochemical Composition of Newly Isolated Indigenous Microalga *Scenedesmus bajacalifornicus* BBKLP-07," *Applied Biochemistry and Biotechnology*, vol. 182, no. 1, pp. 335–348, 2017.
- [2] M. K. Lam, M. I. Yusoff, Y. Uemura et al., "Cultivation of *Chlorella vulgaris* using nutrients source from domestic wastewater for biodiesel production: Growth condition and kinetic studies," *Journal of Renewable Energy*, vol. 103, pp. 197–207, 2017.

- [3] S. R. Medipally, F. M. Yusoff, S. Banerjee, and M. Shariff, "Microalgae as sustainable renewable energy feedstock for biofuel production," *BioMed Research International*, vol. 2015, Article ID 519513, 13 pages, 2015.
- [4] C.-Y. Kao, S.-Y. Chiu, T.-T. Huang, L. Dai, L.-K. Hsu, and C.-S. Lin, "Ability of a mutant strain of the microalga *Chlorella* sp. to capture carbon dioxide for biogas upgrading," *Applied Energy*, vol. 93, pp. 176–183, 2012.
- [5] G. A. Oliveira, E. Carissimi, I. Monje-Ramírez, S. B. Velasquez-Orta, R. T. Rodrigues, and M. T. O. Ledesma, "Comparison between coagulation-flocculation and ozone-flotation for *Scenedesmus* microalgal biomolecule recovery and nutrient removal from wastewater in a high-rate algal pond," *Bioresource Technology*, vol. 259, pp. 334–342, 2018.
- [6] H. A. Abu Hajar, R. G. Riefler, and B. J. Stuart, "Cultivation of *Scenedesmus dimorphus* using anaerobic digestate as a nutrient medium," *Bioprocess and Biosystems Engineering*, vol. 40, no. 8, pp. 1197–1207, 2017.
- [7] L. D. Zhu, Z. H. Li, and E. Hiltunen, "Strategies for Lipid Production Improvement in Microalgae as a Biodiesel Feedstock," *BioMed Research International*, vol. 2016, Article ID 8792548, 8 pages, 2016.
- [8] W. Leong, J. Lim, M. Lam, Y. Uemura, and Y. Ho, "Third generation biofuels: A nutritional perspective in enhancing microbial lipid production," *Renewable & Sustainable Energy Reviews*, vol. 91, pp. 950–961, 2018.
- [9] A. Mohd-Sahib, J. Lim, M. Lam et al., "Mechanistic kinetic models describing impact of early attachment between *Chlorella vulgaris* and polyurethane foam material in fluidized bed bioreactor on lipid for biodiesel production," *Algal Research*, vol. 33, pp. 209–217, 2018.
- [10] X. B. Tan, M. K. Lam, Y. Uemura, J. W. Lim, C. Y. Wong, and K. T. Lee, "Cultivation of microalgae for biodiesel production: A review on upstream and downstream processing," *Chinese Journal of Chemical Engineering*, vol. 26, no. 1, pp. 17–30, 2018.
- [11] M. G. De Moraes and J. A. V. Costa, "Carbon dioxide fixation by *Chlorella kessleri*, *C. vulgaris*, *Scenedesmus obliquus* and *Spirulina* sp. cultivated in flasks and vertical tubular photobioreactors," *Biotechnology Letters*, vol. 29, no. 9, pp. 1349–1352, 2007.
- [12] G. Sibi, V. Shetty, and K. Mokashi, "Enhanced lipid productivity approaches in microalgae as an alternate for fossil fuels—a review," *Journal of the Energy Institute*, vol. 89, no. 3, pp. 330–334, 2016.
- [13] J. Cheng, Y. Huang, J. Feng, J. Sun, J. Zhou, and K. Cen, "Mutate *Chlorella* sp. by nuclear irradiation to fix high concentrations of CO₂," *Bioresource Technology*, vol. 136, pp. 496–501, 2013.
- [14] F. Qi, H. Pei, W. Hu, R. Mu, and S. Zhang, "Characterization of a microalgal mutant for CO₂ biofixation and biofuel production," *Energy Conversion and Management*, vol. 122, pp. 344–349, 2016.
- [15] H. Vigeolas, F. Duby, E. Kaymak et al., "Isolation and partial characterization of mutants with elevated lipid content in *Chlorella sorokiniana* and *Scenedesmus obliquus*," *Journal of Biotechnology*, vol. 162, no. 1, pp. 3–12, 2012.
- [16] K. Anandarajah, G. Mahendrapurumal, M. Sommerfeld, and Q. Hu, "Characterization of microalga *Nannochloropsis* sp. mutants for improved production of biofuels," *Applied Energy*, vol. 96, pp. 371–377, 2012.
- [17] L. Wobbe and C. Remacle, "Improving the sunlight-to-biomass conversion efficiency in microalgal biofactories," *Journal of Biotechnology*, vol. 201, pp. 28–42, 2014.
- [18] X. Zhang, X.-F. Zhang, H.-P. Li et al., "Atmospheric and room temperature plasma (ARTP) as a new powerful mutagenesis tool," *Applied Microbiology and Biotechnology*, vol. 98, no. 12, pp. 5387–5396, 2014.
- [19] F. G. A. Fernández, F. G. Camacho, J. A. S. Pérez, J. M. F. Sevilla, and E. M. Grima, "Modeling of biomass productivity in tubular photobioreactors for microalgal cultures: Effects of dilution rate, tube diameter, and solar irradiance," *Biotechnology and Bioengineering*, vol. 58, no. 6, pp. 605–616, 1998.
- [20] D. J. Bonfil, M. Ronen-Tarazi, D. Sültemeyer, J. Lieman-Hurwitz, D. Schatz, and A. Kaplan, "A putative HCO₃⁻ transporter in the cyanobacterium *Synechococcus* sp. strain PCC 7942," *FEBS Letters*, vol. 430, no. 3, pp. 236–240, 1998.
- [21] A. Demirbaş, "Production of biodiesel from algae oils," *Energy Sources, Part A: Recovery, Utilization, and Environmental Effects*, vol. 31, no. 2, pp. 163–168, 2009.
- [22] F. Han, H. Pei, W. Hu, M. Song, G. Ma, and R. Pei, "Optimization and lipid production enhancement of microalgae culture by efficiently changing the conditions along with the growth-state," *Energy Conversion and Management*, vol. 90, pp. 315–322, 2015.
- [23] L. M. Chng, K. T. Lee, and D. C. Chan, "Evaluation on Microalgae Biomass for Bioethanol Production," *IOP Conference Series: Materials Science and Engineering*, vol. 206, 2017.
- [24] G. Shen, M. Zhao, D. Cao, D. Qu, and M. Li, "Effect of nutrient regimes on desalination rate and lipid production of *scenedesmus obliquus* in saline water," *Desalination and Water Treatment*, vol. 93, pp. 93–99, 2017.
- [25] W. Thiansathit, T. C. Keener, S.-J. Khang, T. Ratpukdi, and P. Hovichitr, "The kinetics of *Scenedesmus obliquus* microalgae growth utilizing carbon dioxide gas from biogas," *Biomass & Bioenergy*, vol. 76, pp. 79–85, 2015.
- [26] Z. Wang, X.-D. Yao, M.-S. Miao et al., "Effects of urban wastewater dilution on growth and biochemical properties of *Scenedesmus* sp.," *Desalination and Water Treatment*, vol. 57, no. 60, pp. 29363–29370, 2016.
- [27] M. Song, H. Pei, W. Hu, and G. Ma, "Evaluation of the potential of 10 microalgal strains for biodiesel production," *Bioresource Technology*, vol. 141, pp. 245–251, 2013.
- [28] F. Qi, H. Pei, G. Ma, S. Zhang, and R. Mu, "Improving productivity and quality of biodiesel from *Chlorella vulgaris* SDEC-3M through customized process designs," *Energy Conversion and Management*, vol. 129, pp. 100–107, 2016.
- [29] E.-H. Chang and S.-S. Yang, "Some characteristics of microalgae isolated in Taiwan for biofixation of carbon dioxide," *Botanical Bulletin of Academia Sinica*, vol. 44, no. 1, pp. 43–52, 2003.
- [30] I. Akkerman, M. Janssen, J. Rocha, and R. H. Wijffels, "Photobiological hydrogen production: photochemical efficiency and bioreactor design," *International Journal of Hydrogen Energy*, vol. 27, no. 11–12, pp. 1195–1208, 2002.
- [31] A. M. Illman, A. H. Scragg, and S. W. Shales, "Increase in *Chlorella* strains calorific values when grown in low nitrogen medium," *Enzyme and Microbial Technology*, vol. 27, no. 8, pp. 631–635, 2000.
- [32] M. Dubois, K. Gilles, J. K. Hamilton, P. A. Rebers, and F. Smith, "A colorimetric method for the determination of sugars," *Nature*, vol. 168, no. 4265, p. 167, 1951.
- [33] AOAC, *Official Methods of Analysis*, AOAC international, Arlington, USA, 15th edition, 1990.
- [34] E. W. Becker, *Microalgae - Biotechnology and Microbiology. Cambridge Studies in Microbiology*, J. Sir Baddiley, N. H. Carey, I. J. Higgins, and W. G. Potter, Eds., vol. 10, Cambridge University Press, 2011.

- [35] S.-N. Mohd-Noor, C.-Y. Wong, J.-W. Lim et al., "Optimization of self-fermented period of waste coconut endosperm destined to feed black soldier fly larvae in enhancing the lipid and protein yields," *Journal of Renewable Energy*, vol. 111, pp. 646–654, 2017.
- [36] J. Degen, A. Uebele, A. Retze, U. Schmid-Staiger, and W. Trösch, "A novel airlift photobioreactor with baffles for improved light utilization through the flashing light effect," *Journal of Biotechnology*, vol. 92, no. 2, pp. 89–94, 2001.
- [37] D. E. Robertson, S. A. Jacobson, F. Morgan, D. Berry, G. M. Church, and N. B. Afeyan, "A new dawn for industrial photosynthesis," *Photosynthesis Research*, vol. 107, no. 3, pp. 269–277, 2011.
- [38] H.-W. Yen, P.-W. Chen, and L.-J. Chen, "The synergistic effects for the co-cultivation of oleaginous yeast-Rhodotorula glutinis and microalgae-Scenedesmus obliquus on the biomass and total lipids accumulation," *Bioresource Technology*, vol. 184, pp. 148–152, 2015.
- [39] F. A. Ansari, S. K. Gupta, M. Nasr, I. Rawat, and F. Bux, "Evaluation of various cell drying and disruption techniques for sustainable metabolite extractions from microalgae grown in wastewater: A multivariate approach," *Journal of Cleaner Production*, vol. 182, pp. 634–643, 2018.
- [40] R. A. E. Hamouda and G. W. Abou-El-Souod, "Influence of various concentrations of phosphorus on the antibacterial, antioxidant and bioactive components of green microalgae scenedesmus obliquus," *International Journal of Pharmacology*, vol. 14, no. 1, pp. 99–107, 2018.
- [41] A. Nakanishi, S. Aikawa, S.-H. Ho et al., "Development of lipid productivities under different CO₂ conditions of marine microalgae *Chlamydomonas* sp. JSC4," *Bioresource Technology*, vol. 152, pp. 247–252, 2014.
- [42] M. J. Griffiths, R. P. Van Hille, and S. T. L. Harrison, "The effect of nitrogen limitation on lipid productivity and cell composition in *Chlorella vulgaris*," *Applied Microbiology and Biotechnology*, vol. 98, no. 5, pp. 2345–2356, 2014.
- [43] K. J. McCree, "The action spectrum, absorptance and quantum yield of photosynthesis in crop plants," *Agricultural and Forest Meteorology*, vol. 9, pp. 191–216, 1971.
- [44] J. R. Bolton and D. O. Hall, "The maximum efficiency of photosynthesis," *Photochemistry and Photobiology*, vol. 53, no. 4, pp. 545–548, 1991.
- [45] L. Brennan and P. Owende, "Biofuels from microalgae—a review of technologies for production, processing, and extractions of biofuels and co-products," *Renewable & Sustainable Energy Reviews*, vol. 14, no. 2, pp. 557–577, 2010.



Hindawi

Submit your manuscripts at
www.hindawi.com

