

## Research Article

# Antioxidant and Tyrosinase Inhibitory Activities of Seed Oils from *Torreya grandis* Fort. ex Lindl.

Hong-Xin Cui <sup>1,2</sup>, Fang-Fang Duan,<sup>3</sup> Shan-Shan Jia,<sup>3</sup> Fang-Rong Cheng,<sup>1</sup> and Ke Yuan <sup>3</sup>

<sup>1</sup>College of Pharmacy, Henan University of Chinese Medicine, Zhengzhou 450046, China

<sup>2</sup>Collaborative Innovation Center for Respiratory Disease Diagnosis and Treatment & Chinese Medicine Development of Henan Province, Zhengzhou 450046, China

<sup>3</sup>Jiyang College of Zhejiang Agriculture and Forestry University, Zhu'ji 311800, China

Correspondence should be addressed to Ke Yuan; yuan\_ke001@126.com

Received 18 May 2018; Revised 14 July 2018; Accepted 28 August 2018; Published 18 September 2018

Academic Editor: Devanand P. Fulzele

Copyright © 2018 Hong-Xin Cui et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Torreya grandis* Fort. ex Lindl. is a plant belonging to the Taxaceae family and *Torreya grandis* cv. Merrillii is the only grafted and thoroughbred species belonging to this species. In this study, we extracted five different seed oils, including *T. grandis* seed oil (TGSO), *T. grandis* “Xiangyafei” seed oil (XYSO), *T. grandis* “Zhimafei” seed oil (ZMSO), *T. grandis* “Majus” seed oil (TGMSO), and *T. grandis* “cunguangfei” seed oil (CGSO) using physical pressure. The resulting extracts were analyzed to determine their fatty acid composition, antioxidant activity, and inhibitory activity towards tyrosinase. The results of the antioxidant activity assays revealed that XYSO and ZMSO exhibited much greater DPPH radical scavenging activity and ferric reducing power than TGSO. Notably, all five of the seed oils showed dose-dependent inhibitory activity towards tyrosinase. XYSO and TGSO gave the highest activities of all of the seed oils tested in the current study against monophenolase and diphenolase, with IC<sub>50</sub> values of 227.0 and 817.5 μg/mL, respectively. The results of this study show that wild TGSOs exhibit strong antioxidant and tyrosinase inhibition activities. These results therefore suggest that wild TGSOs could be used as a potential source of natural antioxidant agents and tyrosinase inhibitors.

## 1. Introduction

Tyrosinase (EC 1.14.18.0) is a copper-containing mixed-function oxidase that is ubiquitously expressed in animals, plants, and microorganisms. Furthermore, tyrosinase is a key rate-limiting enzyme that can catalyze enzyme browning and melanin synthesis. Tyrosinase also exhibits monophenolase and diphenolase activities, which catalyze the hydroxylation of L-tyrosine to L-DOPA and the oxidation of L-DOPA to dopaquinone, which can undergo nonenzymatic polymerization to give dark pigments [1, 2]. In humans, the overexpression of tyrosinase leads to the overproduction of melanin in the skin, which can trigger hyperpigmentation effects such as freckles, melasma, age spots, and melanoma [3]. Free radicals also play an important role in the biosynthesis of melanin, and several studies have shown that free radicals are involved in the catalytic reactions of tyrosinase to give dopaquinone, which can undergo various oxidation

reactions. Increasing in the number or the activity of the free radicals present in living systems can therefore lead to an increase in the production of melanin [4, 5]. It is noteworthy that antioxidants such as vitamin C and Trolox have been reported to exhibit potent inhibitory activities towards tyrosinase and melanin production and have consequently been used to inhibit melanogenesis [6]. Materials capable of inhibiting the activity of tyrosinase have attracted worldwide interest from researchers working in a variety of different fields, including biological medicine, agricultural sciences, chemistry, and pharmacology. Tyrosinase inhibitors capable of inhibiting the biosynthesis of melanin are used currently in various hyperpigmentation and cosmetic agents to control the formation of freckles [7, 8].

From an agricultural perspective, the overexpression of tyrosinase is related to the discoloration of some fruits and vegetables, as well as changes in their flavor and nutritional characteristics, which can have an adverse impact on their

commercial value. The development of new biocompatible tyrosinase inhibitors capable of reducing the production of quinone and hydroquinone to prevent enzymatic browning and improve the color and nutritional characteristics of food is therefore highly desirable [9, 10]. Several synthetic tyrosinase inhibitors have been reported in the literature, such as kojic acid, which is widely used as a skin-whitening agent in cosmetics. However, the use of this compound in this context has been associated with several serious side effects, including erythema and contact dermatitis [11, 12]. Further research is therefore required to identify natural and biocompatible tyrosinase inhibitors that can be used as whitening agents and antibrowning agents in the cosmetic and food industries.

*Torreya grandis* Fort. ex Lindl. are plants belonging to the Taxaceae family, which are also known as “chiguo”. Six different species and two varieties of *Torreya* have been reported to date. Three of these species (*Torreya fargesii*, *Torreya fargesii* var. *yunnanensis*, and *Torreya grandis*) and two varieties (*Torreya jackii* and *Torreya grandis* var. *Jiulongshanensis*) can be found in China; one species (*Torreya nucifera*) can be found in Japan; and two species (*Torreya taxifolia* and *Torreya californica*) can be found in America [13]. *Torreya grandis*, which has been cultivated in China for more than 1500 years, is a rare and unique species, which is mainly distributed across the hilly areas of subtropical China, which is located in Zhejiang, Fujian, Anhui, Jiangxi, Jiangsu, Hunan, Hubei, Guizhou, and other places in China; vast territories were distributed, among them, *Torreya grandis* Fort. ex Lindl is mainly distributed in Zhuji, Zhejiang province. In recent years, a large number of ancient trees of *Torreya grandis* Fort. ex Lindl have been found in this area; among them, there are 42 ancient trees over a thousand years old, and the oldest one has reached 1500 years [14, 15]. *Torreya grandis* cv. *Merrillii* is the only grafted and thoroughbred species of *Torreya grandis* Fort. ex Lindl, which has a wide variety of seedling species because of its broad range of characteristics. *Torreya grandis* Fort. ex Lindl is well known for its edible seeds of *Torreya grandis*, which have a unique nutty flavor and high nutritional value, leading to their use in traditional Chinese medicine [14, 16, 17]. It is noteworthy that there are still four kinds of *Torreya grandis* of wild mutation variety of Zhimafei, Cunguangfei, Xiangyafei, and Dayuanfei, in China, and that the seeds of these wild species have comparable flavor and quality characteristics to those of cultivated *Torreya grandis*. However, the seeds of these wild plants are unsuitable for human consumption, and these species have consequently been abandoned without reasonable utilization. Previous studies have demonstrated that *Torreya grandis* seeds have multiple biological properties including antioxidative, anti-inflammatory, antiatherosclerosis, antiviral, antifungal, anti-tumor and antihelminthic activities, because of their rich nutritional content and numerous bioactive fatty acid, protein, vitamin, and mineral components [16–20]. With this in mind, we investigated the inhibitory activities of TGSO and four wild *Torreya grandis* seed oils (TGSOs) towards tyrosinase to provide a platform for the rational utilization of wild *Torreya grandis*.

## 2. Materials and Methods

**2.1. Chemicals.** 1,1-Diphenyl-2-picrylhydrazyl (DPPH), tripyridyltriazine (TPTZ), 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic, Trolox, and mushroom tyrosinase (3610 U/mg, *Agaricus bisporus*) were purchased from Sigma (St. Louis, MO, USA). Kojic acid, L-ascorbic acid, L-tyrosine, and L-DOPA (L-3,4-dihydroxyphenylalanine) were obtained from Aladdin Co., Ltd (Shanghai, China). All of the other chemicals and solvents used in this study were purchased as the analytical reagent grade from Huadong Medicine Co., Ltd (Zhejiang, China).

**2.2. Collection and Preparation of Plant Material.** Four kinds *T. grandis* of wild mutation variety of Zhimafei, Cunguangfei, Xiangyafei, and Dayuanfei and only one cultivar of *Torreya grandis* after grafting were collected from Zhuji, Zhejiang province, China, in October, 2015. These seeds were subsequently identified and authenticated by Professor Dr. Pinzhang Deng at the Forestry Bureau Zhuji, Zhejiang Province. After authentication, the seeds were cleaned, hulled, and immediately dried in a microwave oven (three space heats for 1 min), before being placed in a drying oven at 65°C for 6 h. Twenty-gram samples of the dried seeds were then placed in a QYZ-230 fully automatic hydraulic oil press (Shandong, China) and pressed to obtain the squeezed seed oils.

**2.3. GC-MS Analyses of Squeezed Seed Oils.** GC-MS analysis was conducted on a Bruker SCION SQ 456 Gas Chromatograph (Bruker Daltonics Inc., Billerica, MA, USA), which was attached to a mass spectrometer. The gas chromatograph system was equipped with a nonpolar, HP-5 capillary column (30 m × 250 μm × 0.25 μm), which was coated with 5% phenyl-95% dimethylpolysiloxane, as well as a flame ionization detector (FID). The oven temperature was programmed as follows: 45°C for 2 min; 45 to 270°C at a rate of 10°C/min; and 270°C for 6 min. Helium was used as the carrier gas at the rate 1 mL/min. All of the oil samples were dissolved in hexane at a concentration of 0.1 mg/μL. The sample injection volume was set to 1 μL with a split ratio of 1:10. L

Mass spectra were recorded in the EI mode at 70 eV with a scan quality range of 50–550 amu. Temperature of the ionization source is 200°C, voltage of the detector is 350 V. Mass spectra data for the different peaks were cross referenced with those found in the NIST27 and WILEY7 libraries to identify the different components. Each mass spectrogram corresponding to each chromatographic peak is qualitatively determined by computer chart-base; the relative content of each component is calculated by the peak area normalization method according to its total ion current chart.

### 2.4. In Vitro Antioxidant Activities

**2.4.1. DPPH Free Radical Scavenging Activity Assay.** The DPPH free radical scavenging activities of the seed oils were measured using a procedure from the literature with minor modifications [21]. The samples were evaluated in

96-well microtiter plates. Stock solutions of the seed oils were prepared at a concentration of 1 mg/mL in dimethyl sulphoxide, DMSO. Standard solutions of Trolox were prepared for concentrations in the range of 0–2.5 mM. Two hundred microliter aliquots of each sample solution were separately added to the 96-well microtiter plate, followed by 50  $\mu$ L of a DPPH solution (1 mM in 50% ethanol), and the resulting mixtures were incubated in the dark for 45 min at room temperature. The absorbance of each well was then measured at 517 nm using an Infinite M 200 Microplate Reader (Tecan, Männedorf, Switzerland). The DPPH free radical scavenging potential was calculated using the following formula:

$$\begin{aligned} & \text{DPPH free radical scavenging activity (I\%)} \\ & = 1 - \frac{(A_p - A_c)}{A_{\max}} \times 100\% \end{aligned} \quad (1)$$

where  $A_p$  is the absorbance of the sample,  $A_c$  is the absorbance of vehicle, and  $A_{\max}$  is the absorbance of the DPPH solution. Each sample was tested three times ( $n=3$ ). The concentration of DPPH was calculated using a Trolox standard curve and expressed as the Trolox equivalent antioxidant capacity (TEAC) with units of mmol Trolox equivalents/g.

**2.4.2. Ferric Reducing Power (FRAP) Assay.** The FRAP assay was performed as described previously [22]. Fresh FRAP reagent was prepared each day by mixing 300  $\mu$ L of acetate buffer (pH 3.6, 0.3 mol/L) with 30  $\mu$ L of TPTZ (10 mmol/L) and 30  $\mu$ L of iron(III) chloride (20 mM). A small sample (10  $\mu$ L) was added to 300  $\mu$ L of the FRAP reagents described above, and the resulting mixture was incubated in the dark for 30 min. The absorbance of the mixture was then measured at 593 nm. Each sample was tested three times ( $n=3$ ), and the results were expressed as TEAC with units of mmol Trolox equivalents/g.

**2.4.3. Total Reducing Power (TRP) Assay.** TRP was determined using a previously published method [23]. Briefly, 500  $\mu$ L of each sample was treated with 2.5 mL of sodium phosphate buffer (0.2 mol/L, pH 6.6) and 2.5 mL of 1% potassium ferricyanide, and the resulting mixture was incubated at 50°C for 30 min. The sample was then treated with 2.5 mL of 10% trichloroacetic acid, and the resulting mixture was centrifuged at 4000  $\times$ g for 10 min. The upper layer (2.5 mL) was collected and mixed with 0.5 mL of 0.1% ferric chloride and 2.5 mL of distilled water. The absorbance of the resulting mixture was then measured at 700 nm against a blank. Several Trolox standards were prepared as calibration solutions with concentrations in the range of 0.04–0.80 mg/mL. Each sample was tested three times ( $n=3$ ), and the results were expressed as TEAC with units of mmol Trolox equivalents/g.

## 2.5. Tyrosinase Inhibitory Activity Assay

**2.5.1. Monophenolase and Diphenolase Activity Assay.** The tyrosinase inhibitory activity was evaluated using a previously

reported method with minor modifications [24]. L-DOPA and L-tyrosine were used as substrates in this assay. Forty microliters of L-DOPA (10 mM, for the diphenolase activity assay) or L-tyrosine (5.0 mM, for the monophenolase activity assay) was mixed with 80  $\mu$ L of phosphate buffer (0.1 M, pH 6.8) in a 96-well microtiter plate, and the resulting mixture was incubated for 10 min at 37°C. Forty microliters of seed oil (50, 100, 200, 400, and 800  $\mu$ g/mL in 50% DMSO) and 40  $\mu$ L of mushroom tyrosinase (250 U/mL, in PBS) were then added to each well on the plate, and the absorbance characteristics of the resulting mixtures were measured at 475 nm (for the diphenolase activity) or 490 nm (for the monophenolase activity) using an ELISA reader (Infinite M 200, Swiss Tecan) at 60-second intervals over a period of 120 min. PBS was added instead of the test sample as a blank control and kojic acid (50  $\mu$ g/mL) and ascorbic acid (50  $\mu$ g/mL) were used as positive controls. The inhibition for each enzyme assay was calculated as follows:

$$\% \text{ Inhibition} = \frac{(\text{Abs}_{\text{control}} - \text{Abs}_{\text{sample}})}{\text{Abs}_{\text{control}}} \times 100 \quad (2)$$

Each experiment was carried out in triplicate ( $n=3$ ). The  $IC_{50}$  value, i.e., the concentration of seed oil, kojic acid, or ascorbic acid required to inhibit the activity of the enzyme by 50%, was calculated from the dose-response curves by nonlinear regression analysis using version 5.0 of the GraphPad Prism software version 5.0 (GraphPad software Inc., San Diego, CA, USA).

**2.5.2. Kinetic Analysis of the Inhibition of Tyrosinase.** The inhibition kinetics of the different seed oils (50, 100, 200, 400, and 800  $\mu$ g/mL) towards tyrosinase were determined using different concentrations (0, 0.5, 1, 1.5, and 2 mM) of L-DOPA and L-tyrosine. The type of inhibition was analyzed using a Lineweaver–Burk plot. All of the experiments were performed in triplicate. The enzyme-inhibitor dissociation constant ( $K_i$ ) and  $K_m$  values were obtained from the slope of each plot and the y-intercept versus the seed oil concentration, respectively [25].


**2.6. Statistical Analyses.** Data were expressed as the mean values  $\pm$  SD. Statistical analyses were performed using SPSS 19.0 (SPSS Inc., Chicago, IL, USA) and version 5.0 of the GraphPad Prism Software (GraphPad).

## 3. Results and Discussion

**3.1. GC-MS Results of the Five Squeezed Seed Oils.** Plant seed oils continue to receive growing interest from researchers working in a variety of different fields, including chemists, pharmacists, and physicians because of their potential applications in the pharmaceutical, cosmetics, and food industries [26]. *Torreya grandis* cv. *Merrillii* is the only grafted and thoroughbred species of *Torreya grandis* Fort. ex Lindl to have been reported in the literature to date and its seeds possess a unique flavor, as well as being rich in nutrients. Despite their huge potential, wild *Torreya grandis* are currently poorly

TABLE 1: Main chemical components of the seed oils obtained from the five different varieties of *T. grandis*.

Compound	TGSO	XYSO	ZMSO	TGMSO	CGSO
oleic acid	30.52%	19.58%	8.07%	12.45%	20.51%
linoleic acid	25.94%	21.67%	7.52%	15.49%	0
palmitic acid	3.23%	8.65%	7.60%	0	7.27%
octadecanal	1.27%	10.77%	9.95%	33.50%	0
linolenic acid(LNA)	5.35%	4.21%	5.79%	0	2.02%
Cis-8,11,14-Eicosatrienoic acid	2.67%	3.24%	3.01%	1.13%	0.86%
Cis-5,11,14-Eicosatrienoic acid	13.93%	11.56%	10.33%	3.93%	3.57%
Oil yield	59.47%	56.99%	51.67%	35.01%	29.89%

FIGURE 1: Total ion chromatograms of the seed oils obtained from the five different varieties of *T. grandis*. 1: TGSO, 2: XYSO, 3: ZMSO, 4: TGMSO, and 5: CGSO.

utilized, especially those that cannot be eaten, with most of these plant resources simply going to waste. To develop a better understanding of the composition of the seed oils derived from these plants, we conducted a series of gas chromatography experiments, and the results are shown in Figure 1 and Table 1. The results showed that the chemical compositions of the seed oils derived from the five different varieties of *Torreya grandis* were roughly the same, although there were some subtle differences. Oleic acid, linolenic acid, and palmitic acid are the main fatty acids compositions in the five seed oils, in particular, the content of conjugate linoleic acid is relatively the highest, and it has been reported that linolenic acid has obvious tyrosinase inhibition activity [27]. The nonedible wild *Torreya grandis* seeds, such as XYSO and ZMSO, showed similar chemical compositions to TGSO. Although it would not be possible to directly commercialize the seed oils obtained from wild *Torreya grandis*, they could potentially be used to develop edible oils.

**3.2. Antioxidant Results of the Five Squeezed Seed Oils.** Oxidation is essential to many living organisms for the production of energy to fuel biological processes. However, free radicals and several other reactive oxygen species are continuously produced *in vivo* as a consequence of oxidation processes, and these reactive species can result in tissue damage and cell death. The oxidative damage caused by free radicals plays a major contributory role in aging and numerous debilitating diseases, including diabetes, cirrhosis, and cancer [28]. Although several enzymes (e.g., superoxide dismutase) and

antioxidant compounds (e.g., ascorbic acid) can protect most organisms against free radical-induced oxidative damage, these systems cannot prevent this damage completely. The use of antioxidant supplements or the consumption of foods containing antioxidants can therefore be used as a strategy to reduce the level of oxidative damage [29]. The antioxidant activities of the seed oils obtained from the five different varieties of *Torreya grandis* evaluated in the current study were measured using DPPH, FRAP, and TRP assays. The DPPH assay, which measures the scavenging ability of an antioxidant towards the stable DPPH free radical, is one of the most commonly used methods for evaluating the radical scavenging activity of an antioxidant. Furthermore, it is well known that radical scavenging ability correlates well with antioxidant activity [30]. Ferric reducing ability (FRAP) and the TRP are commonly used methods for assessing the reducing power of an antioxidant *in vitro*. The research groups of Yen and Siddhuraju reported that there is a close relationship between the reducing power and antioxidant activity of antioxidants, and that the antioxidant capacity generally increases with increasing reducing power [31, 32]. In this way, it is therefore possible to evaluate the antioxidant ability of an antioxidant based on its reducing power. As shown in Table 2, the antioxidant capacities of the seed oils obtained from the five different varieties of *Torreya* were weaker than that of the positive control ascorbic acid. However, the DPPH scavenging capacities of the seed oils obtained from the five different varieties of *Torreya* were similar. The FRAP and TRP properties of the different seed oils also showed a similar trend, and there were significant differences ( $P < 0.05$ ) between the seed oils obtained from the different varieties of *Torreya* in one assay. The DPPH free radical scavenging abilities of the oils tested in the current study were found to be of the following order: ascorbic acid > XYSO > TGSO > ZMSO > TGMSO > CGSO; whereas the FRAP and TRP properties of the five different squeezed oils were of the order: ascorbic acid > TGSO > XYSO > ZMSO > TGMSO > CGSO.

**3.3. Tyrosinase Inhibitory Activities of the Five Squeezed Seed Oils.** The effects of the five different varieties of seed oils on the oxidation reactions of L-DOPA and L-tyrosine using mushroom tyrosinase were determined experimentally. Tyrosinase is an important enzyme that plays a big role in melanogenesis within melanocytes. Furthermore,

TABLE 2: Result of the antioxidant activity assays.

Assays	TGSO	XYSO	ZMSO	TGMSO	CGSO	Ascorbic acid	Kojic acid
DPPH(mgTEs/g oil)	42.12±4.36 <sup>c</sup>	56.73±0.67 <sup>b</sup>	35.35±0.99 <sup>d</sup>	24.87±0.88 <sup>e</sup>	22.34±0.71 <sup>f</sup>	194.56±3.22 <sup>a</sup>	-
FRAP(mgTEs/g oil)	690.32±41.32 <sup>b</sup>	667.89±50.21 <sup>c</sup>	588.91±33.21 <sup>d</sup>	401.67±46.04 <sup>e</sup>	355.56±44.87 <sup>f</sup>	1135.41±55.68 <sup>a</sup>	-
TRP(mgTEs/g oil)	199.76±35.12 <sup>b</sup>	186.77±31.23 <sup>c</sup>	151.03±30.76 <sup>d</sup>	111.43±11.56 <sup>e</sup>	109.61±23.98 <sup>f</sup>	385.98±41.21 <sup>a</sup>	-

TEAC values (mg TEs/g) of the antioxidants derived from the five different varieties of *T. grandis*.

melanogenesis plays an important role in protecting the skin from sun-related injuries and is principally responsible for skin color. However, abnormal hyperpigmentation problems such as freckles, chloasma, and lentigines can have an adverse impact on the physical appearance of skin, representing serious esthetic problems [28]. Tyrosinase activity is one of the major targets used to screen for new inhibitors of melanin biosynthesis. Tyrosinase inhibitors can be used to achieve healthy skin-whitening and have consequently been widely used in cosmetics, medicine, food, and agriculture, where there is a growing interest in the development of safe and effective tyrosinase inhibitors from natural plants [33]. Kojic acid is a metabolite of fungal that exhibits inhibitory activity towards tyrosinase. The inhibitory effects of kojic acid in this context have been attributed to its ability to chelate copper at the active site of this enzyme. For this reason, kojic acid is often used as a positive control to discover new tyrosinase inhibitors, as well as being used as a cosmetic skin-whitening agent and a food additive for preventing enzymatic browning. But, its uses are restricted because of its several side effects [34]. The effects of the five different varieties of squeezed oil on the diphenolase activity of mushroom tyrosinase were assayed using L-DOPA as a substrate with kojic acid as a reference compound. The results revealed that all five of the oils inhibited the oxidation of L-DOPA in a dose-dependent manner (Figure 2). However, the inhibitory effects of the oils were less than that of kojic acid. Among the five oils, TGSO, XYSO, and ZMSO showed the greatest activities at highest concentration (800 µg/mL). However, at the lowest concentration (50 µg/mL), DYSO and CGSO exhibited the best activity. The IC<sub>50</sub> values of the five different oils against tyrosinase were less than 400 µg/mL.

Ascorbic acid is commonly used as a tyrosinase inhibitor because of its antibrowning effect, and this compound has also been used as positive reference compound to evaluate new monophenolase inhibitors [35]. Ascorbic acid can also be used to affect the chemical reduction of dopaquinone, thereby avoiding the formation of dopachrome and melanin by the subsequent reduction of o-dopaquinone to L-DOPA. However, the application of ascorbic acid in this context has been limited by its poor stability [36]. The effects of the five different seed oils on monophenolase activity were studied using L-tyrosine as a substrate and ascorbic acid as a positive control. As shown in Figure 3, all five of the seed oils tested in the current study exhibited concentration-dependent inhibitory activity towards the tyrosinase-mediated oxidation of L-tyrosine. However, the inhibitory activities of all five oils were less than that of ascorbic acid, especially at a high concentration (800 µg/mL). The IC<sub>50</sub> values of the seed oils derived from the five different varieties of *Torreyia* were

greater than 800 µg/mL. Lastly, as is known in Figure 2, the five different seed oils showed slightly better diphenolase activity than their monophenolase activity.

The seed oils from the five different varieties of *Torreyia grandis* all exhibited potent inhibitory activity towards tyrosinase (especially TGSO and XYSO), most likely because of their strong antioxidant activity. It is possible that the free radical scavenging activities of these seed oils prevented the initiation of a radical chain, thereby inhibiting the activity of tyrosinase by reducing the supply of oxygen during the tyrosinase reaction. These could be antioxidant active components present in *Torreyia* seed oils, such as conjugated linoleic acid and Oleic acid, which could reduce the catalytic activity of tyrosinase by preventing the active effect of oxygen on tyrosinase.

**3.4. Inhibition Kinetics of the Five Squeezed Seed Oils.** The inhibition kinetics of the five different oils were analyzed using the Lineweaver–Burk method to determine their inhibitory activities towards diphenolase and monophenolase, and the corresponding plots are shown in Figures 2 and 3. The results revealed that the K<sub>m</sub> values of the oils varied considerably with increasing concentration, whereas their V<sub>max</sub> values remained largely unchanged. This result indicated that all five oils were behaving as competitive inhibitors of diphenolase and monophenolase. The inhibition constants (K<sub>i</sub>) of the different oils are shown in Table 3, together with their K<sub>m</sub> and V<sub>max</sub> values.

## 4. Conclusions

It is generally believed that there are four main methods for inhibiting tyrosinase activity: (1) affecting the chelating activity of copper at the active site of the enzyme, thereby preventing the binding of copper ions to oxygen, leading to the irreversible deactivation of tyrosinase [37, 38]; (2) the introduction of a powerful antioxidant, thereby preventing the activation of oxygen by tyrosinase [39, 40]; (3) the use of a free radical scavenging agent to prevent the formation of melanins [41]; and (4) the use of competitive inhibitors. Compounds with high antioxidant and radical scavenging activities could inhibit tyrosinase [42]. The seed oils derived from the five different varieties of *Torreyia grandis* in the current study all exhibited potent inhibitory activity towards tyrosinase (especially TGSO and XYSO), most likely because of their strong antioxidant activity. It is possible that the free radical scavenging activities of these oils prevented the initiation of a radical chain, thereby inhibiting the activity of tyrosinase by reducing the supply of oxygen during




FIGURE 2: Determination of the mode of tyrosinase inhibition for the seed oils derived from the five different species of *T. grandis*. Lineweaver-Burk plots for the tyrosinase activity (diphenolase activity) using L-DOPA as a substrate with seed oil concentrations of 50 ( $\square$ ), 100 ( $\bullet$ ), 200 ( $\blacksquare$ ), 400 ( $\blacktriangle$ ), and 800 ( $\blacklozenge$ )  $\mu\text{g/ml}$ . (The inset represents the slope ( $K_m/V_{max}$ ) versus seed oil concentration ( $\mu\text{g/ml}$ )). (a) TGSO, (b) XYSO, (c) ZMSO, (d) TGMSO, and (e) CGSO.




FIGURE 3: Determination of the mode of tyrosinase inhibition for the seed oils derived from the five different species of *T. grandis*. Lineweaver-Burk plots for the tyrosinase activity (monophenolase activity) using L-tyrosine as a substrate with seed oil concentrations of 50 ( $\square$ ), 100 ( $\bullet$ ), 200 ( $\blacksquare$ ), 400 ( $\blacktriangle$ ), and 800 ( $\blacklozenge$ )  $\mu\text{g/ml}$  (The inset represents the slope ( $K_m/V_{max}$ ) versus seed oil concentration ( $\mu\text{g/ml}$ ).) (a) TGSO, (b) XYSO, (c) ZMSO, (d) TGMSO, and (e) CGSO.

TABLE 3: The test results of tyrosinase inhibition activity and kinetic parameters.

Name	$V_{max}$ ( $\Delta OD_{475}/min$ )	$IC_{50}$ ( $\mu g/ml$ )	$K_m$ (mM)	$K_i$ ( $\mu g/ml$ )
Diphenolase				
TGSO	0.167	237.42±2.23	1.92	637.79±6.31
XYSO	0.147	227.01±2.68	1.61	587.6±3.87
ZMSO	0.1	277.98±1.38	0.89	297.39±5.61
TGMSO	0.063	244.52±4.15	0.87	317.57±3.48
CGSO	0.083	326.63±2.55	0.77	280.74±4.26
Kojic acid	-	32.65±0.42	-	-
Monophenolase				
TGSO	0.066	849.42±4.37	0.72	481.87±6.84
XYSO	0.05	817.53±3.79	0.44	473.7±4.32
ZMSO	0.04	879.65±5.13	0.37	396.7±5.11
TGMSO	0.036	936.44±6.38	0.14	563.17±3.27
CGSO	0.027	968.28±6.52	0.22	443.43±4.44
Ascorbic acid	-	35.97±0.13	-	-

Summary of the inhibition kinetics of the antioxidants derived from the five different varieties of *T. grandis*. Notes. Data represent the mean values  $\pm$  SD (n=3).

the tyrosinase reaction. There could be several antioxidants present in *Torreya* seed oils, which could reduce the catalytic activity of tyrosinase by preventing the active effect of oxygen on tyrosinase. However, only *in vitro* experiments showed that seed oil of the five *T. grandis* has the potential application as skin-whitening, the research has yet to be validated by animal experiments *in vivo*. Although the diphenolase  $IC_{50}$  are not particularly low in seed oil, it has no powerful inhibition of tyrosinase, but some of their constituents might be more effective and have strong whitening effect. Further in depth studies would be required involving the individual components to accurately determine which compounds are responsible for inhibiting this enzyme. These findings suggest that wild TGSOs (especially XYSO) exhibit significant antioxidant and inhibitory activities towards tyrosinase (especially diphenolase), and that they could therefore be used to develop skin-whitening and/or anti-food-browning agents. Notably, the yields of the wild TGSOs (XYSO and ZMSO were particularly outstanding) were similar to that of TGSO, highlighting the potential for wild seeds to be converted into edible oils.

## Abbreviations

TGSO:	<i>T. grandis</i> seed oil
XYSO:	<i>T. grandis</i> "Xiangyafei" seed oil
ZMSO:	<i>T. grandis</i> "Zhimafei" seed oil
TGMSO:	<i>T. grandis</i> "Majus" seed oil
CGSO:	<i>T. grandis</i> "cunguangfei" seed oil
DPPH:	1,1-Diphenyl-2-picrylhydrazyl
FRAP:	Ferric reducing power
TPTZ:	Tripyridyltriazine
Trolox:	6-Hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic
DMSO:	Dimethyl sulphoxide
TEAC:	Trolox equivalent antioxidant capacity
PBS:	Phosphate buffer
$IC_{50}$ :	50% inhibition concentration

$K_i$ : Enzyme-inhibitor dissociation constant  
 $K_m$ : Michaelis-Menten constant.

## Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

## Conflicts of Interest

The authors declare that they have no conflicts of interest in this research.

## Acknowledgments

This project is sponsored by Zhejiang Provincial Natural Science Foundation of China (LY13H280011).




## References

- [1] J. P. Ebanks, R. R. Wickett, and R. E. Boissy, "Mechanisms regulating skin pigmentation: the rise and fall of complexion coloration," *International Journal of Molecular Sciences*, vol. 10, no. 9, pp. 4066–4087, 2009.
- [2] C. A. Ramsden and P. A. Riley, "Tyrosinase: The four oxidation states of the active site and their relevance to enzymatic activation, oxidation and inactivation," *Bioorganic & Medicinal Chemistry*, vol. 22, no. 8, pp. 2388–2395, 2014.
- [3] Y. Yamaguchi, M. Brenner, and V. J. Hearing, "The regulation of skin pigmentation," *The Journal of Biological Chemistry*, vol. 282, no. 38, pp. 27557–27561, 2007.
- [4] N. Alam, K. N. Yoon, J. S. Lee, H. J. Cho, and T. S. Lee, "Consequence of the antioxidant activities and tyrosinase inhibitory effects of various extracts from the fruiting bodies of *Pleurotus ferulae*," *Saudi Journal of Biological Sciences*, vol. 19, no. 1, pp. 111–118, 2012.
- [5] L.-H. Peng, S. Liu, S.-Y. Xu et al., "Inhibitory effects of salidroside and paeonol on tyrosinase activity and melanin synthesis



- in mouse B16F10 melanoma cells and ultraviolet B-induced pigmentation in guinea pig skin," *Phytomedicine*, vol. 20, no. 12, pp. 1082–1087, 2013.
- [6] G.-H. Wang, C.-Y. Chen, C.-P. Lin et al., "Tyrosinase inhibitory and antioxidant activities of three Bifidobacterium bifidum-fermented herb extracts," *Industrial Crops and Products*, vol. 89, pp. 376–382, 2016.
  - [7] F. Saqib, K. H. Janbaz, and M. K. Sherwani, "In vitro inhibitory potential of methanolic extract of *Celosia argentea* var. *Cristata* on tyrosinase, acetylcholinesterase and butyrylcholinesterase enzymes," *Bangladesh Journal of Pharmacology*, vol. 10, no. 2, pp. 449–454, 2015.
  - [8] J. Solimine, E. Garo, J. Wedler et al., "Tyrosinase inhibitory constituents from a polyphenol enriched fraction of rose oil distillation wastewater," *Fitoterapia*, vol. 108, pp. 13–19, 2016.
  - [9] Y.-S. Lin, S.-H. Chen, W.-J. Huang et al., "Effects of nicotinic acid derivatives on tyrosinase inhibitory and antioxidant activities," *Food Chemistry*, vol. 132, no. 4, pp. 2074–2080, 2012.
  - [10] E. Neeley, G. Fritch, A. Fuller, J. Wolfe, J. Wright, and W. Flurkey, "Variations in IC<sub>50</sub> values with purity of mushroom tyrosinase," *International Journal of Molecular Sciences*, vol. 10, no. 9, pp. 3811–3823, 2009.
  - [11] H. Ando, M. S. Matsui, and M. Ichihashi, "Quasi-drugs developed in Japan for the prevention or treatment of hyperpigmentary disorders," *International Journal of Molecular Sciences*, vol. 11, no. 6, pp. 2566–2575, 2010.
  - [12] J. Ortonne, "Normal and abnormal skin color," *Annales de Dermatologie et de Vénérologie*, vol. 139, pp. S125–S129, 2012.
  - [13] M. K. Saeed, Y. Deng, R. Dai, W. Li, Y. Yu, and Z. Iqbal, "Appraisal of antinociceptive and anti-inflammatory potential of extract and fractions from the leaves of *Torreya grandis* Fort Ex. Lindl.," *Journal of Ethnopharmacology*, vol. 127, no. 2, pp. 414–418, 2010.
  - [14] Q. Ni, Q. Gao, W. Yu, X. Liu, G. Xu, and Y. Zhang, "Supercritical carbon dioxide extraction of oils from two *Torreya grandis* varieties seeds and their physicochemical and antioxidant properties," *LWT- Food Science and Technology*, vol. 60, no. 2, pp. 1226–1234, 2015.
  - [15] Y. Yang, Z. Jin, Q. Jin, and M. Dong, "Isolation and fatty acid analysis of lipid-producing endophytic fungi from wild Chinese *Torreya Grandis*," *Microbiology (Russian Federation)*, vol. 84, no. 5, pp. 710–716, 2015.
  - [16] D. Dong, H. Wang, F. Xu, C. Xu, X. Shao, and H. Li, "Supercritical carbon dioxide extraction, fatty acid composition, oxidative stability, and antioxidant effect of *Torreya grandis* seed oil," *Journal of the American Oil Chemists' Society*, vol. 91, no. 5, pp. 817–825, 2014.
  - [17] L. I. Zhang-Ju, C. F. Luo, X. J. Chen, X. J. Feng, and Y. U. Wei-Wu, "Component analysis and nutrition evaluation of seeds of *Torreya grandis* "Merrillii"," *Journal of Zhejiang Forestry College*, vol. 22, pp. 540–544, 2004.
  - [18] B.-Q. Chen, X.-Y. Cui, X. Zhao et al., "Antioxidative and acute antiinflammatory effects of *Torreya grandis*," *Fitoterapia*, vol. 77, no. 4, pp. 262–267, 2006.
  - [19] Z. He, H. Zhu, W. Li et al., "Chemical components of cold pressed kernel oils from different *Torreya grandis* cultivars," *Food Chemistry*, vol. 209, pp. 196–202, 2016.
  - [20] L. Ni and W.-Y. Shi, "Composition and free radical scavenging activity of kernel oil from *Torreya grandis*, *Carya cathayensis* and *Myrica rubra*," *Iranian Journal of Pharmaceutical Research*, vol. 13, no. 1, pp. 221–226, 2014.
  - [21] H. Cao, X. Chen, and K. Yamamoto, "Bovine serum albumin significantly improves the DPPH free radical scavenging potential of dietary polyphenols and gallic acids," *Anti-Cancer Agents in Medicinal Chemistry*, vol. 12, no. 8, pp. 940–948, 2012.
  - [22] C. Li, J. Feng, W.-Y. Huang, and X.-T. An, "Composition of polyphenols and antioxidant activity of rabbiteye blueberry (*Vaccinium ashei*) in Nanjing," *Journal of Agricultural and Food Chemistry*, vol. 61, no. 3, pp. 523–531, 2013.
  - [23] N. López-Carreras, S. Fernández-Vallinas, R. Hernández, M. Miguel, and A. Aleixandre, "Short-term effect of an aqueous *Fraxinus excelsior* L. seed extract in spontaneously hypertensive rats," *Food Research International*, vol. 53, no. 1, pp. 81–87, 2013.
  - [24] P. Suganya, K. Jeyaprakash, G. R. Mallavarapu, and R. Murugan, "Comparison of the chemical composition, tyrosinase inhibitory and anti-inflammatory activities of the essential oils of *Pogostemon plectranthoides* from India," *Industrial Crops and Products*, vol. 69, pp. 300–307, 2015.
  - [25] R. Matsuura, H. Ukeda, and M. Sawamura, "Tyrosinase inhibitory activity of citrus essential oils," *Journal of Agricultural and Food Chemistry*, vol. 54, no. 6, pp. 2309–2313, 2006.
  - [26] R. D. S. Teixeira, P. R. Rocha, H. C. Polonini, M. A. F. Brandão, M. D. G. A. M. Chaves, and N. R. B. Raposo, "Mushroom tyrosinase inhibitory activity and major fatty acid constituents of Amazonian native flora oils," *Brazilian Journal of Pharmaceutical Sciences*, vol. 48, no. 3, pp. 399–404, 2012.
  - [27] F. Yildiztekin, S. Nadeem, E. Erol, M. Yildiztekin, A. L. Tuna, and M. Ozturk, "Antioxidant, anticholinesterase and tyrosinase inhibition activities, and fatty acids of *Crocus mathewii* – A forgotten endemic angiosperm of Turkey," *Pharmaceutical Biology*, vol. 54, no. 9, pp. 1557–1563, 2016.
  - [28] E. Neagu, G. L. Radu, C. Albu, and G. Paun, "Antioxidant activity, acetylcholinesterase and tyrosinase inhibitory potential of *Pulmonaria officinalis* and *Centarium umbellatum* extracts," *Saudi Journal of Biological Sciences*, vol. 25, no. 3, pp. 578–585, 2018.
  - [29] H. Cao, Y. Xie, and X. Chen, "Type 2 diabetes diminishes the benefits of dietary antioxidants: Evidence from the different free radical scavenging potential," *Food Chemistry*, vol. 186, pp. 106–112, 2015.
  - [30] M. Z. Gul, L. M. Bhakshu, F. Ahmad, A. K. Kondapi, I. A. Qureshi, and I. A. Ghazi, "Evaluation of *Abelmoschus moschatus* extracts for antioxidant, free radical scavenging, antimicrobial and antiproliferative activities using *in vitro* assays," *BMC Complementary and Alternative Medicine*, vol. 11, article 64, 2011.
  - [31] P. Siddhuraju, P. S. Mohan, and K. Becker, "Studies on the antioxidant activity of Indian Laburnum (*Cassia fistula* L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp," *Food Chemistry*, vol. 79, no. 1, pp. 61–67, 2002.
  - [32] G.-C. Yen, P.-D. Duh, and C.-L. Tsai, "Relationship between Antioxidant Activity and Maturity of Peanut Hulls," *Journal of Agricultural and Food Chemistry*, vol. 41, no. 1, pp. 67–70, 1993.
  - [33] R. Biswas, P. K. Mukherjee, M. K. Dalai, P. K. Mandal, and M. Nag, "Tyrosinase inhibitory potential of purpurin in *Rubia cordifolia*-A bioactivity guided approach," *Industrial Crops and Products*, vol. 74, pp. 319–326, 2015.
  - [34] C. L. Burnett, W. F. Bergfeld, D. V. Belsito et al., "Final report of the safety assessment of kojic acid as used in cosmetics," *International Journal of Toxicology*, vol. 29, no. 6, 2010.
  - [35] Y.-H. Jo, G.-U. Seo, H.-G. Yuk, and S.-C. Lee, "Antioxidant and tyrosinase inhibitory activities of methanol extracts from

- Magnolia denudata and Magnolia denudata var. purpurascens flowers,” *Food Research International*, vol. 47, no. 2, pp. 197–200, 2012.
- [36] T.-S. Chang, “An updated review of tyrosinase inhibitors,” *International Journal of Molecular Sciences*, vol. 10, no. 6, pp. 2440–2475, 2009.
- [37] C. Sarikurkcu, G. Zengin, M. Oskay, S. Uysal, R. Ceylan, and A. Aktumsek, “Composition, antioxidant, antimicrobial and enzyme inhibition activities of two *Origanum vulgare* subspecies (subsp. *vulgare* and subsp. *hirtum*) essential oils,” *Industrial Crops and Products*, vol. 70, pp. 178–184, 2015.
- [38] P. M. Souza, S. T. Elias, L. A. Simeoni et al., “Plants from Brazilian Cerrado with Potent Tyrosinase Inhibitory Activity,” *PLoS ONE*, vol. 7, no. 11, 2012.
- [39] K. Boonsiripiphat and C. Theerakulkait, “Extraction of rice bran extract and some factors affecting its inhibition of polyphenol oxidase activity and browning in potato,” *Preparative Biochemistry and Biotechnology*, vol. 39, no. 2, pp. 147–158, 2009.
- [40] D. Fiocco, D. Fiorentino, L. Frabboni et al., “Lavender and peppermint essential oils as effective mushroom tyrosinase inhibitors: A basic study,” *Flavour and Fragrance Journal*, vol. 26, no. 6, pp. 441–446, 2011.
- [41] B. Fatiha, H. Didier, G. Naima et al., “Phenolic composition, in vitro antioxidant effects and tyrosinase inhibitory activity of three Algerian *Mentha* species: *M. spicata* (L.), *M. pulegium* (L.) and *M. rotundifolia* (L.) Huds (Lamiaceae),” *Industrial Crops and Products*, vol. 74, pp. 722–730, 2015.
- [42] G. Nagarani, A. Abirami, and P. Siddhuraju, “A comparative study on antioxidant potentials, inhibitory activities against key enzymes related to metabolic syndrome, and anti-inflammatory activity of leaf extract from different *Momordica* species,” *Food Science & Human Wellness*, vol. 3, pp. 36–46, 2014.



**Hindawi**  
Submit your manuscripts at  
[www.hindawi.com](http://www.hindawi.com)

