Review Article

Clinically Correlated MicroRNAs in the Diagnosis of Non-Small Cell Lung Cancer: A Systematic Review and Meta-Analysis

Min Jiang ^{1,2}, Xuelian Li,² Xiaowei Quan ², Xiaoying Li ² and Baosen Zhou ^{1,2}

¹Department of Clinical Epidemiology and Center of Evidence-Based Medicine, The First Affiliated Hospital, China Medical University, Shenyang, Liaoning 110001, China

²Department of Epidemiology, School of Public Health, China Medical University, Shenyang 110122, China

Correspondence should be addressed to Baosen Zhou; bszhou@cmu.edu.cn

Received 5 February 2018; Revised 30 April 2018; Accepted 7 June 2018; Published 28 June 2018

Academic Editor: Maria L. Tornesello

Copyright © 2018 Min Jiang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

(1) *Background.* Non-small cell lung cancer (NSCLC) has a high mortality rate. MiRNAs have been found to be diagnostic biomarkers for NSCLC. However, controversial results exist. We conducted this meta-analysis to evaluate the diagnostic value of miRNAs for NSCLC. (2) *Methods.* Databases and reference lists were searched. Pooled sensitivity (SEN), specificity (SPE), and area under the curve (AUC) were applied to examine the general diagnostic efficacy, and subgroup analysis was also performed. (3) *Results.* Pooled SEN, SPE, and AUC were 85%, 88%, and 0.93, respectively, for 71 studies. Multiple miRNAs (AUC: 0.96) obtained higher diagnostic value than single miRNA (AUC: 0.86), and the same result was found for Caucasian population (AUC: 0.97) when compared with Asian (AUC: 0.91) and Caucasian/African population (AUC: 0.92). MiRNA had higher diagnostic efficacy when participants contained both smokers and nonsmokers (AUC is 0.95 for imbalanced group and 0.91 for balanced group) than when containing only smokers (AUC: 0.90). Meanwhile, AUC was 0.91 for both miR-21 and miR-210. (4) *Conclusions.* Multiple miRNAs such as miR-21 and miR-210 could be used as diagnostic tools for NSCLC, especially for the Caucasian and nonsmoking NSCLC.

1. Introduction

Lung cancer is the principal cause of cancer-associated deaths among males both in developed and in developing countries, and it has exceeded the breast cancer becoming the major cause of cancer-related deaths in females in the developed countries [1]. Non-small cell lung cancer (NSCLC) is a major type of lung cancer that is responsible for 85% lung cancer-associated deaths. Smoking has been recognized as a primary environmental risk factor of lung cancer. However, only a small number of smokers will develop into lung cancer patients.

MicroRNA is a group of 19–22 nucleotide, small, single-stranded, and conserved noncoding RNA that acts as a regulator of gene expression at both the posttranscriptional and the translational levels through acting on the 3'-untranslated region (UTR) of messenger RNA (mRNA) [2]. MiRNAs play important roles in various biological processes associated with the tumorigenesis such as the cellular proliferation,

differentiation, metabolism, and apoptosis [3, 4]. It is available to isolate the miRNAs from the clinical specimens including the plasma, serum, sputum, and tissue. Meanwhile, it has a high stability. Due to these advantages, the miRNAs are increasingly becoming an ideal tool for the detection of NSCLC.

Recently, a series of articles have shown that different miRNAs might be applied to detect the NSCLC [5–7]. For example, miR-21, an oncogenic miRNA, has been shown to be overexpressed in lung cancer as well as other various human tumors [8]. Upregulation of miR-21 could promote the tumorigenesis of lung cancer through inhibiting the apoptosis process and negatively regulating the Ras/MEK/ERK signal pathway [9]. High miR-210 expression was correlated with the increased lymph node metastasis and a poor prognosis in patients with NSCLC [10]. Both these two, miR-21 and miR-210, have been explored to be used as diagnostic tools for NSCLC, no matter whether they are applied in combination with other miRNAs or alone [11–14]. However,

as a result of the small sample sizes, the different miRNAs profiling, and the differences of the specimen and ethnicity, inconsistencies existed among studies that had examined the diagnostic value of miR-21, miR-210, and other miRNAs for NSCLC. Therefore, a meta-analysis was performed to assess the performance of miRNAs in the detection for NSCLC.

2. Materials and Methods

2.1. Search Strategy. Our meta-analysis was based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA). We searched PubMed, Google Scholar, Chinese National Knowledge Infrastructure (CNKI), Embase, and Medline to find all associated articles in order to investigate the potential utility of miRNAs as diagnostic tools for NSCLC. The combination of the Medical Subject Headings (MeSH) and the keywords ("lung neoplasm" OR "lung malignancy" OR "lung cancer") AND ("miRNA" OR "microRNAs") AND ("ROC curve" OR "sensitivity" OR "specificity" OR "diagnosis") was used (updated to April 5, 2017). The reference lists of the reviews were also searched to obtain all the acceptable articles.

2.2. Study Selection. A series of criteria were applied for study inclusion and exclusion. For inclusion, the criteria were as follows: (1) patients with NSCLC; (2) the type of the controls being healthy controls (HC) or patients with benign pulmonary diseases (BPD); (3) assessing the diagnostic value of the miRNAs; (4) the possibility of extracting or calculating TP, FP, FN, and TN from the articles. For exclusion, the criteria were as follows: (1) studies that were duplicate publications, reviews, or unrelated; (2) studies without complete data.

2.3. Data Collection and Quality Assessment. Two authors collected the data independently as follows: the first author, publication year, and participant demographic characteristics (ethnicity, sample size, mean or median age, smoking status, the types of the controls, and the testing method of controls and cancer); types of the specimen; miRNA profiling and the data used for this meta-analysis (SEN, SPE, TP, FP, FN, and TN). The quality of these articles was assessed with the QUADAS-2 guidelines [15].

2.4. Statistical Analysis. All the statistical analyses were conducted by RevMan 5.3 (version 1.4) software and STATA 11.0 (STATA-Corp, College Station, TX, version 11.0) software. The heterogeneity among the selected studies was assessed through the Q test and the I^2 value [16]. The P value for the Q test being less than 0.05 or the $I^2 \geq 50\%$ demonstrated that there was heterogeneity among the included studies. The pooled SPE [TN/(FP+TN)], SEN [TP/(FN+TP)], diagnostic odds ratio (DOR) [PLR/NLR], the negative likelihood ratio (NLR) [(1-SPE)/SPE], the positive likelihood ratio (PLR) [(SEN/(1-SEN))], and their 95% confidence intervals (95% CIs) were evaluated by a bivariate random-effect-regression model. The SROC curve was constructed and the AUC value was calculated too. A Fagan nomogram was also constructed to evaluate the clinical utility of miRNAs in the

diagnosis of NSCLC. Subgroup analyses (grouped by miRNA profiling: single and multiple; smoking status: only smokers, smokers, and nonsmokers (imbalanced between groups), smokers and nonsmokers (balanced between groups), and unknown smoking status; specimen: serum, plasma, whole blood/blood cell, and not blood; ethnicity: Asian, Caucasian, and Caucasian/African; control-type: BPD, HC, and BPD/HC; stage: early stage and no early stage; and case number: large (≥ 50) and small (< 50)) and meta-regression analysis were used to identify the potential sources of the heterogeneity. The Deeks' funnel plot asymmetry test was also applied to explore the publication bias, with the P value less than 0.01 considered significant [17].

3. Results

3.1. Literature Search and the Studies' Characteristics. As shown in Figure 1(a), 2594 eligible articles were included, of which 2145 articles were removed as unrelated and duplicate articles. And then 370 reviews were also excluded, leaving 79 articles with full texts, and another 21 articles were then removed through carefully reading: 14 articles met the exclusion criteria and 7 articles did not have the complete data. Ultimately, 58 articles [5–7, 11–14, 18–68] with 71 studies published from 2009 to 2017 including 9,099 participants (5111 cases with NSCLC and 3988 controls from the healthy individuals and the patients with the benign pulmonary disease (BPD)) were included. The main characteristics of these 71 studies were shown in Table 1. Wang Y's article [7], Fan LH's article [52], Nadal E's article [45], Tang DF's article [32], Razzak R's article [14], Wang W's article [68], Yu L's article [19], and Xing LX's article [18] included 2 studies. Bediaga's article [28] included 3 studies, Wang C's article [46] included 4 studies, and the remaining articles [5, 6, 11–13, 20–27, 29–31, 33–44, 47–51, 53–67] included 1 study, respectively. Meanwhile, there were 18 studies [13, 14, 28, 31, 33, 45, 46, 48, 54, 56, 63, 65, 66] performed in Caucasian, 11 studies [18, 19, 21, 23, 27, 29, 30, 38, 44] performed in Caucasian/African, and 1 study [26] performed in African populations; the remaining studies were performed in Asian populations. A total of 50 studies detected the miRNAs in blood such as the whole blood, plasma, serum, and peripheral blood mononuclear cells (PBMC) [6, 7, 11, 20–24, 26, 29–35, 37, 39–42, 44–47, 49–54, 56–64, 66–68], while the remaining studies were detected in nonblood samples (7 tissue [5, 25, 28, 55, 68], 1 pleural effusion [43], 12 sputum [12, 14, 18, 19, 27, 36, 38, 48, 65], and 1 BAL [13]). We evaluated 45 studies for assessing the diagnostic value of multiple miRNAs and 26 studies [5, 6, 11, 20, 22, 24–26, 33–37, 39–41, 43, 47, 50, 51, 55, 57, 58, 60, 67, 68] of single miRNA.

The quantitative real-time polymerase chain reaction (qRT-PCR) and digital polymerase chain reaction (digital PCR) were used in these studies to test the expression levels of different miRNAs, and the most common reference miRNAs were RNU6B, miR-39, and miR-16. Quality of the enrolled studies summarized in Figure 1(b) was generally good.

3.2. Pooled Diagnostic Performance. Significant heterogeneity was obtained since I^2 values for SEN and SPE were

TABLE 1: The main features of 71 included studies in this meta-analysis.

Study ID	Ethnicity	Specimen	Case N	Age	Control N	Age	Type of control	Stage	MIRNA profiling	SEN (%)	SPE (%)	Reference miRNA	microRNA assay	Smoking status*
Zhang H 2017	Asian	plasma	129	59.6	83	60.0	HC	I-II	miR-145, miR-20a, miR-21, miR-223 miR-429, miR-205, miR-200b, miR-203, miR-12, miR-34b	81.8	90.1	miR-16	qRT-PCR	3
Halvorsen A 2016	Caucasian	serum	100	62.6	58	57.6	HC	I-IV	20 miRNAs ^a miR-195 miR-182, miR-183, miR-210, miR-126	88.0	71.0	miR-220, miR-19b, U6	qRT-PCR	3
TaiMei C2016	Asian	blood	110	65.0	52	65.7	HC	I-III	miR-195	89.1	100	miR-159a, U6	qRT-PCR	4
Su KL 2016	Asian	plasma	100	NA	100	NA	HC	I-III		78.0	86.0	miR-39	qRT-PCR	3
Zhu WY 2016	Asian	plasma	112	58.5	40	57.9	HC	I-III	miR-182, miR-183, miR-210, miR-126	81.3	100	U6	qRT-PCR	2
Jiang LP 2016	Asian	tissue	154	54.9	63	57.8	BPD	I-IV	miR-26b	79.9	79.4	U6	qRT-PCR	3
Wang Y 2016	Asian	plasma	82	NA	91	NA	HC	I-II	miR-532, miR-628, miR-425	91.5	97.8	miR-39	qRT-PCR	4
Wang Y 2016	Asian	plasma	36	NA	43	NA	HC	I-II	miR-532, miR-628, miR-425	97.2	95.3	miR-39	qRT-PCR	4
Fan LH 2016	Asian	serum	94	60.5	58	58.1	HC	I-III	miR-15b, miR-16, miR-20a	86.2	91.4	NA	qRT-PCR	4
Fan LH 2016	Asian	serum	70	59.7	54	58.0	HC	I-III	miR-15b, miR-16, miR-20a	94.3	94.2	NA	FQDs	4
Sun L 2016	Asian	plasma	87	60.7	96	53.8	HC,BPD	I-IV	miR-30a	61.0	84.3	U6	qRT-PCR	4
Su Y 2016	Asian	sputum	144	66.3	171	65.2	BPD	I	miR-21, miR-31, miR-210	81.5	85.9	U6	qRT-PCR	1
Gao X 2016	Asian	plasma	30	61.1	30	60.2	HC	I	miR-324, miR-1285	93.3	90.0	miR-39	qRT-PCR	4
Wang X 2016	Asian	plasma	59	55.9	59	57.6	BPD	I-III	miR-486	83.1	78.0	miR-16	qRT-PCR	4
Wei J 2016	Asian	plasma	63	61.0	30	57.0	HC	I-IV	miR-21	76.2	70.0	miR-16	qRT-PCR	3
Razzak 2016	Caucasian	sputum	22	68	10	58	HC,BPD	III-IV	miR-21, miR-210, miR-372	64	100	U6	qRT-PCR	4

TABLE 1: Continued.

Study ID	Ethnicity	Specimen	Case N	Age	Control N	Type of control	Stage	MIRNA profiling	SEN (%)	SPE (%)	Reference miRNA	microRNA assay	Smoking status*
Razzak 2016	Caucasian	sputum	21	70	10	HC,BPD	I-II	miR-21, miR-210, miR-372 miR-720,	67	90	U6	qRT-PCR	4
Leidinger P 2016	Caucasian	blood	74	NA	20	HC	I-III	miR-29c, miR-199a, miR-378a,let-7f miR-182, miR-10a, miR-301b, miR-1244, miR-301a, miR-135b, miR-224,	91.0	98.0	U24,U48	qRT-PCR	4
Wang WZ 2016	Asian	tissue	15	57	16	HC	I-IV	miR-21 miR-1244 miR-21, miR-143, miR-155, miR-210, miR-373 miR-483,	93.3	93.8	miR-16	qRT-PCR	4
Wang WZ 2016	Asian	serum	54	NA	15	HC	I-IV	miR-193a, miR-25, miR-214, miR-7	81.5	80	miR-39	qRT-PCR	4
Kim JL O 2015	Caucasian	BAL	21	70	10	HC,BPD	I-II	miR-486 miR-483, miR-193a, miR-25, miR-483,	85.7	100	U6	qRT-PCR	4
Wang C 2015	Asian	serum	19	61.8	19	HC	I-IV	miR-193a, miR-25, miR-214, miR-7	100	84	let-7d/g/i	qRT-PCR	4
Li WS 2015	Asian	plasma	11	59	11	HC	I-III	miR-486 miR-483, miR-193a, miR-25, miR-214, miR-7	90.9	81.8	miR-39, U44	qRT-PCR	4
Wang C 2015	Asian	serum	63	61.9	63	HC	I-IV	miR-483, miR-193a, miR-25, miR-214, miR-7	89.0	68.0	let-7d/g/i	qRT-PCR	4
Wang C 2015	Caucasian	serum	108	67.2	56	BPD	I-IV	miR-483, miR-193a, miR-25, miR-214, miR-7	95.0	95.0	let-7d/g/i	qRT-PCR	4
Wang C 2015	Caucasian	serum	108	67.2	48	HC	I-IV	miR-483, miR-193a, miR-25, miR-214, miR-7	95.0	84.0	let-7d/g/i	qRT-PCR	4
Nadal E 2015	Caucasian	serum	70	67.5	22	HC,BPD	I-III	miR-141, miR-200b, miR-193b miR-141, miR-200b, miR-193b	96.0	95.0	U6	qRT-PCR	2
Nadal E 2015	Caucasian	serum	84	65.5	23	HC,BPD	I-III	miR-141, miR-200b, miR-193b	97.0	96.0	U6	qRT-PCR	2
Guo WG 2015	Asian	plasma	126	NA	50	HC	I-IV	mir-204	76.0	82.0	U6	qRT-PCR	4

TABLE 1: Continued.

Study ID	Ethnicity	Specimen	Case N	Age	Control N	Age	Type of control	Stage	MIRNA profiling	SEN (%)	SPE (%)	Reference miRNA	microRNA assay	Smoking status*
Ma J 2015	Caucasian, African	PBMC	84	64.1	69	62.4	BPD	I-IV	miR-19b, miR-29b, miR-148a, miR-148b, miR-152	72.6	82.6	miR-423-3p	qRT-PCR	2
Li L 2015	Asian	serum	36	56.0	30	58.0	HC,BPD	I-IV	miR-148b, miR-152	72.2	90.0	U6	qRT-PCR	4
Zhang XL 2015	Asian	tissue	125	61.0	125	61.0	HC	I-IV	miR-141	64.8	64.8	miR-191, miR-103	qRT-PCR	3
Zhao W 2015	Asian	serum	80	57.6	60	55.4	HC	NA	miR-21	73.8	71.7	U6	qRT-PCR	4
Wang RJ 2015	Asian	serum	70	64.4	70	63.7	HC	NA	miR-145	92.8	61.4	miR-39	qRT-PCR	3
Yang JS 2015	Asian	serum	152	NA	300	NA	HC	I-IV	miR-152, miR-148a, miR-148b, miR-21	96.0	91.0	U6	qRT-PCR	3
Xing LX 2015	Caucasian	sputum	67	66.4	69	64.9	BPD	I-II	miR-21, miR-31, miR-210	82.1	88.4	U6, miR-16	qRT-PCR	4
Liu CM 2015	Asian	Pleural effusion	61	53.8	70	54.4	BPD	NA	miR-192	61.3	79.5	U6	qRT-PCR	2
Dou HL 2015	Asian	plasma	120	63.2	360	NA	HC	I-IV	miR-152	86.0	81.3	U6	digital PCR	4
Yang YL 2015	Asian	PBMC	74	62.5	52	61.8	HC	I-IV	miR-10b	86.5	76.9	miR-16	qRT-PCR	3
Li N 2014	Caucasian, African	sputum	35	68.9	40	65.7	HC	I	miR-31, miR-210	65.7	85.0	NA	qRT-PCR	4
Zhu W 2014	Asian	serum	70	59.0	48	NA	HC	I-IV	miR-429	54.3	81.2	U6,U48	qRT-PCR	4
LI M 2014	Asian	serum	514	NA	54	NA	HC	I-IV	miR-499	73.7	92.7	miR-39	qRT-PCR	3
Ulivi P 2013	Caucasian	blood	86	68.0	24	65.0	HC	I-II	miR-328	70.0	83.0	U38B,U58A	qRT-PCR	4
Bediaga 2013	Caucasian	tissue	45	66.4	45	66.4	HC	I-IV	8 miRNAs ^b	100	97.8	4miRNAs ^c	qRT-PCR	3
Bediaga 2013	Caucasian	tissue	47	67.8	47	67.8	HC	I-IV	8 miRNAs ^b	97.5	96.3	4miRNAs ^c	qRT-PCR	3
Bediaga 2013	Caucasian	tissue	22	68.4	22	68.4	HC	I-IV	8 miRNAs ^b	100	95.0	4miRNAs ^c	qRT-PCR	3
Anjuman 2013	Caucasian, African	sputum	39	65.6	42	62.3	BPD	I	miR-210, miR-31	61.5	90.5	U6	qRT-PCR	4
Tang DF 2013	Asian	plasma	62	64.8	60	66.0	HC	I-III	miR-21, miR-145, miR-155	69.4	78.3	U6	qRT-PCR	1

TABLE 1: Continued.

Study ID	Ethnicity	Specimen	Case N	Age	Control N	Type of control	Stage	MIRNA profiling	SEN (%)	SPE (%)	Reference miRNA	microRNA assay	Smoking status*
Tang DF 2013	Asian	plasma	34	65.2	32	HC	I-III	miR-21, miR-145, miR-155	76.5	81.3	U6	qRT-PCR	1
Mozzoni 2013	Caucasian	plasma	54	69.1	46	BPD	I-III	miR-21, miR-486	87.0	86.5	miR-16	qRT-PCR	4
ZENG XL 2013	Asian	PBMC	64	58.9	26	HC	I-IV	miR-143	75.0	92.3	U6	qRT-PCR	4
Yang XQ 2013	Asian	sputum	24	60.5	24	BPD	I-IV	let-7a	87.5	83.3	U6	digital PCR	4
Ma J 2013	Caucasian, African	plasma	36	66.7	38	HC	I	miR-21, miR-335	71.8	80.6	NA	qRT-PCR	4
Cazzoli R 2013	Caucasian, African	plasma	50	66.1	30	BPD	I	miR-151a, miR-30a, miR-200b, miR-629, miR-100, miR-154 miR-182	96.0	60.0	let7a	qRT-PCR	2
Abd-E 2013	African	Serum	65	54.1	37	HC	I-II	miR-152, miR-145, miR-199a, miR-24, miR-20a, miR-25 miR-21, miR-143, miR-155, miR-210, miR-372	100	86.5	SNORD68	qRT-PCR	4
Sanfiorenzo C 2013	Caucasian	plasma	52	65.1	10	BPD	I-III	miR-199a, miR-24, miR-20a, miR-25 miR-21, miR-143, miR-155, miR-210, miR-372	90.9	83.3	miR-192, miR-16	qRT-PCR	4
Roa Wilson H 2012	Caucasian	sputum	24	68.8	6	HC	I-II	miR-143, miR-155, miR-210, miR-372	83.3	100	U6	qRT-PCR	4
Li GJ 2012	Asian	plasma	16	NA	14	BPD	I	miR-22, miR-200b	85.3	94.5	18S	qRT-PCR	4
Ma YX 2012	Asian	serum	193	NA	110	HC	I-IV	miR-125b	78.2	66.4	NA	qRT-PCR	4
Hennessey P 2012	Caucasian, African	serum	55	68.2	75	HC	I-IV	miR-15b, miR-27b	100	84.0	miR-16	qRT-PCR	4
ZengXL 2012	Asian	PBMC	34	NA	26	HC	I-IV	miR-150	87.5	69.2	U6	qRT-PCR	4
Zhao M 2012	Asian	tissue	55	NA	55	HC	I-IV	miR-29a	49.1	85.5	U6	qRT-PCR	3
Shen J 2011	Caucasian, African	plasma	34	68.0	29	HC	I-IV	miR-21, miR-126, miR-210, miR-486	91.7	96.6	miR-16	qRT-PCR	1
Jeong H 2011	Asian	blood	35	67.0	30	HC	I-IV	let-7a	90.3	90.3	U6	qRT-PCR	4
Wei J 2011	Asian	plasma	77	59.6	36	HC	I-IV	miR-21	61.0	83.3	miR-16	qRT-PCR	3
Liu S 2011	Asian	plasma	130	53.1	170	HC	I-III	miR-126	46.4	90	NA	qRT-PCR	3

TABLE 1: Continued.

Study ID	Ethnicity	Specimen	Case N	Age	Control N	Age	Type of control	Stage	MIRNA profiling	SEN (%)	SPE (%)	Reference miRNA	microRNA assay	Smoking status*
Yu L 2010	Caucasian, African	sputum	36	68.2	36	66.7	HC	I	miR-486,	80.6	91.7	U6	qRT-PCR	4
									miR-21,					
									miR-200b,					
									miR-375					
Yu L 2010	Caucasian, African	sputum	64	67.0	58	65.0	HC	I-IV	miR-486,	70.3	80.0	U6	qRT-PCR	3
									miR-21,					
									miR-200b,					
									miR-375					
Xing LX 2010	Caucasian, African	sputum	48	67.5	48	65.9	HC	I	miR-205,	73.0	96.0	U6	qRT-PCR	4
									miR-210,					
									miR-708					
									miR-205,					
Xing LX 2010	Caucasian, African	sputum	67	68.0	55	65.0	HC	I-IV	miR-210,	72.0	95.0	U6	qRT-PCR	3
									miR-708					
Keller Andreas 2009	Caucasian	blood	17	64.2	19	37.9	HC	I-III	24miRNAs ^d	92.5	98.1	NA	qRT-PCR	4

^amiR-451, miR-1290, miR-636, miR-30c, miR-22-3p, miR-19b, miR-486-5p, miR-20b, miR-93, miR-34b, miR-185, miR-126-5p, miR-93-3p, miR-1274a, miR-142-5p, miR-628-5p, miR-486-3p, miR-425, miR-645, miR-24; ^bmiR-96, miR-450a, miR-183, miR-9, miR-577, Let-7i, miR-27b and miR-34a; ^cmiR-26a, miR-140-5p, miR-195, miR-30b; ^dmiR-126, miR-423, miR-15a, let-7d, let-7i, miR-22, miR-98, miR-19a, miR-20b, miR-324, miR-574, miR-195, miR-25, let-7e, let-7f, let-7g, miR-140, miR-339, miR-361, miR-1283, miR-18a, miR-26b; * 1: only smokers; 2: smokers and nonsmokers (smoking status was imbalanced between groups); 3: smokers and nonsmokers (smoking status was balanced between groups); 4: unknown smoking status.

N: number; HC: healthy control; BPD: benign pulmonary disease; miR: microRNA; SEN: sensitivity; SPE: specificity; FQDs: fluorescence quantum dots; BAL: bronchoalveolar lavage.

FIGURE 1: Flow chart of this meta-analysis of miRNAs in NSCLC detection (a) and the quality of these included articles according to the QUADAS-2 guidelines: proportion of articles with risk of bias (left) and proportion of articles with concerns regarding applicability (right) (b).

FIGURE 2: Forest plots of SEN and SPE for the NSCLC diagnosis. Both the SEN and SPE of each study were shown by squares with the 95% confidence interval shown by the error bars.

89.05% (95% CI: 87.07-91.03%) and 79.59% (95% CI: 75.18-84.01%), respectively. Therefore, a random-effect model was conducted for this study. Results indicated the pooled SEN and SPE for these 71 studies were 85% (95% CI: 82-88%) and 88% (95% CI: 85-90%), respectively (Figure 2). The PLR and NLR were 6.9 (95% CI: 5.6-8.4) and 0.17 (95% CI:

0.14-0.21), respectively (Figure 3), the DOR was 40 (95% CI: 28-58), and the AUC was 0.93 (95% CI: 0.90-0.95) (Figure 4(a)).

3.3. Publication Bias. Results of the Deeks' funnel plot asymmetry test showed that the publication bias did not exist in

FIGURE 3: Fagan plot of PLR and NLR to evaluate the clinical utility of miRNAs for diagnosis of NSCLC.

FIGURE 4: SROC curve of the miRNAs as diagnostic tools for NSCLC (a) and the Deeks' test for assessing the publication bias for miRNAs in the detection of NSCLC (b).

these studies as the funnel plot was symmetry (Figure 4(b)) and P value equaled 0.12.

3.4. Subgroup Analyses and Meta-Regression Analysis. Results of the meta-regression analysis demonstrated that the heterogeneity might be explained by miRNA profiling

($P < 0.001$) and case number ($P < 0.05$) for SPE and by miRNA profiling ($P < 0.01$) for SEN as described in Figure 5. The subgroup analyses were also conducted and the results were presented in Table 2. For the subgroups of smoking status, compared with the subgroup of only smokers (SEN: 80% (95% CI: 70-87%), SPE: 86% (95% CI: 77-91%), and AUC:

TABLE 2: Subgroup analyses for the selected studies.

Subgroups	No	SEN [95%CI]	SPE [95%CI]	PLR[95%CI]	NLR [95%CI]	DOR[95%CI]	AUC [95%CI]
MiR profiling							
single	26	0.77[0.71-0.82]	0.80[0.77-0.84]	3.9[3.3-4.7]	0.28[0.22-0.36]	14[10-20]	0.86[0.82-0.88]
multiple	45	0.88[0.85-0.91]	0.91[0.88-0.93]	10.0[7.5-13.3]	0.13[0.10-0.17]	79[50-126]	0.96[0.93-0.97]
Smoking status							
only smokers	4	0.80[0.70-0.87]	0.86[0.77-0.91]	5.6[3.2-9.9]	0.23[0.14-0.38]	24[9-66]	0.90[0.87-0.92]
S+NS (imbalanced)*	6	0.88[0.74-0.95]	0.90[0.73-0.97]	9.2[3.0-28.2]	0.13[0.05-0.31]	71[14-360]	0.95[0.93-0.97]
S+NS (balanced) *	18	0.83[0.74-0.90]	0.86[0.80-0.90]	5.9[3.9-8.8]	0.19[0.12-0.32]	30[13-69]	0.91[0.88-0.93]
unknown status	43	0.86[0.82-0.89]	0.88[0.85-0.91]	7.3[5.7-9.4]	0.16[0.12-0.21]	46[30-70]	0.93[0.91-0.95]
Specimen							
plasma	22	0.82[0.76-0.87]	0.87[0.83-0.90]	6.3[4.6-8.5]	0.20[0.15-0.28]	31[18-52]	0.92[0.89-0.94]
serum	19	0.91[0.86-0.95]	0.85[0.79-0.89]	6.1[4.3-8.5]	0.10[0.06-0.17]	60[28-128]	0.94[0.91-0.95]
Whole blood/blood cell	9	0.84[0.78-0.89]	0.92[0.80-0.97]	10.9[3.9-30.3]	0.17[0.11-0.26]	64[17-234]	0.92[0.89-0.94]
not blood	21	0.80[0.72-0.86]	0.89[0.85-0.93]	7.5[4.9-11.7]	0.22[0.16-0.32]	34[16-71]	0.92[0.89-0.94]
Ethnicity							
Asian	41	0.82[0.77-0.85]	0.86[0.82-0.88]	5.7[4.5-7.2]	0.21[0.17-0.27]	27[18-40]	0.91[0.88-0.93]
Caucasian	18	0.91[0.86-0.95]	0.92[0.87-0.96]	12[7.0-20.4]	0.09[0.06-0.15]	127[54-302]	0.97[0.95-0.98]
Caucasian/African	12	0.85[0.72-0.93]	0.87[0.81-0.91]	6.6[4.6-9.4]	0.17[0.09-0.33]	39[17-88]	0.92[0.89-0.94]
Control-type							
BPD	13	0.84[0.77-0.89]	0.84[0.80-0.88]	5.3[4.1-6.8]	0.19[0.13-0.28]	27[16-46]	0.90[0.87-0.92]
HC	50	0.86[0.82-0.89]	0.88[0.85-0.91]	7.4[5.7-9.5]	0.16[0.12-0.21]	47[30-74]	0.94[0.91-0.95]
BPD, HC	8	0.81[0.67-0.90]	0.91[0.79-0.96]	8.8[3.4-22.9]	0.21[0.11-0.40]	42[9-187]	0.93[0.90-0.95]
Stage							
I-II	18	0.84[0.78-0.89]	0.90[0.86-0.93]	8.3[5.8-11.9]	0.17[0.12-0.25]	48[27-87]	0.94[0.91-0.96]
I-IV	50	0.86[0.82-0.89]	0.88[0.84-0.90]	6.5[5.4-8.7]	0.16[0.13-0.22]	42[27-66]	0.93[0.90-0.95]
No. of cases							
small	25	0.88[0.82-0.92]	0.91[0.88-0.94]	10.0[7.1-14.2]	0.14[0.09-0.21]	74[38-143]	0.95[0.93-0.97]
large	46	0.84[0.79-0.87]	0.86[0.82-0.88]	5.8[4.6-7.2]	0.19[0.15-0.24]	31[20-46]	0.91[0.89-0.94]
MiR-210	12	0.77[0.72-0.81]	0.93[0.88-0.96]	11.0[6.2-19.4]	0.25[0.20-0.31]	44[22-87]	0.91[0.88-0.93]
MiR-21	16	0.82[0.77-0.86]	0.87[0.84-0.89]	6.3[5.0-8.1]	0.21[0.15-0.28]	31[19-50]	0.91[0.88-0.93]

No: the number of the studies; HC: healthy control; BPD: benign pulmonary disease; SEN: sensitivity; SPE: specificity; PLR: positive likelihood ratio; NLR:negative likelihood ratio; DOR: diagnostic odds ratio; AUC:area under the curve; no. of case: small (<50) and large (≥50).

* S: smokers; NS: nonsmokers; imbalanced: the smoking status was imbalanced between groups; balanced: the smoking status was balanced between groups.

FIGURE 5: Forest plots for the meta-regression analysis: SEN and SPE. The factors included miRNA profiling, smoking status, specimen, ethnicity, type of control, case number, and stage.

0.90 (95% CI: 0.87-0.92)), miRNAs had a higher diagnostic efficacy in the subgroups of smokers and nonsmokers (SEN: 88% (95% CI: 74-95%), SPE: 90% (95% CI: 73-97%), and AUC: 0.95 (95% CI: 0.93-0.97) for imbalanced groups and SEN: 83% (95% CI: 74-90%), SPE: 86% (95% CI: 80-90%), and AUC: 0.91 (95% CI: 0.88-0.93) for balanced groups). Subgroup analysis by specimen showed that studies with serum samples exhibited higher diagnostic accuracy with SEN: 91% (95% CI: 86-95%), SPE: 85% (95% CI: 79-89%), and AUC: 0.94 (95% CI: 0.91-0.95) than studies with plasma samples with the SEN: 82% (95% CI: 76-87%), SPE: 87% (95% CI: 83-90%), and AUC: 0.92 (95% CI: 0.89-0.94) and not bleeding samples with the SEN: 80% (95% CI: 72-86%), SPE: 89% (95% CI: 85-93%), and AUC: 0.92 (95% CI: 0.89-0.94), respectively. When compared with the large sample size, miRNA might be a better diagnostic tool for small sample size with SEN: 88% (95% CI: 82-92%), SPE: 91% (95% CI: 88-94%), and AUC: 0.95 (95% CI: 0.93-0.97). In the subgroups for the ethnicity, the miRNAs obtained a better diagnostic value in the Caucasian populations with the SEN: 91% (95% CI: 86-95%), SPE: 92% (95% CI: 87-96%), and AUC: 0.97 (95% CI: 0.95-0.98), respectively, when compared with the Asian populations with the SEN: 82% (95% CI: 77-85%), SPE: 86% (95% CI: 82-88%), and AUC: 0.91 (95% CI: 0.88-0.93), respectively, and the Caucasian/African populations with SEN: 85% (95% CI: 72-93%), SPE: 87% (95% CI: 81-91%), and AUC: 0.92 (95% CI: 0.89-0.94), respectively. In the subgroups of the miRNAs profiling, the multiple miRNAs had a higher accuracy for diagnosing the NSCLC with SEN: 88% (95% CI: 85-91%), SPE: 91% (95% CI: 88-93%), and AUC: 0.96 (95% CI: 0.93-0.97), respectively, when compared with the single miRNA with the SEN: 77% (95% CI: 71-82%), SPE: 80% (95% CI: 77-84%), and AUC: 0.86 (95% CI: 0.82-0.88), respectively. miRNAs had a higher value to distinguish the NSCLC patients from healthy individuals with the SEN: 86% (95% CI: 82-89%), SPE: 88% (95% CI: 85-91%), and AUC: 0.94 (95% CI: 0.91-0.95) than controls with benign pulmonary disease with SEN: 84% (95% CI: 77-89%), SPE: 84% (95% CI: 80-88%), and AUC: 0.90 (95% CI: 0.87-0.92). Compared with other miRNAs, miR-210 and miR-21 were more often used as diagnostic tools. However, they were usually associated with other miRNAs. The sensitivity, specificity, and AUC were, respectively, 77% (95% CI: 72-81%), 93% (95% CI: 88-96%), and 0.91 (95% CI: 0.88-0.93) for miR-210 with other miRNAs. The sensitivity, specificity, and AUC of miR-21 with other miRNAs were, respectively, 82% (95% CI: 77-86%), 87% (95% CI: 84-89%), and 0.91 (95% CI: 0.88-0.93).

4. Discussion

Due to the high mortality rate and low survival rate of NSCLC, there is an urgent need for the accurate detection method for the early detection of NSCLC especially for the nonsmoking NSCLC patients. Although miRNAs may have a high diagnostic accuracy according to the previous articles, the clinical utility of the miRNA for diagnosing NSCLC remains controversial. Compared with the previous meta-analyses [69–71], there were more studies and participants included in this meta-analysis. Our analysis showed the

pooled SEN was 85% (95% CI: 82-88%), the pooled SPE was 88% (95% CI: 85-90%), and the AUC was 0.93 (95% CI: 0.90-0.95), suggesting that miRNAs had pretty high diagnostic value for NSCLC. Our results also showed that the pooled DOR was 40 (95% CI: 28-58), indicating that for an individual proved positive by miRNAs the chance of having NSCLC is 40 times higher than the negative ones. For the subgroup analyses, higher accuracy was observed in the multiple miRNA profiling when compared with the single miRNA, which was consistent with the previous conclusions [69–71]. MiRNAs might have a higher diagnostic efficacy for the nonsmoking NSCLC patients compared with the smoking ones. Meanwhile, differences were also observed among the Caucasian, Asian, and Caucasian/African populations. This result could be supported by the Wang H's article [71]. Furthermore, miRNAs from serum samples exhibited higher diagnostic value than miRNAs from other specimen. These results meant that combinations of various miRNAs may be better diagnostic tools than the single miRNA, and miRNA isolated from serum could have a higher diagnostic value for the Caucasian populations when compared with the Asian and Caucasian/African populations. Among the different multiple miRNAs, miR-210 and miR-21 associated with other miRNAs could be used for the detection of NSCLC. However, there were still some limitations that could not be neglected in this meta-analysis such as the heterogeneity among these 71 studies, the different methods in miRNA profiling, the possibility that some articles are missed or not published online.

5. Conclusions

Our meta-analysis showed the practicability of miRNAs for diagnosing NSCLC and demonstrated that the multiple miRNAs might have a relatively high diagnostic value for NSCLC compared with the single miRNA diagnosis. miR-210 and miR-21 could be used as effective tools through combining with other miRNAs. In addition, miRNAs, especially isolated from serum, had a better diagnostic accuracy in Caucasian populations than the Asian populations as well as the Caucasian/African populations. When compared with the smoking NSCLC patients, miRNAs might have a higher diagnostic efficacy for the nonsmoking ones. However, studies on the large samples are still demanded to verify our results.

Abbreviations

miRNA:	MicroRNA
NSCLC:	Non-small cell lung cancer
SEN:	Sensitivity
SPE:	Specificity
SROC:	Summary receiver operating characteristic
AUC:	The area under the SROC curve
mRNA:	Messenger RNA
3'-UTR:	3'-untranslated region
BPD:	Benign pulmonary disease
HC:	Healthy Control
CNKI:	Chinese national knowledge infrastructure

PLR: Positive likelihood ratio
 NLR: Negative likelihood ratio
 DOR: Diagnostic odds ratio
 TP: True positive
 FP: False positive
 FN: False negative
 TN: True negative
 qRT-PCR: Quantitative real-time polymerase chain reaction
 PBMC: Peripheral blood mononuclear cells
 PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses
 BAL: Bronchoalveolar lavage.

Conflicts of Interest

The authors declare that there are no potential conflicts of interest.

Authors' Contributions

Baosen Zhou, Xuelian Li, and Min Jiang conceived and designed this study. Min Jiang and Xiaoying Li searched the literature and analyzed the data. Xiaowei Quan contributed to the analysis tools and the statistical analysis. Min Jiang and Baosen Zhou wrote and revised the paper.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (no. 81502878) and the Doctoral Research Project of Liaoning Province (no. 201601117).

References

- [1] L. A. Torre, F. Bray, R. L. Siegel, J. Ferlay, and J. Lortet-Tieulent, "Global cancer statistics, 2012," *CA: A Cancer Journal for Clinicians*, vol. 65, no. 2, pp. 87–108, 2015.
- [2] M. Lagos-Quintana, R. Rauhut, W. Lendeckel, and T. Tuschl, "Identification of novel genes coding for small expressed RNAs," *Science*, vol. 294, no. 5543, pp. 853–858, 2001.
- [3] S. P. Nana-Sinkam and C. M. Croce, "Non-coding RNAs in cancer initiation and progression and as novel biomarkers," *Molecular Oncology*, vol. 5, no. 6, pp. 483–491, 2011.
- [4] G. A. Calin and C. M. Croce, "MicroRNA signatures in human cancers," *Nature Reviews Cancer*, vol. 6, no. 11, pp. 857–866, 2006.
- [5] X. Zhang, P. Li, and M. Rong, "MicroRNA-141 is a biomarker for progression of squamous cell carcinoma and adenocarcinoma of the lung: clinical analysis of 125 patients," *The Tohoku Journal of Experimental Medicine*, vol. 235, no. 3, pp. 161–169, 2015.
- [6] X. Wang, X. Zhi, Y. Zhang, G. An, and G. Feng, "Role of plasma MicroRNAs in the early diagnosis of non-small-cell lung cancers: a case-control study," *Journal of Thoracic Disease*, vol. 8, no. 7, pp. 1645–1652, 2016.
- [7] Y. Wang, H. Zhao, X. Gao et al., "Identification of a three-miRNA signature as a blood-borne diagnostic marker for early diagnosis of lung adenocarcinoma," *Oncotarget*, vol. 7, no. 18, pp. 26070–26086, 2016.
- [8] B. Wang and Q. Zhang, "The expression and clinical significance of circulating microRNA-21 in serum of five solid tumors," *Journal of Cancer Research and Clinical Oncology*, vol. 138, no. 10, pp. 1659–1666, 2012.
- [9] M. E. Hatley, D. M. Patrick, M. R. Garcia et al., "Modulation of K-Ras-dependent lung tumorigenesis by MicroRNA-21," *Cancer Cell*, vol. 18, no. 3, pp. 282–293, 2010.
- [10] J. Osugi, Y. Kimura, Y. Owada et al., "Prognostic Impact of Hypoxia-Inducible," *Journal of Oncology*, vol. 2015, pp. 1–8, 2015.
- [11] J. Wei, L.-K. Liu, W. Gao et al., "Reduction of plasma MicroRNA-21 is associated with chemotherapeutic response in patients with non-small cell lung cancer," *Chinese Journal of Cancer Research*, vol. 23, no. 2, pp. 123–128, 2011.
- [12] Y. Su, M. A. Guarnera, H. Fang, and F. Jiang, "Small non-coding RNA biomarkers in sputum for lung cancer diagnosis," *Molecular Cancer*, vol. 15, no. 1, article no. 36, 2016.
- [13] J. O. Kim, S. Gazala, R. Razzak et al., "Non-small cell lung cancer detection using microRNA expression profiling of bronchoalveolar lavage fluid and sputum," *Anticancer Research*, vol. 35, pp. 1873–1880, 2015.
- [14] R. Razzak, E. Bédard, J. Kim et al., "MicroRNA expression profiling of sputum for the detection of early and locally advanced non-small-cell lung cancer: a prospective case-control study," *Current Oncology*, vol. 23, no. 2, p. 86, 2016.
- [15] P. F. Whiting, A. W. S. Rutjes, M. E. Westwood et al., "Quadas-2: a revised tool for the quality assessment of diagnostic accuracy studies," *Annals of Internal Medicine*, vol. 155, no. 8, pp. 529–536, 2011.
- [16] J. P. T. Higgins, S. G. Thompson, J. J. Deeks, and D. G. Altman, "Measuring inconsistency in meta-analyses," *British Medical Journal*, vol. 327, no. 7414, pp. 557–560, 2003.
- [17] J. J. Deeks, P. Macaskill, and L. Irwig, "The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed," *Journal of Clinical Epidemiology*, vol. 58, no. 9, pp. 882–893, 2005.
- [18] L. Xing, N. W. Todd, L. Yu, H. Fang, and F. Jiang, "Early detection of squamous cell lung cancer in sputum by a panel of microRNA markers," *Modern Pathology*, vol. 23, no. 8, pp. 1157–1164, 2010.
- [19] L. Yu, N. W. Todd, L. Xing et al., "Early detection of lung adenocarcinoma in sputum by a panel of microRNA markers," *International Journal of Cancer*, vol. 127, no. 12, pp. 2870–2878, 2010.
- [20] H. C. Jeong, E. K. Kim, J. H. Lee, J. M. Lee, H. Nayoo, and J. K. Kim, "Aberrant expression of let-7a miRNA in the blood of non-small cell lung cancer patients," *Molecular Medicine Reports*, vol. 4, no. 2, pp. 383–387, 2011.
- [21] J. Shen, Z. Liu, N. W. Todd et al., "Diagnosis of lung cancer in individuals with solitary pulmonary nodules by plasma microRNA biomarkers," *BMC Cancer*, vol. 11, article 374, 2011.
- [22] S. Liu and J. Yang, "Application of plasma circulation microRNA - 126 to diagnosis of non - small cell lung cancer," *Medical Journal of National Defending Forces in Southwest China*, vol. 21, pp. 1280–1283, 2011.
- [23] P. T. Hennessey, T. Sanford, A. Choudhary et al., "Serum microRNA biomarkers for detection of non-small cell lung cancer," *PLoS ONE*, vol. 7, no. 2, Article ID e32307, 2012.
- [24] M. Yuxia, T. Zhennan, and Z. Wei, "Circulating miR-125b is a novel biomarker for screening non-small-cell lung cancer and predicts poor prognosis," *Journal of Cancer Research and Clinical Oncology*, vol. 138, no. 12, pp. 2045–2050, 2012.

- [25] M. Zhao, X. Shen, P. Zhan, T. Lv, H. Liu, and Y. Song, "Expression of miRNA-29a in non-small cell lung cancer and clinical significance," *Chinese Clinical Oncology*, vol. 17, pp. 428–432, 2012.
- [26] A. A. Abd-El-Fattah, N. A. H. Sadik, O. G. Shaker, and M. L. Aboulftouh, "Differential microRNAs expression in serum of patients with lung cancer, pulmonary tuberculosis, and pneumonia," *Cell Biochemistry and Biophysics*, vol. 67, no. 3, pp. 875–884, 2013.
- [27] N. Anjuman, N. Li, M. Guarnera, S. A. Stass, and F. Jiang, "Evaluation of lung flute in sputum samples for molecular analysis of lung cancer," *linical and Translational Medicine*, vol. 2, article 15, 2013.
- [28] N. G. Bediaga, M. P. A. Davies, A. Acha-Sagredo et al., "A microRNA-based prediction algorithm for diagnosis of non-small lung cell carcinoma in minimal biopsy material," *British Journal of Cancer*, vol. 109, no. 9, pp. 2404–2411, 2013.
- [29] R. Cazzoli, F. Buttitta, M. di Nicola et al., "MicroRNAs derived from circulating exosomes as noninvasive biomarkers for screening and diagnosing lung cancer," *Journal of Thoracic Oncology*, vol. 8, no. 9, pp. 1156–1162, 2013.
- [30] J. Ma, N. Li, M. Guarnera, and F. Jiang, "Quantification of Plasma miRNAs by Digital PCR for Cancer Diagnosis," *Biomarker Insights*, vol. 8, p. BMI.S13154, 2013.
- [31] P. Mozzoni, I. Banda, M. Goldoni et al., "Plasma and EBC microRNAs as early biomarkers of non-small-cell lung cancer," *Biomarkers*, vol. 18, no. 8, pp. 679–686, 2013.
- [32] D. Tang, Y. Shen, M. Wang et al., "Identification of plasma microRNAs as novel noninvasive biomarkers for early detection of lung cancer," *European Journal of Cancer Prevention*, vol. 22, no. 6, pp. 540–548, 2013.
- [33] P. Ulivi, G. Foschi, M. Mengozzi et al., "Peripheral blood miR-328 expression as a potential biomarker for the early diagnosis of NSCLC," *International Journal of Molecular Sciences*, vol. 14, no. 5, pp. 10332–10342, 2013.
- [34] X. L. Zeng, S. Y. Zhang, J. F. Zheng, H. Yuan, and Y. Wang, "Altered miR-143 and miR-150 expressions in peripheral blood mononuclear cells for diagnosis of non-small cell lung cancer," *Chinese Medical Journal*, vol. 126, pp. 4510–4516, 2013.
- [35] Z. Wang, X. Zhang, H. Bai et al., "EML4-ALK Rearrangement and Its Clinical Significance in Chinese Patients with Advanced Non-Small Cell Lung Cancer," *Oncology*, vol. 83, no. 5, pp. 248–256, 2012.
- [36] X. Yang, Y. Zhang, B. Sun et al., "Diagnostic value of the detection of MicroRNAs in sputum of patients with non-small cell lung cancer," *Journal of Clinical Pulmonary Medicine*, vol. 18, pp. 226–229, 2013.
- [37] M. Li, Q. Zhang, L. Wu et al., "Serum miR-499 as a novel diagnostic and prognostic biomarker in non-small cell lung cancer," *Oncology Reports*, vol. 31, no. 4, pp. 1961–1967, 2014.
- [38] N. Li, J. Ma, M. A. Guarnera, H. Fang, L. Cai, and F. Jiang, "Digital PCR quantification of miRNAs in sputum for diagnosis of lung cancer," *Journal of Cancer Research and Clinical Oncology*, vol. 140, no. 1, pp. 145–150, 2014.
- [39] W. Zhu, J. He, D. Chen et al., "Expression of miR-29c, miR-93, and miR-429 as potential biomarkers for detection of early stage non-small lung cancer," *PLoS ONE*, vol. 9, no. 2, Article ID e87780, 2014.
- [40] H. Dou, Y. Wang, G. Su, and S. Zhao, "Decreased plasma let-7c and miR-152 as noninvasive biomarker for non-small-cell lung cancer," *International Journal of Clinical and Experimental Medicine*, vol. 8, pp. 9291–9298, 2015.
- [41] W. Guo, Y. Zhang, Y. Zhang et al., "Decreased expression of MIR-204 in plasma is associated with a poor prognosis in patients with non-small cell lung cancer," *International Journal of Molecular Medicine*, vol. 36, no. 6, pp. 1720–1726, 2015.
- [42] L. Li, Y. Y. Chen, Li. SQ, C. Huang, and Y. Z. Qin, "Expression of miR-148/152 family as potential biomarkers in non-small-cell lung cancer," *Medical Science Monitor*, vol. 21, pp. 1155–1161, 2015.
- [43] C. Liu, C. Wang, F. Chen, H. Yu, J. Wang, and J. Yang, "Combined detection of pleural effusion cell DNA aneuploidy and microRNA-192 and its related factors in diagnosis of non-small cell lung cancer," *China Journal of Modern Medicine*, vol. 25, pp. 26–30, 2015 (Chinese).
- [44] J. Ma, Y. Lin, M. Zhan, D. L. Mann, S. A. Stass, and F. Jiang, "Differential miRNA expressions in peripheral blood mononuclear cells for diagnosis of lung cancer," *Laboratory Investigation*, vol. 95, no. 10, pp. 1197–1206, 2015.
- [45] E. Nadal, A. Truini, A. Nakata et al., "A Novel Serum 4-microRNA Signature for Lung Cancer Detection," *Scientific Reports*, vol. 5, Article ID 12464, 2015.
- [46] C. Wang, M. Ding, M. Xia et al., "A Five-miRNA Panel Identified From a Multicentric Case-control Study Serves as a Novel Diagnostic Tool for Ethnically Diverse Non-small-cell Lung Cancer Patients," *EBioMedicine*, vol. 2, no. 10, pp. 1377–1385, 2015.
- [47] R. J. Wang, Y. H. Zheng, P. Wang, and J. Z. Zhang, "Serum miR-125a-5p, miR-145 and miR-146a as diagnostic biomarkers in non-small cell lung cancer," *International Journal of Clinical and Experimental Pathology*, vol. 8, pp. 765–771, 2015.
- [48] L. Xing, J. Su, M. A. Guarnera et al., "Sputum microRNA biomarkers for identifying lung cancer in indeterminate solitary pulmonary nodules," *Clinical Cancer Research*, vol. 21, no. 2, pp. 484–489, 2015.
- [49] J.-S. Yang, B.-J. Li, H.-W. Lu et al., "Serum miR-152, miR-148a, miR-148b, and miR-21 as novel biomarkers in non-small cell lung cancer screening," *Tumor Biology*, vol. 36, no. 4, pp. 3035–3042, 2015.
- [50] Y.-L. Yang, L.-P. Xu, F.-L. Zhuo, and T.-Y. Wang, "Prognostic value of microRNA-10b overexpression in peripheral blood mononuclear cells of nonsmall-cell lung cancer patients," *Tumor Biology*, vol. 36, no. 9, pp. 7069–7075, 2015.
- [51] W. Zhao, J. J. Zhao, L. Zhang et al., "Serum miR-21 level: a potential diagnostic and prognostic biomarker for non-small cell lung cancer," *International Journal of Clinical and Experimental Medicine*, vol. 8, pp. 14759–14763, 2015.
- [52] L. Fan, H. Qi, J. Teng et al., "Identification of serum miRNAs by nano-quantum dots microarray as diagnostic biomarkers for early detection of non-small cell lung cancer," *Tumor Biology*, vol. 37, no. 6, pp. 7777–7784, 2016.
- [53] X. Gao, Y. Wang, H. Zhao et al., "Plasma miR-324-3p and miR-1285 as diagnostic and prognostic biomarkers for early stage lung squamous cell carcinoma," *Oncotarget*, vol. 7, no. 37, 2016.
- [54] A. R. Halvorsen, M. Bjaanæs, M. LeBlanc et al., "A unique set of 6 circulating microRNAs for early detection of non-small cell lung cancer," *Oncotarget*, vol. 7, no. 24, pp. 37250–37259, 2016.
- [55] L.-P. Jiang, Z.-T. Zhu, and C.-Y. He, "Expression of miRNA-26b in the diagnosis and prognosis of patients with non-small-cell lung cancer," *Future Oncology*, vol. 12, no. 9, pp. 1105–1115, 2016.
- [56] P. Leidinger, T. Brefort, C. Backes et al., "High-throughput qRT-PCR validation of blood microRNAs in non-small cell lung cancer," *Oncotarget*, vol. 7, no. 4, pp. 4611–4623, 2016.

- [57] K. Su, T. Zhang, Y. Wang, and G. Hao, "Diagnostic and prognostic value of plasma microRNA-195 in patients with non-small cell lung cancer," *World Journal of Surgical Oncology*, vol. 14, no. 1, 2016.
- [58] L. Sun, Y. Chen, Q. Su et al., "Increased plasma miRNA-30a as a biomarker for non-small cell lung cancer," *Medical Science Monitor*, vol. 22, pp. 647–655, 2016.
- [59] M. C. Tai, K. Yanagisawa, M. Nakatochi et al., "Blood-borne miRNA profile-based diagnostic classifier for lung adenocarcinoma," *Scientific Reports*, vol. 6, no. 1, 2016.
- [60] J. Wei, W. Gao, C.-J. Zhu et al., "Identification of plasma microRNA-21 as a biomarker for early detection and chemosensitivity of non-small cell lung cancer," *Chinese Journal of Cancer*, vol. 30, no. 6, pp. 407–414, 2011.
- [61] W. Zhu, K. Zhou, Y. Zha et al., "Diagnostic value of serum miR-182, miR-183, miR-210, and miR-126 levels in patients with early-stage non-small cell lung cancer," *PLoS ONE*, vol. 11, no. 4, Article ID e0153046, 2016.
- [62] H. Zhang, F. Mao, T. Shen et al., "Plasma miR-145, miR-20a, miR-21 and miR-223 as novel biomarkers for screening early-stage non-small cell lung cancer," *Oncology Letters*, vol. 13, no. 2, pp. 669–676, 2017.
- [63] A. Keller, P. Leidinger, A. Borries et al., "MiRNAs in lung cancer - Studying complex fingerprints in patient's blood cells by microarray experiments," *BMC Cancer*, vol. 9, article no. 1471, p. 353, 2009.
- [64] G. Li, Y. Huang, Y. He et al., "Study of miRNA Signatures as Biomarkers for Early Stage Nonsmoking Female Lung Cancer of Xuanwei Country of Yunnan Province," *Cancer prevention and control research*, vol. 39, pp. 802–806, 2012.
- [65] W. H. Roa, J. O. Kim, R. Razzak et al., "Sputum microRNA profiling: a novel approach for the early detection of non-small cell lung cancer," *Clinical & Investigative Medicine*, vol. 35, no. 5, pp. E271–E281, 2012.
- [66] C. Sanfiorenzo, M. I. Ilie, A. Belaid et al., "Two panels of plasma microRNAs as non-invasive biomarkers for prediction of recurrence in resectable NSCLC," *PLoS ONE*, vol. 8, no. 1, Article ID e54596, 2013.
- [67] W. Li, Y. Wang, Q. Zhang et al., "MicroRNA-486 as a biomarker for early diagnosis and recurrence of non-small cell lung cancer," *PLoS ONE*, vol. 11, no. 1, 2016.
- [68] W. Wang, W. Li, M. Ding et al., "Identification of miRNAs as non-invasive biomarkers for early diagnosis of lung cancers," *Tumor Biology*, vol. 37, no. 12, pp. 16287–16293, 2016.
- [69] L. Chen and H. Jin, "MicroRNAs as novel biomarkers in the diagnosis of non-small cell lung cancer: a meta-analysis based on 20 studies," *Tumor Biology*, vol. 35, no. 9, pp. 9119–9129, 2014.
- [70] Y. Huang, Q. Hu, Z. Deng, Y. Hang, J. Wang, and K. Wang, "MicroRNAs in body fluids as biomarkers for non-small cell lung cancer: a systematic review," *Technology in Cancer Research & Treatment*, vol. 13, no. 3, pp. 277–287, 2014.
- [71] H. Wang, S. Wu, L. Zhao, J. Zhao, J. Liu, and Z. Wang, "Clinical use of microRNAs as potential non-invasive biomarkers for detecting non-small cell lung cancer: A meta-analysis," *Respirology*, vol. 20, no. 1, pp. 56–65, 2015.