Research Article

Validation of Reference Genes for Gene Expression Normalization in RAW264.7 Cells under Different Conditions

Zhenzhen Bao,^{1,2} Yanli Huang,² Jiyu Chen,² Zhenglong Wang,²
Jiang Qian,² Jiyang Xu ,² and Yucheng Zhao ^{2,3}

¹School of Pharmacy, Jiangsu Health Vocational College, Nanjing, Jiangsu, China

²School of Life Science and Technology, China Pharmaceutical University, Nanjing, Jiangsu, China

³Jiangsu Key Laboratory of Bioactive Natural Product Research and State Key Laboratory of Natural Medicines, School of Traditional Chinese Pharmacy, China Pharmaceutical University, Nanjing, Jiangsu, China

Correspondence should be addressed to Jiyang Xu; jiyangx@126.com and Yucheng Zhao; zhaoyucheng1986@126.com

Received 6 November 2018; Revised 3 March 2019; Accepted 3 April 2019; Published 16 May 2019

Academic Editor: Salvatore Gallone

Copyright © 2019 Zhenzhen Bao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

RAW264.7 is a macrophage strain derived from mice tumour and shows a significant ability in antigen uptake. Real-time quantitative PCR (RT-qPCR) is one of the most commonly used methods in gene studies and requires suitable reference genes to normalize and quantitate the expression of gene of interest with sensitivity and specificity. However, suitable reference genes in RAW264.7 cells have not yet been identified for accurate gene expression quantification. In the current study, we evaluated expression levels of ten candidate reference genes in RAW264.7 cells under different conditions. RT-qPCR results indicated significant differences in the expression levels among the ten reference genes. Statistical analyses were carried out using geNorm, NormFinder, and BestKeeper software to further investigate the stability of the reference genes. Integrating the results from the three analytical methods, cytochrome c-1 and hydroxymethylbilane synthase were found to be the most stable and therefore more suitable reference genes, while ribosomal protein L4 and cyclophilin A were the least stable. This study emphasises the importance of identifying and selecting the most stable reference genes for normalization and provides a basis for future gene expression studies using RAW264.7 cells.

1. Introduction

Reverse transcription quantitative real-time PCR (RT-qPCR) is an important method for gene expression studies [1, 2]. This technology has become a very popular method owing to its high speed, high sensitivity, and high-throughput capabilities [3–5]. However, the results are inevitably affected by sample variation and PCR efficiency, which could lead to erroneous interpretations. Hence, in order to ensure accurate measurement of the expression levels of genes in various conditions, normalization of target gene expression with that of a proper reference gene is absolutely necessary when using RT-qPCR [6–8]. Several reference genes, including glyceraldehyde-3-phosphate dehydrogenase (GAPDH), beta-2-microglobulin (B2M), and β -actin (ACTB), have been reported for their stable expression in all tissues and cells [9, 10]. Considering that gene expression levels may vary among cells or tissues

and may also change under certain circumstances, geNorm [11], NormFinder [12], and BestKeeper [13] analytical software have been specially designed for screening of reference gene stability.

Macrophages are important immune cells and play a critical role in anti-infection, antitumour, and immune regulation processes. A large number of cell lines of murine macrophages, such as P338D1 and J774A1 cells, have been commonly used in the study of microbiology and immunology [14, 15]. In contrast, RAW264.7 cells, macrophages derived from mice tumour, have rarely been examined for suitable candidate reference gene expression in a variety of drug treatments [16]. Hence, in this study, we chose RAW264.7 cells to evaluate the expression levels of reference genes for reliable normalization under different conditions. Conversely, a number of studies have shown that no single reference gene exists that can be expressed stably under any

TABLE 1: Information of the ten reference genes used in the real time quantitative PCR.

Gene	Description	Primer: forward/reverse (5'-3')	Length (bp)	Accession number
ACTB	β -actin	F: CAGGTCATCACTATTGGCAA R: AGGTCTTTACGGATGTCAAC	143	NM_007393
GAPDH	Glyceraldehyde-3-phosphate dehydrogenase	F: TGCTGAGTATGTCGTGGAGT R: GTTCACACCCATCACAAACA	136	NM_001289726
RPL4	Ribosomal protein L4	F: GGAAGTTGGATGAGCTGTAT R: TCAAGATTCTGCTAAGGTCG	108	NM_024212
HPRT1	Hypoxanthine phosphoribosyltransferase-1	F: TAGTGAAACTGGAAAAGCCA R: AAGCTTTACTAGGCAGATGG	135	NM_013556
PPIA	Cyclophilin A	F: CGGTTCCCAGTTTATCT R: ATGGCTTCCACAATGTTTCAT	102	NM_008907
CYC1	Cytochrome c-1	F: CTAACCCTGAGGCTGCAAGA R: GCCAGTGAGCAGGAAAATA	113	NM_025567
HMBS	Hydroxymethyl-bilane synthase	F: ATGGCTCAGATAGCATGCAA R: GGGCTCCTCTTGAATGTTA	126	NM_013551
Eef1a1	Eukaryotic translation elongation factor 1 alpha 1	F: CCTACCACCAACTCGTCCAA R: AAAGGTAACCACCATGCCAG	136	NM_010106
GUSB	Glucuronidase β	F: ATGTCCTGCTGAGAGGTGTC R: TCCAGCCTCTCACCAGTAGC	121	NM_010368
LDHA	lactate dehydrogenase A	F: AACATCTCACTCCCCACAGC R: CTCACAGGGGTAATCGAAGC	128	NM_010699

experimental condition [17]. Therefore, ten reference genes, ACTB, GAPDH, ribosomal protein L4 (RPL4), hypoxanthine phosphoribosyltransferase-1 (HPRT1), cyclophilin A (PPIA), Cytochrome c-1 (CYC1), hydroxymethylbilane synthase (HMBS), eukaryotic translation elongation factor 1 alpha 1 (Eef1a1), glucuronidase β (GUSB), and lactate dehydrogenase A (LDHA), were selected and analysed for suitability under different conditions [9, 10, 18–24].

This study utilized the three analytical methods mentioned above to examine the stability of multiple commonly used reference genes in RAW264.7 cells. To the best of our knowledge, this is the first study to systematically evaluate the expression stability of candidate reference genes in RAW264.7 cells under different conditions (various drugs and concentrations). We believe that the present work will provide a substantial foundation for future research in mouse or human cells.

2. Materials and Methods

2.1. Cell Culture and Treatment. Murine macrophage cells, RAW264.7, were grown in DMEM (Gibco, USA) supplemented with 100 U/mL penicillin-streptomycin and 10% foetal bovine serum (FBS, Bioind) and incubated at 37°C in a 5% CO₂ humidified atmosphere. Upon confluence, cells were trypsinised and treated, respectively, with various drugs at multiple concentrations: high-glucose (HG; 50 mM, 100 mM, 200 mM), hydrogen peroxide (H₂O₂; 50 μ M, 100 μ M, 200 μ M), lipopolysaccharide (LPS; 0.1, 0.5, 1 μ g/mL), cobalt chloride (CoCl₂; 50 μ M, 100 μ M, 200 μ M), and palmitic acid (PA; 50 μ M, 100 μ M, 200 μ M). Cells without treatment (WT) acted as the control

group. Three biological repeats were used for each condition.

2.2. PCR Primers for Reference Genes. Ten reference genes (ACTB, GAPDH, RPL4, HPRT1, PPIA, CYC1, HMBS, Eef1a1, GUSB, and LDHA) of RAW264.7 cells were selected for evaluation based on the high frequency of their use in other related studies [9, 10, 18–24]. Primers used for determining the expression of the ten reference genes are listed in Table 1. According to the manufacturer, these primers were designed and optimized using Primer 5 as follows: primer length, 18–22 bp; GC content, 40%–60%; amplification length, 100–150 bp, to avoid the formation of primer-dimer during the reaction.

2.3. RNA Extraction and cDNA Synthesis. RNA was extracted from the treated and control RAW264.7 cells using RNAiso Plus total RNA kit (TransGen Biotech, Dalian, China). DNase I (Takara, Dalian, China) treatment was carried out to purify RNA. The RNA samples with OD₂₆₀/OD₂₈₀ ratios between 1.8 and 2.0 and a total amount of 0.5 μ g were used for cDNA synthesis to ensure the precision of the trial. Additionally, the purification of the RNA was confirmed by agarose gel electrophoresis. The cDNA was synthesised from the RNA using reverse transcriptase and the HiScript® Q RT SuperMix for qPCR Kit (Vazyme, Nanjing, China), by following manufacturer's instructions, and stored at –20°C until use for subsequent reaction.

2.4. Quantitative Real-Time PCR Analysis. The RT-qPCR reactions were set up using cDNA, forward/reverse primer,

and Hieff™ qPCR SYBR® Green Master Mix (Yeasen, Shanghai, China), according to manufacturer's instructions. The reactions were carried out in 96-well PCR reaction plates in a LightCycler 480 system (Roche Molecular Biochemicals, Mannheim, Germany).

2.5. Data Analysis. The stability of reference genes was analysed using the three statistical software programs: geNorm [11], NormFinder [12], and BestKeeper [13]. The geNorm measures gene expression stability according to the values of M , which were calculated from the cycle threshold (C_t) values obtained from RT-qPCR. M values indicated the pairwise variation between individual gene and the other reference genes; lower values of M represented higher expression stabilities, while a higher M value indicated a more unstable reference gene. In addition, geNorm could also determine the optimal number of candidate reference genes required for normalization based on pairwise variation. Like geNorm, NormFinder tended to rank the stability of reference genes according to the M values, with the lowest M value indicating the most stable gene. BestKeeper, an Excel-based statistical method that analyses expression variability of reference genes, ranked the stability of reference genes from most to least based on the key factors of standard deviation (SD) and coefficient of variance (CV).

3. Results

3.1. Validation of Primer Specificity. As shown in Supplementary Figure 1, PCR and subsequent agarose gel electrophoresis were used to identify the specificity of the designed primers. The single band and peak of melting curve indicated that the primers possessed good specificity (Supplementary Figure 2).

3.2. The Expression Levels of the Reference Genes. The C_t value generated from the RT-qPCR is the fluorescence threshold for each primer pair and reflects the expression levels of the reference genes; a low C_t value indicates high expression [4–6]. The overall mean C_t values for the ten reference genes are shown in Figure 1. Among the ten genes, there was an obvious difference in expression levels as seen from the C_t values, which ranged from 15 to 30. $CYC1$ was the least expressed reference gene with the highest mean C_t value, while $Eef1a1$, $GAPDH$, and $ACTB$ were the three most stably expressed genes having lower C_t values. In addition, $LDHA$ had a narrow range of C_t values, indicating that the variability of the expression level was constant under different conditions and it might be the best reference gene with a relatively high expression level. Similarly, $ACTB$ and $HMBS$ could be considered as suitable choices for stable reference genes owing to their relatively narrow C_t ranges. In contrast, $RPL4$ showed a large range of C_t values and would be unsuitable as a reference gene. To further systematically assess the stability of the ten reference genes under various treatments, the C_t values were transformed to analyse expression levels using the three software programs: geNorm, NormFinder, and BestKeeper.

FIGURE 1: Expression levels of the ten reference genes ($ACTB$, $GAPDH$, $RPL4$, $HPRT1$, $PPIA$, $CYC1$, $HMBS$, $Eef1a1$, $GUSB$, and $LDHA$) in RAW264.7 cells. Squares of the box indicate the means; the lines indicate the median and whiskers indicate the highest and lowest values.

3.3. GeNorm Analysis. GeNorm used the equation $2^{-\Delta\Delta C_t}$ (C_t values were collected from different experimental conditions), to transform the C_t values into relative quantification data to analyse the stability of reference genes, which were ranked from the most to the least stable based on their M values. The M values were calculated by pairwise variation analysis and were considered reliable to select stable reference genes when the M values were greater than 1.5. A lower M value reflected a higher stability [11]. As shown in Figure 2, the stability rank of the ten reference genes was different under different conditions, indicating that different concentrations or drug treatments affected the gene expression. For the control group, the M values of the ten reference genes were ranked as follows: $RPL4 > LDHA > PPIA > HPRT1 > GUSB > ACTB > HMBS > Eef1a1 > CYC1 > GAPDH$, revealing that $GAPDH$ was the most stable reference gene, while $RPL4$ was the least stable. However, for other groups, the ten reference genes had different stability ranks. For instance, the rank of the reference genes after LPS treatment ($0.5 \mu\text{g/mL}$) was as follows: $PPIA > LDHA > ACTB > GUSB > RPL4 > Eef1a1 > CYC1 > HMBS > GAPDH > HPRT1$, indicating that, under these conditions, $HPRT1$ ranked the most stable reference gene, while $PPIA$ was the least stable. In addition, different concentrations of the same drug resulted in the same gene to be ranked as the most stable reference gene; for example, $ACTB$ was the most stable reference gene after treatment with both low and high concentrations of HG. Moreover, different drug treatments showed the same reference gene to be the most stable at similar doses of treatment. For example, $HMBS$ was ranked the most stable candidate gene in H_2O_2 , LPS, and $CoCl_2$ treatments, respectively, but only upon treatment with high concentrations of the drugs. Taken together, these results indicated that the stability of the reference genes was not constant under different conditions. Notably, geNorm analysis indicated that $GAPDH$ might be the most stable

FIGURE 2: Expression stability of the reference genes analysed by geNorm. M values represent the average expression stability; the stability is ranked from left to right, indicating the stability is inversely related to the M values. The treatments of drugs and concentrations are listed in the figure correspondingly.

reference genes under various experimental conditions (drug treatments and concentrations).

3.4. NormFinder Analysis. The NormFinder is a statistical algorithm that calculates the M values, which are then

used to rank the stability of the reference genes. Similar to geNorm, the most stable gene is associated with the smallest M value [12]. As shown in Table 2, the ten reference genes in the control group were ranked based on stability as follows: GAPDH, HMBS, CYC1, GUSB, ACTB, HPRT1,

TABLE 2: Expression stability values of ten reference genes in RAW264.7 cells calculated by NormFinder analysis.

Rank	WT	HG	HG	H ₂ O ₂	H ₂ O ₂	LPS	LPS	LPS	CoCl ₂	CoCl ₂	CoCl ₂	PA	PA	PA
		50 mM	100 mM	200 mM	50 μM	100 μM	200 μM	0.1 μg/mL	0.5 μg/mL	1 μg/mL	50 μM	100 μM	200 μM	200 μM
1	RPL4	EF1a1	EF1a1	GAPDH	RPL4	RPL4	LDHA	LDHA	LDHA	LDHA	LDHA	EF1a1	EF1a1	GUSB
	0.048	0.073	0.067	0.039	0.051	0.043	0.053	0.046	0.046	0.056	0.048	0.058	0.069	0.042
2	LDHA	GUSB	RPL4	PPIA	LDHA	EF1a1	EF1a1	PPIA	PPIA	GUSB	PPIA	GUSB	LDHA	PPIA
	0.046	0.057	0.042	0.037	0.049	0.040	0.050	0.036	0.036	0.050	0.048	0.048	0.065	0.034
3	PPIA	RPL4	LDHA	EF1a1	PPIA	LDHA	PPIA	GUSB	GUSB	EF1a1	RPL4	LDHA	GUSB	EF1a1
	0.030	0.053	0.037	0.036	0.042	0.039	0.047	0.035	0.035	0.049	0.045	0.044	0.048	0.034
4	EF1a1	PPIA	GUSB	RPL4	HPRT1	ACTB	RPL4	ACTB	ACTB	PPIA	ACTB	RPL4	PPIA	LDHA
	0.027	0.041	0.036	0.031	0.029	0.035	0.033	0.030	0.030	0.048	0.031	0.027	0.046	0.028
5	HPRT1	HPRT1	GAPDH	LDHA	EF1a1	PPIA	GAPDH	GUSB	GAPDH	GAPDH	GAPDH	CYC1	RPL4	ACTB
	0.024	0.040	0.031	0.027	0.025	0.025	0.027	0.027	0.027	0.036	0.029	0.026	0.039	0.028
6	ACTB	GAPDH	ACTB	CYC1	GUSB	GUSB	ACTB	RPL4	RPL4	RPL4	EF1a1	PPIA	HPRT1	GAPDH
	0.023	0.039	0.028	0.020	0.023	0.016	0.023	0.022	0.022	0.032	0.028	0.031	0.028	0.024
7	GUSB	LDHA	HMBS	HPRT1	GAPDH	CYC1	CYC1	GAPDH	GAPDH	ACTB	GAPDH	ACTB	GAPDH	RPL4
	0.023	0.035	0.026	0.020	0.017	0.016	0.022	0.016	0.016	0.013	0.027	0.021	0.022	0.021
8	CYC1	HMBS	CYC1	HMBS	HMBS	HMBS	GUSB	HPRT1	HPRT1	CYC1	HPRT1	EF1a1	CYC1	HMBS
	0.016	0.030	0.025	0.020	0.014	0.015	0.014	0.012	0.012	0.006	0.022	0.019	0.018	0.020
9	HMBS	ACTB	HPRT1	GUSB	CYC1	GAPDH	HPRT1	HMBS	HMBS	HMBS	HMBS	HPRT1	ACTB	CYC1
	0.015	0.019	0.023	0.020	0.012	0.012	0.008	0.010	0.010	0.006	0.014	0.015	0.017	0.015
10	GAPDH	CYC1	PPIA	ACTB	ACTB	HPRT1	HMBS	GAPDH	HPRT1	HPRT1	CYC1	HMBS	HMBS	HPRT1
	0.011	0.018	0.020	0.018	0.010	0.009	0.006	0.010	0.010	0.002	0.007	0.001	0.006	0.015

FIGURE 3: Determination of the optimal number of reference genes for normalization using geNorm analysis. Pairwise variation ($V_n/n+1$) of reference genes under different conditions are listed. WT, H50 (H100, H200), L0.1 (L0.5, L1), C50 (C100, C200), HG50 (HG100, HG200), and PA50 (PA100, PA200), respectively, were the abbreviation for the control group; H_2O_2 , 50 μM (100 μM , 200 μM); lipopolysaccharide, 0.1 $\mu g/mL$ (0.5, 1 $\mu g/mL$); $CoCl_2$, 50 μM (100 μM , 200 μM); high-glucose, 50 mM (100 mM, 200 mM); and palmitic acid, 50 μM (100 μM , 200 μM).

Eef1a1, PPIA, LDHA, and RPL4. Thus, GAPDH was the most expressed stable reference gene, followed by HMBS; RPL4 was the least expressed one. These results were similar to those obtained with geNorm data analysis of the control group. Interestingly, under some conditions, more than one reference gene showed the same M value, and therefore identical stability; for instance, CYC1 and HPRT1 showed the same M value and were determined to be the most stably expressed genes upon treatment with 200 μM PA. Consistent with the results of geNorm analysis, the HMBS gene seemed to be the most stable reference gene, as it showed the lowest M value among the most stable genes under different conditions. However, differences were also noted between the geNorm and NormFinder analyses; for instance, CYC1 was ranked as the most stable gene by NormFinder, while geNorm ranked it as the fourth.

3.5. BestKeeper Analysis. Distinct from the geNorm and NormFinder methods, BestKeeper determines the gene stability based on the SD and the CV. The SD of the reference genes is a key factor to identify the most stable gene in the BestKeeper analysis; the higher the CV and SD values, the more unstable the reference gene [13]. Supplementary Table 1 shows the $CV \pm SD$ values, arranged in the decreasing order

of their value from top to bottom, representing the stability rank of the reference gene from least to most. Reference genes with $SD > 1.5$ were eliminated for further analysis. Specifically, PPIA ($SD = 1.53$) in 50 μM $CoCl_2$, LDHA ($SD = 1.56$) in 200 μM $CoCl_2$, and RPL4 ($SD = 2.50$) were excluded to be able to accurately select the best normalization gene from the candidate genes. GUSB and HMBS frequently appeared as the most stable reference genes. In contrast, Eef1a1 and RPL4 often ranked as the top-most unstable genes. Notably, in the control group, GUSB had the smallest $CV \pm SD$ values of 0.34 ± 0.07 , followed by CYC1 with 1.30 ± 0.36 , and RPL4 had the highest expression variation with $CV \pm SD$ value of 4.13 ± 1.19 . Together, the above findings indicated that BestKeeper analysis differed significantly from that of geNorm and NormFinder analysis.

3.6. Optimal Number of Reference Genes for Normalization. GeNorm can select the most stable expression gene and also determine the optimal number of reference genes for accurate normalization according to the pairwise variation ($V_n/n+1$), which is calculated from the two sequential normalization factors (NFn) and uses $V_n/n+1 < 0.15$ as a criterion to confirm the optimal number of reference genes for accurate analysis. The results of $V_n/n+1$ are shown in Figure 3. Most of the

V2/3 values were greater than 0.15, indicating that using only two most stably expressed genes would not improve the accuracy under all conditions. According to the cut-off value of 0.15, the control group, H_2O_2 (50 μM , 100 μM , 200 μM), LPS (0.1, 0.5, 1 $\mu\text{g/mL}$), 100 μM CoCl_2 , and 200 μM PA could be assessed using only two suitable reference genes, while three reference genes were required under conditions of 200 μM CoCl_2 and 100 μM PA, and four were required under conditions of HG (50 mM, 100 mM, 200 mM), 50 μM CoCl_2 , and 50 μM PA.

4. Discussion

It is critical to select the proper reference genes for normalization when using RT-qPCR quantification for analysing gene expression [25–27]. Our study evaluated the stability of ten reference genes in RAW264.7 cells by geNorm [11], NormFinder [12], and BestKeeper [13] programs, by comparing their expression levels under different conditions. The Ct values representing expression levels of the reference genes ranged from 15 to 30, in accordance with the results of most reports [4].

Based on the principle that lower Ct values indicate higher expression profile [4, 6], our results indicated that *Eef1a1*, *GAPDH*, and *ACTB* were the three most highly expressed reference genes, while *CYCl* was the least expressed gene with Ct values being nearly 30 under conditions. In addition, owing to a narrow Ct range, which represents a stable expression profile, *LDHA*, *ACTB*, and *HMBS* were considered as the most stable reference genes, while *RPL4* was the least stable. Our findings showed some differences between geNorm, NormFinder, and BestKeeper analyses, indicating that it is necessary to systematically analyse expression stability combining various analysis methods, as also suggested by previous reports [28–30].

GeNorm analysis indicated that *GAPDH*, *HMBS*, and *HPRT1* were the three most stable genes under different experimental conditions. NormFinder ranked *HMBS*, *CYCl*, *HPRT1*, and *GAPDH* on the basis of high expression stability, while BestKeeper indicated that *GUSB* and *HMBS* could be considered as the most suitable normalization genes. Thus, the stability ranking of the candidate genes was not consistent across different analyses. Nonetheless, on comparing the outcomes of the three analytical programs, *HMBS* was the only one that was commonly identified by all three programs to be highly and stably expressed. Additionally, only slight differences were observed between the stability of the ten reference genes when comparing results of NormFinder and geNorm, likely due to the similar method adopted by both programs to evaluate gene expression stability. Integrating results of all three analyses, *CYCl* and *HMBS* were considered to be the most stably expressing genes suitable for normalizing qPCR data in RAW264.7 cells.

GAPDH is a frequently used reference gene, although a number of studies have reported that it does not maintain constant expression level under different experimental conditions, and thus it is unsuitable for normalizing RT-qPCR data [31–34]. However, *GAPDH* was recognized as the most stable gene with a lower M value by geNorm analysis in

our study, consistent with previous studies in mouse uterus cells and J774A1 murine macrophage cells [15, 35]. *RPL4* and *PPIA* significantly ranked low, indicating that they were not suitable as normalizing genes. However, previous studies have reported that *RPL4* was a highly stable gene, as its expression was not influenced by some experimental conditions [36–38]. Meanwhile, *PPIA* has been considered a highly expressed reference gene for normalization in macrophages [33, 37].

Considering that using a single reference gene could lead to inaccuracies during RT-qPCR quantification [39, 40], it is necessary to determine the optimal number of reference genes to be used for gene expression studies. In the current study, a pairwise variation ($V_n/n+1$) indicated that most groups required only two reference genes for normalization, as the V2/3 values were lower than the cut-off value of 0.15. However, the $V_n/n+1$ values of certain groups exceeded 0.15, suggesting the need for additional reference genes (three or four) to improve the accuracy of normalization.

5. Conclusions

In summary, we evaluated the stability of ten reference genes in RAW264.7 cells using three analytical approaches, namely, geNorm, NormFinder, and BestKeeper, to determine the most stable reference genes under different conditions. Our findings reveal that two reference genes are sufficient for accurate normalization in most conditions, although some cases might require more than two reference genes for accurate evaluation of gene expression levels. Importantly, our data indicate that *CYCl* and *HMBS* are the most suitable reference genes, while *RPL4* and *PPIA* are the most variable and unsuitable genes for normalization in RAW264.7 cells. These findings might play a crucial role in the selection of appropriate reference genes in further studies.

Data Availability

The data used to support the findings of this study are included within the article.

Conflicts of Interest

The authors declare no conflicts of interest.

Authors' Contributions

Zhenzhen Bao, Yanli Huang, and Jiyu Chen contributed equally to this work.

Acknowledgments

This project is funded by National Natural Science Foundation of China (81703637, 81573564), China Postdoctoral Science Foundation (2016M601922, 2018T110577), and Natural Science Fund in Jiangsu Province (BK20170736). This work is also funded by teaching and research project of Jiangsu Health Vocational College (JKKYTD201701, JKA201812).

Supplementary Materials

Supplementary 1. Supplementary Figure 1: Agarose gel (1%) electrophoresis of the ten reference genes. 1-10 represent ACTB, GAPDH, RPL4, HPRT1, PPIA, CYC1, HMBS, Eef1a1, GUSB, and LDHA, respectively.

Supplementary 2. Supplementary Figure 2: Melt curves of the ten reference genes.

Supplementary 3. Supplementary Table 1: Expression stability values of the ten reference genes calculated by BestKeeper in RAW264.7 cells.

References

- [1] C. Caldana, W.-R. Scheible, B. Mueller-Roeber, and S. Ruzicic, "A quantitative RT-PCR platform for high-throughput expression profiling of 2500 rice transcription factors," *Plant Methods*, vol. 3, no. 1, p. 7, 2007.
- [2] M. W. Pfaffl, "The ongoing evolution of qPCR," *Methods*, vol. 50, no. 4, pp. 215–216, 2010.
- [3] X. Li, D. Zhang, H. Li et al., "Characterization of reference genes for RT-qPCR in the desert moss *Syntrichia caninervis* in response to abiotic stress and desiccation/rehydration," *Frontiers in Plant Science*, vol. 6, p. 38, 2015.
- [4] S. A. Bustin, V. Benes, J. A. Garson et al., "The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments," *Clinical Chemistry*, vol. 55, no. 4, pp. 611–622, 2009.
- [5] H. Cassan-Wang, M. Soler, H. Yu et al., "Reference genes for high-throughput quantitative reverse transcription-PCR analysis of gene expression in organs and tissues of eucalyptus grown in various environmental conditions," *Plant & Cell Physiology (PCP)*, vol. 53, no. 12, pp. 2101–2116, 2012.
- [6] Q. Gao, X.-Y. Wang, J. Fan et al., "Selection of reference genes for real-time PCR in human hepatocellular carcinoma tissues," *Journal of Cancer Research and Clinical Oncology*, vol. 134, no. 9, pp. 979–986, 2008.
- [7] S. M. Gresner, E. Golanska, D. Kulczycka-Wojdala, D. J. Jaskolski, W. Papierz, and P. P. Liberski, "Selection of reference genes for gene expression studies in astrocytomas," *Analytical Biochemistry*, vol. 408, no. 1, pp. 163–165, 2011.
- [8] H. Zhuang, Y. Fu, W. He, L. Wang, and Y. Wei, "Selection of appropriate reference genes for quantitative real-time PCR in *Oxytropis ochrocephala* Bunge using transcriptome datasets under abiotic stress treatments," *Frontiers in Plant Science*, vol. 6, p. 475, 2015.
- [9] Y. Higashimura, Y. Nakajima, R. Yamaji et al., "Up-regulation of glyceraldehyde-3-phosphate dehydrogenase gene expression by HIF-1 activity depending on Sp1 in hypoxic breast cancer cells," *Archives of Biochemistry and Biophysics*, vol. 509, no. 1, pp. 1–8, 2011.
- [10] F. Wang, J. Wang, D. Liu, and Y. Su, "Normalizing genes for real-time polymerase chain reaction in epithelial and nonepithelial cells of mouse small intestine," *Analytical Biochemistry*, vol. 399, no. 2, pp. 211–217, 2010.
- [11] J. Vandesompele, K. De Preter, F. Pattyn et al., "Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes," *Genome Biology*, vol. 3, no. 7, Article ID RESEARCH0034, 2002.
- [12] C. L. Andersen, J. L. Jensen, and T. F. Ørntoft, "Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets," *Cancer Research*, vol. 64, no. 15, pp. 5245–5250, 2004.
- [13] M. W. Pfaffl, A. Tichopad, C. Prgomet, and T. P. Neuvians, "Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: bestKeeper—excel-based tool using pair-wise correlations," *Biotechnology Letters*, vol. 26, no. 6, pp. 509–515, 2004.
- [14] D. M. Segal, S. K. Dower, and J. A. Titus, "The FcR-mediated endocytosis of model immune complexes by cells from the P338D1 mouse macrophage line. I. Internalization of small, nonaggregating oligomers of IgG," *The Journal of Immunology*, vol. 130, no. 1, pp. 130–137, 1983.
- [15] F. Ferraz and J. Fernandez, "Selection and validation of reference house-keeping genes in the J774A1 macrophage cell line for quantitative real-time PCR," *Genetics and Molecular Research*, vol. 15, no. 1, Article ID 15017720, 2016.
- [16] D. L. Taylor, P. C. Thomson, K. de Silva, and R. J. Whittington, "Validation of endogenous reference genes for expression profiling of RAW264.7 cells infected with *Mycobacterium avium* subsp. *paratuberculosis* by quantitative PCR," *Veterinary Immunology and Immunopathology*, vol. 115, no. 1–2, pp. 43–55, 2007.
- [17] K. Dheda, J. F. Huggett, J. S. Chang et al., "The implications of using an inappropriate reference gene for real-time reverse transcription PCR data normalization," *Analytical Biochemistry*, vol. 344, no. 1, pp. 141–143, 2005.
- [18] Z. Kolkova, A. Arakelyan, B. Casslén, S. Hansson, and E. Kriegova, "Normalizing to GAPDH jeopardises correct quantification of gene expression in ovarian tumours - IPO8 and RPL4 are reliable reference genes," *Journal of Ovarian Research*, vol. 6, no. 1, p. 60, 2013.
- [19] A. S. Stephens, S. R. Stephens, and N. A. Morrison, "Internal control genes for quantitative RT-PCR expression analysis in mouse osteoblasts, osteoclasts and macrophages," *BMC Research Notes*, vol. 4, no. 1, p. 410, 2011.
- [20] O. J. Harrison, N. Moorjani, C. Torrens, S. K. Ohri, and F. R. Cagampang, "Endogenous reference genes for gene expression studies on bicuspid aortic valve associated aortopathy in humans," *PLoS ONE*, vol. 11, no. 10, Article ID e0164329, 2016.
- [21] W. Zhu, Y. Lin, H. Liao, and Y. Wang, "Selection of reference genes for gene expression studies related to intramuscular fat deposition in *Capra hircus* skeletal muscle," *PLoS ONE*, vol. 10, no. 3, Article ID e0121280, 2015.
- [22] B. R. Everaert, G. A. Boulet, J.-P. Timmermans, and C. J. Vrints, "Importance of suitable reference gene selection for quantitative real-time PCR: special reference to mouse myocardial infarction studies," *PLoS ONE*, vol. 6, no. 8, Article ID e23793, 2011.
- [23] T. Fink, P. Lund, L. Pilgaard, J. G. Rasmussen, M. Duroux, and V. Zachar, "Instability of standard PCR reference genes in adipose-derived stem cells during propagation, differentiation and hypoxic exposure," *BMC Molecular Biology*, vol. 9, p. 98, 2008.
- [24] S. Ren, F. Zhang, C. Li et al., "Selection of housekeeping genes for use in quantitative reverse transcription PCR assays on the murine cornea," *Molecular Vision*, vol. 16, pp. 1076–1086, 2010.
- [25] D. G. Ginzinger, "Gene quantification using real-time quantitative PCR: an emerging technology hits the mainstream," *Experimental Hematology*, vol. 30, no. 6, pp. 503–512, 2002.

- [26] V. J. Gadkar and M. Filion, "Validation of endogenous reference genes in *Buglossoides arvensis* for normalizing RT-qPCR-based gene expression data," *SpringerPlus*, vol. 4, no. 1, p. 178, 2015.
- [27] D. V. Maltseva, N. A. Khaustova, N. N. Fedotov et al., "High-throughput identification of reference genes for research and clinical RT-qPCR analysis of breast cancer samples," *Journal of Clinical Bioinformatics*, vol. 3, no. 1, p. 13, 2013.
- [28] Y. Zhao, J. Luo, S. Xu et al., "Selection of reference genes for gene expression normalization in *Peucedanum praeruptorum* Dunn under abiotic stresses, hormone treatments and different tissues," *PLoS ONE*, vol. 11, no. 3, Article ID e0152356, 2016.
- [29] F. Almeida-Oliveira, J. G. Leandro, P. Ausina, M. Sola-Penna, and D. Majerowicz, "Reference genes for quantitative PCR in the adipose tissue of mice with metabolic disease," *Biomedicine & Pharmacotherapy*, vol. 88, pp. 948–955, 2017.
- [30] B. Żyżyńska-Granica and K. Koziak, "Identification of suitable reference genes for real-time PCR analysis of statin-treated human umbilical vein endothelial cells," *PLoS ONE*, vol. 7, no. 12, Article ID e51547, 2012.
- [31] J. Gimeno, N. Eattock, A. Van Deynze, and E. Blumwald, "Selection and validation of reference genes for gene expression analysis in switchgrass (*Panicum virgatum*) using quantitative real-time RT-PCR," *PLoS ONE*, vol. 9, no. 3, Article ID e91474, 2014.
- [32] P. Matouskova, H. Bartikova, I. Bousova, V. Hanusova, B. Szotakova, and L. Skalova, "Reference genes for real-time PCR quantification of messenger rnas and micrnas in mouse model of obesity," *PLoS ONE*, vol. 9, no. 1, Article ID e86033, 2014.
- [33] F. Jacob, R. Guertler, S. Naim et al., "Careful selection of reference genes is required for reliable performance of RT-qPCR in human normal and cancer cell lines," *PLoS ONE*, vol. 8, no. 3, Article ID e59180, 2013.
- [34] R. D. Barber, D. W. Harmer, R. A. Coleman, and B. J. Clark, "GAPDH as a housekeeping gene: analysis of GAPDH mRNA expression in a panel of 72 human tissues," *Physiological Genomics*, vol. 21, no. 3, pp. 389–395, 2005.
- [35] P. Lin, X. Lan, F. Chen, Y. Yang, Y. Jin, and A. Wang, "Reference gene selection for real-time quantitative PCR analysis of the mouse uterus in the peri-implantation period," *PLoS ONE*, vol. 8, no. 4, Article ID e62462, 2013.
- [36] M. U. Cinar, M. A. Islam, M. Pröll et al., "Evaluation of suitable reference genes for gene expression studies in porcine PBMCs in response to LPS and LTA," *BMC Research Notes*, vol. 6, no. 1, p. 56, 2013.
- [37] S.-J. Park, S. G. Kwon, J. H. Hwang, D. H. Park, T. W. Kim, and C. W. Kim, "Selection of appropriate reference genes for RT-qPCR analysis in Berkshire, Duroc, Landrace, and Yorkshire pigs," *Gene*, vol. 558, no. 1, pp. 152–158, 2015.
- [38] C. Zeng, L. He, W. Peng et al., "Selection of optimal reference genes for quantitative RT-PCR studies of boar spermatozoa cryopreservation," *Cryobiology*, vol. 68, no. 1, pp. 113–121, 2014.
- [39] K. E. Reid, N. Olsson, J. Schlosser, F. Peng, and S. T. Lund, "An optimized grapevine RNA isolation procedure and statistical determination of reference genes for real-time RT-PCR during berry development," *BMC Plant Biology*, vol. 6, p. 27, 2006.
- [40] A. R. Paolacci, O. A. Tanzarella, E. Porceddu, and M. Ciaffi, "Identification and validation of reference genes for quantitative RT-PCR normalization in wheat," *BMC Molecular Biology*, vol. 10, p. 11, 2009.