

Research Article

Phytochemical Analysis, Anti-inflammatory, and Antioxidant Activities of *Dendropanax dentiger* Roots

Li Yang,¹ Yiwei Fang,² Ronghua Liu ,¹ and Junwei He ³

¹Key Laboratory of Modern Preparation of TCM, Ministry of Education, Jiangxi University of Traditional Chinese Medicine, Nanchang 330004, China

²First Affiliated Hospital of Gannan Medical University, Ganzhou 341000, China

³Research Center of Natural Resources of Chinese Medicinal Materials and Ethnic Medicine, Jiangxi University of Traditional Chinese Medicine, Nanchang 330004, China

Correspondence should be addressed to Ronghua Liu; rhliu@163.com and Junwei He; hjwjn2008@163.com

Received 6 July 2020; Revised 4 November 2020; Accepted 10 November 2020; Published 21 November 2020

Academic Editor: Ahmet Özer Sehirli

Copyright © 2020 Li Yang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dendropanax dentiger root is a traditional medicinal plant in China and used to treat inflammatory diseases for centuries, but its phytochemical profiling and biological functions are still unknown. Thus, a rapid, efficient, and precise method based on ultra high-performance liquid chromatography coupled with quadrupole time-of-flight tandem mass spectrometry (UHPLC-Q-TOF-MS/MS) was applied to rapidly analyse the phytochemical profiling of *D. dentiger* with anti-inflammatory and antioxidant activities *in vitro*. As a result, a total of 78 chemical compositions, including 15 phenylpropanoids, 15 alkaloids, 14 flavonoids, 14 fatty acids, 7 phenols, 4 steroids, 4 cyclic peptides, 3 terpenoids, and 2 others, were identified or tentatively characterized in the roots of *D. dentiger*. Moreover, alkaloid and cyclic peptide were reported from *D. dentiger* for the first time. In addition, the ethanol crude extract of *D. dentiger* roots exhibited remarkable anti-inflammatory activity against cyclooxygenase- (COX-) 2 inhibitory and antioxidant activities *in vitro*. This study is the first to explore the phytochemical analysis and COX-2 inhibitory activity of *D. dentiger*. This study can provide important phytochemical profiles and biological functions for the application of *D. dentiger* roots as a new source of natural COX-2 inhibitors and antioxidants in pharmaceutical industry.

1. Introduction

Over the past few years, secondary metabolites from natural products play an important role in the development of new drugs [1]. Higher plants represent sources of abundant phytochemicals with a wide range of biological effects and have attracted more attention in the past decades [2–6]. Consequently, most medicinal plants belong to higher plants have been widely to treat many human diseases in traditional folk medicine [1, 7–9]. Although numerous studies on the medicinal plants used as traditional Chinese medicines (TCMs), problems of chemical compositions and biological properties remained the main barriers in the development of modern traditional medicines or new drugs.

The genus *Dendropanax* (Araliaceae), known as “Shushen” in Chinese, comprises about 80 known species

in tropical America and eastern Asia. In China, 16 native species have been found, which were widely cultivated in parks and/or used as folk medicine [10]. *D. dentiger* (Harms) Merr. is native to China and widely distributed in Guangxi, Jiangxi, Yunnan, and Guangdong provinces. In TCM, the roots of *D. dentiger* have been used as an important folk medicine for the treatment of inflammatory diseases [11]. Due to its potential pharmaceutical industry promoting effects, *D. dentiger* afforded structurally diverse and biologically active compounds, such as steroids, alkaloids, flavonoids, and monoterpenes; some of them showed potential anti-inflammatory, cytotoxic, and antioxidant activities [12, 13]. Although lots of chemical compositions report on *D. dentiger*, the full chemical profiling and COX-2 inhibitory activity of this plant have not yet been studied so far.

This study was the first time to determine the phytochemical profiling and COX-2 inhibitory activity. In addition, it was also to evaluate the antioxidant activity, including DPPH and ABTS assays *in vitro*. This finding may contribute to the processing and utility of *D. dentiger*.

2. Materials and Methods

2.1. Chemicals and Reagents. The COX-2 inhibitor screening assay kit was purchased from Beyotime Biotechnology (Shanghai, China). 2,2-Diphenyl-1-picrylhydrazyl (DPPH), 2,2'-azinobis-(3-ethylbenzthiazoline-6-sulphonate) (ABTS), and celecoxib were purchased from Sigma-Aldrich (St. Louis, MO, USA). L-Ascorbic acid (Vc) was purchased from Aladdin (Shanghai, China). Acetonitrile and formic acid (LC-MS grade) were purchased from Fisher Scientific (Pittsburgh, PA, USA). HPLC grade water was deionized using a Milli-Q ultrapure water system (Merck Millipore, Milford, MA, USA).

2.2. Plant Material. The roots of *D. dentiger* were collected in the town of Baidu, Baise City, Guangxi, China, in October 2016. A botanical voucher specimen of this plant (No. DD20161022) was deposited at authors' laboratory and was identified by one of the authors Ronghua Liu [10].

2.3. Extraction Procedure. The dried and powdered roots of *D. dentiger* (10.0 kg) were extracted with 95% EtOH (60 L × 3) and subsequently 50% EtOH (60 L × 3) by maceration at room temperature for seven days. All filtrates were combined and evaporated under reduced pressure (EYELA, Tokyo, Japan) to obtain the ethanol crude extract of *D. dentiger* (DD, 1275 g, 12.75%).

2.4. UHPLC-Q-TOF-MS/MS. The UHPLC-Q-TOF-MS/MS was provided in our previously published article [14]. Chromatographic separation was conducted on a Luna Omega C18 (100 × 2.1 mm, 1.6 μm, Phenomenex Inc., CA, USA) keeping at 40°C. 0.1% aqueous formic acid (*v/v*, A) and acetonitrile (B) were used as mobile phases. The gradient elution with the flow rate of 0.3 mL/min was performed as follows: 0-15 min, 25% B; 15-18 min, 25%-55% B; 18-40 min, 55%-95% B; 40-42 min, 95% B for column cleaning, and a conditioning cycle time of 3 min with the same initial conditions of 5% B. The sample inject volume was 3 μL.

2.5. COX-2 Inhibitory Assay. The anti-inflammatory effect of the sample against COX-2 inhibition was determined using colorimetric COX-2 inhibitor screening assay kit (no. S0168) and using celecoxib as the positive drug [2-4]. Briefly, 75 μL of assay buffer, 5 μL of cofactor working solution, and 5 μL of working solution were mixed with 5 μL of the sample at different concentrations and then incubated at 37°C. After 10 min, 5 μL of probe and 5 μL of substrate were added in all wells and then incubated at 37°C for 5 min, and the absorbance was determined (A_{sample}). The absorbance of a blank (A_{blank}) and control (A_{control}) composed of only the sample and COX-2 enzyme solutions was also determined, respectively. The COX-2 inhibitory activity (%) = $[(A_{\text{control}} - A_{\text{sample}})/(A_{\text{control}} - A_{\text{blank}})] \times 100\%$.

2.6. Antioxidant Assay

2.6.1. DPPH Radical Scavenging Activity. The DPPH radical scavenging activity of the sample was provided in our previously published articles [2-4]. Briefly, 150 μL of DPPH solution (dissolved 0.2 mM in methanol) was mixed with 50 μL of the sample at different concentrations. The mixture was stirred and incubated in the dark at 30°C for 30 min, and the absorbance was determined at 517 nm (A_{sample}). The absorbance of a blank (A_{blank}) and negative control (A_{control}) composed of only the sample and DPPH solutions was also determined, respectively. The DPPH radical scavenging activity of the sample was calculated by the following equation: DPPH scavenging activity = $[1 - (A_{\text{sample}} - A_{\text{blank}})/A_{\text{control}}] \times 100\%$. Vc was used as a positive control in this experiment.

2.6.2. ABTS Radical Scavenging Activity. The ABTS radical scavenging activity of the sample was carried out using the method reported by Sun et al. with minor modification [15]. Briefly, 1.76 mL of K₂S₂O₈ (140 mM) and 100 mL of ABTS solution (7 mM) were mixed and stored in the dark at 25°C for 12 h. Then, the ABTS stock solution was diluted with PBS (0.1 M, pH 7.4) until an absorbance value of 0.70 ± 0.02 was reached at 734 nm to obtain the diluted ABTS⁺ radical solution. Subsequently, 10 μL of the sample was mixed with 195 μL the diluted ABTS⁺ radical solution and incubated in the dark at 25°C for 106 min, and the absorbance of the sample at 734 nm (A_{sample}) was measured. The absorbance of a blank (A_{blank}) and negative control (A_{control}) composed of only the sample and diluted ABTS⁺ radical solutions was also determined, respectively. The ABTS radical scavenging activity of the sample was calculated by the following equation: ABTS scavenging activity = $[1 - (A_{\text{sample}} - A_{\text{blank}})/A_{\text{control}}] \times 100\%$. Vc was used as a positive control in this experiment.

2.7. Statistical Analysis. Graphpad Prism 6 was used for statistical analysis, and the data were presented as the means ± standard deviation (SD). One-way analysis of variance (ANOVA) and Tukey's test were used for comparison of differences in groups. Differences with $p < 0.05$ indicated statistical significance.

3. Results and Discussion

3.1. Identification of Main Constituents in *D. dentiger* Root Extract. In the present study, the phytochemical compositions were identified using UHPL-Q-TOF-MS/MS based on the existing literatures and public databases, including ChemSpider, Massbank PubChem, and mzCloud, and summarized and described in Table 1 [16-44]. The base peak chromatograms of *D. dentiger* roots extract in positive and negative ion modes were presented in Figure 1. A total of 78 compounds, including 15 phenylpropanoids, 15 alkaloids, 14 flavonoids, 14 fatty acids, 7 phenols, 4 steroids, 4 cyclic peptides, 3 terpenoids, and 2 others, were identified. The molecular formula was accurately assigned within mass error of 5 ppm. Then, the fragment ions were

TABLE 1: Compounds identified from the roots of *D. dentiger* by UHPLC-Q-TOF-MS/MS in positive/negative ion mode.

| No. | RT (min) | Ion mode | Molecular weight | Measured mass | Error (ppm) | Molecular formula | Fragments | Identification | Reference |
|------------------|----------|----------------------|------------------|---------------|-------------|---|---|---|-----------|
| Phenylpropanoids | | | | | | | | | |
| 17 | 5.03 | [M-H] ⁻ | 342.09508 | 341.08806 | 0.7 | C ₁₅ H ₁₈ O ₉ | 191.0568 ^a , 179.0351, 173.0458, 135.0467, 93.0375 | Caffeoyl hexose | 16 |
| 18 | 5.23 | [M-H] ⁻ | 354.09508 | 353.08806 | 0.7 | C ₁₆ H ₁₈ O ₉ | 353.0872, 191.0571 ^a , 179.0356, 161.0254 | Chlorogenic acid | 16 |
| 19 | 5.59 | [M-H] ⁻ | 326.10017 | 325.09321 | 1 | C ₁₅ H ₁₈ O ₈ | 176.0497, 163.0405, 145.0302, 119.0524 ^a , 114.0389, 59.0167 | Coumaric acid glucoside | 17 |
| 21 | 5.91 | [M-H] ⁻ | 180.04226 | 179.03629 | 4.3 | C ₉ H ₈ O ₄ | 135.0461 ^a , 134.0386, 89.0430 | Caffeic acid | 18 |
| 22 | 6.89 | [M-H] ⁻ | 338.10017 | 337.0932 | 0.9 | C ₁₆ H ₁₈ O ₈ | 191.0574 ^a , 173.0460, 163.0410, 119.0529, 93.0385 | 3-p-COQA | 19 |
| 26 | 7.93 | [M-H] ⁻ | 368.11073 | 367.10358 | 0.3 | C ₁₇ H ₂₀ O ₉ | 193.0511, 191.0567 ^a , 134.0389, 93.0377 | Methyl 4-caffeoylquininate | 19 |
| 27 | 8.12 | [M-H] ⁻ | 504.18429 | 503.17658 | -0.9 | C ₂₂ H ₃₂ O ₁₃ | 503.1761 ^a , 341.1243, 221.0673, 161.0474, 101.0269 | Tinoscorside D | 20 |
| 31 | 9.56 | [M-H] ⁻ | 742.26842 | 741.26178 | 0.9 | C ₃₄ H ₄₆ O ₁₈ | 417.1571 ^a , 402.1310, 181.0524, 166.0289 | Syringaresinol-4,4'-bis-O-β-D-glucopyranoside | 21 |
| 36 | 10.77 | [M-H] ⁻ | 472.19446 | 471.18683 | -0.8 | C ₂₂ H ₃₂ O ₁₁ | 189.0565, 163.0773, 134.0381, 105.0375 ^a , 89.0250, 71.0169 | Eugenol rutinoside | 22 |
| 38 | 11.00 | [M-H] ⁻ | 624.20542 | 623.19799 | -0.3 | C ₂₉ H ₃₆ O ₁₅ | 623.1974 ^a , 461.1662, 179.0354, 161.0254 ^a , 135.0459 | Verbascoside | 23 |
| 40 | 12.49 | [M-H] ⁻ | 580.21559 | 579.20843 | 0.2 | C ₂₈ H ₃₆ O ₁₃ | 417.1550, 402.1323, 387.1019, 181.0516 ^a , 166.0279, 151.0047 | Syringaresinol-4'-O-β-D-glucopyranoside | 21 |
| 42 | 13.18 | [M-H] ⁻ | 516.12678 | 515.11912 | -0.7 | C ₂₅ H ₂₄ O ₁₂ | 353.0878, 191.0566, 179.0359, 173.0466 ^a , 161.0255, 135.0467 | 3,5-di-O-Caffeoylquinic acid | 24 |
| 50 | 15.03 | [M-H] ⁻ | 500.13186 | 499.12418 | -0.8 | C ₂₅ H ₂₄ O ₁₁ | 353.0878, 337.0925, 191.0569, 179.0361, 173.0467 ^a , 135.0463 | 4-PCO-5-CQA | 19 |
| 54 | 16.76 | [M-H] ⁻ | 208.07356 | 207.06732 | 5 | C ₁₁ H ₁₂ O ₄ | 161.0258, 135.0466, 133.0313 ^a | Caffeic acid ethyl ester | 21 |
| 55 | 17.52 | [M-H] ⁻ | 524.22576 | 523.21801 | -0.9 | C ₂₆ H ₃₆ O ₁₁ | 361.1659 ^a , 346.1430, 317.1404, 231.0668, 161.0257 | Secoisolariciresinol hexose | 16 |
| Alkaloids | | | | | | | | | |
| 2 | 1.51 | [M + H] ⁺ | 135.0545 | 136.06184 | 0.5 | C ₅ H ₅ N ₅ | 136.0627, 119.0356 ^a , 107.0489, 92.0260, 91.0554 ^a , 65.0414 | Adenine | 18 |
| 3 | 1.54 | [M + H] ⁺ | 181.07389 | 182.08102 | -0.8 | C ₉ H ₁₁ NO ₃ | 119.0492, 91.0561 ^a , 77.0414, 65.0423 | Tyrosine | 25 |
| 4 | 1.59 | [M + H] ⁺ | 267.09675 | 268.10379 | -0.9 | C ₁₀ H ₁₃ N ₅ O ₄ | 136.0622 ^a , 119.0360 | Adenosine | 18 |
| 5 | 1.65 | [M + H] ⁺ | 283.09167 | 284.09882 | -0.5 | C ₁₀ H ₁₃ N ₅ O ₅ | 152.0569 ^a , 135.0308, 110.0364 | Guanosine | 25 |
| 6 | 1.71 | [M + H] ⁺ | 131.09463 | 132.10211 | 1.5 | C ₆ H ₁₃ NO ₂ | 86.0987 ^a , 72.9415, 69.0718, 57.0693, 55.9383, 55.0230 | Isoleucine | 25 |

TABLE 1: Continued.

| No. | RT (min) | Ion mode | Molecular weight | Measured mass | Error (ppm) | Molecular formula | Fragments | Identification | Reference |
|------------|----------|----------------------|------------------|---------------|-------------|--|---|---------------------------------------|-----------|
| 7 | 1.72 | [M + H] ⁺ | 275.13689 | 276.14402 | -0.5 | C ₁₂ H ₂₁ NO ₆ | 276.1431, 258.1327, 230.1383 ^a , 212.1256, 87.0330 | Glutaryl carnitine | 26 |
| 9 | 2.55 | [M + H] ⁺ | 165.07898 | 166.08619 | -0.4 | C ₉ H ₁₁ NO ₂ | 120.0811, 13.0550, 77.0412 ^a | Phenylalanine | 27 |
| 23 | 7.16 | [M + H] ⁺ | 341.16271 | 342.17031 | 1 | C ₂₀ H ₂₃ NO ₄ | 342.1718, 297.1133, 282.0899, 265.0867 ^a , 222.0899, 58.0696 | Magnoflorine | 26 |
| 24 | 7.31 | [M + H] ⁺ | 327.14706 | 328.15459 | 0.8 | C ₁₉ H ₂₁ NO ₄ | 328.1557, 178.0862 ^a , 163.0627, 151.0759 | Stepholidine | 28 |
| 37 | 10.85 | [M + H] ⁺ | 277.11028 | 278.11756 | 0 | C ₁₈ H ₁₅ NO ₂ | 278.1183, 263.0948, 220.1129, 204.0813 ^a | Dehydrooemerine | 27 |
| 44 | 13.79 | [M + H] ⁺ | 589.2887 | 590.29653 | 0.9 | C ₃₁ H ₄₃ NO ₁₀ | 590.2971 ^a , 572.2870, 558.2670, 540.2608, 508.2340 | Benzoylmesaconine | 26 |
| 48 | 14.94 | [M + H] ⁺ | 603.30435 | 604.31125 | -0.6 | C ₃₂ H ₄₅ NO ₁₀ | 604.3141 ^a , 554.2761, 242.1194 | Benzoylaconine | 26 |
| 49 | 15.00 | [M + H] ⁺ | 335.11576 | 336.1233 | 0.8 | C ₂₀ H ₁₇ NO ₄ | 336.1244, 321.1012, 320.0937 ^a , 306.0774, 292.0981, 278.0822 | Berberine | 26 |
| 52 | 15.83 | [M + H] ⁺ | 573.29378 | 574.30155 | 0.9 | C ₃₁ H ₄₃ NO ₉ | 574.3045 ^a , 542.2772, 510.2458, 105.0336 | Benzoylhypaconine | 26 |
| 57 | 17.95 | [M + H] ⁺ | 615.30435 | 616.31177 | 0.2 | C ₃₃ H ₄₅ NO ₁₀ | 616.3162 ^a , 556.2911, 524.2635, 338.1757, 161.0597 | Hypaconitine | 26 |
| Flavonoids | | | | | | | | | |
| 25 | 7.41 | [M-H] ⁻ | 594.15847 | 593.15094 | -0.4 | C ₂₇ H ₃₀ O ₁₅ | 593.1523, 503.1208, 473.1098, 383.0778, 353.0675 ^a , 297.0772 | Apigenin-6,8-di-C-glucoside | 17 |
| 29 | 8.66 | [M-H] ⁻ | 564.14791 | 563.14072 | 0.2 | C ₂₆ H ₂₈ O ₁₄ | 563.1414, 545.1304, 503.1196, 473.1092, 443.0980, 383.0781, 353.0676 ^a , 297.0786, 173.0466, 93.0369 | Schaftoside | 19 |
| 30 | 8.91 | [M-H] ⁻ | 448.10056 | 447.09339 | 0.2 | C ₂₁ H ₂₀ O ₁₁ | 357.0627, 339.0514, 327.0523 ^a , 299.0569, 297.0410, 133.0307 | Isoorientin | 19 |
| 32 | 10.07 | [M-H] ⁻ | 432.10565 | 431.09862 | 0.6 | C ₂₁ H ₂₀ O ₁₀ | 431.0991, 341.0666, 311.0567, 293.0454, 283.0616 ^a , 161.0247, 117.0365 | Vitexin | 19 |
| 33 | 10.11 | [M-H] ⁻ | 610.15339 | 609.14652 | 0.7 | C ₂₇ H ₃₀ O ₁₆ | 609.1490, 447.1153, 301.0363, 300.0284 ^a , 271.0259, 255.0310, 161.0249 | Rutin | 19 |
| 34 | 10.15 | [M + H] ⁺ | 464.09548 | 465.10317 | 0.9 | C ₂₁ H ₂₀ O ₁₂ | 303.0504 ^a , 257.0444, 201.0546, 85.0309 | Myricitrin | 29 |
| 35 | 10.34 | [M-H] ⁻ | 580.14282 | 579.13562 | 0.1 | C ₂₆ H ₂₈ O ₁₅ | 579.1365, 285.0411 ^a | Luteolin-7-O-xylosyl-glucoside | 19 |
| 39 | 11.76 | [M-H] ⁻ | 462.11621 | 461.10885 | -0.2 | C ₂₂ H ₂₂ O ₁₁ | 461.1063, 299.0560 ^a , 284.0334, 256.0375 | 5,7,2'-Trihydroxy-6-methoxyflavone | 30 |
| 45 | 14.18 | [M-H] ⁻ | 492.12678 | 491.11956 | 0.1 | C ₂₃ H ₂₄ O ₁₂ | 491.1189 ^a , 459.0923, 323.0771, 315.0732, 314.0442, 179.0361, | Tricin 5-glucoside/tricin 7-glucoside | 17 |

TABLE 1: Continued.

| No. | RT (min) | Ion mode | Molecular weight | Measured mass | Error (ppm) | Molecular formula | Fragments | Identification | Reference |
|-------------|----------|----------------------|------------------|---------------|-------------|---|--|---------------------------------|-----------|
| 47 | 14.88 | [M-H] ⁻ | 446.08491 | 445.07734 | -0.7 | C ₂₁ H ₁₈ O ₁₁ | 175.0398, 160.0178, 152.0134, 153.0208, 132.0219, 108.0286 269.0451 ^a | Baicalin | 30 |
| 53 | 16.06 | [M-H] ⁻ | 286.04774 | 285.04067 | 0.7 | C ₁₅ H ₁₀ O ₆ | 285.0414, 257.0410, 243.0281, 217.0511, 201.0205, 187.0389, 175.0420, 151.0053, 133.0311 ^a , 132.0218, 105.0355, 83.0196, 65.0104 | Luteolin | 17 |
| 56 | 17.84 | [M-H] ⁻ | 300.06339 | 299.05669 | 1.9 | C ₁₆ H ₁₂ O ₆ | 284.0313 ^a , 256.0382, 227.0343, 151.0067 | Diosmetin | 19 |
| 61 | 19.12 | [M + H] ⁺ | 402.13147 | 403.13883 | 0.2 | C ₂₁ H ₂₂ O ₈ | 403.1397, 388.1151, 373.0922 ^a , 342.1110 | Nobiletin | 31 |
| 66 | 20.54 | [M-H] ⁻ | 270.05282 | 269.04576 | 0.8 | C ₁₅ H ₁₀ O ₅ | 269.0467 ^a , 241.0515, 225.0568, 213.0563, 197.0617, 181.0660, 171.0458 | Apigenin | 32 |
| Fatty acids | | | | | | | | | |
| 16 | 4.98 | [M-H] ⁻ | 176.06847 | 175.06192 | 4.1 | C ₇ H ₁₂ O ₅ | 115.0411, 113.0637, 85.0689 ^a | Hydroxy-methylglutaric acid | 33 |
| 58 | 18.28 | [M-H] ⁻ | 228.13616 | 227.12906 | 0.8 | C ₁₂ H ₂₀ O ₄ | 183.1407 ^a , 165.1305 | Dihydroxy dodecadienoic acid | 34 |
| 59 | 18.40 | [M-H] ⁻ | 329.23349 | 329.2336 | 0.8 | C ₁₈ H ₃₄ O ₅ | 329.2329, 229.1451, 211.1348, 209.1191, 171.1038 ^a , 139.1141, 127.1144 | Trihydroxy-octadecaenoic acid | 35 |
| 62 | 19.55 | [M-H] ⁻ | 310.21441 | 309.20744 | 1 | C ₁₈ H ₃₀ O ₄ | 209.1192 ^a , 207.1408, 185.1193, 163.1139, 137.0985, 125.0991, 99.0849, 97.0682, 57.0403 | Dihydroxy-octadecatrienoic acid | 35 |
| 65 | 20.54 | [M-H] ⁻ | 314.24571 | 313.23872 | 0.9 | C ₁₈ H ₃₄ O ₄ | 313.2389, 277.2178, 201.1139 ^a , 199.0981, 171.1029, 127.1142, 125.0980 | Dihydroxy-octadecaenoic acid | 35 |
| 67 | 20.72 | [M-H] ⁻ | 312.23006 | 311.22326 | 1.5 | C ₁₈ H ₃₂ O ₄ | 311.2234, 293.2129, 275.2005, 211.1353, 201.1141, 185.1188, 171.1040 ^a , 139.1145, 127.1155 | Dihydroxy-octadecadienoic acid | 35 |
| 68 | 21.47 | [M + H] ⁺ | 352.26136 | 353.26862 | -0.1 | C ₂₁ H ₃₆ O ₄ | 261.2204, 187.1461, 145.1020, 131.0859, 107.0870, 93.0718, 81.0721 ^a , 67.0580 | Glyceryl linolenate | 36 |
| 69 | 23.75 | [M-H] ⁻ | 294.2195 | 293.21295 | 2.5 | C ₁₈ H ₃₀ O ₃ | 293.2114 ^a , 249.2220, 197.1180, 185.1186, 125.0981, 113.0987 | Hydroxy-octadecatrienoic acid | 35 |
| 71 | 28.55 | [M + H] ⁺ | 354.27701 | 355.28463 | 1 | C ₂₁ H ₃₈ O ₄ | 337.2750, 263.2377, 245.2258, 175.1478, 161.1327, 147.1159, 109.1017, 95.0862, 81.0720 ^a | Glyceryl linoleate | 36 |

TABLE 1: Continued.

| No. | RT (min) | Ion mode | Molecular weight | Measured mass | Error (ppm) | Molecular formula | Fragments | Identification | Reference |
|-----------------|----------|-------------------------|------------------|---------------|-------------|---|--|---------------------------------------|-----------|
| 72 | 29.08 | [M-H] ⁻ | 356.29266 | 355.28536 | -0.1 | C ₂₁ H ₄₀ O ₄ | 355.2758, 355.2918, 293.2837 ^a , 295.0241, 240.9931 | Monoolein | 22 |
| 74 | 30.78 | [M + H] ⁺ | 330.27701 | 331.28412 | -0.5 | C ₁₉ H ₃₈ O ₄ | 313.2738, 109.1018, 95.0863, 85.1026, 81.0716, 71.0880, 57.0740 ^a | Glycerin palmitate | 36 |
| 75 | 37.15 | [M-H] ⁻ | 284.27153 | 283.2645 | 0.9 | C ₁₈ H ₃₆ O ₂ | 283.2631 ^a , 282.3573, 265.2527, 199.8494 | Tearic acid or its isomer | 35 |
| 77 | 40.40 | [M-H] ⁻ | 312.30283 | 311.29527 | -0.9 | C ₂₀ H ₄₀ O ₂ | 311.2948 ^a , 311.1673, 293.2749, 184.0163, 183.0121 | Arachidic acid/eicosanoic acid | 37 |
| 78 | 43.36 | [M-H] ⁻ | 340.33413 | 339.32722 | 1.1 | C ₂₂ H ₄₄ O ₂ | 339.1992, 184.0199, 183.0127 ^a , 119.0516 | Behenic acid | 22 |
| Phenols | | | | | | | | | |
| 8 | 1.93 | [M-H] ⁻ | 170.02152 | 169.0152 | 4.6 | C ₇ H ₆ O ₅ | 125.0254 ^a , 124.0202, 97.0341, 79.0239, 69.0410 | Gallic acid | 38 |
| 10 | 2.80 | [M-H] ⁻ | 316.07943 | 315.07253 | 1.2 | C ₁₃ H ₁₆ O ₉ | 153.0201, 152.0127, 108.0246 ^a | 3-Carboxy-4-hydroxy-phenoxy glucoside | 19 |
| 11 | 2.94 | [M-H] ⁻ | 330.09508 | 329.08808 | 0.8 | C ₁₄ H ₁₈ O ₉ | 167.0359, 152.0128 ^a , 123.0474, 108.0251 | Vanillic acid hexose | 16 |
| 12 | 3.47 | [M + H] ⁺ | 198.05282 | 199.05998 | -0.6 | C ₉ H ₁₀ O ₅ | 181.0440, 140.0465, 125.0234 ^a , 97.0297, 77.0406 | Syringic acid | 25 |
| 13 | 3.50 | [M-H] ⁻ | 360.10565 | 359.0987 | 0.9 | C ₁₅ H ₂₀ O ₁₀ | 197.0465, 182.0234, 167.0000, 138.0344 ^a , 123.0112, 95.0176 | Methoxypolygoacetophenoxide | 33 |
| 14 | 3.82 | [M-H] ⁻ | 256.0583 | 255.05159 | 2.2 | C ₁₁ H ₁₂ O ₇ | 225.0399, 196.0402, 181.0514, 163.0405, 148.0190 ^a , 135.0469, 120.0222, 109.0389, 95.0205 | Piscidic acid | 39 |
| 28 | 8.55 | [M + H] ⁺ | 182.05791 | 183.06503 | -0.8 | C ₉ H ₁₀ O ₄ | 140.0463, 125.0236, 95.0504, 77.0414 ^a , 65.0421 | Syringaldehyde | 25 |
| Steroids | | | | | | | | | |
| 63 | 19.56 | [M + HCOO] ⁻ | 738.41904 | 783.41695 | 1 | C ₃₉ H ₆₂ O ₁₃ | 783.4257, 737.4121 ^a | 25(27)-ene-Timosaponin AIII | 35 |
| 64 | 19.91 | [M + HCOO] ⁻ | 1050.52469 | 1095.52283 | 0.9 | C ₅₀ H ₈₂ O ₂₃ | 1049.5222 ^a , 917.4771, 887.4673, 593.3684 | F-Gitonin | 35 |
| 73 | 30.23 | [M-H] ⁻ | 340.24023 | 339.23303 | 0.2 | C ₂₃ H ₃₂ O ₂ | 339.2323, 163.1143 ^a | Dimethisterone | 35 |
| 76 | 38.29 | [M + H] ⁺ | 412.37052 | 413.37809 | 0.7 | C ₂₉ H ₄₈ O | 413.3786, 395.3701, 255.2108, 213.1639, 173.1328, 159.1170 ^a , 145.1017, 109.0658 | α-Spinasterol | 40 |
| Cyclic peptides | | | | | | | | | |
| 41 | 12.63 | [M + H] ⁺ | 678.50438 | 679.51224 | 0.8 | C ₃₆ H ₆₆ N ₆ O ₆ | 679.5139 ^a , 661.5044, 566.4337, 548.4185, 486.2149, 435.3332, 322.2482, 209.1660, 114.0932 | Cyclo hexaleucyl (or isoleucyl) | 35 |

TABLE 1: Continued.

| No. | RT (min) | Ion mode | Molecular weight | Measured mass | Error (ppm) | Molecular formula | Fragments | Identification | Reference |
|------------|----------|----------------------|------------------|---------------|-------------|---|--|---------------------------------|-----------|
| 43 | 13.71 | [M + H] ⁺ | 791.58845 | 792.59618 | 0.6 | C ₄₂ H ₇₇ N ₇ O ₇ | 792.5976 ^a , 774.5873, 679.5137, 661.5069, 566.4308, 548.4199, 453.3648, 435.3336, 340.2605, 322.2510, 227.1767, 209.1654 | Cyclo hetaleucyl (or isoleucyl) | 35 |
| 46 | 14.58 | [M + H] ⁺ | 904.67251 | 905.68109 | 1.4 | C ₄₈ H ₈₈ N ₈ O ₈ | 905.6811 ^a , 887.6715, 774.5847, 548.4196, 435.3353 | Cyclo octaleucyl (or isoleucyl) | 35 |
| 51 | 15.33 | [M + H] ⁺ | 1017.75658 | 1018.76517 | 1.3 | C ₅₄ H ₉₉ N ₉ O ₉ | 1018.7684 ^a , 1000.7590, 887.6637 | Cyclo nonaleucyl (or isoleucyl) | 35 |
| Terpenoids | | | | | | | | | |
| 15 | 4.70 | [M-H] ⁻ | 376.13695 | 375.12984 | 0.5 | C ₁₆ H ₂₄ O ₁₀ | 375.1307, 213.0777 ^a , 169.0879, 151.0776, 113.0265 | Mussaenosidic acid | 18 |
| 20 | 5.79 | [M-H] ⁻ | 390.11621 | 389.10865 | -0.7 | C ₁₆ H ₂₂ O ₁₁ | 389.1098, 345.1181, 209.0459, 183.0691, 165.0554, 139.0781, 121.0672, 95.0526, 69.0405 ^a | Oleoside | 41 |
| 60 | 18.59 | [M-H] ⁻ | 822.40379 | 821.39586 | -0.8 | C ₄₂ H ₆₂ O ₁₆ | 821.3970 ^a , 351.0564 | Glycyrrhizin | 42 |
| Others | | | | | | | | | |
| 1 | 0.90 | [M-H] ⁻ | 342.11621 | 341.10951 | 1.7 | C ₁₂ H ₂₂ O ₁₁ | 179.0573, 161.0463, 119.0376, 113.0272, 89.0287, 71.0195, 59.0207 ^a | Sucrose | 43 |
| 70 | 24.37 | [M + H] ⁺ | 278.15181 | 279.1589 | -0.7 | C ₁₆ H ₂₂ O ₄ | 149.0233 ^a , 121.0287 | Dibutyl phthalate | 44 |

^aBase peak. RT: retention time; 3-p-COQA: 3-O-trans-coumaroylquinic acid; 4-PCO-5-CQA: 4-O-feruloyl-5-coumaroylquinic acid.

used to further confirm the chemical structure. Furthermore, the fragmentation pathways of some representative compounds were proposed in order to facilitate structural identification. Among them, compounds 7, 10, 11, 13-32, 34-39, 41, 43-46, 48, 50-59, 61-65, 67-69, and 71-77 were reported for the first time in the Araliaceae family. Moreover, this is the first report on compounds 1-3, 5, 6, 9, 12, 40, 42, 47, 49, 60, 66, and 78 from the genus *Dendropanax* and compounds 4, 8, 33, and 70 from *Dendropanax denticiger* [12, 13].

3.1.1. Phenylpropanoids. Phenylpropanoids were widely distributed in medicinal plants and its structures containing one or more C₆-C₃ units, which include three structure types, including simple phenylpropanoids, coumarins, and lignans [14]. A total of 15 phenylpropanoids in the roots of *D. denticiger* extract were identified in negative ion mode, including 12 simple phenylpropanoids and 3 lignans (Figure 2).

Compounds 17-19, 21, 22, 26, 27, 36, 38, 42, 50, and 54 were simple phenylpropanoids, while compounds 27 and 36 were phenylpropanol and phenylpropene, respectively. Moreover, compounds 17, 18, 21, 26, 38, 42, 50, and 54 were caffeic acid derivatives, including 4 caffeoylquinic acid derivatives (18, 26, 42, 50). They combine by the quinic acid and

caffeic acid with esteratic linkage and have similar cleavage pathways. The typical neutral losses of caffeoyl, quinine, H₂O, and CO₂ were the major cleavage pathway of such compounds. Taking compound 18 as an example, it gave the same MS² base peak at *m/z* 191.0571 due to the loss of caffeic acid and a relatively intense secondary ion at *m/z* 179.0356, while the ion at *m/z* 161.0254 was produced by continuous loss of H₂O, allowing the assignment of chlorogenic acid as reported by the reference data. The possible fragmentation mechanism was depicted in Figure S1. Besides, compound 42 also has the same fragmentation pathways.

Three compounds (31, 40, 55) belonging to the lignan, which contain two or more C₆-C₃ units. Compound 55 with a deprotonated molecule at *m/z* 523.21801 showed a base peak at *m/z* 361.1659 resulting from the loss of a hexosyl residue and was tentatively assigned as secoisolariciresinol hexose.

3.1.2. Alkaloids. In this study, a total of 15 alkaloids (Figure 3) in the roots of *D. denticiger* extract were identified, including 4 diterpenoid alkaloids (44, 48, 52, and 57), 4 isoquinoline alkaloids (23, 24, 37, and 49), 3 purine alkaloids (2, 4, and 5), 3 amino acid (3, 6, and 9), and 1 other alkaloid (7).




FIGURE 1: The base peak chromatograms of the *D. dentiger* root extract by UPLC-Q-TOF-MS/MS in negative and positive ion modes.

Compounds **44**, **48**, **52**, and **57** were diterpenoid alkaloids, which were belonging to aconitum alkaloids. In tandem mass spectrum of aconitum alkaloids commonly observe the neutral losses of H₂O (18 Da), MeOH (32 Da), CO₂ (44 Da), and PhCOOH (122 Da). Take the case of the **52**, it gave fragment ions at *m/z* 574.3045, 542.2772, 510.2458, and 105.0336 in the positive mode were corresponding to [M+H]⁺, [M+H-CH₃OH]⁺, [M+H-2CH₃OH]⁺, and [M+H-C₂₄H₃₉NO₈]⁺, respectively. Compared with literature data, compound **52** was identified as benzoylhypaconine, and the possible fragmentation mechanism was depicted in Figure S2.

Compounds **23**, **24**, **37**, and **49** were isoquinoline alkaloids, which were widely distributed in medicinal plants and have high medicinal value. Compound **23** gave fragment ions at *m/z* 342.1718, 297.1133, 282.0899, and 265.0867 in the positive mode were corresponding to [M+H]⁺, [M+H-C₂H₆N]⁺, [M+H-C₂H₆N-CH₃]⁺, and [M+H-C₂H₆N-CH₃OH]⁺, respectively, of which ring B lost a C₂H₆N by a-cleavage and formed a Cp-ring; then, the ring A lost a methoxy at C-6 and formed an epoxy between C-6 and C-7. The tandem mass pattern of this compound was similar with magnoflorine. Thus, it could be identified as magnoflorine. The ESI-MS spectra of compound **24** exhibited similar quasi-molecular ions peak [M+H]⁺ at *m/z* 328.1557; their MS² generated fragments at *m/z* 178.0862 and *m/z* 151.0759 by splitting of RDA on C-ring. Hence, compound **24** was tentatively identified as stepholidine. The [M+H]⁺ ion of compound **37** at *m/z* 278.1183 had a similar mass and fragmentation pathway to the dehydrooemerine, according to the characteristic ions at *m/z* 263.0948 [M+H-CH₃]⁺, *m/z* 220.1129 [M+H-CH₂O-CO]⁺, and *m/z* 204.0813 [M+H-CH₄-CH₂O-CO]⁺. For compound **49**, the positive mode MS spectrum showed the parent ion at *m/z* 336.1233 [M+H]⁺, and MS² spectrum showed the fragment ions at *m/z* 321.1012 [M+H-CH₃]⁺, 320.0937 [M+H-CH₄]⁺, 306.0774 [M+H-2CH₃]⁺, 292.0981 [M+H-CH₄-CO]⁺, and 278.0822 [M+H-2CH₃-CO]⁺. Compared with literature data, compound

49 was identified as berberine, and the possible fragmentation mechanism was depicted in Figure S3.

Moreover, other 7 alkaloid compounds **2**, **3**, **4**, **5**, **6**, **7**, and **9** were identified as adenine, tyrosine, adenosine, guanosine, isoleucine, glutaryl carnitine, and phenylalanine, respectively. To the best of our knowledge, alkaloid was reported from *D. dentiger* for the first time.

3.1.3. Flavonoids. The mass spectra fragmentation patterns were widely used to provide the structural characterization of flavonoids in relation to the flavonoid aglycone and flavonoids glycoside. Moreover, the identification of the flavonoid aglycone was based on fragmentations, which related to the lost small neutral molecules and radicals (CH₃, H₂O, CO, and CO₂), as well as the loss of a glucuronic acid (176 Da), hexose residue (162 Da), and apiose residue (132 Da) for flavonoids glycoside [14].

In this study, 5 flavonoids (**39**, **53**, **56**, **61**, and **66**) and 9 flavonoid glycosides (**25**, **29**, **30**, **32**, **33**, **34**, **35**, **45**, and **47**) in the roots of *D. dentiger* extract were identified based on the molecular weight and fragmentation information (Figure 4). Compounds **39**, **53**, **56**, **61**, and **66** were belonging to flavonoids, which considered as 5,7,2'-trihydroxy-6-methoxyflavone, luteolin, diosmetin, nobiletin, and apigenin, respectively. The MS² spectrum of **66** shown in Figure S4 was a representative example, which showed a [M-H]⁻ ion at *m/z* 269.0467, in accordance with the elemental composition of C₁₅H₁₀O₅⁻.

Compounds **25**, **29**, **30**, **32**, **33**, **34**, **35**, **45**, and **47** were flavonoid glycosides, which considered as apigenin-6,8-di-C-glucoside, schaftoside, isoorientin, vitexin, rutin, myricitrin, luteolin-7-O-xylosyl-glucoside, tricetin 5-glucoside, and baicalin, respectively. Compound **29** was C-glycosides, which the disaccharide substitution continuously loses 60, 90, and 120 Da fragment ions. Compound **29** showed a [M-H]⁻ ion at *m/z* 563.1414, C-6 substituted hexose broke up in ^{0,4}X₀, ^{0,3}X₀, and ^{0,2}X₀ to obtain 503.1196, 473.1092, and 443.0980 fragment ions, respectively, after that C-8 site pentose fractured at ^{0,3}X₁ and ^{0,2}X₁ to get 383.0781 and 353.0676, because




FIGURE 2: Chemical structures of phenylpropanoids from *D. dentiger* roots.




FIGURE 3: Chemical structures of alkaloids from *D. dentiger* roots.




FIGURE 4: Chemical structures of flavonoids from *D. dentiger* roots.

the C-6 substituent glycosyl groups were superior to the C-8 replacement fracture. According to the characteristics of the fragment ions, compound **29** was easily confirmed as schaftoside. However, compound **33** was O-glycosides, which typically lost the entire sugar neutral molecule with significant loss of 132, 146, 162, and 192 Da fragments. Compound **33** $[M-H]^{-609.1490}$ was rutin, which could be detected aglycone ion $[Y_0]^{-301.0363}$ and radical aglycone ion $[Y_0-H]^{-300.0284}$ after losing carbohydrate continuously in the negative ion mode, and the possible fragmentation mechanism was depicted in Figure S5.

3.1.4. Fatty Acids. In our study, a total of 14 fatty acids (peaks **16**, **58-59**, **62**, **65**, **67-69**, **71**, **72**, **74**, **75**, **77**, and **78**) were identified based on the reference mass spectra and databases.

3.1.5. Phenols. A total of 7 phenols were identified in this study. Compounds **8**, **10**, **11**, **12**, **13**, **14**, and **28** were considered as gallic acid, 3-carboxy-4-hydroxy-phenoxy glucoside,

vanillic acid hexose, syringic acid, methoxypolyacetophenone, picidic acid, and syringaldehyde, respectively.

Take compound **11** as an example, its fragment ions at m/z 167.0359, 152.0128, 123.0474, and 108.0251 were identified as vanillic acid hexose. The ion at m/z 167.0359 was obtained by the loss of hexose, while the ion at m/z 123.0474 was produced by continuous loss of CO_2 . Meanwhile, the ion at m/z 152.0128 was obtained by the loss of CH_3 from the precursor ion at m/z 167.0359, while the ion at m/z 108.0251 was produced by continuous loss of CO_2 . Based on the above fragment ions, which was obtained in the MS^2 spectrum, the structure of compound **11** was easily confirmed as vanillic acid hexose.

3.1.6. Steroids. Four steroids (peaks **63**, **64**, **73**, and **76**) were identified in this study. Peaks **63** and **64** generated $[M+HCOO]^-$ ions at m/z 783.41695 and 1095.52283 in negative mode were unequivocally determined to be 25(27)-ene-timosaponin AIII and F-gitonin by comparison with the reference data. Peak **73** produced $[M-H]^-$ ions at

m/z 339.2323 in ESI⁻ mode. By comparing the quasi-molecular ions and fragmentations with MassBank and reference mass spectra data, peak **73** was tentatively identified as dimethisterone. Compound **76** had $[M+H]^+$ ion at m/z 413.37809, and its fragments were at m/z 395.3701 $[M+H-H_2O]^+$, 255.2108 $[M+H-C_{10}H_{20}-H_2O]^+$, 213.1639 $[M+H-C_{10}H_{20}-H_2O-C_3H_4]^+$, 173.1328 $[M+H-C_{10}H_{20}-H_2O-2C_3H_4]^+$, and 159.1170 $[M+H-C_{10}H_{20}-H_2O-2C_3H_4-CH_2]^+$, and was identified as α -spinasterol.

3.1.7. Cyclic Peptides. A total of 4 cyclic peptides (peaks **41**, **43**, **46**, and **51**) were identified in this study, and the compounds **41**, **43**, **46**, and **51** showed $[M+H]^+$ ion at m/z 679.51224, 792.59618, 905.68109, and 1018.76517, respectively. They have similar fragmentation pathways [14]. This is the first time to report the cyclic peptide from *D. dentiger*.

3.1.8. Terpenoids. In current work, 3 terpenoids (peaks **15**, **20**, and **60**) were identified in negative ion mode.

Compound **15** had $[M-H]^-$ ion at m/z 375.1307, and its fragments were at m/z 213.0777 $[M-H-Glc]^-$, 169.0879 $[M-H-Glc-CO_2]^-$, and 151.0776 $[M-H-Glc-CO_2-H_2O]^-$. Its fragmentation process was the same as the literature. Therefore, compound **15** was identified as mussaenosidic acid. Compound **20** exhibited a pseudomolecular ion at m/z 389.1098 $[M-H]^-$ and fragment ions at m/z 345.1181 $[M-H-CO_2]^-$ corresponding to decarboxylation and m/z 165.0554 $[M-H-CO_2-C_6H_{12}O_6]^-$ corresponding to the cleavage of elenolic acid moieties. Meanwhile, fragment ion at m/z 69.0405 was corresponding to the propiolic acid. Compound **20** was tentatively identified as oleoside. Compound **60** showed $[M-H]^-$ ion at m/z 821.3970 and the fragment ions at m/z 351.0564 were corresponding to $[2GluA-H_2O]^-$, which the fragmentation pathways were similar with glycyrrhizin.

3.1.9. Others. Compounds **1** and **70** were given $[M+H]^+$ ions at m/z 341.10951 and 279.1589 and identified as sucrose and dibutyl phthalate, respectively, by comparing with literature.

3.2. COX-2 Inhibitory Assay. COX-2 is one of the most important proinflammatory enzyme of action for anti-inflammatory drugs, and celecoxib was a COX-2 selective inhibitor in clinical practice [2]. As observed in Table 2, the ethanol crude extract of *D. dentiger* roots showed significant COX-2 inhibitory effect with an IC_{50} value of $77.2 \pm 4.2 \mu\text{g/mL}$; however, there was an indicated remarkable difference ($p < 0.01$) in comparison with that of celecoxib with an IC_{50} value of $22.4 \pm 1.4 \text{ ng/mL}$. To the best of our knowledge, this study was the first time to determine the COX-2 inhibitory activity for *D. dentiger* [12, 13].

3.3. Antioxidant Activity. The DPPH and ABTS free radical scavenging activity assays were mostly used to evaluate the antioxidant effect of natural antioxidants [2]. Hence, the antioxidant activity of the *D. dentiger* roots ethanol crude extract was evaluated using ABTS and DPPH assays, and the results are shown in Table 2. The ethanol crude extract of *D. dentiger* roots showed the outstanding antioxidant

TABLE 2: IC_{50} values of *D. dentiger* root extract (DDR) and standards in COX-2, DPPH, and ABTS assays.

| Samples | IC_{50} ($\mu\text{g/mL}$) | | |
|-----------------------------|---------------------------------|-----------------------|----------------------|
| | COX-2 assay | DPPH assay | ABTS assay |
| DDR | $77.2 \pm 4.2^{**}$ | $255.8 \pm 10.3^{**}$ | $151.9 \pm 6.5^{**}$ |
| Celecoxib ^a | $(22.4 \pm 1.4) \times 10^{-3}$ | NT | NT |
| V _c ^a | NT | 6.0 ± 0.2 | 1.2 ± 0.1 |

^aPositive drug. NT: not tested. Data are shown as mean \pm SD ($n=3$). Differences were analyzed using ANOVA by Tukey's test. $^{**}p < 0.01$ compared with the positive drug.

activity, with IC_{50} values of $255.8 \pm 10.3 \mu\text{g/mL}$ for DPPH assay and $151.9 \pm 6.5 \mu\text{g/mL}$ for ABTS assay; however, there were exhibited significant differences ($p < 0.01$) comparable to those of the positive control V_c with IC_{50} values of 6.0 ± 0.2 and $1.2 \pm 0.1 \mu\text{g/mL}$, respectively.

To date, only one paper was reported the antioxidant activity of *D. dentiger*, and its ethyl acetate and n-butanol fractions showed significant against DPPH free radical scavenging activity [45]. Moreover, 7 phenolic compounds were isolated from the extract of *D. dentiger* and showed moderate or significant against DPPH free radical scavenging activity, with IC_{50} values of 0.038-0.741 μM , comparable to that of V_c with an IC_{50} value of 0.059 μM [45]. Therefore, this observed antioxidant activity could be due to the greater presence of secondary bioactive metabolites belonging to the flavonoids or phenolics noticed in ethanol crude extract of *D. dentiger* roots.

4. Conclusions

To summarize our findings, this study revealed that the root of *D. dentiger* was rich in phenylpropanoids, alkaloids, and flavonoids by UHPLC-Q-TOF-MS/MS and showed significant anti-inflammatory and antioxidant activities. This is the first study to describe the phytochemical profiling and COX-2 inhibitory activity of this plant [12, 13]. This study can provide important chemical information for the application of *D. dentiger* as a new source of natural COX-2 inhibitors and antioxidants in health food and pharmaceutical industry.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflict of interest.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No. 81760705), the Natural Science Foundation of Jiangxi Province (No. 20192BBHL80008), the Research Project of Jiangxi Health Department (No.

2016A038), and the Jiangxi University of Traditional Chinese Medicine (No. JXSYLXK-ZHYA0031).

Supplementary Materials

Fig. S1: tandem mass spectra and its fragmentation of chlorogenic acid in negative ion mode. Figure S2: tandem mass spectra and its fragmentation of benzoylhyaconine in positive ion mode. Figure S3: tandem mass spectra and its fragmentation of berberine in positive ion mode. Figure S4: tandem mass spectra and its fragmentation of apigenin in positive ion mode. Figure S5: tandem mass spectra and its fragmentation of rutin in positive ion mode. (*Supplementary Materials*)

References

- [1] D. D. Newman and G. M. Cragg, "Natural products as sources of new drugs over the nearly four decades from 01/1981 to 09/2019," *Journal of Natural Products*, vol. 83, no. 3, pp. 770–803, 2020.
- [2] J. W. He, L. Yang, Z. Q. Mu et al., "Anti-inflammatory and antioxidant activities of flavonoids from the flowers of *Hosta plantaginea*," *RSC Advances*, vol. 8, no. 32, pp. 18175–18179, 2018.
- [3] J. W. He, X. Y. Huang, Y. Q. Wang et al., "A new flavonol glycoside from the flowers of *Hosta plantaginea* with cyclooxygenase-1/2 inhibitory and antioxidant activities," *Natural Product Research*, vol. 33, no. 11, pp. 1599–1604, 2019.
- [4] L. Yang, S. T. Jiang, Q. G. Zhou, G. Y. Zhong, and J. W. He, "Chemical constituents from the flower of *Hosta plantaginea* with cyclooxygenase inhibition and antioxidant activities and their chemotaxonomic significance," *Molecules*, vol. 22, no. 11, p. 1825, 2017.
- [5] L. Yang, Y. M. Lin, Z. W. He et al., "Hostaflavanol A, a new anti-inflammatory and antioxidant activities flavanol from the flowers of *Hosta plantaginea*," *Medicinal Chemistry Research*, vol. 29, no. 3, pp. 426–430, 2020.
- [6] L. Yang, Y. Y. Zhu, Z. W. He et al., "Plantanone D, a new rare methyl-flavonoid from the flowers of *Hosta plantaginea* with anti-inflammatory and antioxidant activities," *Natural Product Research*, pp. 1–7, 2020.
- [7] J. W. He, L. B. Zeng, R. R. Wei et al., "*Lagopsis supina* exerts its diuretic effect via inhibition of aquaporin-1, 2 and 3 expression in a rat model of traumatic blood stasis," *Journal of Ethnopharmacology*, vol. 231, pp. 446–452, 2019.
- [8] Z. Y. Liu, L. Yang, R. X. Li et al., "Diuretic and Antidiuretic activities of ethanol extract and fractions of *Lagopsis supina* in normal rats," *BioMed Research International*, vol. 2019, Article ID 6927374, 8 pages, 2019.
- [9] L. Yang and J. W. He, "*Hosta plantaginea* (Lam.) Aschers (Yuzan): an overview on its botany, traditional use, phytochemistry, quality control and pharmacology," *RSC Advances*, vol. 9, no. 60, pp. 35050–35058, 2019.
- [10] L. Yang, S. Z. Liu, R. H. Liu, and J. W. He, "Bioassay-guided isolation of cyclooxygenase-2 inhibitory and antioxidant phenylpropanoid derivatives from the roots of *Dendropanax dentiger*," *Bioorganic Chemistry*, vol. 104, p. 104211, 2020.
- [11] State Administration of Traditional Chinese Medicine, *Zhonghua Bencao (Chinese Materia Medica) vol. 5*, Shanghai Science and Technology Press, Shanghai, China, 1999.
- [12] L. Yang, Y. Q. Wang, S. C. Liu, and J. W. He, "Research progress of chemical constituents and pharmacological activities from three commonly used Ban-feng-he medicinal plants," *Chinese Journal of Experimental Traditional Medical Formulae*, vol. 22, no. 22, pp. 191–196, 2016.
- [13] X. M. Wang, L. Yang, J. W. He, G. Ren, M. Yang, and G. Y. Zhong, "Research progress of chemical constituents and pharmacological activities of genus *Dendropanax*," *Chinese Journal of Experimental Traditional Medical Formulae*, vol. 21, no. 24, pp. 229–234, 2015.
- [14] L. Yang, R. H. Liu, and J. W. He, "Rapid analysis of the chemical compositions in *Semiliquidambar cathayensis* roots by ultra high-performance liquid chromatography and quadrupole time-of-flight tandem mass spectrometry," *Molecules*, vol. 24, no. 22, p. 4098, 2019.
- [15] Y. J. Sun, J. Rukeya, W. T. Tao, P. L. Sun, and X. Q. Ye, "Bioactive compounds and antioxidant activity of wolfberry infusion," *Scientific Reports*, vol. 7, no. 1, 2017.
- [16] C. Jiménez-Sánchez, J. Lozano-Sánchez, C. Rodríguez-Pérez, A. Segura-Carretero, and A. Fernández-Gutiérrez, "Comprehensive, untargeted, and qualitative RP-HPLC-ESI-QTOF/MS² metabolite profiling of green asparagus (*Asparagus officinalis*)," *Journal of Food Composition and Analysis*, vol. 46, pp. 78–87, 2016.
- [17] T. T. Xing, *Study on chemical compositions and metabonomics of the peels of mandarins and oranges by UPLC-QTOF-MS*, Master degree of Southwest University, Chongqing, China, 2019.
- [18] F. Zhou, M. S. Lin, R. Li et al., "Analysis of chemical constituents in baihe dihuangtang by UPLC-Q-TOF-MS," *Chinese Journal of Experimental Traditional Medical Formulae*, vol. 26, no. 7, pp. 15–22, 2020.
- [19] M. Liu, M. Z. He, H. W. Gao et al., "Strategy for rapid screening of antioxidant and anti-inflammatory active ingredients in *Gynura procumbens* (Lour.) Merr. based on UHPLC-Q-TOF-MS/MS and characteristic ion filtration," *Biomedical Chromatography*, vol. 33, no. 11, p. e4635, 2019.
- [20] P. van Kiem, C. van Minh, N. T. Dat et al., "Aporphine alkaloids, clerodane diterpenes, and other constituents from *Tinospora cordifolia*," *Fitoterapia*, vol. 81, no. 6, pp. 485–489, 2010.
- [21] X. M. Luo, M. F. Su, X. Y. Chang et al., "Qualitative and quantitative analysis of main chemical constituents in *Eucommia ulmoides* by LC-MS," *Modern Chinese Medicine*, vol. 21, no. 8, pp. 1029–1040, 2019.
- [22] L. P. Zheng, Z. G. He, Z. J. Wu, and C. A. Zhang, "Chemical constituents from *Dendropanax dentiger*," *Chemistry of Natural Compounds*, vol. 8, pp. 883–885, 2012.
- [23] X. X. Gao, N. Wang, J. P. Jia, P. Y. Wang, A. R. Zhang, and X. M. Qin, "Chemical profiling of Dingkun Dan by ultra high performance liquid chromatography Q exactive orbitrap high resolution mass spectrometry," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 177, p. 112732, 2020.
- [24] X. H. Yang, X. L. Cheng, B. Qin et al., "Ultra-high performance liquid chromatography coupled with quadrupole/time of flight mass spectrometry based chemical profiling approach for the holistic quality control of complex Kang-Jing formula preparations," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 124, pp. 319–336, 2016.
- [25] M. H. Liu, X. Tong, J. X. Wang, W. Zou, H. Cao, and W. W. Su, "Rapid separation and identification of multiple constituents in traditional Chinese medicine formula Shenqi Fuzheng Injection by ultra-fast liquid chromatography combined with

- quadrupole-time-of-flight mass spectrometry,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 74, pp. 141–155, 2013.
- [26] D. Gao, B. J. Wang, Z. P. Huo et al., “Analysis of chemical constituents in an herbal formula Jitong Ning Tablet,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 140, pp. 301–312, 2017.
- [27] S. Duan, W. Qi, S. Zhang, K. Huang, and D. Yuan, “Ultra high performance liquid chromatography coupled with electrospray ionization/quadrupole time-of-flight mass spectrometry for the rapid analysis of constituents in the traditional Chinese medicine formula Wu Ji Bai Feng Pill,” *Journal of Separation Science*, vol. 40, no. 20, pp. 3977–3986, 2017.
- [28] J. Xiao, N. N. Song, T. Lu et al., “Rapid characterization of TCM Qianjinteng by UPLC-QTOF-MS and its application in the evaluation of three species of *Stephania*,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 156, pp. 284–296, 2018.
- [29] M. M. Yan, M. Chen, F. Zhou et al., “Separation and analysis of flavonoid chemical constituents in flowers of *Juglans regia* L. by ultra-high-performance liquid chromatography-hybrid quadrupole time-of-flight mass spectrometry,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 164, pp. 734–741, 2019.
- [30] X. Qiao, R. Li, W. Song et al., “A targeted strategy to analyze untargeted mass spectral data: rapid chemical profiling of *Scutellaria baicalensis* using ultra-high performance liquid chromatography coupled with hybrid quadrupole orbitrap mass spectrometry and key ion filtering,” *Journal of Chromatography A*, vol. 1441, pp. 83–95, 2016.
- [31] L. G. Faqueti, L. P. Sandjo, and M. W. Biavatti, “Simultaneous identification and quantification of polymethoxyflavones, coumarin and phenolic acids in *Ageratum conyzoides* by UPLC-ESI-QToF-MS and UPLC-PDA,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 145, pp. 621–628, 2017.
- [32] J. M. Li, M. Z. He, H. Ouyang et al., “Rapid identification of chemical constituents of *Folium hibisci Mutabilis* by UHPLC-Q-TOF-MS/MS,” *Chinese Pharmaceutical Journal*, vol. 51, no. 14, pp. 1162–1168, 2016.
- [33] L. L. Wang, M. M. Sang, E. W. Liu et al., “Rapid profiling and pharmacokinetic studies of major compounds in crude extract from *Polygonum multiflorum* by UHPLC-Q-TOF-MS and UPLC-MS/MS,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 140, pp. 45–61, 2017.
- [34] S. T. Sakna, A. Mocan, H. N. Sultani, N. M. El-fiky, L. A. Wessjohann, and M. A. Farag, “Metabolites profiling of *Ziziphus* leaf taxa via UHPLC/PDA/ESI-MS in relation to their biological activities,” *Food Chemistry*, vol. 293, pp. 233–246, 2019.
- [35] D. Ji, Z. Y. Huang, C. H. Fei, W. W. Xue, and T. L. Lu, “Comprehensive profiling and characterization of chemical constituents of rhizome of *Anemarrhena asphodeloides* Bge,” *Journal of Chromatography B*, vol. 1060, pp. 355–366, 2017.
- [36] X. Y. Zhai, L. Zhang, B. T. Li, G. L. Feng, S. L. Xu, and C. J. Yang, “Chemical components in *Pinelliae Rhizoma* by UPLC-Q-TOF-MS/MS,” *Chinese Journal of Experimental Traditional Medical Formulae*, vol. 25, no. 7, pp. 173–183, 2019.
- [37] X. C. Liu, X. Y. Fan, X. Wang, R. N. Liu, C. F. Meng, and C. Y. Wang, “Structural characterization and screening of chemical markers of flavonoids in *Lysimachiae Herba* and *Desmodii Styracifolii* Herba by ultra high-performance liquid chromatography quadrupole time-of-flight tandem mass spectrometry based metabolomics approach,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 171, pp. 52–64, 2019.
- [38] C. M. Zhang, Y. He, T. T. Li et al., “Analysis of chemical components in herbal formula Qi Bai Granule by UPLC-ESI-Q-TOF-MS,” *Natural Product Research*, vol. 33, no. 15, pp. 2271–2275, 2019.
- [39] V. Miranda, C. D. Maycock, and M. R. Ventura, “A Stereoselective Synthesis of (+)-Piscidic Acid and Cimicifugic Acid L,” *European Journal of Organic Chemistry*, vol. 2015, no. 34, pp. 7529–7533, 2015.
- [40] Y. J. Chen, W. Hao, Z. Y. Wei et al., “Identification of chemical constituents in *Aster tataricus* by UHPLC-Q-TOF-MS,” *Acta Pharmaceutica Sinica*, vol. 54, no. 9, pp. 1645–1654, 2019.
- [41] A. K. Kiss, B. Michalak, A. Patyra, and M. Majdan, “UHPLC-DAD-ESI-MS/MS and HPTLC profiling of ash leaf samples from different commercial and natural sources and their in vitro effects on mediators of inflammation,” *Phytochemistry Analysis*, vol. 31, no. 1, pp. 57–67, 2019.
- [42] S. Z. Li, S. Liu, Z. F. Pi, F. R. Song, Y. R. Jin, and Z. Q. Liu, “Chemical profiling of Fufang-Xialian-Capsule by UHPLC-Q-TOF-MS and its antioxidant activity evaluated by in vitro method,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 138, pp. 289–301, 2017.
- [43] T. H. Wang, J. Zhang, X. H. Qiu, J. Q. Bai, Y. H. Gao, and W. Xu, “Application of ultra-high-performance liquid chromatography coupled with LTQ-Orbitrap mass spectrometry for the qualitative and quantitative analysis of *Polygonum multiflorum* Thumb. and Its Processed Products,” *Molecules*, vol. 21, no. 1, p. 40, 2016.
- [44] H. Yan, Y. M. Zuo, E. Yuan, Y. F. Wang, and Z. L. Zhang, “Analysis of chemical constituents from *Gardeniae Flos* by UHPLC-Q-TOF-MS,” *Journal of Chinese Medicinal Materials*, vol. 41, no. 6, pp. 1359–1364, 2018.
- [45] L. P. Zheng and Z. G. He, “Antioxidant activity of phenolic compounds from *Dendropanax dentiger* (Harms.) Merr,” *Asian Journal of Chemistry*, vol. 25, no. 14, pp. 7809–7812, 2013.