



Review on Analytical and Biological applications of Hydrazones and their Metal Complexes

LAKSHMI NARAYANA SUVARAPU^{1*}, YOUNG KYO SEO¹, SUNG OK BAEK¹,
VARADA REDDY AMMIREDDY²

¹Department of Environmental Engineering, Yeungnam University, Gyeongsan-si Republic of Korea -712 749.

²Analytical Division, Department of Chemistry, S.V. University, Tirupati, India – 517 502.

lakshminarayana_chem@rediffmail.com

Received 03 October 2011; Accepted 31 December 2011

Abstract: Hydrazones are very important group of analytical reagents for the determination of various metal ions by using various analytical techniques. Besides this use of hydrazones are also having biological activities also. In this paper we first discussed about the chemical nature of hydrazones and their biological activities. We mainly focused on the papers which were published during 1980-2011 on analytical applications (spectrophotometric and spectrofluorimetric) of hydrazones. We gave the total established conditions for the determination of various metal ions with hydrazones.

Keywords: Hydrazones, metal ions, Spectrophotometry, Spectrofluorimetry, Biological applications.

Introduction

Hydrazones as Analytical Reagents

Many organic compounds react with metal ions and form colored precipitates or solutions. Hence, they are extensively used as analytical reagents, even though it is difficult to predict with certainty which organic compound is suitable for the analysis of a particular metal ion. Yoe¹ gave a list of more than twenty ways in which they are used. It has been observed that the reactivity of organic reagents with metal ions in the use of the former as analytical reagents requires the presence of certain acidic or basic groupings² and coordinating atoms.

The aim of research in this field is the discovery of compounds possessing a high degree of selectivity and identification of the causes underlying such selectivity. While most of the reagents are not selective, various means are known where by the selectivity of a

reagent may be improved. These include adjustment of the pH, and the use of masking agents which form complexes with the interfering elements in the determination of the test ion. Within the organic reagent molecule, there is generally a single acidic or basic group, or a combination of these two, which is the key to the reactivity of the reagent.

Literature survey has revealed that organic compounds capable of forming chelates or inner complex salts give better results than those containing only acidic or basic groupings, in the field of inorganic analysis. The element in the organic molecule through which the metal is bonded is generally oxygen or nitrogen, less usually it is sulphur. The oxygen containing groups most often met in organic reagents were -OH, -CHO, -COOH, >CO. The nitrogen containing groups (-NH₂, =NH, heterocyclic N) met with in general functional groups are amines (usually aliphatic), heterocyclic rings (usually pyridine), oximes (in which bonding tends to be coordinated to the nitrogen instead of replacement of hydrogen) and azo groupings. The aromatic orthohydroxy carbonyl compounds form stable six membered rings with the metal ions. Hydroxy carbonyl compounds derived from benzene and naphthalene, hydroxy quinones of naphthalene and anthracene series have been introduced as analytical reagents.

It is observed, that in many cases the formation of a precipitate or a soluble colored product is dependent on the presence of definite atomic groupings. Such compounds are therefore designated as metal binding groups with specific or selective action. But, it cannot be over looked that the reaction conditions play a definite role in this direction. Hence, proper choice of solvent and other factors are to be given equal importance. Organic compounds can undergo extensive alteration in their structure as a result of condensation and substitution reactions. These may bring in the useful changes in the reagent to make them better organic reagents. A survey of literature³ shows, that organic compounds containing a phenolic or enolic group and a coordinating group containing nitrogen, oxygen or sulphur forms a variety of complexes with different metal ions.

It is found that -SH group of an organic compound exhibits a higher acidic character than similarly bound -OH group. Thio-keto group (>C=S) also plays an important role compared to its counterpart, keto group (>C=O)⁴.

It is observed that aromatic compounds containing nitroso (-N=O) as well as phenolic -OH groups are also useful as analytical reagents. It is clear from the above brief review presented, that many organic compounds containing acidic or basic groups, besides the coordinating groups form chelates easily and have been used extensively as analytical reagents. However, these investigations reveal that sensitivity and selectivity of the reagent should be established, even though a few general guidelines are available to predict the potentialities of a reagent for the said purpose. In view of large and varied demand for the new methods to determine the metal ions, under specific conditions, the search for new reagents is a continuous process. This exercise of finding new and novel reagents as well as methods for inorganic analysis has a special significance in these days in view of the alarming and complex problem of environmental pollution.

Isonicotinoylhydrazones of carbonyl compounds act as good analytical reagents, but they have not been fully exploited. Hence, in the present investigation a detailed study of these reagents has been made with a view to find out their potentialities in inorganic analysis. Hydrazones are usually named after the carbonyl compounds from which they are obtained. Isonicotinoyl hydrazones are the condensation products of isonicotinic acid hydrazide and the carbonyl compounds. These isonicotinoyl hydrazones are prepared by refluxing a mixture of isonicotinic acid hydrazide and the desired carbonyl compound for 2-3 h in slightly alkaline medium. The compound usually crystallizes out on cooling.

Many of physiologically active hydrazones find application⁵ in the treatment of diseases like tuberculosis, leprosy and mental disorders. Hydrazones also act as herbicides,

insecticides, nematocides, rodenticides and plant growth regulators. Isonicotinic acid hydrazide (INH) is an important antitubercular agent and has potential sites for formation of complexes with metal ions. It is also observed, that isonicotinoyl hydrazones and their metal complexes possess higher activity and lower resistivity to tuberculosis bacteria. These reagents, apart from those specified above are also potential analytical reagents for the determination of several metal ions by different physico-chemical techniques, of which the spectrophotometric determination occupies a special place.

The analytical applications of hydrazones have reviewed by Singh⁶ and Katyal⁷. The latest review was published in this area on 1982. In this, the author reviewed the papers published on analytical potentialities of hydrazones up to 1980. After 1980 so many researchers have worked on the analytical potentialities of hydrazones. Hydrazones have both analytical and biological applications, which attract so many researchers. Because of so much of work published in this area, we are interested to review the papers published on analytical applications of hydrazones from 1980 to 2011. We have so much confidence that this three decades survey is very useful to the scientists who are working in this area.

Biological Applications of Hydrazones and Their Metal Complexes

Interest in the study of hydrazones has been growing because of their antimicrobial, antituberculosis and antitumor activities^{8,9}. Hydrazones play an important role in inorganic chemistry, as they easily form stable complexes with most of the transition metal ions. The development of the field of bioinorganic chemistry has increased the interest in hydrazone complexes, since it has been recognized that many of these complexes may serve as models for biologically important species¹⁰. Coordination compounds derived from arylhydrazones have been reported to act as enzyme inhibitors and are useful due to their pharmacological applications^{11,12}. Hydrazones possessing an azomethine -NHN=CH- proton constitute an important class of compounds for new drug development. Therefore, many researchers have synthesized these compounds as target structures and evaluated their biological activities. Hydrazones and their metal complexes are biologically very active compounds. For example Ragavendran et al¹³ have reported anticonvulsant activity of 4-aminobutyric acid hydrazide, Abdel-Aal et al¹⁴ have reported antiviral activity of N-arylaminoacetyl hydrazones against *Herpes simplex virus-1* and *Hepatitis-A virus (HAV)*, Walcourt et al¹⁵ have reported antimalarial activity of 2-hydroxy-1-naphth-aldehyde isonicotinoyl hydrazone and Savini et al¹⁶ have reported antitumor activity of 3- and 5-methyl -thiophene-2-carboxaldehyde α -(N)-heterocyclic hydrazone derivatives.

Some important hydrazones and their complexes with antibacterial activity reported earlier are discussed here under.

Singh et al¹⁷ have reported the antibacterial activity of cobalt(II), nickel(II), zinc(II), copper(II) and cadmium(II) complexes of acetophenone-4-aminobenzoylhydrazone and 4-hydroxyacetophenone-4-aminobenzoylhydrazone. The authors have evaluated the antibacterial activity of these complexes against *Escherichia coli* and *Aspergillus niger*. They reported that at each concentration, copper(II) complexes are more active than zinc(II) complexes with both the hydrazones. The antifungal activity of the metal complexes is less than their parent ligands. Nora H. Al-Sha'alan¹⁸ has reported the antibacterial activity of 7-chloro-4-(benzylidenehydrazo)quinoline and its complexes with copper(II), nickel(II), cobalt(II), manganese(II) and iron(III). The author has evaluated the antibacterial activity against gram positive bacteria (*Staphylococcus aureus*), gram negative bacteria (*Escherichia coli*) and antifungal activity against *Candida albicans*. The ligand is also proved to be biologically active. The author reported that chelation tends to make ligand to act as more powerful and potent bactericidal agent.

Kucukguzel et al¹⁹ have reported antibacterial activity of 2,3,4-pent aneotriene-3[4-[(5-nitro-2-furyl)methylenehydrazide]carbonyl]phenyl]-hydrazone against *Staphylococcus aureus* and *Mycobacterium tuberculosis* at a concentration of $3.13 \mu\text{g mL}^{-1}$. Gulerman et al²⁰ have reported the in vivo metabolism of 4-fluorobenzoic acid (5-nitro-2-furyl)methylene hydrazide which is effective against *Staphylococcus aureus*.

Sah and Peoples²¹ synthesized hydrazones by reacting isonicotinic acid hydrazide (INH) with various aldehydes and ketones. These compounds are reported to have inhibitory activity in mice infected by various strains of *Mycobacterium tuberculosis*. Ersan S et al²² have reported antimicrobial activity of N-[(α -methyl benzilidene)-(3-substituted 1,2,4-triazol-5-yl-thio) acetohydrazides. The authors synthesized eighteen new hydrazones by reacting ortho- or para- substituted acetophenones with (3-substituted-1,2,4-triazol-5-ylthio)acetohydrazide in ethanol.

The synthesized hydrazones are tested for antimicrobial activity. These compounds exhibit only poor activity against gram positive bacteria (*Staphylococcus aureus* and *Enterococcus faecalis*) and gram negative bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*). Moderate activity is observed against fungi *Candida albicans*, *Candida parapsilosis* and *Candida krusei*. Ulusoy et al²³ have reported the synthesis and antimicrobial activity of N²-substituted alkylidene /arylidene-6-phenylimidazothiazole-3-acetic acid hydrazides. These compounds show antibacterial activity against *Staphylococcus aureus* and *Staphylococcus epidermidis*. (MIC is $25.0 \mu\text{g mL}^{-1}$). Kucukguzel et al²⁴ have reported the synthesis and antibacterial activity of ethyl-2-arylhydrazones-3-oxobutyrate. These compounds show significant activity against *Staphylococcus aureus*.

Rodriguez-Arguelles et al²⁵ have reported the antibacterial activities of cobalt(II), copper(II), nickel(II) and zinc(II) complexes of 2-thiophenecarbonyl hydrazone and isonicotinoylhydrazones of 3-(N-methyl)isatin. 2-thiophene carbonyl hydrazone metal complexes exhibit a strong inhibition on the growth of *Haemophilus influenzae* (MIC is $0.15\text{--}1.50 \mu\text{g mL}^{-1}$) and good antibacterial activity towards *Bacillus subtilis* (MIC is $3.0\text{--}25.0 \mu\text{g mL}^{-1}$). The authors have reported that the antimicrobial activity of thiophene carbonyl hydrazone derivatives is greater than that of the isonicotinoyl hydrazone derivatives. Loncle et al²⁶ have reported the antifungal activity of tosylhydrazone-cholesterol derivatives against *Candida albicans* at a concentration of $1.0\text{--}5.0 \mu\text{g mL}^{-1}$. Recently critical reviews have been published by various authors on antibacterial activity of hydrazones^{27,28}.

Analytical Applications of Hydrazones

Spectrophotometric Applications

Katyer et al²⁹ employed Orthohydroxybenzaldehyde isonicotinoylhydrazone for its chelatometric properties with several metal ions. It precipitates nickel(II), zinc(II), palladium(II), cerium(II) and copper(II) and also forms soluble complexes with lead(II), iron(II), cobalt(II), tin(II), vanadium(II), stibium(III), aluminium(III), iron(III), zirconium(IV) and thorium(IV).

Salicylaldehyde isonicotinoylhydrazone³⁰ is used for the spectrophotometric determination of gallium(III) and indium(III). Dimethylaminobenzaldehyde isonicotinoylhydrazone³¹ was employed for its selective detection of mercury(I) and mercury(II). All the reagents which were used for the spectrophotometric determination of metal ions and the established conditions like λ_{max} , pH, validity of Beer's law, molar absorptivity and the composition are presented in Table 1³²⁻¹⁰³. In this table we described about the reagents which were used in the period of 1980 - 2011. Kavrentis employed Phthalaldehyde bisguanyldihydrazone¹⁰⁴ for the determination of Co(II), Cu(II) and Ni(II), 4-Dimethylaminobenzaldehyde isonicotinoylhydrazone¹⁰⁵ for Arsenic(III) and Antimony(III), Salicylaldehyde isonicotinoylhydrazone¹⁰⁶ for Molybdenum(VI). Hydrazones were

extensively used for the spectrophotometric determination of metal ions during this period. We gave the total description about all the papers published during this period about the spectrophotometric determination of metal ions by using hydrazones as analytical reagents.

Table 1. Analytical applications of Hydrazones in the determination of metal ions by using Spectrophotometer.

Reagent	Metal ion	λ_{max} (nm)	pH	Beer's law ppm	$\epsilon \times 10^4$	M:L	Solvent	Ref
2-Aceto-1-naphthal-N-salicylhydrazone	Mn(II)	-	-	-	-	-	-	32
α -(Benzimidazolyl)- α' , α'' -(N-5- nitro-2-pyridylhydrazone)-toulene	Cu(II)	410	6.0	0-2.50	3.81	1:2	-	33
2-Benzimidazolyl-3-sulphophenyl methanone-5-nitro-2- pyridyl-hydrazone	Co(II)	517	2.7-9.4	0.02-1.00	6.65	1:2	-	34
Benzyl-2-pyridylketone-2-quinolyl hydrazone	Hg(II)	475	9.0-10.4	-	5.01	1:2	-	35
Biacetyl mono (2-pyridyl) hydrazone	Zn(II)	440	10.0	0.04-1.20	5.20	-	-	36
2,2'- Bipyridyl -2-pyridylhydrazone	Co(II)						Iso amyl alcohol	37
	Pd(II)	585	0.5-1.5H Cl	0.5-1.5	1.90	1:1	chloroform	38
2,2'- Dipyridyl-2-quinolylhydrazone	Pd(II)	-	-	-	-	-	-	39
Bis (isonicotinoylhydrazone)	Zr(IV)	410	2.8	-	-	-	-	40
N-cyanoacylacetaldehydehydrazone	Mo(VI)	790	Phosphoric acid	-	-	1:1	-	41
1,2-Cyclohexanedione bis(benzoyl) hydrazone	Ti(IV)	477	1.75 - 3.00	-	1.05	1:2	-	42
1,2-Cyclohexanedione-2-oxime-1-guanylhhydrazone	Cu(II)	-	-	-	0.52	-	-	43

Diacetylmonoxime- p-hydroxybenzoyl hydrazone	Sn(II)	430	-	0.25- 2.76	3.2	1:2	-	44
2,6- Diacetylpyridine bis (furoylhydrazone)	U(VI)	400	3.0	0.72- 10.80	1.5	-	-	45
2,6- Deacetylpyridine bis(benzoyl hydrazone)	U(VI)	420	-	-	-	-	Dichloro methane	46
2,6- Diacetylpyridine bis (benzoylhydrazone)	V(V)	335	2.6- 4.0	-	2.74	1:1	-	47
2,4- Dihydroxyacetophen one benzoylhydrazone	Mo(VI),	440	3-6	0.5-8.0	0.9	1:2	-	48
	V(V)	410	3-6	0.22- 4.9	0.77	1:1	-	
3,4- Dihydroxybenzaldehy de guanylhyazone	Mo(VI)	550	5.0	0.3-9	0.88	-	-	
	Fe(III)	355	3-6	0.03- 1.1	4.13	-	-	49
	Co(II)	355	-	0.5- 16.00	0.29	-	-	
2,4- Dihydroxybenzaldehy de isonicotinoylhydrazo ne	Ti(IV)	430	1.5	0.09- 2.15	1.35	1:2	-	50
3,4- Dihydroxybenzaldehy de isonicotinoylhydrazo ne	V(V)	360	5.5	0.5-5.3	1.29	1:1	-	51
	Cr(VI)	400	5.5	0.5-7.7	1.35	1:1	-	52
	Pd(II)	380	3.0	0.5- 20.0	0.53	1:1	-	53
2,4- Dihydroxybenzaldehy de isonicotinoylhydrazo ne	Th(IV)	390	2.0- 8.0	0.30- 7.00	2.20	1:1	-	54
	Zn(II)	390	4.0- 10.0	0.10- 1.50	3.55	1:1	-	55
2,4- Dihydroxybenzophe	Ce(IV)	400	10.0	0.7-7.0	2.0	1:1	-	56

none								
benzoichydrazone								
Diphenylglyoxal	Ca(II)	432	7.80	-	1.76	1:1		
bis(2-hydroxybenzoylhydrazone)							-	57
	Sn(II)	455	3.5-7.0		2.25	-	MIBK	58
	Ti(IV)	500	0.1H ₂ SO ₄	-	1.5	1:3	Benzyl al	59
2,2'-Dipyridyl-2-benzothiazolylhydrazone	Fe(II)	427	4.5-8.4	-	3.41	1:3	-	60
		615	3.0-9.6		1.23	-		
Di-2-pyridylketone benzoylhydrazone	Pd(II)	455	-	0.00-10.00	0.94	-	Benzen e	61
	Fe(II)	379	4.7-6.0	0.00-30.00	1.59	-	Benzen e	62
	Ni(II)	-	-	-	-	-	-	63
Di-2-pyridylketone benzylhydrazone	Fe(II), Fe(III)	-	-	-	-	-	-	64
	Co(II)	-	-	-	-	-	-	65
	Ni(II)	-	-	-	-	-	-	66
Di-2-pyridylketone-2-furan carbothiohydrazone	Re(VII)	546	-	0.70-14.00	1.64	-	MIBK	67
2,2'-Dipyridylketone hydrazone	Cu(II), Hg(II)	-	-	-	-	-	-	68
Di-2-pyridylketone isonicotinoylhydrazone	Fe(II)	-	-	-	-	-	-	69
Di-2-pyridylketone-2-pyridylhydrazone	Pd(II)	-	-	-	-	-	-	70
	V(II)	-	-	-	-	-	-	71
Di-2-pyridylketone salicyloylhydrazone	Fe(II), Fe(III)	-	-	-	-	-	-	72
Di-2-pyridylketone thiophenylhydrazone	U(VI)	-	-	-	-	-	-	73
Di-2-pyridylmethanone-2-(5-nitro)pyridylhydrazone	Pd(II)	560	-	-	3.78	-	1,2-dichloro ethane	74
	Fe(II)	505	2.0-7.5	0.00-0.84	5.83	1:2	-	75
2,2'-Dipyridyl-2-pyridylhydrazone	Fe(II), Fe(III)	535	-	0-30 mg/L	-	-	-	76
				0-50				

				mg/L				
	Pd(II)	560	-	Upto 12 mg/L	-	-	Chloroform	77
2-Hydroxybenzaldehyde-5-nitropyridylhydrazone	Co(II)	470	-	0.02-1.50 mg/L	6.5	-	-	78
2-Hydroxy-1-naphthaldehyde benzoylhydrazone	V(V), Fe(III)	465 540	5.0	0.12-2.50 0.14-4.20	-	-	-	79
Isoniazid-p-diethylaminosalicylaldehyde hydrazone	Fe(III)	-	-	-	-	-	-	80
5-Nitro-2-pyridinecarbaldehyde-5-nitro-2-pyridylhydrazone	Ni(II)	561	5.5-10	-	13.5	1:2	benzene	81
N,N'-oxalyl-bis(salicylaldehyde) hydrazone	Cu(II)	422	2.0	0.4-1.8	0.92	1:1	-	82
Orthohydroxypropio phenone isonicotinoylhydrazone	U(VI)	380	3.0	0.47-17	1.15	1:1	-	83
p-Methylisonitrosoacetophenonehydrazone	Co(II)	520	7.5	0.2-6	1.83	-	Chloroform	84
p-Methylisonitrosoacetophenonehydrazone	Cu(II)	510	7.0	0.1-1.0	0.63	-	Chloroform	85
Phenyl-2-picolylketone-2-pyridyl hydrazone	Pd(II)	-	-	-	-	-	-	86
Pyridine-2-acetaldehydesalicylhydrazone	Pd(II)	425	2-4.5		1.3	1:2	Chloroform	87
Pyridine-2-acetaldehydesalicylhydrazone	Pb(II)	380	8.6-9.3	1.5-6.2	1.93	1:2	Chloroform	88
2-Pyridinecarbaldehyde-3,5-dinitro-2-pyridylhydrazone	Ni(II)	-	-	-	-	-	-	89
Di-2-pyridylmethanone-1-phthalazinyl-	Ni(II)	500	4.3-10.7	0.1-1.0	5.40	1:2	Benzene	90

hydrazone								
Pyridine-2-carboxaldehyde-2-hydroxybenzoylhydrazone	U(VI)	375	3.5-4.6	1.0-5.6	4.74	1:2	MIBK	91
Pyridine-2-carboxaldehydeisonicotinoyl hydrazone	Fe(II)	-	-	-	-	-	-	92
Pyridoxalsalicylalhydrazone	Ti(IV)	440	0.9-2.5		0.55	1:1		93
2-Pyridylketonebenzoylhydrazone	Fe(II), Fe(III)	-	-	-	-	1:1	-	94
1-(2-Pyridylmethylidene-5-salicylidene)thiocarbohydrazone	Pd(II)	505	8.3	-	1.65	-	-	95
2-Pyridyl-3'-sulphophenylmethanone-2-pyrimidylhydrazone	Fe(II)	379 580	7.3-10 -	- -	4.79 1.18	1:1 -	- -	96
Pyrvaldehyde-2-benzothiazolylhydrazone	Cd(II)	-	-	-	-	-	-	97
Salicylaldehydecarbohydrazone	Zn(II)	445	5.4	-	-	1:1	-	98
Salicylaldehydeguanylhydrazone	Mn(II)	505	-	8-80µg/l	-	-	-	99
2-(3-Sulfobenzoyl)pyridinebenzoylhydrazone	Fe(II)	-	7-11	0-4	-	1:2	-	100
Thiazole-2-carbaldehyde-2-quinolylhydrazone	Pd(II)	625	acidic	0-4.2	1.93	1:1	benzene	101
2-Thiophenaldehyde-2-pyridylhydrazone	Cu(II)	435	8.5	0.017-1.2	-	1:3	-	102
2,3,4-Trihydroxyacetophenonephenylhydrazone	Th(IV)	370	3.5	-	32.4 6	1:1	-	103

Spectrofluorimetric Applications

Most of the researchers were used the hydrazones as analytical reagents for the determination of metal ions by using spectrophotometer, but few researchers considered these hydrazones as spectrofluorimetric reagents also. β -cyclodextrin-o-vanilinfurfural hydrazone¹⁰⁷ was used as spectrofluorimetric reagent for the determination of traces of cadmium(II). Di-2-pyridylketone-2-furoylhydrazone was used for the determination of gallium(III) by Salgado et al¹⁰⁸. Orthovanillin furfural -hydrazone¹⁰⁹ and

Orthovanilinfuroylhydrazone¹¹⁰ were used for the spectrofluorimetric determination of Os(VIII) and Cd(II), respectively.

Conclusions

Based on the above information about the biological and analytical applications of hydrazones it is concluded that the hydrazones are precious reagents in the determination of metal ions in various environmental matrices. We believe that the three decades survey is very useful to the future studies in this respect.

References

1. Yoe JH. 1949. Recent advances in Analytical Chemistry, (Inter Science Publishers) New York.
2. Feigl F, Organic reagents in Inorganic analysis. *Analytical Chemistry* 1936, **8**, 3. 401- 410.
3. Feigl F, Beitr ige zur qualitativen Mikroanalyse. *Mikrochemie* 1924, **2**, 186-188.
4. Clark R E D, The colorimetric determination of tin by means of Toluene -3:4-dithiol (Dithiol). *Analyst* 1937, **62**, 661.
5. Yu Kitaev P, Buzykin B I and Troepolskaya T V, The structure of hydrazones. *Russian Chemical Reviews* 1970, **39**, 441-456.
6. Singh R B, Hydrazones as analytical reagents: A Review. *Talanta* 1982, **29**, 77-84.
7. Katyal M and Dutt Y, Analytical applications of hydrazones. *Talanta* 1975, **22**, 151 -166.
8. Jeewoth T, Li Kam Wah H, Bhowon M G, Ghoorohoo D and Babooram K, Synthesis and anti bacterial/catalytical properties of Schiff bases and Schiff base metal complexes derived form 2,3-diaminopyridine. *Synthesis and Reactivity in Inorganic and Metal Organic Chemistry* 2000, **30**, 1023-1038.
9. Jayabalakrishnan C and Natarajan K, Synthesis, characterization and biological activities of Ruthenium(II) carbonyl complexes containing bifunctional tridentate Schiff bases. *Synthesis and Reactivity in Inorganic and Metal Organic Chemistry* 2001, **31**, 983-995.
10. Dharamaraj N, Viswanathamurthi P and Natarajan K, Ruthenium (II) complexes containing bidentate Schiff bases and their antifungal activity. *Transition Metal Chemistry* 2001, **26**, 105-109.
11. Savini L, Chiasserini L, Gaeta A and Pellerano C, Synthesis and anti-tubercular evaluation of 4-Quinolylhydrazones. *Bioorganic and Medicinal Chemistry* 2002, **10**, 2193-2198.
12. Agarwal R K, Singh L, Sharma D K and Sing R, Synthesis, spectral and thermal investigations of some oxovanadium(IV) complexes of hydrazones of Isonicotinic acid hydrazide. *Turkish Journal of Chemistry* 2005, **29**, 309-316.
13. Ragavendran JV, Sriram D, Patel S K, Reddy I V, Bharathwajan N, Stables J and Yogeeswari P, Design and synthesis of anticonvulsants from a combined phthalimide-GABA-anilide and hydrazone pharmacophore. *European Journal of Medicinal Chemistry* 2007, **42**, 146-151.
14. Abdel-Aal T M, El-Sayed W A and El-Ashry E S H, Synthesis and antiviral evaluation of some sugar Arylglycinoylhydrazones and their oxadiazoline derivatives. *Archiv der Pharmazie Chemistry in Life Sciences* 2006, **339**, 656-663.
15. Walcourt A, Loyevsky M, Lovejoy D B, Gofdeuk V R and Richardson D R, Novel aroyl hydrazone and thiosemicarbazone iron chelators with anti-malarial activity against chloroquine-resistant and -sensitive parasites. *The International Journal of Biochemistry and Cell Biology* 2004, **36**, 401-407.

17. Savini L, Chiasserini L, Travagli V, Pellerano C, Novellino E, Cosentino S and Pisano M B, New α -(N)-heterocyclic hydrazones: evaluation of anticancer, anti-HIV and antimicrobial activity. *European Journal of Medicinal Chemistry* 2004, **39**, 113-122.
18. Singh B, Narang K K and Srivastava R, Studies of complexes of Co(II), Ni(II), Cu(II) and Cd(II) with Acetophenone and 4-hydroxyacetophenone, 4-aminobenzoyl hydrazones. *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry* 2003, **33**, 1025-1036.
19. Nora H. Al-Sha'alan, Antimicrobial activity and spectral, magnetic and thermal studies of some transition metal complexes of a Schiff base hydrazone containing a Quinoline moiety. *Molecules* 2007, **12**, 1080-1091.
20. Kucukguzel S G, Rollas S, Erdeniz H and Kiraz M, Synthesis, characterization and antimicrobial evaluation of ethyl 2-arylhydrazono-3-oxobutyrate. *European Journal of Medicinal Chemistry* 1999, **34**, 153-160.
21. Gulerman N N, Oruc E E, Kartal F and Rollas S, In vivo metabolism of 4-fluoro benzoic acid [(5-nitro-2-furanyl)methylene] hydrazide in rats. *European Journal of Drug Metabolism and Pharmacokinetics* 2000, **25**, 103-108.
22. Sah P P T and Peoples S A, Isonicotinoyl hydrazones as antitubercular agents and derivatives for identification of Aldehydes and ketones. *Journal of American Pharmacists Association* 1954.43, 513-524.
23. Ersan S, Nacak S and Berkem R, Synthesis and antimicrobial activity of N-[(α -methyl)benzylidene]-(3-substituted-1,2,4-triazol-5-yl-thio)acetohydrazides. *IL Farmaco* 1998, **53**, 773-776.
24. Ulsoy N, Capan G, Otuk G and Kiraz M, Synthesis and antimicrobial activity of new 6-phenylimidazo[2,1-b]thiazole derivatives. *Bollettino Chimico Farmaceutico* 2000, **139**(4), 167-172.
25. Kucukguzel S G, Rollas S, Kucukguzel I and Kiraz M, Synthesis and antimicrobial activity of some coupling products from 4-aminobenzoic acid hydrazones. *European Journal of Medicinal Chemistry* 1999, **34**, 1093-1100.
26. Rodriguez-Arguelles M C, Vazquez S M, Touceda P T, Matalobos J S, Deibe A M G, Ferrari M B, Pelosi G, Pelizzi C and Zanil F, Complexes of 2-thio phenyl carbonyl and isonicotinoyl hydrazones of 3-(N-methyl)isatin. A study of their antimicrobial activity. *Journal of Inorganic Biochemistry* 2007, **101**, 138-147.
27. Loncle C, Brunel J M, Vidal N, Dherbomez M and Letourneux Y, Synthesis and antifungal activity of cholesterol-hydrazone derivatives. *European Journal of Medicinal Chemistry* 2004, **39**, 1067-1071.
28. Nayyar, Amit, Jain and Rahul, Recent advances in new structural classes of anti-tuberculosis agents. *Current Medicinal Chemistry* 2005, **12**, 1873-1886.
29. Scior T and Disele G S J, Isoniazid is not a lead compound for its pyridyl ring derivatives, isonicotinoyl amides, hydrazides, and hydrazones: A critical review. *Current Medicinal Chemistry* 2006, **13**, 2205-2219.
30. Katiyar T S and Tandon S N, 1-Isonicotinoyl-2-salicylidenehydrazine as a new chelatometric reagent. *Talanta* 1964, **11**, 892-894.
31. Kouimtzis G S and Vasiliades V C, Spectrophotometric and solvent extraction study of o-hydroxybenzaldehyde isonicotinoyl hydrazone complexes with gallium and Indium. *Microchemical Journal* 1975, **20**, 173-179.
32. Vasilikiotis G S, Analytical applications of isonicotinoyl hydrazones: I. A new selective reagent for mercury. *Microchemical Journal* 1968, **13**, 526-528.
33. Khattab M A, El-Enany G and Soliman N E, Structural studies of 2-Aceto-1-Naphthol-N-Salicylhydrazone complexes of Cu(II), Co(II), Ni(II), Mn(II), Ce(III) and UO_2^{2+} . *Pakistan Journal of Scientific and Industrial Research* 1984, **27**, 81.

34. Park C I, Kim H S and Cha K W, Spectrophotometric determination of copper after selective extraction with α -(2-Benzimidazolyl)- α' , α'' -(N-5-nitro-2-pyridylhydrazone)-toluene in the presnence of Brij 58. *Bulletin of Korean Chemical Society* 1999, **20**(3), 352-354.
35. Ishii H, Odashima T and Kawamonzen Y, Spectrophotometric studies on complexation reactions of some water-soluble 5-nitro-2-pyridylhydrazones with cobalt. Spectrophotometric and derivative spectrophotometric determination of trace cobalt with 2-benzimidazolyl-3-sulphophenylmethanone-5-nitro-2-pyridyl hydrazone. *Analytica Chimica Acta* 1991, **244**, 223-229.
36. Medinilla J, Ales F and Sanchez F G, Spectrophotometric and second-derivative spectrophotometric determination of mercury in organomercurials by means of benzyl-2-pyridylketone-2-quinolylhydrazone. *Talanta* 1986, **33**, 329-334.
37. Rosales D, Ariza J L G and Asuero A G, Spectrophotometric determination of palladium in catalysts and carbenicillin with 1-(2-pyridylmethylidene)-5-thiocarbohydrazone. *Analyst* 1986, **111**, 449-453.
38. Kouimtzis Th A, Apostolopoulou C and Staphilakis I. Spectrophotometric determination of cobalt in sea water and brines by solvent extraction with 2,2'-dipyridyl-2''-pyridyl hydrazone. *Analytica Chimica Acta* 1980, **113**, 185-188.
39. Stratis J A, Anthemidis A N and Vasilikiotis G S, Solvent extraction and spectrophotometric determination of palladium(II) with 2,2'-bipyridyl 2-pyridylhydrazone (DPPH). *Analyst* 1984, **109**, 373-376.
40. Otomo M, 2,2'-Dipyridyl-2-quinolylhydrazone as a reagent for the spectro -photometric determination of metals: The extractive spectrophotometric determination of palladium(II). *Analytica Chimica Acta* 1980, **116**, 161-168.
41. Garcia-Villanova R J and Garcia-Villanova R, Biacetyl bis(isonicotinoyl hydra zone) complexes with Zirconium(IV). I. Spectrophotometric study. *Anales De Quimica Serie B* 1983, **79**, 401-405.
42. Kabil M A, Hafez M A H, El-Asmy A A and Mostafa M M, Spectrophotometric determination of Molybdenum with N-Cyanoacetylacetaldehyde hydrazone. *Journal of the Chinese Chemical Society* 1986, **33**, 183-187.
43. Garcia-Varagas M, Trevilla S and Milla M, Synthesis and characterization of 1,2-cylco hexanedione bis-benzoylhydrazone and its application to the determination of ti in minerals and rocks. *Talanta* 1986, **33**, 209-214.
44. Salinas F, Nevado J J B and Valiente P, Kinetic spectrophotometric determina -tion of nanogram levels of manganese by use of the salicylaldehyde guanyl hydrazone-hydrogen peroxide system. *Talanta* 1987, **34**, 321-324.
45. Varghese A and Khadar A M A, Highly selective derivative spectrophotometric determination of Tin(II) in alloy samples in the presence of Cetylpyridinium chloride. *Acta Chimica Slovenica* 2006, **53**, 374-380.
46. Alvarez M B, Palmieri M, Davis D and Fritz J S, 2,6-Diacetylpyridine bis(arylhydra - zone) derivatives as analytical reagents for uranium. *Talanta* 1987, **34**, 473-477.
47. Casoli A, Mangia A, Mori G and Predieri G, Liquid-liquid extraction and determination of uranium(IV) with 2,6-diacetylpyridine bis (benzoylhydrazone). *Analytica Chimica Acta* 1986, **186**, 283-287.
48. Sasaki Y, 2,6-Diacetylpyridine bis(benzoylhydrazone) and 2,6-diacetylpyridine bis(2-hydroxybenzoylhydrazone) as spectrophotometric reagents for determina -tions of Iron(III) and Vanadium(V). *Bulletin of the Institute for Chemical Research, Kyoto University* 1986, **64**, 140-149.

49. Mallikarjuna Rao K, Sreenivasulu Reddy T and Brahmaji Rao S, Spectrophotometric determination of molybdenum(VI) and vanadium(V) using resacetophenone isonicotinoyl hydrazone. *Journal of Indian Institute of Science* 1986, **66**, 39-41.
50. Kavlentis E, Synthesis and analytical properties of 3,4-dihydroxy benzaldehyde Guanyldihydrazone (3,4-DBGH). *Mikrochimica Acta* 1986, **88**, 27-32.
51. Babaiah O, Kesava Rao C P, Reddy T S and Reddy V K, Rapid, selective, direct and derivative spectrophotometric determination of titanium with 2,4-dihydroxybenzaldehyde isonicotinoylhydrazone. *Talanta* 1996, **43**, 551-558.
52. Lakshmi Narayana S, Janardhan Reddy K, Adi Narayana Reddy S, Sarala Y, and Varada Reddy A, A highly sensitive spectrophotometric determination of micro amounts of vanadium(V) in environmental and alloy samples by using 3,4-dihydroxybenzaldehydeisonicotinoylhydrazone. *Environmental Monitoring and Assessment* 2008, **144**, 341-349.
53. Lakshmi Narayana S, Adi Narayana Reddy S, Subbarao Y, Hwang I and Varada Reddy A, A simple and highly sensitive spectrophotometric determination of Cr(VI) in food samples by using 3,4-dihydroxy benzaldehyde isonicotinoyl hydrazone (3,4-DHBINH). *Food Chemistry* 2010, **121**, 1269-1273.
54. Lakshmi Narayana S, Ramachandraiah C, Varada Reddy A, Lee D and Shim J, Determination of traces of Pd(II) in spiked samples by using 3,4-dihydroxy benzaldehyde isonicotinoyl hydrazone as a chelating agent with UV visible spectrophotometer. *E- Journal of Chemistry* 2011, **8**, 217-225.
55. Sivaramaiah S, Raveendra Reddy P, Krishna Reddy V and Sreenivasulu Reddy T, Direct and derivative spectrophotometric determination of thorium with 2,4-dihydroxybenzaldehyde isonicotinoylhydrazone. *Journal of Radioanalytical and Nuclear Chemistry* 2000, **245**, 367-370.
56. Sivaramaiah S and Raveendra Reddy P, Direct and derivative spectrophotometric determination of Zinc with 2,4-dihydroxybenzaldehyde isonicotinoyl hydrazone in potable water and pharmaceutical samples. *Journal of Analytical Chemistry* 2005, **60**, 828-832.
57. Kesava Rao C, Krishna Reddy V and Reddy T S, Rapid and sensitive spectrophotometric determination of cerium(IV) with 2,4-dihydroxybenzophenone benzoichydrazone. *Talanta* 1994, **41**, 237-241.
58. Silva M and Valcarcel M, Spectrophotometric determination of microgram amounts of calcium in waters and foods using diphenylglyoxal bis(2-hydroxy benzoylhydrazone). *Analyst* 1980, **105**, 193-202.
59. Silva M and Valcarcel M, Extraction and spectrophotometric determination of trace amounts of tin(II) with diphenylglyoxal bis(2-hydroxybenzoyl hydrazone). *Microchemical Journal* 1980, **25**, 289-294.
60. Silva M and Valcarcel M, Spectrophotometric method for the determination of microgram amounts of titanium using an extraction technique with diphenylglyoxalbis(2-hydroxybenzoylhydrazone). *Microchemical Journal* 1980, **25**, 117-123.
61. Singh RB, Odashima T and Ishii H, Spectrophotometric and analogue derivative spectrophotometric determination of micro-amounts of iron with 2,2'-dipyridyl-2-benzothiazolylhydrazone. *Analyst* 1983, **108**, 1120-1127.
62. Nakanishi T and Otomo M, Extraction-spectrophotometric determination of palladium(II) with Di-2-pyridyl ketone benzolyhydrazone. *Analytical Sciences* 1985, **1**, 161-163.
63. Nakanishi T and Otomo M, Solvent extraction and spectrophotometric determination of iron(II) with di-2-pyridyl ketone benzolyhydrazone. *Microchemical Journal* 1986, **33**, 172-178.

64. Avila Terra L H S, Cunha areias M C da, Gaubeur I, Encarnacion M and Suarez-Iha V, Solvent extraction-spectrophotometric determination of nickel(II) in natural waters using Di-2-pyridyl ketone benzoylhydrazine. *Spectroscopy Letters* 1999, **32**, 257-271.
65. Zatar N A, Abu-zuhri A Z, Al-nuri Md. A, Mahmoud F M, Abu-obaid A A, Simultaneous spectrophotometric determination of iron(II) and iron(III) mixtures using Di-2-pyridyl ketone benzoylhydrazine. *Spectroscopy Letters* 1989, **22**, 1203-1214.
66. Rossi M V, Suarez-Iha M E V and Hoffman M R, Spectrophotometric determination of stability constants of Co(II) and DPKBH (Di-2-pyridyl ketone benzoylhydrazine). *Spectroscopy Letters* 1995, **28**, 1153-1166.
67. Terra S A L H, Encarnacion M and Suarez-Iha V, A new spectrophotometric determination with Di-2-pyridyl ketone benzoylhydrazine for determination of nickel(II). *Spectroscopy Letters* 1997, **30**, 625-639.
68. Kettrup A, Seshadri T and Jakobi F, Spectrophotometric determination of rhenium with di(2-pyridyl) ketone-2-furancarbothiohydrazine and other thio- hydrazones. *Analytica Chimica Acta* 1980, **115**, 383-388.
69. Grases F, Garcia Sanchez F and Valcarcel M, Fluorimetric determination of copper and mercury based on their catalytic effects on the autoxidation of 2,2'-dipyridyl ketone hydrazine. *Analytica Chimica Acta* 1980, **119**, 359-365.
70. Bernhardt P V, Caldwell L M, Chaston T B, Chin P and Richardson D R, Cyto -toxic iron chelators: characterization of the structure, solution chemistry and redox activity of ligands and iron complexes of the di-2-pyridyl ketone isonicotinoylhydrazine analogues. *Journal of Inorganic Biological Chemistry* 2003, **8**, 866-880.
71. Kavlentis E, Selective extractive spectrophotometric determination of palladium (II) with Di-2-pyridyl ketone 2-pyridyl-hydrazine. *Analytical Letters* 1988, **21**, 1681-1688.
72. Kavlentis E, Selective spectrophotometric determination of vanadium(II) in the presence of 35 common cations by means of Di-2-pyridyl ketone 2-pyridyl hydrazine (DPPH). *Analytical Letters* 1989, **22**, 2083-2089.
73. Cunha Areias M C da, Avila-Terra L H S, Gaubeur I, Suarez-Iha M E V, A new simultaneous spectrophotometric determination method for determination of iron(II) and iron(III) in natural waters. *Spectroscopy Letters* 2001, **34**, 289-300.
74. Abu-Eid M, Zatar N A, Al-Nuri Md A, Khamis M and Khalaf S, Spectrophotometric determination of uranium in ores using Di-2-pyridyl ketone hydrazine derivatives. *Spectroscopy Letters* 1995, **25**, 585-592.
75. Kanetake T and Otomo M. Extractive spectrophotometric determination of palladium with Di-2-pyridylmethanone 2-(5-nitro)pyridylhydrazine. *Analytical Sciences* 1988, **4**, 411-415.
76. Takaoka T, Taya T and Otomo M, Extractive spectrophotometric determination of trace iron(II) with di-2-pyridylmethanone-2-(5-nitro)pyridylhydrazine. *Talanta* 1992, **39**, 77-80.
77. Themelis D G, Tzanavaras P D, Kika F S and Sofoniou M C, Flow-injection manifold for the simultaneous spectrophotometric determination of Fe(II) and Fe(III) using 2,2'-dipyridyl-2-pyridylhydrazine and a single-line double injection approach. *Fresenius Journal of Analytical Chemistry* 2001, **371**, 364-368.
78. Anthemidis A N, Themelis D G and Stratis J A, Stopped-flow injection liquid -liquid extraction spectrophotometric determination of palladium in airborne particulate matter and automobile catalysts. *Talanta* 2001, **54**, 37-43.
79. Park C I and Cha K W, Spectrophotometric determination of trace amounts of cobalt with 2-hydroxybenzaldehyde-5-nitro-pyridylhydrazine in presence of surfactant after separation with amberlite IRC-718 resin. *Talanta* 1998, **46**, 1515-1523.

80. Krishna Reddy V, Mutta Reddy S, Chennaiah A, Reddy P R and Sreenivasulu Reddy T, First order derivative spectrophotometric determination of vanadium (V) and iron(III) individually and simultaneously. *Journal of Analytical Chemistry* 2003, **58**, 442-446.
81. Richardson D R, Tran E H and Ponka P, The potential of iron chelators of the pyridoxal isonicotinoylhydrazone class as effective antiproliferative agents. *Blood* 1995, **86**, 4295-4306.
82. Odashima T, Kohata K, Yagi K and Ishii H. Extraction spectrophotometric determination of trace nickel with 5-nitro-2-pyridinecarbaldehyde 5-nitro-2-pyridyl hydrazone. *Bunseki Kagaku* 1995, **44**, 135-138.
83. Kim Y N, Choi K S, Lee I H, Bark K M and Chung R J, Spectrophotometric determination of copper with N,N'-Oxalylbis (salicylaldehyde hydrazone). *Journal of the Korean Chemical Society* 1992, **36**, 95-99.
84. Ramachandraiah C, Vijaya Kumari D and Lakshminarayana K, A sensitive spectrophotometric method for the determination of trace amounts of uranium (VI) using o-hydroxypropionophenone isonicotinoylhydrazone. *Journal of Radio Analytical and Nuclear Chemistry* 1993, **175**, 185-190.
85. Lokhande R S, Jaywant A S and Barhate V D, Extractive spectrophotometric determination of cobalt(II) with p-methylisonitrosoacetophenonehydrazone. *Asian Journal of Chemistry* 1996, **8**, 610-612.
86. Lokhande R S and Jaywant A S, Extractive spectrophotometric determination of copper with p-methylisonitrosoacetophenonehydrazone. *Asian Journal of Chemistry* 1999, **11**, 1040-1042.
87. Zatar N A, Al-Nuri Md A, Abu-Eid M, Hannoun Md, Abu-Zuhri A Z, Khalaf S and Khamis M, Spectrophotometric determination of cobalt with Di-2-pyridyl ketone benzoylhydrazone. *Spectroscopy Letters* 1991, **24**, 1145-1152.
88. Sinha S H and Sawant A D, Extraction and spectrophotometric determination of palladium with Pyridine-2-acetaldehyde salicyloylhydrazone. *Bulletin of the Chemical Society of Japan* 1992, **65**, 1622-1625.
89. Bale M N, Dave D P and Sawant A D, Extraction and spectrophotometric determination of lead(II) with pyridine-2-acetaldehyde salicyloylhydrazone. *Talanta* 1995, **42**, 1291-1296.
90. Odashima T and Ishii H, Synthesis and properties of hydrazones from 3- and/or 5-nitro-2-pyridylhydrazine and heterocyclic Aldehydes, characterization of their complexes and extraction-spectrophotometric determination of traces of nickel with 2-pyridinecarbaldehyde 3,5-dinitro-2-pyridylhydrazone. *Analytica Chimica Acta* 1993, **277**, 79-88.
91. Odashima T, Yamada M, Yonemori N and Ishii H, Synthesis and chromogenic properties of some Phthalazinylhydrazones and extractive spectrophotometric and analogue-derivative spectrophotometric determination of trace amounts of nickel with Di-2-pyridylmethanone-1-phthalazinylhydrazone. *Bulletin of the Chemical Society of Japan* 1987, **60**, 3225-3231.
92. Bale M N and Sawant A D, Solvent extraction and spectrophotometric determination of uranium(VI) with pyridine-2-carboxaldehyde 2-hydroxy benzoyl hydrazone. *Journal of Radioanalytical and Nuclear Chemistry* 2001, **247**, 531-534.
93. Chaston T B and Richardson Des R, Interactions of the pyridine-2-carboxaldehyde isonicotinoylhydrazone class of chelators with iron and DNA: implications for toxicity in the treatment of iron overload disease. *Journal of Biological Inorganic Chemistry* 2003, **8**, 427-438.

94. Gallego M, Valcarcel M and Garcia-Vergas M, Pyridoxal salicyloylhydrazone as a chromogenic reagent for the determination of titanium. *Microchimica Acta* 1983, **79**, 289-297.
95. Suarez Iha M E V, Pehkonen S O and Hoffmann M R, Stability, stoichiometry, and structure of Fe(II) and Fe(III) complexes with Di-2-pyridyl ketone benzoyl hydrazone: Environmental applications. *Environmental Science and Technology* 1994, **28**, 2080-2086.
96. Rosales D, Ariza J L G and Asuero A G, Spectrophotometric determination of palladium in catalysts and carbenicillin with 1-(2-pyridylmethylidene)-5-thiocarbohydrazone. *Analyst* 1986, **111**, 449-453.
97. Aita T, Odashima T and Ishii H, Synthesis and chromogenic properties of water-soluble hydrazones, spectrophotometric and analogue derivative spectro-photometric determination of trace amounts of iron with 2- pyridyl-3'-sulpho-phenylmethanone 2-pyrimidylhydrazone (PSPmH) and kinetic studies of the complex formation of iron(II) with PSPmH. *Analyst* 1984, **109**, 1139-1145.
98. Roquette-Pinto C L S and Libergott E K, Extraction-spectrophotometric determination of cadmium with pyruvylidine-2-hydrazinobenzothiazole in benzene. *Analytica Chimica Acta* 1980, **117**, 349-351.
99. Pozo M E U, Torres A G D and Pavon J M C. Use of salicylaldehyde carbo hydrazone as a reagent for the determination of trace amounts of zinc in biological samples and alloys. *Analyst* 1988, **113**, 547-549.
100. Salinas F, Nevado J J B and Valiente P, Kinetic spectrophotometric determination of nanogram levels of manganese by use of the salicylaldehyde guanylhydrazone-hydrogen peroxide system. *Talanta* 1987, **34**, 321-324.
101. Nakanishi T and Otomo M, Spectrophotometric determination of iron(II) with 2-(3'-sulphobenzoyl) pyridine benzoylhydrazone. *Microchemical Journal* 1987, **36**, 128-134.
102. Nakagawa T, Doi K and Otomo M, Extraction-spectrophotometric determination of palladium(II) with thiazole-2-carbaldehyde-2-quinolylhydrazone. *Analyst* 1985, **110**, 387-390.
103. Sanchez F G and Lopez M H, Determination of copper in biological tissue by means of photochemically generated (z)-2-thiophenylaldehyde 2-pyridyl hydra zone. *Analyst* 1986, **111**, 445-448.
104. Chalapathi P V, Prathima B, Subba Rao Y, Janardhan Reddy K, Ramesh G N , Venkata Ramana Reddy D and Varada Reddy A. A chelating reagent 2,3,4-Trihydroxy acetophenone phenylhydrazone (THAPPH) used for selective and sensitive kinetic spectrophotometric determination of Thorium(IV) from ores. *Research Journal of Chemistry and Environment* 2011, **15**, 579-585.
105. Kavlentis E, Synthesis of Phthalaldehyde bisguanylhydrazone (PABGH) -Spectrophotometric analysis of Co(II), Cu(II), Ni(II) mixtures in the presence of several cations. *Analytical Letters* 1988, **21**, 689-697.
106. Kavlentis E, Spectrophotometry determination of Arsenic (III) and Antimony (III) by means of Isonicotinoylhydrazones of 4-Dimethyl amino benzaldehyde and 2-hydroxynaphthaldehyde. *Analytical Letters* 1987, **20**, 2043-2047.
107. Kavlentis E, Salicylaldehyde isonicotinoylhydrazone as a specific analytical reagent for the selective extractive spectrophotometric determination of Molybdenum(VI) in presence of several cations. *Analytical Letters* 1988, **21**, 107-113.
108. Tang B, Zhang L, Zhang J, Chen Z Z and Wang Y, Synthesis of a novel host molecule of cross-linking-polymeric-beta-cyclodextrin-o-vanillin furfural hydrazone and spectro fluorimetric analysis of its identifying cadmium. *Spectrochimica Acta Part A* 2004, **60**, 2425-2431.

109. Salgado M, Ojeda C B, Torres A G D and Pavon J M C, Di-2-pyridyl ketone 2-furoylhydrazone as a reagent for the fluorimetric determination of low concentrations of gallium and its application to biological samples. *Analyst* 1988, **113**, 1283-1285.
110. Tang B, Zhang H and Wang Y, Flow injection spectrofluorimetric determination of trace amounts of osmium. *Spectrochimica Acta Part A* 2005, **61**, 2239-2244.
111. Tang B, Yue T, Zhang L, Wu J and Chen Z, Flow injection spectro fluorimetric method for the determination of Cadmium. *Microchimica Acta* 2004, **148**, 71-76.

