Management of perianal Crohn's disease

MRB KEIGHLEY, MS, FRCS

MRB KEIGHLEY, MS, FRCS. Management of perianal Crohn's disease. Can J Gastroenterol 1993;7(2):266-271. Most perianal Crohn's disease is asymptomatic and may well resolve given adequate medical treatment for more proximal lesions. This will consist, in the main, of sulphasalazine and prednisolone, both of which are known to be effective in the treatment of Crohn's disease. Should the local perianal disease become more severe, a trial of oral metronidazole may be worthwhile. The development of an ischiorectal or perianal abscess is indication for the simplest surgical drainage procedure, and rectal strictures resulting from the healing of perianal fissures may be gently dilated. Further progression of disease may be treated by diversion of the fecal stream, but this will stand a greater chance of success in patients without florid rectal Crohn's disease. Diversion is also indicated as a preliminary to the repair of rectovaginal fistulae. Patients with florid perianal Crohn's disease and severe anorectal disease will probably come to proctocolectomy, but initial defunctioning of the colon will make the operative procedure easier, may facilitate perineal healing, and in some patients, subsequent proctectomy with the risk of a persistent perineal sinus may be avoided.

Key Words: Fecal diversion, Perianal Crohn's disease, Proctocolectomy

Traitement de la maladie de Crohn périnéale

RÉSUMÉ: Dans la plupart des cas, la maladie de Crohn périnéale est asymptomatique et peut rentrer dans l'ordre avec un traitement médicamenteux adéquat pour les lésions plus proximales. Il consistera principalement de salazopyrine et de prednisolone, tous deux connus pour leur efficacité dans le traitement de la maladie de Crohn. Si la maladie périnéale locale s'aggrave, il peut être approprié d'essayer le métronidazole par voie orale. Le développement d'un abcès ischiorectal ou périnéale est une indication pour le drainage chirurgical le plus simple et le rétrécissement rectal qui résulte de la cicatrisation d'une fissure périnéale peut être dilaté en douceur. La progression de la maladie peut être traitée par déviation de l'écoulement fécal, mais cette intervention aura de meilleures chances de succès chez les patients sans maladie de Crohn rectale

Department of Surgery, Queen Elizabeth Hospital, Birmingham, United Kingdom

Correspondence: Professor MRB Keighley, Department of Surgery, Queen Elizabeth Hospital, Birmingham, B15 2TH, United Kingdom

PERIANAL PROBLEMS ARE COMMON among patients with Crohn's disease. Such lesions may precede the occurrence of proximal gastrointestinal Crohn's disease by as much as 22 years (1). They occur more commonly in association with colonic rather than small bowel disease (2,3) but the exact incidence of perianal Crohn's disease is difficult to assess because estimates vary widely between series. An incidence of between 20 and 80% is reported for perianal disease in Crohn's patients (3-5), the problem being to decide what constitutes perianal Crohn's disease? Some series consider only symptomatic disease (6), some include cases with skin tags (7) while others do not (8). Such differences probably account for the observed variations in frequency, rather than a truly altered incidence of perianal disease between series. The perianal lesions of Crohn's disease are often asymptomatic but they can become both painful and disabling. In this case they will require energetic treatment.

The treatment of any pathological condition depends upon the cause of that condition, but the cause of Crohn's disease remains unknown. It is proposed by Hughes (9) that perianal Crohn's disease takes its origin in certain primary lesions which occur in

manifeste. La déviation est également indiquée comme mesure préliminaire avant la réparation d'une fistule rectovaginale. Les patients porteurs d'une maladie de Crohn périnéale manifeste et d'une maladie anorectale grave devront probablement subir une proctocolectomie, mais une iléostomie initiale rendra l'intervention chirurgicale plus facile, aidera probablement la circulation périnéale et chez certains patients, la proctectomie subséquente avec risque de sinus périnéal persistant, sera ainsi évitée.

relation to the anorectal mucosa. These primary lesions include mucosal fissures, ulcerated edematous piles and cavitating ulcers. Fissuring of the anal margin heals by fibrosis and may subsequently cause the anal stenosis which is commonly seen in these patients (10). Some edematous piles resolve to form skin tags, but it is the cavitating ulcers which progress to form the majority of the tissue destruction. These ulcers penetrate through the full thickness of the rectal mucosa, spreading sepsis into the perirectal tissues. Their propensity to spread through the rectal musculature might partly be due to obstruction of forward flow of rectal contents by the anal sphincters. This may encourage fecal material under pressure to discharge down the perirectal tracts. Once outside the wall of the anorectum, sepsis may spread for quite long distances before emerging on the perianal skin to form a fistula. If such a fistula tract becomes obstructed, extensive anorectal sepsis may result.

Such a pathological sequence of events seems dire, and certainly perianal Crohn's disease can look alarming. However, the temptation to treat alarming appearances with aggressive surgery must be resisted if the situation is not to be made worse by iatrogenic morbidity. Careful studies of the natural history of perianal Crohn's disease reveal that it is surprisingly indolent. Of particular interest are the 109 patients reported by Fielding to have perianal Crohn's disease in 1968 (6). Patients with skin tags alone were excluded from this study and only those patients with either a definite fistula or fissure were included. This cohort of patients was re-examined in 1978 to determine the outcome of the perianal Crohn's disease. During follow-up, 10 of the original patients in the study

came to proctectomy, although in only five was this done directly because of perianal disease. Fourteen patients died from coincidental causes or from complications of Crohn's disease. Twenty-four were completely asymptomatic but refused to be reassessed proctologically in 1978. The remaining 61 patients had their perianal Crohn's disease thoroughly evaluated both in 1968 and in 1978 (11). Of 54 patients with an anal fissure in 1968, 17 had healed spontaneously by 1978. Twenty-seven further patients had also healed their anal fissures but with some minor degree of narrowing of the anal canal. Two patients with no fissure at the beginning of the study developed one over the follow-up decade.

In 1968, 21 patients had an anal fistula. Eight patients who had no surgical treatment healed their fistulae spontaneously. Seven were treated surgically by laying open a low fistula, but one of these recurred. The remainder with fistulae were unhealed but as they were asymptomatic no operation intervention was necessary. These findings underline the generally benign nature of perianal Crohn's disease. When planning treatment for such lesions it must be against the background of a strong tendency for spontaneous resolution of the lesion and the risk of incontinence by surgical intervention (23).

MEDICAL TREATMENT OF PERIANAL CROHN'S DISEASE

Before any form of treatment is started, it is important to establish the diagnosis of the lesion as accurately as possible. Any anal lesion in a patient who is known to be suffering from Crohn's disease is likely to be perianal Crohn's disease. However, if the perianal disease is the first lesion to

manifest itself – as is the case in 30% of the author's patients – diagnosis may be more difficult. In this situation the clinician may consider the perianal disease to be a nonspecific lesion. However, the relative lack of pain, multiplicity of the lesions, or the undermined nature of the fissures on the lateral margin of the anus, may alert the observer that this is in fact perianal Crohn's disease. Further criteria for differentiation may be a visually active proctitis at sigmoidoscopy. Biopsy of the rectal mucosa for histological examination together with similar biopsies of any skin tag or fistula track may allow a firm histological diagnosis to be made. Other pathologies which may be confused with perianal Crohn's disease include tuberculosis, venereal proctitis, bilharzia and hidradenitis suppurativa. In seriously ill patients with sudden unexplained florid perianal disease, leukemia or myeloma should be considered.

There is evidence that perianal lesions may heal if active Crohn's disease elsewhere in the gut is successfully treated (12,13). Such treatment may include resection of a proximal lesion. For example Heuman and colleagues (3) found that in a series of 15 patients with perianal lesions before a resection for more proximal Crohn's disease, 80% of the perianal lesion healed following resection (2). Similar results are reported by Hellers et al (14) who found that 47% of their patients with anal fistula healed spontaneously after intestinal resection. There are no prospective, controlled trials of drug therapy in perianal Crohn's disease alone, but bed-rest, correction of fluid depletion, as well as sulphasalazine (4, 15-17) and methylprednisolone (4,15, 16) are known to be effective treatment for Crohn's colitis. The immunosuppressive drug azathioprine and its metabolite 6-mercaptopurine are also effective in the treatment of colonic Crohn's disease. There is some evidence that they may be useful in healing perirectal fistulae (18,19). Against this, the National Cooperative Crohn's Disease Study (NCCDS) in the United States did not find azathioprine of value in healing Crohn's disease (15).

This difference may in part be due to the shorter follow-up period in the NCCDS compared with the studies of Brooke et al (18) and Presant et al (19).

There is mounting evidence that antibiotics are effective in controlling perianal Crohn's disease. In 1977 Allan and Cooke (20) reported striking improvement in two patients with perianal Crohn's disease treated by metronidazole. This report was followed by further studies (21,22); in the study of Brandt and colleagues (22) 26 patients with perianal Crohn's disease were treated with 20 mg/kg/day metronidazole orally. Ten patients progressed to complete healing of perianal disease while on the drug. The remaining 16 patients underwent partial healing. Unfortunately the drug was required for many months and in one case for three years. Such prolonged treatment was associated with paresthesia in 50% of patients, which developed a mean of six months after onset of treatment. Most paresthesia was bilateral and pedal. However, Ambrose and others found the response to metronidazole to be disappointing (24).

Further evidence that Crohn's colitis might respond to metronidazole therapy is provided by the trial carried out by Ursing et al (17) who demonstrated that oral metronidazole therapy was as effective as sulphasalazine in controlling Crohn's colitis. A prospective, randomized, cross-over trial of metronidazole therapy in Crohn's disease showed clinical improvement in six patients with Crohn's colitis but not in a further 16 patients with small bowel disease (25). How metronidazole brings about this improvement is uncertain. It seems likely that it exerts its action through its antimicrobial effect on the bowel microflora and as a systemic antibiotic. Ursing and colleagues (17) found that metronidazole reduces the *Bacteroides* content of the colon after only one week and this often lasted until the drug was stopped. Besides its antimicrobial effect metronidazole is known to influence leucocyte chemotaxis (26) as well as being an immunosuppressive agent (27). Both or either of these actions could contribute to its efficacy.

SURGICAL TREATMENT FOR PERIANAL CROHN'S DISEASE

The potential benefits of surgical treatment for perianal Crohn's disease must be carefully weighed against the possibility that spontaneous resolution will occur in the perianal disease. A recent study from Birmingham (23) reviewed 202 Crohn's disease patients. Their notes were examined to assess evidence of perianal Crohn's disease at some time during their previous illness. The most common manifestation of perianal disease was skin tags in 75 patients. Most skin tags were asymptomatic and had disappeared by the time of review in 1984. There were 38 patients with documented evidence of a fissure-in-ano but 29 of these had also disappeared spontaneously by the time of review. Low lying fistula-in-ano had been present in 40 patients but 14 had spontaneously resolved on review. Thirteen were still present but caused little trouble, while 10 patients with a low fistula required proctectomy. Twelve patients with a high fistula-in-ano were identified, of whom seven had come to proctectomy and the remaining had troublesome symptoms. A similar pattern was observed in six patients with a rectovaginal fistula, four of whom had come to proctectomy by the time of review. Thus, this study showed a strong tendency for perianal lesions in Crohn's disease to remit spontaneously, similar to results reported by Buchmann et al (1).

Of the patients in the Birmingham series (26), only a small number received operative treatment for perianal disease. In those who did come to surgery, there was a high incidence of complications together with little clinical benefit. Gentle anal dilation achieved healing in only four of seven patients with a fissure and one patient was rendered temporarily incontinent of feces afterwards. Laying open of a low fistula-in-ano achieved healing in only one of 12 fistulas but six developed impairment of continence afterwards. This finding is in contrast with a report from St Mark's Hospital, which relates to 49 low fistulae treated along conventional lines; 25 healed successfully (28). No patient with a high fistula was

treated by local surgery; the majority were advised to undergo a proctectomy. It is of great interest that the presence of perianal disease has a marked effect on the healing of proctectomy wounds after rectal excision. Of 27 patients with perianal disease who came to proctectomy, 19 were associated with a persistent perianal sinus, compared with complete perianal healing in all patients undergoing a rectal excision in the absence of perianal disease (23).

The most pressing indication for local surgery for perianal Crohn's disease is the drainage of a perianal or ischiorectal abscess. This is a surgical emergency as for other forms of pyogenic abscess. If an abscess is drained and a fistula results, then unless this is very low and symptomatic there is no further indication for surgery.

On the current available evidence, anal fissures, which are often painless, usually heal spontaneously. They do not, therefore, require surgical treatment. During the healing process they may produce some degree of fibrosis with consequent anal narrowing. Such narrowing is of short length and usually responds well to gentle dilation with bougies. These short superficial strictures are in contrast to the long, craggy, fibrous anorectal strictures that sometimes occur as a consequence of cavitating ulceration of the anorectum in Crohn's proctitis. Such strictures may respond to balloon dilation (29) but more commonly require a proctectomy if they become symptomatic (30).

Low anal fistulae in perianal Crohn's disease are often asymptomatic and many heal spontaneously. There are few indications to lay them open, but if a subcutaneous fistula is associated with an abscess in the absence of active proximal disease, it is relatively safe to lay it open while draining the abscess. Some surgeons recommend the use of partial internal sphincterotomy (31,32) for low intersphincteric fistula. The present author has not found this to be a useful operation in perianal Crohn's disease. Patients with a high fistula and perianal Crohn's disease often come to proctectomy (23,28). This is because such fistulae often arise in the rectum as a result of cavitating

ulceration. Enterovaginal fistulae may occur in association with perianal Crohn's disease. Givel and colleagues (33) report their experience of 13 such fistulas. Seven came to proctectomy with perineal healing in five, and a persistent vaginocutaneous fistula in one. In a further two patients, attempts at local repair of the fistula without a covering colostomy were unsuccessful. In a further patient, a local repair of the rectovaginal fistula was successful following creation of a proximal loop ileostomy. The ileostomy was subsequently closed with no ill effects. One further patient developed an adenocarcinoma of the fistula track after 10 years of medical treatment. Provided the patient does not require a proctectomy for coincidental severe rectal disease, a local repair of the rectovaginal fistula should be attempted following creation of a proximal loop ileostomy. Very low anovaginal fistulae may be safely laid open.

Coexisting hemorrhoids are not common in patients with perianal Crohn's disease (10) and their treatment in patients with active disease is not to be recommended. In one study, six of 20 patients with hemorrhoids and active Crohn's disease who underwent treatment for their hemorrhoids came to proctectomy for complications apparently dating from the treatment of their hemorrhoids (34).

In summary, the indications for local surgery in perianal Crohn's disease are limited to the drainage of pus, gentle dilation of short low strictures, and a very occasional laying open of a symptomatic low anal fistula. Some enterovaginal fistulae should also be repaired but if so a preliminary loop ileostomy should be raised; however, a large proportion of these patients have few symptoms and, if so, the fistula should be left alone.

FECAL DIVERSION

Whether fecal diversion has a place in the management of intractable perianal Crohn's disease remains debatable. It is possible to divert the fecal stream in one of two ways. Recently there has been much interest in the use of an elemental diet to treat

acute exacerbations of Crohn's disease by reducing the fecal stream. Most reports do not specifically mention the effect of such diets on perianal manifestations of Crohn's disease, but use of an elemental diet was shown to be as effective as prednisolone in the treatment of large and small bowel Crohn's disease (35) and remission of disease can be achieved more rapidly when an elemental diet is combined with a non-absorbable antibiotic (36). The elemental diet is of theoretical value for the treatment of perianal disease but as yet not much information is available concerning its therapeutic efficacy.

An alternative approach uses a loop (37) or split ileostomy (38) to defunction the colon. This method has been extensively tested with regard to its effect in treating Crohn's colitis, and some information is available concerning perianal Crohn's disease. The two largest series of patients with Crohn's colitis treated by this technique are those from Oxford (39) and the Cleveland Clinic (40). The Oxford group has shown that when patients with Crohn's colitis are treated by a split ileostomy, 25% relapse while the colon is defunctioned and a further 25% once the ileostomy is closed (41). The Oxford group reviewed 102 patients who had undergone a split ileostomy for Crohn's disease, 29 of whom had symptomatic perianal Crohn's disease at the time of fecal diversion. Eleven of 19 patients with fistulae-in-ano improved with diversion, six healed partially and five healed completely. In addition, 12 women had rectovaginal fistulae; these five fistulae persisted but remained asymptomatic, two healed spontaneously following defunction, and of these one relapsed when the ileostomy was closed. A further five healed after local surgery while the colon remained defunctioned and of these, two relapsed when the ileostomy was closed. Overall, the perianal disease improved or healed after 72% of the split ileostomies. The best response to diversion appeared to be in patients without rectal involvement and whose intestinal disease was quiescent. Harper (39) also noted that the reduction in inflamma-

tion and improvement in the perineum greatly facilitated the perineal dissection in patients requiring proctectomy. Furthermore, it has been shown that healing of the perineal wound was less certain following proctectomy in patients with Crohn's proctitis, with a high fistula-in-ano, or with a rectovaginal fistula (42).

Zelas and Jagelman (40) reviewed the result of a loop ileostomy in 79 patients with severe Crohn's colitis or ileocolitis. Twenty-three of the patients had severe perianal Crohn's disease at the time of defunction, and 22 improved following diversion of the fecal stream as assessed by clinical and serological parameters. Of these, six remained well with an ileostomy alone. Nine required definitive surgery a mean of nine months following ileostomy formation and six had a relapse of symptoms requiring definitive surgery 11 months after ileostomy formation. Six remained well with an ileostomy alone when followed for between three and five years. Other smaller series of patients treated by defunctioning ileostomy for perianal Crohn's disease testify to the value of this method of management (43,44). Not all surgeons agree, however (2), and several reports of the application of defunctioning ileostomy (primarily to treat Crohn's colitis but including some patients with perianal Crohn's disease) revealed disappointing results. For example the series of Berman et al (45) includes 10 patients with perianal Crohn's disease. Among the 10 patients before diversion, two had a fissure-in-ano, seven had a fistula-in-ano and one had both. After diversion over a mean follow-up period of 25 months, all the fissures healed as did one fistula, but in six patients the fistula-in-ano persisted. Three patients developed a rectal stricture while defunctioned (also noted by other surgeons [41]) and in one case this was the main indication for a subsequent proctectomy. Four further patients developed perianal Crohn's disease following defunction. McIlrath (46) reports similar disappointing results with only four of 10 patients showing any improvement of perianal Crohn's disease following fecal diver-

sion, and in a further four cases worsening of the anorectal disease led to proctectomy.

MANAGEMENT OF COMPLICATIONS OF PERIANAL CROHN'S DISEASE

The most serious complications of perianal Crohn's disease are incontinence of feces or the development of carcinoma. Alexander-Williams (47)

reported 16 patients with incontinence of feces subsequent to perianal Crohn's disease. Six of these patients were completely incontinent of feces and 10 were incontinent only when they had diarrhea. Four of the less severe group had undergone no previous surgical treatment but the other six all had some form of anal surgery. Results such as these have led to the view that incontinence subsequent to anorectal

Crohn's disease is due to aggressive surgeons and not to progressive disease. Adenocarcinoma may develop in the track of a chronic fistula and one such is reported in relation to a longstanding rectovaginal fistula in Crohn's disease (48) which had been present for 20 years. This is such a rare occurrence that it should not influence the treatment policy in perianal Crohn's disease.

REFERENCES

- Baker WN, Milton-Thompson GJ. The anal lesion as a sole presenting symptoms of intestinal Crohn's disease. *Gut* 1971;12:A865.
- Heuman R, Bolin T, Sjodahl R, Tagesson C. The incidence and cause of perianal complications and arthralgia after intestinal resection with restoration of continuity for Crohn's disease. *Br J Surg* 1981;68:528-30.
- Homan WP, Tang CK, Thorburnasson B. Anal lesions complicating Crohn's disease. *Arch Surg* 1976;111:1333-5.
- Rankin GB, Watt D, Melyk CS. National cooperative Crohn's disease study. Extraintestinal manifestations and perianal complications. *Gastroenterol* 1979;77:914-20.
- Lockhart-Mummery HE. Crohn's disease: Anal lesions. *Dis Colon Rectum* 1975;18:200.
- Loygue J, Huguier M. Le traitement chirurgical des localisations anorectales de la maladie de Crohn. *Arch Fr Mal App Dis* 1971;60:29.
- Keighley MRB, Allan RN. Current status and influence of operation on perianal Crohn's disease. *Int J Colorectal Dis*. (In press)
- Fielding JF. Perianal lesions in Crohn's disease. *J Roy Coll Surg Edin* 1972;32:1717.
- Hughes LE, Jones IRG. Perianal lesions in Crohn's disease. In: Allan RN, Keighley MRB, Hawkins C, Alexander-Williams J, eds. *Inflammatory Bowel Diseases*. Churchill Livingstone 1982.
- Alexander-Williams J, Buchmann P. Perianal Crohn's disease. *World J Surg* 1980;4:203-8.
- Buchmann P, Keighley MRB, Allan RN, Thompson H, Alexander-Williams J. Natural history of perianal Crohn's disease. Ten year follow-up: A plea for conservatism. *Am J Surg* 1980;140:642-4.
- Lockhart-Mummery HE, Morson BC. Crohn's disease of the large bowel. *Gut* 1964;5:493-509.
- Garlock JH. Surgery of the Alimentary Tract. London: Butterworths, 1967
- Hellers G, Bergstrand O, Ewerth S. Occurrence and outcome of the primary treatment of anal fistulae in Crohn's disease. *Gut* 1980;21:525-7.
- Summers RW, Switz DM, Sessions JT. National cooperative Crohn's disease study: Results of drug treatment. *Gastroenterol* 1979;77:847-96.
- Melchow H, Ewe K, Branders JW, et al. European cooperative Crohn's disease study (ECCDS). Results of drug treatment. *Gastroenterology* 1984;86:249-66.
- Ursing B, Alm T, Baramy F. A comparative study of metronidazole and sulfasalazine for active Crohn's disease. The cooperative Crohn's disease study in Sweden. II: Results. *Gastroenterology* 1984;83:550-62.
- Brooke BN, Cave DR, King DW. The place of azathioprine for Crohn's disease. *Lancet* 1976; i: 1041-2.
- Present DM, Korelitz BI, Wisch N, et al. Treatment of Crohn's disease with 6-mecaptopurine. *N Engl J Med* 1980;302:981-7.
- Allan RN, Cooke TW. Evaluation of metronidazole in the management of Crohn's disease. *Gut* 1977;18:A422.
- Bernstein LH, Frank MS, Brandt LJ. Healing of perianal Crohn's disease with metronidazole. *Gastroenterology* 1982;79:357-65.
- Brandt LJ, Bernstein LH, Bolye JS, Frank MS. Metronidazole therapy for perianal Crohn's disease: A follow-up study. *Gastroenterology* 1982;83:383-7.
- Keighley MRB, Allan RN. Current status and influence of operation on perianal Crohn's disease. *Int J Colorectal Dis* 1986;1:104-7.
- Ambrose NS, Allan RN, Keighley MRB, et al. *Dis Colon Rectum* 1984;28:81-5.
- Blichfeldt P, Blomhoff JP, Myhre E, Gjone E. Metronidazole in Crohn's disease: a double blind cross-over clinical trial. *Scand J Gastroenterol* 1978;13:123-7.
- Gnarpe H, Persson S, Belsheim J. Influence of metronidazole and tinidazole on leucocyte chemotaxis in Crohn's disease. *Infection* 1978;6(Suppl):107-9.
- Grove DI, Mahmoud AAF, Warren KS. Suppression of cell mediated immunity by metronidazole. *Int Arch Allergy Appl Immunol* 1977;54:422-7.
- Marks CG, Ritchie JK, Lockhart-Mummery HE. Anal fistula in Crohn's disease. *Br J Surg* 1981;68:525-7.
- Alexander-Williams J, Allan A, Morel P, et al. Therapeutic coaxial balloon dilatation of Crohn's stricture. *Ann R Coll Surg Eng* 1986;68:95-7.
- Lockhart-Mummery LE. Perianal Crohn's disease: Invited commentary. *World Journal Surgery* 1980;4:208.
- Parks AG. Pathogenesis and treatment of fistula-in-ano. *Br Med J* 1961;1:463-9.
- Sohn N, Korelitz B, Weinstein MA. Anorectal Crohn's disease: Definitive surgery for fistulas and recurrent abscess. *Am J Surg* 1980;139:394-7.
- Givel JC, Hawker P, Allan RN, Alexander-Williams J. Enterovaginal fistulas associated with Crohn's disease. *Surg Gynecol Obstet* 1982;155:494-6.
- Jeffrey PJ, Ritchie JK, Parks AG. Treatment of haemorrhoids in patients with inflammatory bowel disease. *Lancet* 1977;i:1084-5.
- O'Morain C, Segal AW, Levi AJ. Elemental diet as a primary treatment of acute Crohn's disease: a controlled trial. *Br Med J* 1984;288:1859-63.
- Saverymutu S, Hodgson HJF, Chadwick VS. Controlled trial comparing prednisolone with an elemental diet plus non-absorbable antibiotics in active Crohn's disease. *Gut* 1985;26:994-8.
- Alexander-Williams J. Loop ileostomy and colostomy for faecal diversion. *Ann R Coll Surg Eng* 1974;54:141-8.
- Lee E. Split ileostomy in the treatment of Crohn's disease of the colon. *Ann R Coll Surg Engl* 1975;56:94-102.
- Harper PH, Kettlewell MGW, Lee E. The effect of split ileostomy on perianal Crohn's disease. *Br J Surg* 1982;69:608-10.
- Zelas P, Jagelman DG. Loop ileostomy

- in the management of Crohn's colitis in the debilitated patient. *Ann Surg* 1980;191:164-8.
41. Harper PH, Truelove SC, Lee ECG, Kettlewell MGW, Jewell DP. Split ileostomy and ileocolectomy for Crohn's disease of the colon and ulcerative colitis: a 20 year survey. *Gut* 1983;24:106-13.
 42. Scammell BE, Keighley MRB. Delayed perineal wound healing after proctectomy for Crohn's colitis. *Br J Surg* 1986;73:150-2.
 43. Waterman IT, Pena AS. The place of split ileostomy in the management of Crohn's disease. In: *The Management of Crohn's Disease*. Amsterdam-Oxford: Excerpta Med, 1976:224.
 44. Oberhelman HA. The effect of intestinal diversion by ileostomy on Crohn's disease of the colon. In: *The Management of Crohn's Disease*. Amsterdam-Oxford: Excerpta Med, 1976:216.
 45. Berman JH, Thompson H, Cooke WT, Alexander-Williams J. The effect of diversion of intestinal contents on the progress of Crohn's disease of the large bowel. *Gut* 1971;12:11-15.
 46. McIlrath DC. Divertin ileostomy or colostomy in the management of Crohn's disease of the colon. *Arch Surg* 1971;103:308-16.
 47. Alexander-Williams J. Perianal Crohn's disease. Amsterdam-Oxford: Excerpta Med, 1975:43.
 48. Buchmann P, Allan RB, Thompson H, Alexander-Williams J. Carcinoma in a recto-vaginal fistula in a patient with Crohn's disease. *Am J Surg* 1980;140:462.
-