Diagnosis?

Lillian Sung MD, Nicole Le Saux MD, Vivian Frenkel MD

A seven-year-old black boy born in Canada with perinatally acquired human immunodeficiency virus (HIV) infection presented with recurrent nausea, diarrhea and intolerance of gastric tube feeding. Additionally, there was a history of a 3 kg weight loss over the two weeks before admission and a history of persistent fatigue over the four weeks before admission.

Significant past medical history revealed cytomegalic virus (CMV)-retinitis diagnosed and treated three months previously, chronic sinusitis and moderate HIV encephalopathy. Approximately three months before admission, his CD4 count was $41/\text{mm}^3$, and his viral load was $4.6 \log_{10}$ HIV RNA copies/mL of plasma. His medications included stavudine, lamivudine, saquinovir, granulocyte colony-stimulating factor, clarithromycin, fluconazole, intravenous ganciclovir, and monthly pentamidine and immune globulin.

On physical examination, he was a thin boy in the 25th percentile for weight. He was afebrile. His examination was remarkable for the absence of oral candidiasis, the presence of diffuse palpable lymph nodes less than 0.5 cm in the cervical, axillary and inguinal areas, and the absence of skin lesions. His abdominal exam revealed no masses, no tenderness and a palpable nontender liver edge 1 cm below the costal margin.

Blood for CMV polymerase chain reaction (PCR) was positive, while Epstein-Barr virus (EBV) PCR was negative; EBV se-

Figure 1) Cytology (Papanicolaou stain) from fine needle aspiration of liver lesion

rology (immunoglobulin [Ig] G) for capsid antigen was positive; EBV IgM was negative. A chest radiograph was normal. Examination of the stool was positive for adenovirus but negative for *Clostridium difficile* toxin, ova and parasites. Gastric and colonic biopsies were negative for parasites, mycobacteria or viral changes. An ultrasound of the liver demonstrated two well defined hypoechoic liver lesions, each measuring approximately 13 mm, located in the right lobe of the liver. A fine needle aspiration of one of the hepatic lesions was performed (Figure 1).

What is your diagnosis?

Continued on page 293

Correspondence: Dr N Le Saux, Children's Hospital of Eastern Ontario, 401 Smythe Road, Ottawa, Ontario K1H 8L1. Telephone 613-737-7600, fax 613-738-4832, e-mail LESAUX@CHEO.on.ca