Diagnosis?

T Lambert MD¹, JM Langley MD MSc FRCPC², A Issekutz MD FRCPC¹

A 15-month-old African-Canadian male was referred for consultation because of a recent febrile illness associated with abdominal distension, decreased appetite, fever and lethargy for one week. His mother noted that he had not been himself for about six months, over which time he became increasingly irritable, lost interest in play and had a weight loss of 3 kg. He was constipated, and defecation was painful. There was no history of sweats, cough, rhinorrhea, discharge from eyes or ears, rash, arthralgia or arthritis, or urinary symptoms. He was unable to walk or crawl but could sit unsupported and scoot around the floor. He had a five-word vocabulary. He was born at 40 weeks gestation by induced vaginal delivery. There was history of oligohydramnios, but the pregnancy, labour and delivery were otherwise unremarkable. He was on no medications and his immunizations were up to date. He had a history of three skin infections of the diaper, scalp and periorbital area as well as pimples that were treated effectively with topical fucidin. He had been hospitalized for treatment of cervical adenitis at age three months, and for gastroenteritis and dehydration at 14 months.

The boy lived with his family in a small town in the Maritimes. They had travelled to Barbados in the previous year while the father engaged in agricultural employment. Animal

exposures included domestic cats, a dog and a goldfish. The family history was notable only for irritable bowel syndrome and hypertension.

On examination, he looked chronically ill. Temperature was 38.3°C, pulse 170 beats/min and respiration 55 breaths/min. Blood pressure was normal. Height was 75 cm (fifth to 10th percentile), the weight 7.8 kg (less than fifth percentile) and the head circumference 47.5 cm (25th to 50th percentile). His abdomen was protuberant and tympanic, and bowel sounds were heard. The liver was palpable 3 cm below the costal margin; there was no splenomegaly or adenopathy. The rest of his examination was unremarkable. A complete blood count revealed a hypochromic, microcytic anemia (hemoglobin 68 g/L) with a normal total white and platelet count. Aspartate aminotransferase was slightly elevated at 78 U/L. An abdominal ultrasound revealed three cystic masses in the liver of mixed echogenicity; the largest was 7×6 cm and had a necrotic centre. Numerous smaller lesions were also present. Radiological appearance was consistent with a hepatoblastoma. Stools for culture, ova and parasite examination, and occult blood were negative.

What is the diagnosis?

Continued on page 365

¹Department of Pediatrics, Queen's University, Kingston, Ontario; ²Department of Pediatrics, Dalhousie University, Halifax, Nova Scotia
Correspondence and reprints: Dr Joanne Langley, IWK-Grace Health Centre for Children, Women and Families, Dalhousie University, 8th Floor Labs, 5850 University Avenue, Halifax, Nova Scotia B3J 3G9. Telephone 902-428-8498, fax 902-428-3217, e-mail joanne.langley@dal.ca