Diagnosis?

Caroline Quach MD, Dorothy Moore MD, Earl E Rubin MD

A 20-year-old male receiving chemotherapy for an osteosarcoma of the lower limb was admitted to hospital with a two-day history of painful oral ulcers and fever. His last course of therapy (VP-16, ifosfamide) finished 11 days before admission.

Past medical history was significant for an episode of pre-septal cellulitis 15 months earlier, an episode of fever and neutropenia nine months previously, and a left maxillary sinusitis four months earlier, each without positive cultures. All of these episodes responded completely to empirical antibiotic therapy. He was sexually active but had no new partners recently, and had no prior history of sexually transmitted diseases. He lived and worked on a farm, and there was no disease among the livestock. His vaccinations were up-to-date, and he had no history of any recent trauma.

At the time of the present admission, he reported seeing some blood in his saliva, which he felt was coming from his nose or the back of his throat. He denied any cough. The remainder of his review of systems revealed no abnormalities. On physical examination, temperature was 38.9°C, and there were discrete oral ulcers throughout the mouth. The remain-

der of the examination was normal. A complete blood count revealed profound neutropenia (absolute neutrophil count of 20/mm³). Following appropriate cultures, empirical treatment with parenteral piperacillin/tazobactam, tobramycin and acyclovir was initiated.

Within hours of his presentation, his mother noted that his left eyelid was red and swollen, and, by 12 h after admission, the process extended down the left side of his face to his neck. As well, within 12 h of admission, his overall status had deteriorated, and he became unresponsive and hypotensive. In addition, he developed fulminant disseminated intravascular coagulation. He was intubated and transferred to the intensive care unit. An infused computed tomographic scan of the head revealed a diffuse cerebral edema with air extending from the left maxillary sinus to the orbit. In addition, a mass was suspected to be present in the middle of the maxillary sinus.

Despite maximal general support and the addition of clindamycin and amphotericin B, the patient died 30 h after admission. Approximately 18 h after admission, an initial blood culture was noted to be positive.

What is the diagnosis?

continued on page 300

Montreal Children's Hospital and McGill University, Montreal, Quebec

Correspondence and reprints: Dr Earl Rubin, Montreal Children's Hospital, 2300 Tupper Street, Montreal, Quebec H3H 1P3.